

HIW_Indicators

Version 1.5.2.1

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
10th graders disapproval of heavy drinking (percent)	Percent of adolescents who disapprove of having one or two alcoholic drinks nearly every day-10th graders	Yes	No	No	No	1445
10th graders disapproval of trying marijuana (percent)	Percent of adolescents who disapprove of trying marijuana or hashish once or twice-10th graders	Yes	No	No	No	1448
10th graders steroid use (percent)	Percent of 10th graders using steroids	Yes	No	No	No	1441
8th graders disapproval of heavy drinking (percent)	Percent of adolescents who disapprove of having one or two alcoholic drinks nearly every day-8th graders	Yes	No	No	No	1444
8th graders disapproval of trying marijuana (percent)	Percent of adolescents who disapprove of trying marijuana or hashish once or twice-8th graders	Yes	No	No	No	1447
8th graders steroid use (percent)	Percent of 8th graders using steroids	Yes	No	No	No	1440
Abuse of prescription pain relievers (percent)	Percent of person 12 years and older with nonmedical use of prescription pain relievers in the past year	Yes	No	No	No	1461
Abuse of prescription psychotherapeutic drug (percent)	Percent of persons 12 years and older with nonmedical use of any prescription psychotherapeutic drug in the past year	Yes	No	No	No	1465
Abuse of prescription sedatives (percent)	Percent of persons 12 years and older with nonmedical use of prescription sedatives in the past year	Yes	No	No	No	1464
Abuse of prescription stimulants (percent)	Percent of persons 12 years and older with nonmedical use of prescription stimulants in the past year	Yes	No	No	No	1463
Abuse of prescription tranquilizers (percent)	Percent of persons 12 years and older with nonmedical use of prescription tranquilizers in the past year	Yes	No	No	No	1462
Access to online health information (percent)	Percent of online health information seekers who report easily accessing health information	Yes	No	No	No	865
Access to school athletic facilities (percent)	Percent of the Nation's public and private schools that provide access to their physical activity spaces and facilities for all persons outside of normal school hours (that is, before and after the school day, on weekends, and during summer and other vacations)	Yes	No	No	No	1318

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Accidental puncture or laceration rate (per 100,000 beneficiaries)	Patient Safety Indicators: Accidental puncture or laceration rate (observed, among all surgical discharges based on secondary diagnosis field) among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2037
Activity limitation due to arthritis among adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who experience a limitation in activity due to arthritis or joint symptoms	Yes	No	No	No	419
Activity limitation due to chronic back conditions (per 1,000)	Activity limitation due to chronic back conditions	Yes	No	No	No	428
Acute hospital readmissions (number)	Total inpatient readmissions within 30 days of an acute hospital stay	Yes	Yes	No	Yes	278
Acute hospital readmissions (percent)	Inpatient readmissions within 30 days of an acute hospital stay	Yes	Yes	No	Yes	279
Acute inpatient PPS Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who had covered acute inpatient prospective payment system services	Yes	Yes	No	Yes	167
Acute inpatient PPS Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered acute inpatient prospective payment system services	Yes	Yes	No	Yes	168
Added sugar consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake provided by added sugars for persons aged 2 years and older	Yes	No	No	No	1237
Admissions for injection drug use (number)	Number of admissions to substance abuse treatment for injection drug use	Yes	No	No	No	1453
Adolescent artificial UV tanning (percent)	Percent of adolescents in grades 9 through 12 who report using artificial sources of ultraviolet light for tanning	Yes	No	No	No	501
Adolescent deaths aged 10 - 14 years (per 100,000)	Rates of adolescent and young adult deaths-Adolescents aged 10 to 14 years	Yes	No	No	No	1126

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Adolescent deaths aged 15 - 19 years (per 100,000)	Rates of adolescent and young adult deaths-Adolescents aged 15 to 19 years	Yes	No	No	No	1127
Adolescent participation in daily school physical education (percent)	Percent of adolescents who participate in daily school physical education	Yes	Yes	No	No	1317
Adolescent skin cancer protective measures (percent)	Percent of adolescents in grades 9 through 12 who follow protective measures that may reduce the risk of skin cancer	Yes	No	No	No	503
Adolescent suicide attempts requiring medical attention (per 100)	Rate of suicide attempts by adolescents that require medical treatment	Yes	Yes	No	No	1106
Adolescent wellness checkup (percent)	Percent of adolescents who have had a wellness checkup in the past 12 months	Yes	No	No	No	405
Adolescent-adult connection-discuss problems (percent)	Percent of adolescents who have an adult in their lives with whom they can talk about serious problems	Yes	No	No	No	409
Adolescent-adult connection-parents attend activities (percent)	Percent of parents who attend events and activities in which their adolescents participate	Yes	No	No	No	410
Adolescents binge drinking (percent)	Percent of persons engaging in binge drinking during the past month-Adolescents aged 12 to 17 years	Yes	No	No	No	1438
Adolescents females receiving formal instruction on abstinence (percent)	Percent of adolescents females who received formal instruction on abstinence before they were 18 years old	Yes	No	No	No	802
Adolescents females receiving formal instruction on birth control methods (percent)	Percent of adolescents females who received formal instruction on birth control methods before they were 18 years old	Yes	No	No	No	804
Adolescents females receiving formal instruction on HIV/AIDS prevention (percent)	Percent of adolescents females who received formal instruction on HIV/AIDS prevention before they were 18 years old	Yes	No	No	No	806
Adolescents females receiving formal instruction on STDs (percent)	Percent of adolescents females who received formal instruction on STDs before they were 18 years old	Yes	No	No	No	808
Adolescents females who talked to a parent or guardian about abstinence (percent)	Percent of adolescents females who talked to a parent or guardian about abstinence before they were 18 years old	Yes	No	No	No	810

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Adolescents females who talked to a parent or guardian about birth control methods (percent)	Percent of adolescents females who talked to a parent or guardian about birth control methods before they were 18 years old	Yes	No	No	No	812
Adolescents females who talked to a parent or guardian about HIV/AIDS prevention (percent)	Percent of adolescents females who talked to a parent or guardian about HIV/AIDS prevention before they were 18 years old	Yes	No	No	No	814
Adolescents females who talked to a parent or guardian about STDs (percent)	Percent of adolescents females who talked to a parent or guardian about STDs before they were 18 years old	Yes	No	No	No	816
Adolescents inhalant use (percent)	Percent of adolescents who use inhalants	Yes	No	No	No	1443
Adolescents males receiving formal instruction on abstinence (percent)	Percent of adolescents males who received formal instruction on abstinence before they were 18 years old	Yes	No	No	No	803
Adolescents males receiving formal instruction on birth control methods (percent)	Percent of adolescents males who received formal instruction on birth control methods before they were 18 years old	Yes	No	No	No	805
Adolescents males receiving formal instruction on HIV/AIDS prevention (percent)	Percent of adolescents males who received formal instruction on HIV/AIDS prevention before they were 18 years old	Yes	No	No	No	807
Adolescents males receiving formal instruction on STDs (percent)	Percent of adolescents males who received formal instruction on STDs before they were 18 years old	Yes	No	No	No	809
Adolescents males who talked to a parent or guardian about abstinence (percent)	Percent of adolescents males who talked to a parent or guardian about abstinence before they were 18 years old	Yes	No	No	No	811
Adolescents males who talked to a parent or guardian about birth control methods (percent)	Percent of adolescents males who talked to a parent or guardian about birth control methods before they were 18 years old	Yes	No	No	No	813
Adolescents males who talked to a parent or guardian about HIV/AIDS prevention (percent)	Percent of adolescents males who talked to a parent or guardian about HIV/AIDS prevention before they were 18 years old	Yes	No	No	No	815

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Adolescents males who talked to a parent or guardian about STDs (percent)	Percent of adolescents males who talked to a parent or guardian about STDs before they were 18 years old	Yes	No	No	No	817
Adolescents never using alcohol (percent)	Percent of at risk adolescents aged 12 to 17 years who, in the past year, refrained from using alcohol for the first time	Yes	No	No	No	1428
Adolescents never using marijuana (percent)	Percent of at risk adolescents aged 12 to 17 years who, in the past year, refrained from using marijuana for the first time	Yes	No	No	No	1429
Adolescents perception of risk associated with binge drinking (percent)	Percent of adolescents aged 12 to 17 years perceiving great risk associated with substance abuse-Consuming five or more alcoholic drinks at a single occasion once or twice a week	Yes	No	No	No	1450
Adolescents perception of risk associated with cocaine (percent)	Percent of adolescents aged 12 to 17 years perceiving great risk associated with substance abuse-Using cocaine once per month	Yes	No	No	No	1452
Adolescents perception of risk associated with marijuana (percent)	Percent of adolescents aged 12 to 17 years perceiving great risk associated with substance abuse-Smoking marijuana once per month	Yes	No	No	No	1451
Adolescents recent use of alcohol or illicit drugs (percent)	Percent of adolescents reporting use of alcohol or any illicit drugs during the past 30 days	Yes	No	No	No	1432
Adolescents recent use of marijuana (percent)	Percent of adolescents reporting use of marijuana during the past 30 days	Yes	No	No	No	1433
Adolescents riding with a driver who has been drinking (percent)	Percent of adolescents who report that they rode, during the previous 30 days, with a driver who had been drinking alcohol	Yes	Yes	No	No	1427
Adolescents with care in family-centered, comprehensive, coordinated systems (percent)	Percent of children with special health care needs who receive their care in family-centered, comprehensive, coordinated systems-Children aged 12 to 17 years	Yes	No	No	No	1155
Adult artificial UV tanning (percent)	Percent of adults aged 18 and older who report using artificial sources of ultraviolet light for tanning	Yes	No	No	No	502

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Adult asthma admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Adult asthma admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2045
Adult recent use of illicit drugs (percent)	Percent of adults reporting use of any illicit drug during the past 30 days	Yes	No	No	No	1434
Adult skin cancer protective measures (percent)	Percent of adults aged 18 years and older who follow protective measures that may reduce the risk of skin cancer	Yes	No	No	No	504
Adults binge drinking (percent)	Percent of persons engaging in binge drinking during the past month-Adults aged 18 years and older	Yes	No	No	No	1437
Aerobic physical activity among adolescents (percent)	Percent of adolescents who meet current Federal physical activity guidelines for Aerobic physical activity	Yes	Yes	No	No	1323
Aerobic physical activity and muscle-strengthening activity among adults (percent)	Percent of adults who meet the objectives for aerobic physical activity and for muscle-strengthening activity	Yes	No	No	No	1322
Aerobic physical activity: >150 min/week moderate or >75 min/week vigorous (percent)	Percent of adults who engage in aerobic physical activity of at least moderate intensity for at least 150 minutes/week, or 75 minutes/week of vigorous intensity, or an equivalent combination	Yes	No	No	No	1319
Aerobic physical activity: >300 min/week moderate or >150 min/week vigorous (percent)	Percent of adults who engage in aerobic physical activity of at least moderate intensity for more than 300 minutes/week, or more than 150 minutes/week of vigorous intensity, or an equivalent combination	Yes	No	No	No	1320
African American Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months in who are African American	Yes	Yes	No	Yes	312
Age-appropriate child restraint use, 1-3 years (percent)	Percent of age-appropriate vehicle restraint system use in children: Children aged 1 to 3 years	Yes	No	No	No	1055
Age-appropriate child restraint use, 4-7 years (percent)	Percent of age-appropriate vehicle restraint system use in children: Children aged 4 to 7 years	Yes	No	No	No	1056

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Age-appropriate child restraint use, 8-12 years (percent)	Percent of age-appropriate vehicle restraint system use in children: Children aged 8 to 12 years	Yes	No	No	No	1057
Age-appropriate child restraint use, birth - 12 months (percent)	Percent of age-appropriate vehicle restraint system use in children: Birth to 12 months	Yes	No	No	No	1054
AIDS among adolescents and adults (per 100,000)	Rate of new AIDS cases among adolescents and adults	Yes	No	No	No	929
AIDS among heterosexuals (number)	Number of new AIDS cases among adolescent and adult heterosexuals	Yes	No	No	No	943
AIDS among injection drug users (number)	Number of new AIDS cases among adolescents and adults who inject drugs	Yes	No	No	No	931
AIDS among men who have sex with men (number)	Number of new AIDS cases among adolescent and adult men who have sex with men	Yes	No	No	No	930
AIDS, perinatally acquired (number)	Number of new cases of perinatally acquired AIDS	Yes	No	No	No	939
Air Quality Index (AQI) exceeding 100 (number)	Number of days the Air Quality Index (AQI) exceeds 100	Yes	No	No	No	665
Airborne toxins - area sources (number)	Air toxic emissions to decrease the risk of adverse health effects caused by airborne toxics - area sources	Yes	No	No	No	671
Airborne toxins - mobile sources (number)	Air toxic emissions to decrease the risk of adverse health effects caused by airborne toxics - mobile sources	Yes	No	No	No	670
Airborne toxins -major sources (number)	Air toxic emissions to decrease the risk of adverse health effects caused by airborne toxics - major sources	Yes	No	No	No	672
Alcohol and/or illicit drug treatment receipt (percent)	Percent of persons who need alcohol and/or illicit drug treatment and received specialty treatment for abuse or dependence in the past year-Alcohol and/or illicit drug treatment	Yes	No	No	No	1455
Alcohol sold (number)	Average annual alcohol consumption rate	Yes	No	No	No	1439
Alcohol treatment receipt (percent)	Percent of persons who need alcohol and/or illicit drug treatment and received specialty treatment for abuse or dependence in the past year-Alcohol abuse or dependence	Yes	No	No	No	1456

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Alcohol-attributable deaths (number)	Number of deaths attributable to alcohol	Yes	No	No	No	1467
Alcohol-impaired driving deaths (per 100 vehicle miles)	Alcohol-impaired driving (08+ blood alcohol content [BAC]) fatality rate	Yes	No	No	No	1466
Alcohol: Excessive drinking (percent)	Percent of adults that report excessive drinking	No	Yes	Yes	No	150
Allergic reactions to food - adults (percent)	Percent of adults with a physician-diagnosed food allergy with a severe allergic reactions to food	Yes	No	No	No	820
Alternative modes of transportation - bicycling (percent)	Percent of trips to work using of alternative modes of transportation - bicycling	Yes	No	No	No	666
Alternative modes of transportation - mass transit (percent)	Percent of trips to work using of alternative modes of transportation - mass transit	Yes	No	No	No	668
Alternative modes of transportation - telecommute (percent)	Percent of trips to work using of alternative modes of transportation - telecommute	Yes	No	No	No	669
Alternative modes of transportation - walking (percent)	Percent of trips to work using of alternative modes of transportation - walking	Yes	No	No	No	667
Ambulatory care sensitive conditions (per 1,000 Medicare fee-for service aged enrollees)	Discharge rate per 1,000 Medicare fee-for-service enrollees for ambulatory care sensitive conditions	No	Yes	Yes	No	27
Ambulatory surgery center Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who used ambulatory surgery center services	Yes	Yes	No	Yes	233
Ambulatory surgery center Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used ambulatory surgery center services	Yes	Yes	No	Yes	234
Ambulatory Surgical Centers (number)	Number of Medicare participating Ambulatory Surgical Centers	No	Yes	No	Yes	10033
Amelioration of balance and dizziness problems of adults (percent)	Percent of adults with balance and dizziness problems in the past 12 months who reported their condition got worse or did not improve	Yes	No	No	No	766
American Indian or Alaska Native Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months in who are American Indian or Alaska Native	Yes	Yes	No	Yes	315

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Amputation among diabetics rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Rate of lower extremity amputation among patients with diabetes among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2046
Anencephaly (per 100,000 live births)	Rate of anencephaly	Yes	No	No	No	1170
Annual dental examinations - Persons with diabetes (percent)	Percent of persons with diagnosed diabetes who have at least an annual dental examination	Yes	No	No	No	553
Annual dilated eye examinations - Persons with diabetes (percent)	Percent of adults with diabetes who have an annual dilated eye examination	Yes	No	No	No	551
Annual foot examinations - Persons with diabetes (percent)	Percent of adults with diabetes who have at least an annual foot examination	Yes	No	No	No	552
Annual urinary microalbumin measurement - Medicare beneficiaries with diabetes (percent)	Percent of persons with diagnosed diabetes who obtain an annual urinary microalbumin measurement	Yes	No	No	No	549
Antibiotics misuse for common cold (per 100,000)	Rate of courses of antibiotics prescribed for the sole diagnosis of the common cold	Yes	No	No	No	975
Antibiotics misuse for ear infections (percent)	Rate of physician visits by children less than 5 years of age diagnosed with an ear infection that have an antibiotic course ordered, supplied, administered, or continued	Yes	No	No	No	974
Antimicrobial resistance - Ampicillin (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Non-typhoidal Salmonella isolates from humans that are resistant to ampicillin	Yes	No	No	No	832
Antimicrobial resistance - Ceftriaxone (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Non-typhoidal Salmonella isolates from humans that are resistant to Ceftriaxone (third-generation cephalosporin)	Yes	No	No	No	830

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Antimicrobial resistance - Erythromycin (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Campylobacter jejuni isolates from humans that are resistant to Erythromycin	Yes	No	No	No	834
Antimicrobial resistance - Gentamicin (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Non-typhoidal Salmonella isolates from humans that are resistant to gentamicin	Yes	No	No	No	831
Antimicrobial resistance - Nalidixic acid (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Non-typhoidal Salmonella isolates from humans that are resistant to nalidixic acid (quinolone)	Yes	No	No	No	829
Antimicrobial resistance - three or more classes (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant to antimicrobial drugs: Non-typhoidal Salmonella isolates from humans that are resistant to three or more classes of antimicrobial agents	Yes	No	No	No	833
Apparent noise-induced hearing loss among adolescents (per 1,000)	Rate of adolescents who have elevated hearing thresholds, or audiometric notches, in high frequencies (3, 4, or 6 kHz) in both ears, signifying noise-induced hearing loss	Yes	No	No	No	754
Apparent noise-induced hearing loss among adults (per 1,000)	Rate of adults who have elevated hearing thresholds, or audiometric notches, in high frequencies (3, 4, or 6 kHz) in both ears, signifying noise-induced hearing loss	Yes	No	No	No	755
Appropriate asthma medication regimens (percent)	Percent of persons with current asthma who do not use more than one canister of short-acting inhaled beta agonist per month	Yes	No	No	No	1416
Arteriovenous fistula use for vascular access (percent)	Percent of adult hemodialysis patients who use arteriovenous fistulas or have a maturing fistula as the primary mode of vascular access at the start of renal replacement therapy	Yes	No	No	No	471

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Arthritis education among adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who have had effective, evidence-based arthritis education as an integral part of the management of their condition	Yes	No	No	No	426
ASC Medicare service events (per 1,000 beneficiaries)	Rate of ambulatory surgery center service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2020
Asian American or Pacific Islander Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months in who are Asian American or Pacific Islander	Yes	Yes	No	Yes	314
Assistance in reducing exposure to environmental risks for asthma (percent)	Percent of persons with current asthma who have been advised by a health professional to change things in their home, school, and work environments to reduce exposure to irritants or allergens to which they are sensitive	Yes	No	No	No	1417
Assistive and adaptive devices use (percent)	Percent of persons aged 18 years and over with visual impairment who use assistive and adaptive devices	Yes	No	No	No	1575
Assistive listening device use among older adults (per 1,000)	Rate of adults aged 70 years and older with hearing loss who use assistive listening devices	Yes	No	No	No	747
Associate degrees in public health (percent)	Percent of 2-year colleges that offer public health or related Associate degrees	Yes	No	No	No	1388
Asthma activity limitations (percent)	Percent of adults with current asthma who have activity limitations	Yes	No	No	No	1404
Asthma deaths, ages 35-64 years (per 1,000,000)	Asthma death rate: Adults aged 35 to 64 years old	Yes	No	No	No	1408
Asthma deaths, ages 65 years and older (per 1,000,000)	Asthma death rate: Adults aged 65 years and older	Yes	No	No	No	1409
Asthma deaths, ages under 35 years (per 1,000,000)	Asthma death rate: Children and adults under age 35 years	Yes	No	No	No	1407
Asthma emergency department visits, ages 5-64 years (per 10,000)	Rate of hospital emergency department visits for asthma: Children and adults aged 5 to 64 years	Yes	No	No	No	1402

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Asthma emergency department visits, ages 65 years and older (per 10,000)	Rate of hospital emergency department visits for asthma: Adults aged 65 years and older	Yes	No	No	No	1403
Asthma emergency department visits, ages under 5 years (per 10,000)	Rate of hospital emergency department visits for asthma: Children under age 5 years	Yes	No	No	No	1401
Asthma hospitalizations, ages 5-64 years (per 10,000)	Rate of hospitalizations for asthma: Children and adults aged 5 to 64 years	Yes	No	No	No	1399
Asthma hospitalizations, ages 65 years and older (per 10,000)	Rate of hospitalizations for asthma: Adults aged 65 years and older	Yes	No	No	No	1400
Asthma hospitalizations, ages under 5 years (per 10,000)	Rate of hospitalizations for asthma: Children under age 5 years	Yes	No	No	No	1398
Asthma patient education (percent)	Percent of persons with current asthma who receive formal patient education	Yes	No	No	No	1412
Atrial fibrillation Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have atrial fibrillation	Yes	Yes	No	Yes	285
Atrial fibrillation Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have atrial fibrillation	Yes	Yes	No	Yes	286
Autism spectrum disorder screening (percent)	Percent of young children who are screened for an autism spectrum disorder (ASD) and other developmental delays by 24 months of age	Yes	No	No	No	1188
Average age of Medicare beneficiaries 65+ (mean)	Average age of Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	308
Average total hierarchical condition category score for Medicare beneficiaries 65+ (mean)	Average hierarchical condition category score of Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	318
Awareness of early warning symptoms of and response to heart attack (percent)	Percent of adults aged 20 years and older who are aware of the early warning symptoms and signs of a heart attack and the importance of accessing rapid emergency care by calling 9-1-1 or another emergency number	Yes	No	No	No	894

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Awareness of early warning symptoms of heart attack (percent)	Percent of adults aged 20 years and older who are aware of the early warning symptoms and signs of a heart attack	Yes	No	No	No	895
Awareness of HIV infection status (percent)	Percent of HIV-infected persons who are aware of their infection	Yes	No	No	No	942
Awareness of response to early warning symptoms of heart attack (percent)	Percent of adults aged 20 years and older who are aware of the importance of accessing rapid emergency care by calling 9-1-1 or another emergency number	Yes	No	No	No	896
Bacterial pneumonia admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Bacterial pneumonia admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2043
Beach closings (percent)	Percent of days that beaches are open and safe for swimming	Yes	No	No	No	674
Binge drinking: adults (percent)	Percent of adults 18 years and over that report binge drinking	No	Yes	Yes	No	17
Birth dose of hepatitis B vaccination (percent)	Percent of young children who have received universally recommended vaccines-A birth dose of hepatitis B vaccine (0 to 3 days between birth date and date of vaccination, reported by annual birth cohort)	Yes	No	No	No	1001
Births at 32 to 33 weeks of gestation (percent)	Percent of live births at 32 to 33 weeks of gestation	Yes	Yes	Yes	No	1139
Births to females 15-19 years (per 1,000)	Rate of births to females 15-19 years of age	Yes	Yes	Yes	No	22
Births to mothers 40-54 (percent)	Percent of all births to mothers 40-54 years of age	Yes	Yes	Yes	No	98
Births to mothers under 18 (percent)	Percent of all births to mothers less than 18 years of age	Yes	Yes	Yes	No	97
Births to unmarried mothers (percent)	Percent of all births to mothers who report not being married	Yes	Yes	Yes	No	99
Births, moderately preterm (32-36 wk.) (percent)	Percent of live births that are moderately preterm (32-36 weeks gestation)	Yes	Yes	Yes	No	10008
Blood cholesterol levels (mean)	Mean total blood cholesterol levels among adults	Yes	No	No	No	884

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Blood cholesterol screening (percent)	Percent of adults who have had their blood cholesterol checked within the preceding 5 years	Yes	No	No	No	886
Blood donations (percent)	Percent of persons who donate blood	Yes	No	No	No	436
Blood foliate concentrations in reproductive-aged women (percent)	Percent of women of childbearing potential who have low red blood-cell foliate concentrations	Yes	No	No	No	1172
Blood lead levels in children - elevated level (percent)	Percent of children with elevated blood lead levels	Yes	No	No	No	675
Blood lead levels in children - mean level (mean (average))	Mean blood lead levels in children	Yes	No	No	No	676
Blood pressure screening (percent)	Percent of adults who have had their blood pressure measured within the preceding 2 years and can state whether their blood pressure was normal or high	Yes	No	No	No	883
Blood pressure under control--persons with diabetes (percent)	Percent of adults with diagnosed diabetes whose blood pressure is under control	Yes	No	No	No	560
Breast cancer deaths (per 100,000)	Breast cancer death (female) rate	Yes	Yes	Yes	No	488
Breast cancer Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have breast cancer	Yes	Yes	No	Yes	299
Breast cancer Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have breast cancer	Yes	Yes	No	Yes	300
Breast cancer screening (percent)	Percent of women who receive a breast cancer screening based on the most recent guidelines	Yes	No	No	No	507
Breastfeeding - at 1 year (percent)	Percent of infants who are breastfed-At 1 year	Yes	No	No	No	1149
Breastfeeding - at 6 months (percent)	Percent of infants who are breastfed-At 6 months	Yes	No	No	No	1148
Breastfeeding - ever (percent)	Percent of infants who are breastfed-Ever	Yes	No	No	No	1147
Breastfeeding - exclusively through 3 months (percent)	Percent of infants who are breastfed-Exclusively through 3 months	Yes	No	No	No	1150
Breastfeeding - exclusively through 6 months (percent)	Percent of infants who are breastfed-Exclusively through 6 months	Yes	No	No	No	1151

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Broadband access to internet (percent)	Percent of persons with broadband access to the Internet	Yes	No	No	No	858
Bullying among adolescents (percent)	Percent of bullying among adolescents	Yes	Yes	No	No	1102
Calcium consumption (milligrams)	Mean total daily calcium intake (in mg) by persons aged 2 years and older	Yes	No	No	No	1223
Cancer survival (percent)	Percent of cancer survivors who are living 5 years or longer after diagnosis	Yes	No	No	No	498
Cardiac surgery patients with controlled 6 A.M. postoperative blood glucose (percent)	Hospital Compare: Cardiac surgery patients with controlled 6 A.M postoperative blood glucose	Yes	Yes	No	Yes	339
Cardiovascular death in dialysis patients (per 1,000)	Cardiovascular death rate for persons on dialysis	Yes	No	No	No	465
Cardiovascular death in transplant patients (per 1,000)	Cardiovascular death rate for persons with a functioning kidney transplant	Yes	No	No	No	467
Catheter use for vascular access (percent)	Percent of adult hemodialysis patients who use catheters as the only mode of vascular access	Yes	No	No	No	470
Central venous catheter-related bloodstream infections rate (per 100,000 beneficiaries)	Patient Safety Indicators: Central venous catheter-related bloodstream infections rate (observed, among medical and surgical discharges based on secondary diagnosis field) among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2034
Certificate programs in public health (percent)	Percent of 2-year colleges that offer public health or related Certificate programs	Yes	No	No	No	1389
Cervical cancer screening (percent)	Percent of women who receive a cervical cancer screening based on the most recent guidelines	Yes	No	No	No	505
Cesarean births, first (percent)	Percent of live births that are cesarean deliveries among low-risk (full-term, singleton, vertex presentation) women who have never previously had a cesarean birth	Yes	No	No	No	1133

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Cesarean births, repeat (percent)	Percent of live births that are cesarean among low-risk (full-term, singleton, vertex presentation) women who have had a prior cesarean birth	Yes	No	No	No	1134
Child deaths aged 1 - 4 years (per 100,000)	Rate of child deaths-Children aged 1 to 4 years	Yes	No	No	No	1124
Child deaths aged 5 - 9 years (per 100,000)	Rate of child deaths-Children aged 5 to 9 years	Yes	No	No	No	1125
Child maltreatment deaths (per 100,000)	Child maltreatment death rate	Yes	No	No	No	1085
Children in poverty (percent)	Percent of children under 18 years of age living in poverty	No	Yes	Yes	No	36
Children who have a medical home (percent)	Percent of children who have access to a medical home	Yes	No	No	No	1152
Children with care in family-centered, comprehensive, coordinated systems (percent)	Percent of children with special health care needs who receive their care in family-centered, comprehensive, coordinated systems-Children aged 0 to 11 years	Yes	No	No	No	1154
Children with cerebral palsy born as low birth weight infants (percent)	Percent of children with cerebral palsy born as low birth weight infants (less than 2,500 grams)	Yes	No	No	No	1168
Children with special needs who have a medical home (percent)	Percent of children with special health care needs who have access to a medical home	Yes	No	No	No	1153
Children's exposure to violence (percent)	Percent of children who are exposed to violence	Yes	No	No	No	1104
Children's Hospitals (number)	Number of Children's Hospitals participating in the Medicare program	No	Yes	No	Yes	10034
Chlamydia (per 100,000)	Chlamydia rate	No	Yes	Yes	No	20
Chlamydia among females 15 to 24 years attending family planning clinics (percent)	Percent of adolescents and young adults with Chlamydia trachomatis infections-Among females aged 15 to 24 years attending family planning clinics	Yes	No	No	No	1475

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Chlamydia among females 24 years enrolled in a national job training program (percent)	Percent of adolescents and young adults with Chlamydia trachomatis infections-Among females aged 24 years and under enrolled in a National Job Training Program	Yes	No	No	No	1476
Chlamydia among males 24 years enrolled in a national job training program (percent)	Percent of adolescents and young adults with Chlamydia trachomatis infections-Among males aged 24 years and under enrolled in a National Job Training Program	Yes	No	No	No	1477
Chlamydia cases (number)	Number of reported cases of chlamydia	No	Yes	Yes	No	10064
Chlamydia screening: 16-20 year old females with commercial health insurance (percent)	Percent of sexually active females aged 16 to 20 years enrolled in commercial health insurance plans who are screened for genital Chlamydia infections during the measurement year	Yes	No	No	No	1488
Chlamydia screening: 16-20 year old females with Medicaid (percent)	Percent of sexually active females aged 16 to 20 years enrolled in Medicaid plans who are screened for genital Chlamydia infections during the measurement year	Yes	No	No	No	1486
Chlamydia screening: 21-24 year old females with commercial health insurance (percent)	Percent of sexually active females aged 21 to 24 years enrolled in commercial health insurance plans who are screened for genital Chlamydia infections during the measurement year	Yes	No	No	No	1489
Chlamydia screening: 21-24 year old females with Medicaid (percent)	Percent of sexually active females aged 21 to 24 years enrolled in Medicaid plans who are screened for genital Chlamydia infections during the measurement year	Yes	No	No	No	1487
Chronic hepatitis B in infants and young children (number)	Number of chronic hepatitis B virus infections in infants and young children (perinatal infections)	Yes	No	No	No	962
Chronic kidney disease (percent)	Percent of the adults with chronic kidney disease	Yes	No	No	No	483
Chronic kidney disease deaths (per 100)	Death rate among adults 18 years or older with chronic kidney disease	Yes	No	No	No	484
Chronic kidney disease Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have chronic kidney disease	Yes	Yes	No	Yes	287

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Chronic kidney disease Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have chronic kidney disease	Yes	Yes	No	Yes	288
Chronic kidney disease with knowledge of impaired renal function (percent)	Percent of adults 45 years and older with chronic kidney disease (CKD) who know they have impaired renal function	Yes	No	No	No	485
Chronic obstructive pulmonary disease admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Chronic obstructive pulmonary disease admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2039
Cigar smoking - Adults (percent)	Percent of adults aged 18 years and over smoking cigars	Yes	No	No	No	1500
Cigarette smoking - Adults (percent)	Percent of adults aged 18 years and over smoking cigarettes	Yes	No	No	No	1498
Cirrhosis deaths (per 100,000)	Cirrhosis death rate	Yes	No	No	No	1425
Clinical Laboratory Improvement Act (CHIA) Providers (number)	Number of Clinical Laboratory Improvement Act (CHIA) providers participating in the Medicare program	No	Yes	No	Yes	10035
Cochlear implant surgeries (per 10,000)	Rate of persons who are deaf or very hard of hearing and who have new cochlear implants	Yes	No	No	No	745
College degree, adults 25+ (percent)	Percent of population age 25+ with 4-year college degree or higher	No	Yes	Yes	No	34
College students binge drinking (percent)	Percent of students engaging in binge drinking during the past 2 weeks-College students	Yes	No	No	No	1436
Colorectal cancer deaths (per 100,000)	Colorectal cancer death rate per 100,000	Yes	Yes	Yes	No	86
Colorectal cancer deaths, Healthy People coded (per 100,000)	Colon cancer death rate, using the ICD codes defined by the Healthy People initiative	Yes	No	No	No	490
Colorectal cancer Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have colorectal cancer	Yes	Yes	No	Yes	301
Colorectal cancer Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have colorectal cancer	Yes	Yes	No	Yes	302

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Colorectal cancer screening (percent)	Percent of adults who receive a colorectal cancer screening based on the most recent guidelines	Yes	No	No	No	506
Community Mental Health Centers (number)	Number of Community Mental Health Centers participating in the Medicare program	No	Yes	No	Yes	10036
Community-based primary prevention services - Chronic diseases (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in chronic disease programs	Yes	No	No	No	602
Community-based primary prevention services - Injury (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in injury	Yes	No	No	No	596
Community-based primary prevention services - Mental illness (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in mental illness	Yes	No	No	No	598
Community-based primary prevention services - Nutrition (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in nutrition	Yes	No	No	No	603
Community-based primary prevention services - Physical activity (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in physical activity	Yes	No	No	No	604

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Community-based primary prevention services - Substance abuse (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in substance abuse	Yes	No	No	No	600
Community-based primary prevention services - Tobacco use (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in tobacco	Yes	No	No	No	599
Community-based primary prevention services - Unintended pregnancy (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in unintended pregnancy	Yes	No	No	No	601
Community-based primary prevention services - Violence (percent)	Percent of community-based organizations (including local health departments, tribal health services, nongovernmental organizations, and State agencies) providing population-based primary prevention services in violence	Yes	No	No	No	597
Competencies for public health professionals: Local public health agencies (percent)	Percent of Federal, Tribal, State, and local public health agencies that incorporate core competencies for public health professionals into job descriptions and performance evaluations-Local public health agencies	Yes	No	No	No	1372
Complete vaccination among children (percent)	Percent of children aged 19 to 35 months who receive the recommended doses of DTaP, polio, MMR, Hib, hepatitis B, varicella and PCV	Yes	No	No	No	1008
Comprehensive Outpatient Rehabilitation Facilities (number)	Number of Comprehensive Outpatient Rehabilitation (CORF) Facilities participating in the Medicare program	No	Yes	No	Yes	10037
Condom use among unmarried females (percent)	Percent of sexually active persons who use condoms-Unmarried females aged 15 to 44 years	Yes	No	No	No	940

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Condom use among unmarried males (percent)	Percent of sexually active persons who use condoms- Unmarried males aged 15 to 44 years	Yes	No	No	No	941
Congenital Rubella Syndrome (number)	Number of Congenital Rubella Syndrome (CRS) among children aged 1 year and under (U.S.-acquired cases)	Yes	No	No	No	978
Congenital syphilis (per 100,000)	Congenital syphilis rate	Yes	No	No	No	1474
Congestive heart failure admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Congestive heart failure admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2041
Congregate care - adults (number)	Number of adults with disabilities (aged 22 years and older) living in congregate care facilities with 16 beds or more	Yes	No	No	No	524
Congregate care - children and youth (number)	Number of children and youth with disabilities (aged 21 years and under) living in congregate care facilities with 16 beds or more	Yes	No	No	No	525
Consumer food safety practices - Clean hands (percent)	Percent of consumers who follow key food safety practices-- Clean: Wash hands and surfaces often	Yes	No	No	No	835
Consumer food safety practices - Cook (percent)	Percent of consumers who follow key food safety practices-- Cook: Cook to proper temperatures	Yes	No	No	No	837
Consumer food safety practices - Refrigerate (percent)	Percent of consumers who follow key food safety practices-- Chill: Refrigerate promptly	Yes	No	No	No	838
Consumer food safety practices - Separate food (percent)	Percent of consumers who follow key food safety practices-- Separate: Don't cross-contaminate	Yes	No	No	No	836
Contraceptive use at most recent sexual intercourse (percent)	Percent of females or their partners at risk of unintended pregnancy who used contraception at most recent sexual intercourse	Yes	No	No	No	785
COPD activity limitations (percent)	Percent of adults with chronic obstructive pulmonary disease (COPD) who have activity limitations	Yes	No	No	No	1405
COPD deaths, adults aged 45+ (per 100,000)	Death rate from chronic obstructive pulmonary disease (COPD) among adults 45 years and over	Yes	No	No	No	1406

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
COPD emergency department visits (per 10,000)	Rate of hospital emergency department visits for chronic obstructive pulmonary disease (COPD)	Yes	No	No	No	1423
COPD hospitalizations (per 10,000)	Hospitalization rate for chronic obstructive pulmonary disease (COPD)	Yes	No	No	No	1422
COPD Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have chronic obstructive pulmonary disease	Yes	Yes	No	Yes	289
COPD Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have chronic obstructive pulmonary disease	Yes	Yes	No	Yes	290
Coronary heart disease deaths (per 100,000)	Coronary heart disease death rate	Yes	Yes	Yes	No	877
Counseling for physical activity among adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who receive Health Care provider counseling-For physical activity or exercise	Yes	No	No	No	422
Counseling for weight reduction among adults with arthritis who are overweight (percent)	Percent of adults with doctor-diagnosed arthritis who receive Health Care provider counseling-For weight reduction among overweight and obese persons	Yes	No	No	No	421
Critical Access Hospital Medicare admissions (per 1,000 beneficiaries)	Rate of critical access hospital (CAH) admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2003
Critical Access Hospital Medicare days (per 1,000 beneficiaries)	Average number of critical access hospital (CAH) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2011
Critical access hospital services Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who had covered critical access hospital services	Yes	Yes	No	Yes	174
Critical access hospital services Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered critical access hospital services	Yes	Yes	No	Yes	175
Critical Access Hospitals (number)	Number of Critical Access Hospitals participating in the Medicare program	No	Yes	No	Yes	10038

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Cumulative health instruction meets time standards--elementary school (percent)	Percent of schools that require cumulative instruction in health education that meet the US National Health Education Standards for elementary, middle, and senior high schools-Elementary schools	Yes	No	No	No	734
Cumulative health instruction meets time standards--high school (percent)	Percent of schools that require cumulative instruction in health education that meet the US National Health Education Standards for elementary, middle, and senior high schools-High schools	Yes	No	No	No	736
Cumulative health instruction meets time standards--middle school (percent)	Percent of schools that require cumulative instruction in health education that meet the US National Health Education Standards for elementary, middle, and senior high schools-Middle schools	Yes	No	No	No	735
D.O. training in counseling for health promotion and disease prevention (percent)	Percent of inclusion of core clinical prevention and population health content-Counseling for health promotion and disease prevention in health professions training-DO-granting medical schools	Yes	No	No	No	612
D.O. training in cultural diversity (percent)	Percent of inclusion of core clinical prevention and population health content-Cultural diversity in health professions training-DO-granting medical schools	Yes	No	No	No	613
D.O. training in environmental health (percent)	Percent of inclusion of core clinical prevention and population health content-Environmental health in health professions training-DO-granting medical schools	Yes	No	No	No	615
D.O. training in evaluation of health sciences literature (percent)	Percent of inclusion of core clinical prevention and population health content-Evaluation of health sciences literature in health professions training-DO-granting medical schools	Yes	No	No	No	614
D.O. training in global health (percent)	Percent of inclusion of core clinical prevention and population health content-Global health in health professions training-DO-granting medical schools	Yes	No	No	No	617

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
D.O. training in public health systems (percent)	Percent of inclusion of core clinical prevention and population health content-Public health systems in health professions training-DO-granting medical schools	Yes	No	No	No	616
Death in transplant patients (per 1,000)	Total death rate for persons with a functioning kidney transplant	Yes	No	No	No	466
Deaths among infants with Down syndrome (per 1,000)	One-year mortality rate for infants with Down syndrome	Yes	No	No	No	1187
Deaths due to poor water quality, sanitation, and hygiene worldwide (number)	Global burden of disease resulting in death due to poor water quality, sanitation, and insufficient hygiene	Yes	No	No	No	720
Deaths from falls (per 100,000)	Fall-related death rate: All persons	Yes	Yes	Yes	No	1079
Deaths from falls, ages 65 years and older (per 100,000)	Fall-related death rate: Adults aged 65 years and older	Yes	No	No	No	1080
Deaths, all causes (per 100,000)	Death rate per 100,000 population from all causes of death	Yes	Yes	Yes	No	71
Dehydration admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Dehydration admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2042
Dental caries experience in adolescents (percent)	Percent of adolescents aged 13 to 15 years with dental caries experience in their permanent teeth	Yes	No	No	No	1271
Dental caries experience in children (percent)	Percent of children aged 6 to 9 years with dental caries experience in their primary and permanent teeth	Yes	No	No	No	1270
Dental caries experience in young children (percent)	Percent of young children aged 3 to 5 years with dental caries experience in their primary teeth	Yes	No	No	No	1269
Dental sealants in adolescents (percent)	Percent of adolescents aged 13 to 15 years who have received dental sealants on one or more of their permanent molar teeth	Yes	No	No	No	1283
Dental sealants in children (percent)	Percent of children aged 6 to 9 years who have received dental sealants on one or more of their permanent first molar teeth	Yes	No	No	No	1282

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Dental sealants in young children (percent)	Percent of children aged 3 to 5 years who have received dental sealants on one or more of their primary molar teeth	Yes	No	No	No	1281
Dental services for low-income children (percent)	Percent of low-income children and adolescents who received any preventive dental service during the past year	Yes	No	No	No	1267
Dentist rate (per 100,000 population)	Dentists per capita	No	No	Yes	No	130
Depression Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have depression	Yes	Yes	No	Yes	291
Depression Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have depression	Yes	Yes	No	Yes	292
Depression screening for adults by primary care providers (percent)	Percent of primary care physician office visits that screen adults aged 19 years and older for depression	Yes	No	No	No	1120
Depression screening for youth by primary care providers (percent)	Percent of primary care physician office visits that screen youth aged 12 to 18 years for depression	Yes	No	No	No	1121
Destructive periodontal disease (percent)	Percent of adults aged 45 to 74 years with periodontitis	Yes	No	No	No	1280
Diabetes (percent)	Adults that report having diabetes	No	Yes	Yes	No	125
Diabetes education (percent)	Percent of adults with diagnosed diabetes who receive formal diabetes education	Yes	No	No	No	545
Diabetes long-term complications admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Diabetes long-term complications admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2038
Diabetes Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have diabetes	Yes	Yes	No	Yes	293
Diabetes Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have diabetes	Yes	Yes	No	Yes	294
Diabetes self-management benefits receipt among older adults (percent)	Percent of older adults who receive Diabetes Self-Management Benefits	Yes	No	No	No	1263

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Diabetes-related deaths (per 100,000)	Diabetes death rate	Yes	No	No	No	547
Diagnosed diabetes (percent)	Percent of adults aged 20 years and older with diabetes whose condition has been diagnosed	Yes	No	No	No	546
Diagnosis of von Willebrand Disease (vWD) in women (percent)	Percent of women with von Willebrand disease (vWD) who are timely and accurately diagnosed	Yes	No	No	No	443
Dilated eye examinations (percent)	Percent of adults who have a comprehensive eye examination, including dilation, within the past 2 years	Yes	No	No	No	1566
Disorders diagnosed through newborn blood spot screening (percent)	Percent of children diagnosed with a disorder through newborn blood spot screening who experience developmental delay requiring special education services	Yes	No	No	No	1167
DME Medicare service events (per 1,000 beneficiaries)	Rate of durable medical equipment service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2026
Doctor visits with nutrition counseling (percent)	Percent of physician visits made by all child or adult patients that include counseling about nutrition or diet	Yes	No	No	No	1227
Doctor visits with nutrition counseling: patients with CVD, diabetes, or hyperlipidemia (percent)	Percent of physician office visits made by patients with a diagnosis of cardiovascular disease, diabetes, or hyperlipidemia that include counseling or education related to diet and nutrition	Yes	No	No	No	1225
Doctor visits with weight loss, nutrition or physical activity counseling: obese patients (percent)	Percent of physician office visits made by adult patients who are obese that include counseling or education related to weight reduction, nutrition, or physical activity	Yes	No	No	No	1226
Doctors address parents' developmental concerns --parents of child aged 0-5 years (percent)	Percent of parents who receive information from their doctors or other Health Care professionals when they have a concern about their children's learning, development, or behavior	Yes	No	No	No	741
Domestic water withdrawals (gallons)	Per capita domestic water withdrawals with respect to use and conservation	Yes	No	No	No	673
Drownings (per 100,000)	Drowning death rate	Yes	Yes	Yes	No	1081

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Drug poisoning deaths (per 100,000)	Rate per 100,000 of deaths with underlying causes of drug related poisonings	Yes	Yes	Yes	No	10016
Drug-induced deaths (per 100,000)	Drug-induced death rate	Yes	No	No	No	1426
DTaP vaccination (percent)	Percent of young children who have received universally recommended vaccines-4 doses diphtheria-tetanus-acellular pertussis (DTaP) vaccine by age 19 to 35 months	Yes	No	No	No	993
DTaP vaccination among kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-DTaP vaccine (4 or more doses)	Yes	No	No	No	1003
Durable medical equipment Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who used durable medical equipment services	Yes	Yes	No	Yes	257
Durable medical equipment Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used durable medical equipment services	Yes	Yes	No	Yes	258
Early death in dialysis patients (per 1,000)	Death rate in dialysis patients within the first 3 months of initiation of renal replacement therapy	Yes	No	No	No	464
Early infant diagnostic hearing evaluation (percent)	Percent of infants who did not pass the hearing screening that received audiologic evaluation at no later than age 3 months	Yes	No	No	No	757
Early intervention services for infants with confirmed hearing loss (percent)	Percent of infants with confirmed hearing loss enrolled for intervention services no later than age 6 months	Yes	No	No	No	758
Early intervention services, children under 2 with disabilities (percent)	Percent of children with disabilities, birth through age 2 years, who receive early intervention services in home or community-based settings	Yes	No	No	No	539
Eating disorders (percent)	Percent of adolescents who engage in disordered eating behaviors in an attempt to control their weight	Yes	Yes	No	No	1108
Elder abuse, neglect, and exploitation-States providing information (number)	Number of States-and the District of Columbia-that collect and make publicly available information on the characteristics of victims, perpetrators, and cases of elder abuse, neglect, and exploitation	Yes	No	No	No	1256

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Electronic health records in medical practices (percent)	Percent of medical practices that use electronic health records	Yes	No	No	No	872
Electronic surveillance of rabies (number)	Number of States that use electronic data from rabies animal surveillance to inform public health prevention programs	Yes	No	No	No	1024
Elevated blood lead from work exposures (per 100,000)	Rate of elevated blood lead concentrations from work exposures among adults	Yes	No	No	No	1300
Elevated blood pressure in persons with chronic kidney disease (percent)	Percent of adults with chronic kidney disease who have elevated blood pressure	Yes	No	No	No	480
Elevated lipid levels in chronic kidney disease patients (percent)	Percent of adults with chronic kidney disease who have elevated lipid levels	Yes	No	No	No	481
Emergency department visit rate (per 1,000 beneficiaries)	Rate among beneficiaries of inpatient or hospital outpatient emergency department visits	Yes	Yes	No	Yes	282
Emergency department visits (number)	Total inpatient or hospital outpatient emergency department visits	Yes	Yes	No	Yes	281
Emergency personnel response time (number)	The time necessary to activate designated personnel in response to public health emergency in minutes	Yes	No	No	No	1394
Employment among adults with disabilities (percent)	Employment rates among people with disabilities	Yes	No	No	No	516
Employment of persons with serious mental illness (percent)	Percent of persons with serious mental illness (SMI) who are employed	Yes	No	No	No	1116
End Stage Renal Disease Facilities (number)	Number of End Stage Renal Disease (ESRD) facilities participating in the Medicare program	No	Yes	No	Yes	10039
End-stage kidney failure due to diabetes (per 1,000,000)	Incidence rate of kidney failure due to diabetes	Yes	No	No	No	475
End-stage kidney failure in diabetics (per 1,000,000)	Incidence rate of kidney failure due to diabetes among persons with diabetes	Yes	No	No	No	476
End-stage renal disease (per 1,000,000)	Incidence rate of end-stage renal disease (ESRD)	Yes	No	No	No	462

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
End-stage renal disease deaths (per 1,000)	Total death rate for persons on dialysis	Yes	No	No	No	463
Enforcement of illegal tobacco sales to minors laws - States (number)	Illegal tobacco sales rate of <=5% to minors through enforcement of laws prohibiting the sale of tobacco products to minors-number of States and the District of Columbia	Yes	No	No	No	1493
Enforcement of illegal tobacco sales to minors laws - Territories (number)	Illegal tobacco sales rate of <=5% to minors through enforcement of laws prohibiting the sale of tobacco products to minors-number of Territories	Yes	No	No	No	1494
Epidemiology services: Local public health agencies (percent)	Percent of Tribal, State, and local public health agencies that provide or assure comprehensive epidemiology services to support essential public health services-Local public health agencies	Yes	No	No	No	1363
Epidemiology services: State public health agencies (percent)	Percent of Tribal, State, and local public health agencies that provide or assure comprehensive epidemiology services to support essential public health services-State public health agencies	Yes	No	No	No	1362
Estimated Poverty (percent)	Small area estimates of population living below the poverty line	Yes	Yes	Yes	No	10004
Evaluation for obstructive sleep apnea (percent)	Percent of persons with symptoms of obstructive sleep apnea who seek medical evaluation	Yes	No	No	No	1469
Evaluation of children with Autism spectrum disorders (percent)	Percent of children with an ASD with a first evaluation by 36 months of age	Yes	No	No	No	1189
Excessive drinking (percent)	Percent of adults who drank excessively in the previous 30 days	Yes	No	No	No	1458
Exposure to environmental chemicals - arsenic (ug/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: arsenic	Yes	No	No	No	695

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Exposure to environmental chemicals - BDE 47 (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: BDE 47, (2,2',4,4'-tetrabromodiphenyl ether)	Yes	No	No	No	712
Exposure to environmental chemicals - beta-HCH (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: beta-hexachlorocyclohexane or beta-HCH	Yes	No	No	No	702
Exposure to environmental chemicals - bisphenol A (ug/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Bisphenol A	Yes	No	No	No	709
Exposure to environmental chemicals - cadmium (ug/L)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: cadmium	Yes	No	No	No	696
Exposure to environmental chemicals - chlordane (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Chlordane (Oxychlordane)	Yes	No	No	No	700
Exposure to environmental chemicals - chlorpyrifos (ug/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: 3,4,6-trichloro-2-pyridinol (chlorpyrifos)	Yes	No	No	No	704
Exposure to environmental chemicals - DDT (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: DDT (DDE)	Yes	No	No	No	701
Exposure to environmental chemicals - dioxin-like PCBs (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: PCB 126, representative of dioxin-like PCBs	Yes	No	No	No	707

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Exposure to environmental chemicals - dioxins (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: 1,2,3,6,7,8-hexachlorodibenzo-p-dioxin, representative of the dioxin class	Yes	No	No	No	708
Exposure to environmental chemicals - lead (ug/dL)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: lead	Yes	No	No	No	697
Exposure to environmental chemicals - mercury (children) (ug/L)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Mercury, children aged 1 to 5 years	Yes	No	No	No	698
Exposure to environmental chemicals - mercury (females) (ug/L)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Mercury, females aged 16 to 49 years	Yes	No	No	No	699
Exposure to environmental chemicals - nondioxin-like PCBs (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: PCB 153, representative of nondioxin-like PCBs	Yes	No	No	No	706
Exposure to environmental chemicals - para-Nitrophenol (ug/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Paranitrophenol (methyl parathion and parathions)	Yes	No	No	No	703
Exposure to environmental chemicals - perchlorate (ug/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: perchlorate	Yes	No	No	No	710
Exposure to environmental chemicals - phthalates (ug/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: Mono-n-butyl phthalate	Yes	No	No	No	711

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Exposure to environmental chemicals - pyrethroids (ug/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations of the substances or their metabolites: 3-phenoxybenzoic acid	Yes	No	No	No	705
Exposure to secondhand smoke--nonsmokers aged 12 to 17 years (percent)	Percent of nonsmokers exposed to secondhand smoke--Adolescents aged 12 to 17 years	Yes	No	No	No	1518
Exposure to secondhand smoke--nonsmokers aged 18 years and over (percent)	Percent of nonsmokers exposed to secondhand smoke--Adults aged 18 years and older	Yes	No	No	No	1519
Exposure to secondhand smoke--nonsmokers aged 3 to 11 years (percent)	Percent of nonsmokers exposed to secondhand smoke--Children aged 3 to 11 years	Yes	No	No	No	1517
Extracurricular and out-of-school activities (percent)	Percent of adolescents who participate in extracurricular and out-of-school activities	Yes	No	No	No	404
Fair or poor health: adults (percent)	Percent of adults that report fair or poor health	No	Yes	Yes	No	5
Falling of adults with balance and dizziness problems (percent)	Percent of adults with balance and dizziness problems who have fallen in the past 5 years while experiencing symptoms of dizziness, vertigo, or imbalance	Yes	No	No	No	769
Falls among older adults: Emergency department visits (per 100,000)	Rate of emergency department visits due to falls among older adults	Yes	No	No	No	1259
Federal and State tax increases on tobacco products-Cigarettes (number)	Number of States, DC, and Federal government that have increased tax on cigarettes by \$1.50	Yes	No	No	No	1548
Federal and State tax increases on tobacco products-Smokeless tobacco products (number)	Number of States, DC, and Federal government that have increased tax on smokeless tobacco products by \$1.50	Yes	No	No	No	1549
Federal Qualified Health Centers with oral health components (percent)	Percent of Federally Qualified Health Centers that have an oral health care program	Yes	No	No	No	1287

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Federally Qualified Health Centers (percent)	Number of Federally Qualified Health Centers (FQHC) participating in the Medicare program	No	Yes	No	Yes	10040
Female Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months in who are female	Yes	Yes	No	Yes	309
Female teens using a condom & hormonal or intrauterine contraception: first intercourse (percent)	Percent of sexually active females aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at first intercourse	Yes	No	No	No	798
Female teens using a condom & hormonal or intrauterine contraception: last intercourse (percent)	Percent of sexually active females aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at last intercourse	Yes	No	No	No	800
Females aged 15 to 17 years who have never had sexual intercourse (percent)	Percent of female adolescents aged 15 to 17 years who have never had sexual intercourse	Yes	No	No	No	790
Females aged 15 years and under who have never had sexual intercourse (percent)	Percent of female adolescents aged 15 years and under who have never had sexual intercourse	Yes	No	No	No	792
Females receiving publicly supported contraceptive services and supplies (percent)	Percent of females in need of publicly supported contraceptive services and supplies who receive those services and supplies	Yes	No	No	No	819
Fetal alcohol syndrome (per 10,000 live births)	Rate of fetal alcohol syndrome (FAS)	Yes	No	No	No	1146
Fetal deaths (per 1,000 live births plus fetal deaths)	Rate of fetal deaths at 20 or more weeks of gestation	Yes	No	No	No	1156
Few fruits/vegetables (percent)	Adults that report fewer than 5 servings of fruits/vegetables per day	No	Yes	Yes	No	121
Firearm-related deaths (per 100,000)	Firearm-related death rate	Yes	Yes	Yes	No	1042
Flu vaccination, adults (percent)	Percent of adults age 65+ that report having Influenza vaccine in past 12 months	No	Yes	Yes	No	119
Fluoridation of community water (percent)	Percent of the U.S. population served by community water systems with optimally fluoridated water	Yes	Yes	No	No	1265

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Folic acid intake in reproductive-aged women (percent)	Percent of women of childbearing potential with intake of at least 400 µg of folic acid from fortified foods or dietary supplements	Yes	No	No	No	1171
Follow-up to newborn screening (percent)	Percent of screen-positive children who receive follow-up testing within the recommended time period	Yes	No	No	No	1174
Formula supplementation in breastfed newborns (percent)	Percent of breastfed newborns who receive formula supplementation within the first 2 days of life	Yes	No	No	No	1185
Free school lunch eligibility (percent)	Percent of total students eligible to participate in the National School Lunch Program under the Free Lunch Act	No	Yes	Yes	No	10005
Fruit consumption (cup equivalents per 1,000 calories)	Mean daily intake of fruit (in cup equivalents per 1,000 calories) by persons aged 2 years and older	Yes	No	No	No	1217
Functional limitations in older adults (percent)	Percent of older adults 65 years and over who have moderate to severe functional limitations	Yes	No	No	No	1245
Genetic counseling for women with a family history of breast and/or ovarian cancer (percent)	Percent of women with a family history of breast and/or ovarian cancer who receive genetic counseling	Yes	No	No	No	850
Genital herpes simplex type 2 (percent)	Percent of young adults with genital herpes infection due to herpes simplex type 2	Yes	No	No	No	1482
Geriatric dentists (percent)	Percent of the health care workforce with geriatric certification-Dentists	Yes	No	No	No	1253
Geriatric dieticians (percent)	Percent of the health care workforce with geriatric certification-Registered dieticians	Yes	No	No	No	1255
Geriatric nurses (percent)	Percent of the health care workforce with geriatric certification-Registered nurses	Yes	No	No	No	1252
Geriatric physical therapists (percent)	Percent of the health care workforce with geriatric certification-Physical therapists	Yes	No	No	No	1254
Geriatric physicians (percent)	Percent of the health care workforce with geriatric certification-Physicians	Yes	No	No	No	1250

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Geriatric psychiatrists (percent)	Percent of the health care workforce with geriatric certification-Geriatric psychiatrists	Yes	No	No	No	1251
Giardiasis (number)	Number of reported cases of giardiasis	No	Yes	Yes	No	10021
Gini coefficient (ratio)	Gini coefficient of household income inequality	No	Yes	Yes	No	37
Global Disease Detection (GDD) Regional Centers worldwide (number)	Number of Global Disease Detection (GDD) Regional Centers worldwide to detect and contain emerging health threats	Yes	No	No	No	846
Global Disease Detection (GDD): diagnostic testing capacity (number)	Diagnostic testing capacity in host countries and regionally through the Global Disease Detection (GDD) Regional Centers	Yes	No	No	No	848
Global Disease Detection (GDD): public health professionals trained (number)	Number of public health professionals trained by Global Disease Detection: Global Disease Detection (GDD) programs worldwide	Yes	No	No	No	847
Gonorrhea among Females 15 to 44 years (per 100,000)	Gonorrhea rate among females aged 15 to 44 years	Yes	No	No	No	1478
Gonorrhea among males 15 to 44 years (per 100,000)	Gonorrhea rate among males aged 15 to 44 years	Yes	No	No	No	1479
Gonorrhea rate (per 100,000)	Rate of gonorrhea cases per 100,000 population	No	Yes	Yes	No	10062
Graduate with High School Diploma on time (percent)	Percent of public school students who graduate with a regular diploma 4 years after starting 9th grade	Yes	No	No	No	394
Group B streptococcal disease among newborns (per 1,000)	Early onset group B streptococcal disease rate	Yes	No	No	No	973
HbA1c greater than 9 percent--persons with diabetes (percent)	Percent of adults with diagnosed diabetes with an HbA1c value greater than 9 percent	Yes	No	No	No	558
HbA1c less than 7 percent--persons with diabetes (percent)	Percent of adults with diagnosed diabetes with an HbA1c value less than 7 percent	Yes	No	No	No	559
HbA1c Test, at least two times a year --persons with diabetes (percent)	Percent of adults with diabetes who have a glycosylated hemoglobin measurement at least twice a year	Yes	No	No	No	550

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
HbA1c test: diabetic Medicare beneficiaries (percent)	Percent of diabetic Medicare enrollees receiving a hemoglobin A1c test in the past year	No	Yes	Yes	No	29
Health (medical) insurance (percent)	Percent of persons under age 65 years with health (medical) insurance	Yes	No	No	No	368
Health Care evaluation of adults with balance or dizziness problems (percent)	Percent of adults with balance or dizziness problems in the past 12 months who have ever seen a Health Care provider about their balance or dizziness problems	Yes	No	No	No	762
Health care evaluation of adults with tinnitus (percent)	Percent of adults bothered by tinnitus in the past 12 months who have seen a doctor	Yes	No	No	No	759
Health care organizations monitoring and analyzing adverse events (percent)	Percent of health care organizations that are monitoring and analyzing adverse events associated with medical therapies within their systems	Yes	No	No	No	1196
Health care provider explaining things (percent)	Percent of persons who report that their Health Care provider always explained things so they could understand them	Yes	No	No	No	854
Health care provider listening skills (percent)	Percent of persons who report that their Health Care provider always listened carefully to them	Yes	No	No	No	853
Health care provider referrals to hearing specialists (per 1,000)	Rate of persons who are referred by their primary care physician or other health care provider for hearing evaluation and treatment	Yes	No	No	No	751
Health Care provider showing respect (percent)	Percent of persons who report that their Health Care provider always showed respect for what they had to say	Yes	No	No	No	855
Health care provider spending enough time with patient (percent)	Percent of persons who report that their health care provider always spent enough time with them	Yes	No	No	No	856
Health education certification requirements--elementary school (percent)	Percent of schools that require newly hired staff who teach required health instruction to be certified, licensed, or endorsed by the State in health education-Elementary schools	Yes	No	No	No	731

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Health education certification requirements--high school (percent)	Percent of schools that require newly hired staff who teach required health instruction to be certified, licensed, or endorsed by the State in health education-High schools	Yes	No	No	No	733
Health education certification requirements--middle school (percent)	Percent of schools that require newly hired staff who teach required health instruction to be certified, licensed, or endorsed by the State in health education-Middle schools	Yes	No	No	No	732
Health education training requirements--elementary school (percent)	Percent of schools that require newly hired staff who teach required health education to have undergraduate or graduate training in health education-Elementary schools	Yes	No	No	No	728
Health education training requirements--high school (percent)	Percent of schools that require newly hired staff who teach required health education to have undergraduate or graduate training in health education-High schools	Yes	No	No	No	730
Health education training requirements--middle school (percent)	Percent of schools that require newly hired staff who teach required health education to have undergraduate or graduate training in health education-Middle schools	Yes	No	No	No	729
Health Professions Shortage Area (HPSA) Indicator (yes/no)	Indicator for single county designated Health Professional Shortage Area	No	No	Yes	No	132
Health promotion programs for caregivers (States and DC) (number)	States and the District of Columbia with health promotion programs for caregivers	Yes	No	No	No	529
Health promotion programs for people with disabilities (States and DC) (number)	States and the District of Columbia with health promotion programs for people with disabilities	Yes	No	No	No	527
Health Websites meeting three or more evaluation criteria (percent)	Percent of health-related Websites that meet three or more evaluation criteria disclosing information that can be used to assess information reliability	Yes	No	No	No	863
Health-risk behavior information in higher education - Alcohol and drug use (percent)	Percent of college and university students who receive information from their institution on alcohol and other drug use	Yes	No	No	No	580

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Health-risk behavior information in higher education - HIV and STD (percent)	Percent of college and university students who receive information from their institution on HIV, AIDS and STD infection	Yes	No	No	No	582
Health-risk behavior information in higher education - Injury (percent)	Percent of college and university students who receive information from their institution on unintentional injury	Yes	No	No	No	576
Health-risk behavior information in higher education - Nutrition (percent)	Percent of college and university students who receive information from their institution on unhealthy dietary patterns	Yes	No	No	No	583
Health-risk behavior information in higher education - Physical activity (percent)	Percent of college and university students who receive information from their institution on inadequate physical activity	Yes	No	No	No	584
Health-risk behavior information in higher education - Suicide (percent)	Percent of college and university students who receive information from their institution on suicide	Yes	No	No	No	578
Health-risk behavior information in higher education - Tobacco use (percent)	percent of college and university students who receive information from their institution on tobacco use and addiction	Yes	No	No	No	579
Health-risk behavior information in higher education - Unintended pregnancy (percent)	Percent of college and university students who receive information from their institution on unintended pregnancy	Yes	No	No	No	581
Health-risk behavior information in higher education - Violence (percent)	Percent of college and university students who receive information from their institution on violence	Yes	No	No	No	577
Healthy food outlets (percent)	Percent of zip codes in county with a healthy food outlet	No	Yes	Yes	No	48
Healthy weight in adults (percent)	Percent of adults who are at a healthy weight	Yes	No	No	No	1207
Healthy weight prior to pregnancy (percent)	Percent of women delivering a live birth who had a healthy weight prior to pregnancy	Yes	No	No	No	1182
Hearing aid use among adults (per 1,000)	Rate of adults aged 20 to 69 years with hearing loss who have ever used a hearing aid	Yes	No	No	No	744
Hearing aid use among older adults (per 1,000)	Rate of adults aged 70 years and older with hearing loss who have ever used a hearing aid	Yes	No	No	No	746

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Heart attack Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have had a heart attack	Yes	Yes	No	Yes	283
Heart attack Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have had a heart attack	Yes	Yes	No	Yes	284
Heart attack patients given aspirin at hospital arrival (percent)	Hospital Compare: Heart attack patients given aspirin at hospital arrival	Yes	Yes	No	Yes	319
Heart attack patients prescribed angiotensin converting enzyme inhibitor or angiotensin receptor blocker at hospital discharge (percent)	Hospital Compare: Heart attack patients prescribed angiotensin converting enzyme inhibitor or angiotensin receptor blocker at hospital discharge	Yes	Yes	No	Yes	321
Heart attack patients with aspirin prescribed at hospital discharge (percent)	Hospital Compare: Heart attack patients with aspirin prescribed at hospital discharge	Yes	Yes	No	Yes	320
Heart attack patients with beta blocker prescribed at hospital discharge (percent)	Hospital Compare: Heart attack patients with beta blocker prescribed at hospital discharge	Yes	Yes	No	Yes	323
Heart attack patients with fibrinolytic received within 30 minutes of hospital arrival (percent)	Hospital Compare: Heart attack patients with fibrinolytic received within 30 minutes of hospital arrival	Yes	Yes	No	Yes	324
Heart attack patients with percutaneous coronary intervention within 90 minutes of hospital arrival (percent)	Hospital Compare: Heart attack patients with percutaneous coronary intervention within 90 minutes of hospital arrival	Yes	Yes	No	Yes	325
Heart attack patients with smoking cessation counseling during hospital stay (percent)	Hospital Compare: Heart attack patients with smoking cessation counseling during hospital stay	Yes	Yes	No	Yes	322
Heart disease deaths (per 100,000)	Heart disease death rate	Yes	Yes	Yes	No	83
Heart failure hospitalizations, adults aged 65-74 years (per 1,000)	Rate of hospitalizations of older adults with heart failure as the principal diagnosis-Adults aged 65 to 74 years	Yes	No	No	No	891

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Heart failure hospitalizations, adults aged 75-84 years (per 1,000)	Rate of hospitalizations of older adults with heart failure as the principal diagnosis-Adults aged 75 to 84 years	Yes	No	No	No	892
Heart failure hospitalizations, adults aged 85 years and older (per 1,000)	Rate of hospitalizations of older adults with heart failure as the principal diagnosis-Adults aged 85 years and older	Yes	No	No	No	893
Heart failure Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have heart failure	Yes	Yes	No	Yes	295
Heart failure Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have heart failure	Yes	Yes	No	Yes	296
Heart failure patients prescribed angiotensin converting enzyme inhibitor or angiotensin receptor blocker at hospital discharge (percent)	Hospital Compare: Heart failure patients prescribed angiotensin converting enzyme inhibitor or angiotensin receptor blocker at hospital discharge	Yes	Yes	No	Yes	328
Heart failure patients with discharge instructions (percent)	Hospital Compare: Heart failure patients with discharge instructions	Yes	Yes	No	Yes	326
Heart failure patients with evaluation of left ventricular systolic function (percent)	Hospital Compare: Heart failure patients with evaluation of left ventricular systolic function	Yes	Yes	No	Yes	327
Heart failure patients with smoking cessation counseling (percent)	Hospital Compare: Heart failure patients with smoking cessation counseling	Yes	Yes	No	Yes	329
Hep B vaccination (percent)	Percent of young children who have received universally recommended vaccines-3 doses hepatitis B (hep B) vaccine by age 19 to 35 months	Yes	No	No	No	995
Hepatitis A (per 100,000)	Hepatitis A rate	Yes	No	No	No	967
Hepatitis A vaccination (percent)	Percent of young children who have received universally recommended vaccines-2 doses hepatitis A vaccine by age 19 to 35 months	Yes	No	No	No	1000
Hepatitis A, acute (number)	Number of reported cases of acute Hepatitis A	No	Yes	Yes	No	10022
Hepatitis B among adults (per 100,000)	Rate of new hepatitis B infections in adults aged 19 and older	Yes	No	No	No	988

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Hepatitis B among injection drug users (number)	Number of new hepatitis B infections among high-risk populations: Injection drug users	Yes	No	No	No	989
Hepatitis B among MSM (number)	Number of new hepatitis B infections among high-risk populations: Men who have sex with men	Yes	No	No	No	991
Hepatitis B vaccination among health care workers (percent)	Percent of Hepatitis B vaccine coverage among high-risk populations-Health care personnel	Yes	No	No	No	1016
Hepatitis B vaccination among kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-Hepatitis B vaccine (3 or more doses)	Yes	No	No	No	1006
Hepatitis B, acute (number)	Number of reported cases of acute Hepatitis B	No	Yes	Yes	No	10023
Hepatitis C (per 100,000)	Rate of new hepatitis C infections	Yes	No	No	No	992
Hepatitis C infection status awareness (percent)	Percent of persons aware they have a hepatitis C infection	Yes	No	No	No	1022
Hib invasive disease (number)	Number of serotype b cases of Haemophilus influenzae (Hib) invasive disease (children aged 5 years and under)	Yes	No	No	No	979
Hib vaccination (percent)	Percent of young children who have received universally recommended vaccines-3 doses or more Haemophilus influenzae type b (Hib) vaccine by age 19 to 35 months	Yes	No	No	No	994
High blood cholesterol levels (percent)	Percent of adults with high total blood cholesterol levels	Yes	No	No	No	885
High blood pressure (percent)	Adults that report high blood pressure	No	Yes	Yes	No	123
High blood pressure control among adults (percent)	Percent of adults with hypertension whose blood pressure is under control	Yes	No	No	No	882
High school completion (percent)	Percent of persons aged 18-24 that completes high school education	Yes	No	No	No	565
High school diploma and disabilities (percent)	Percent of students who are served under the Individuals with Disabilities Education Act who graduate high school with a diploma	Yes	No	No	No	395
High school on-time graduation (percent)	Percent of public high school students who graduate with a regular diploma 4 years after starting 9th grade	No	Yes	Yes	No	33

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
High school seniors binge drinking (percent)	Percent of students engaging in binge drinking during the past 2 weeks-High school seniors	Yes	No	No	No	1435
High school seniors disapproval of heavy drinking (percent)	Percent of adolescents who disapprove of having one or two alcoholic drinks nearly every day-12th graders	Yes	No	No	No	1446
High school seniors disapproval of trying marijuana (percent)	Percent of adolescents who disapprove of trying marijuana or hashish once or twice-12th graders	Yes	No	No	No	1449
High school seniors never using alcohol (percent)	Percent of high school seniors never using substances-Alcoholic beverages	Yes	No	No	No	1430
High school seniors never using illicit drugs (percent)	Percent of high school seniors never using substances-Illicit drugs	Yes	No	No	No	1431
High school seniors steroid use (percent)	Percent of 12th graders using steroids	Yes	No	No	No	1442
Hip fractures-Females 65+ (per 100,000)	Rate of hip fractures among older adults: Females aged 65 years and older	Yes	No	No	No	429
Hip fractures-Males 65+ (per 100,000)	Rate of hip fractures among older adults: Males aged 65 years and older	Yes	No	No	No	430
Hispanic Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months in who are Hispanic	Yes	Yes	No	Yes	313
HIV deaths (per 100,000)	HIV infection mortality rate	Yes	Yes	Yes	No	935
HIV prevalence (per 100,000)	Rate of adolescents and adults aged 13 years and over living with HIV	No	Yes	Yes	No	10018
HIV testing among adolescents and adults (percent)	Percent of adolescents and adults who have been tested for HIV in the past 12 months	Yes	No	No	No	944
HIV testing among adolescents and young adults (percent)	Percent of adolescents and young adults who have been tested for HIV in the past 12 months	Yes	No	No	No	947
HIV testing among pregnant women (percent)	Percent of pregnant women who have been tested for HIV during their pregnancy	Yes	No	No	No	946

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
HIV testing in TB patients (percent)	Percent of adults with tuberculosis (TB) who have been tested for HIV	Yes	No	No	No	934
HIV transmission rate (per 100)	Rate of HIV transmission per 100 persons with HIV among adolescents and adults	Yes	No	No	No	959
HIV/AIDS education in substance abuse treatment programs (percent)	Percent of substance abuse treatment facilities that offer HIV/AIDS education, counseling, and support	Yes	No	No	No	933
Home Health Agencies (number)	Number of Home Health Agencies participating in the Medicare program	No	Yes	No	Yes	10041
Home health Medicare episodes (per 1,000 beneficiaries)	Average number of home health episodes among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2018
Home health Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who used home health services	Yes	Yes	No	Yes	221
Home health Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used home health services	Yes	Yes	No	Yes	222
Home health Medicare visits (per 1,000 beneficiaries)	Average number of home health visits among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2017
Homes with lead-based paint - dust-lead hazards. (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US homes that have dust-lead hazards	Yes	No	No	No	723
Homes with lead-based paint - lead-based paint (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US homes that are found to have lead-based paint	Yes	No	No	No	721
Homes with lead-based paint - paint-lead hazards. (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US homes that have paint-lead hazards	Yes	No	No	No	722
Homes with lead-based paint - soil-lead hazards. (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US homes that have soil-lead hazards	Yes	No	No	No	724
Homicide deaths (per 100,000)	Homicide mortality rate	Yes	Yes	Yes	No	1041

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Hospice Facilities (number)	Number of Hospice Facilities participating in the Medicare program	No	Yes	No	Yes	10042
Hospice Medicare admissions (per 1,000 beneficiaries)	Rate of hospice admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2007
Hospice Medicare days (per 1,000 beneficiaries)	Average number of hospice days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2015
Hospice Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who had covered hospice services	Yes	Yes	No	Yes	227
Hospice Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered hospice services	Yes	Yes	No	Yes	228
Hospital 30-day death (mortality) for heart attack patients (percent)	Hospital Compare: Hospital 30-day death (mortality) rates for heart attack patients	Yes	Yes	No	Yes	343
Hospital 30-day death (mortality) for heart failure patients (percent)	Hospital Compare: Hospital 30-day death (mortality) rates for heart failure patients	Yes	Yes	No	Yes	344
Hospital 30-day death (mortality) for pneumonia patients (percent)	Hospital Compare: Hospital 30-day death (mortality) rates for pneumonia patients	Yes	Yes	No	Yes	345
Hospital 30-day readmission for heart attack patients (percent)	Hospital Compare: Hospital 30-day readmission rates for heart attack patients	Yes	Yes	No	Yes	346
Hospital 30-day readmission for heart failure patients (percent)	Hospital Compare: Hospital 30-day readmission rates for heart failure patients	Yes	Yes	No	Yes	347
Hospital 30-day readmission for pneumonia patients (percent)	Hospital Compare: Hospital 30-day readmission rates for pneumonia patients	Yes	Yes	No	Yes	348
Hospital inpatient Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who had covered hospital inpatient services	Yes	Yes	No	Yes	160
Hospital inpatient Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered hospital inpatient services	Yes	Yes	No	Yes	161
Hospital IP Medicare admissions (per 1,000 beneficiaries)	Rate of hospital inpatient admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2001

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Hospital IP Medicare days (per 1,000 beneficiaries)	Average number of hospital inpatient (IP) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2009
Hospital OP Medicare visits (per 1,000 beneficiaries)	Rate of hospital outpatient (OP) visits among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2019
Housing units with physical problems (percent)	Percent of occupied housing units that have moderate or severe physical problems	Yes	No	No	No	663
HPV vaccination among adolescents (percent)	Percent of female adolescents who have 3 doses Human papillomavirus vaccine (HPV) by age 13 to 15 years	Yes	No	No	No	1013
Hypertension admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Hypertension admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2040
Hypertension medication compliance (percent)	Percent of adults with hypertension who are taking the prescribed medications to lower their blood pressure	Yes	No	No	No	912
Hypertension, adults (percent)	Percent of adults with hypertension	Yes	No	No	No	897
Hypertension, children and adolescents (percent)	Percent of children and adolescents with hypertension	Yes	No	No	No	898
Iatrogenic pneumothorax rate (per 100,000 beneficiaries)	Patient Safety Indicators: Iatrogenic pneumothorax rate (observed, among medical and surgical discharges based on secondary diagnosis field) among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2033
Identification of "people with disabilities" in data systems (number)	Percent of major applicable Healthy People 2020 data systems with a standardized set of questions that identify "people with disabilities"	Yes	No	No	No	513
IHS programs with a dental professional directing their dental program (number)	Number of Indian Health Service Areas and Tribal health programs that serve jurisdictions of 30,000 or more persons with a dental public health program directed by a dental professional with public health training	Yes	No	No	No	1292

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Illegal drugs on school property (percent)	Percent of adolescents who have been offered, sold, or given an illegal drug on school property	Yes	Yes	No	No	411
Illicit drug treatment receipt (percent)	Percent of persons who need illicit drug treatment and received specialty treatment for abuse or dependence in the past year-Illicit drug treatment	Yes	No	No	No	1454
Imaging Medicare service events (per 1,000 beneficiaries)	Rate of imaging service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2023
Imaging Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who used imaging services	Yes	Yes	No	Yes	251
Imaging Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used imaging services	Yes	Yes	No	Yes	252
Immunization information system participation by States (number)	Number of States that have 80 percent of adolescents with two or more age-appropriate immunizations recorded in immunization information systems (adolescents aged 11 to 18 years)	Yes	No	No	No	1025
Impaired fecundity - women (percent)	Percent of women aged 18 to 44 years who have impaired fecundity	Yes	No	No	No	1193
Impairment due to cataract (per 1,000)	Rate of visual impairment due to cataract	Yes	No	No	No	1570
Impairment due to diabetic retinopathy (per 1,000)	Rate of visual impairment due to diabetic retinopathy	Yes	No	No	No	1568
Impairment due to glaucoma (per 1,000)	Rate of visual impairment due to glaucoma	Yes	No	No	No	1569
Impairment due to macular degeneration (per 1,000)	Rate of visual impairment due to age-related macular degeneration (AMD)	Yes	No	No	No	1571
Impairment due to refractive errors (per 1,000)	Rate of visual impairment due to uncorrected refractive error	Yes	No	No	No	1567
Improvement plans (number)	Time reported for State public health agencies to establish improvement plans following responses to public health emergencies and exercises in minutes	Yes	No	No	No	1397

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Inability to obtain or delay in obtaining dental care (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary dental care	Yes	No	No	No	381
Inability to obtain or delay in obtaining medical care (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary medical care	Yes	No	No	No	380
Inability to obtain or delay in obtaining medical care, dental care, or prescription medicines (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary medical care, dental care, or prescription medicines	Yes	No	No	No	379
Inability to obtain or delay in obtaining prescription medicines (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary prescription medicines	Yes	No	No	No	382
Inclusion of children and youth with disabilities in regular education programs (percent)	Percent of children and youth with disabilities in classroom settings who spend at least 80 percent of their time in regular education programs	Yes	No	No	No	517
Income eligibility level for Medicaid-covered family planning services (number)	Number of States that set the income eligibility level for Medicaid-covered family planning services to at least the same level used to determine eligibility for Medicaid-covered, pregnancy-related care	Yes	No	No	No	818
Increasing physical activity--persons with pre-diabetes (percent)	Percent of persons at high risk for diabetes (with pre-diabetes) who report increasing their levels of physical activity	Yes	No	No	No	562
Indoor allergen levels - cockroach (units of allergen/grams of settled dust)	Indoor allergen levels: cockroach	Yes	No	No	No	679
Indoor allergen levels - mouse (micrograms of allergen/grams of settled dust)	Indoor allergen levels: mouse	Yes	No	No	No	680
Indoor worksite policies that prohibit smoking (percent)	Percent of persons covered by indoor worksite policies that prohibit smoking	Yes	No	No	No	1492
Infant deaths due to birth defects (per 1,000 live births)	Rate of infant deaths related to birth defects (all birth defects)	Yes	No	No	No	1161

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Infant deaths due to congenital heart defects (per 1,000 live births)	Rate of infant deaths related to birth defects (congenital heart defects)	Yes	No	No	No	1162
Infant deaths from sudden infant death syndrome (per 1,000 live births)	Rate of infant deaths from sudden infant death syndrome (SIDS)	Yes	No	No	No	1163
Infant deaths from sudden unexpected/unexplained causes (per 1,000 live births)	Rate of infant deaths from sudden unexpected/unexplained causes (includes SIDS, Unknown Cause, Accidental Suffocation, and Strangulation in Bed)	Yes	No	No	No	1164
Infants put to sleep on their backs (percent)	Percent of infants who are put down to sleep on their backs	Yes	No	No	No	1141
Infections caused by c. Listeria monocytogenes (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Reduction in infections caused by Listeria monocytogenes	Yes	No	No	No	824
Infections caused by Campylobacter species (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Campylobacter species	Yes	No	No	No	822
Infections caused by Escherichia coli (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Reduction of Shiga toxin-producing Escherichia coli (STEC) O157	Yes	No	No	No	823
Infections caused by Postdiarrheal hemolytic-uremic syndrome (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Postdiarrheal hemolytic uremic syndrome (HUS) in children less than 5 years of age	Yes	No	No	No	826
Infections caused by Salmonella species (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Reduction in infections caused by Salmonella species	Yes	No	No	No	825
Infections caused by Vibrio species (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Vibrio species	Yes	No	No	No	827
Infections caused by Yersinia species (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Yersinia species	Yes	No	No	No	828
Influenza-virus resistance to antiviral agents monitoring (number)	Number of public health laboratories monitoring influenza-virus resistance to antiviral agents	Yes	No	No	No	1028

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Information systems used for environmental health (number)	Quality, utility, awareness, and use of existing information systems for environmental health among states	Yes	No	No	No	664
Initiation of cigar use- adolescents (percent)	Percent of children and adolescents aged 12 to 17 years initiating cigar use in past 12 months	Yes	No	No	No	1508
Initiation of cigar use- young adults (percent)	Percent of young adults aged 18 to 25 years initiating cigar use in the past 12 months	Yes	No	No	No	1512
Initiation of cigarette use - adolescents (percent)	Percent of children and adolescents aged 12 to 17 years initiating cigarette use in past 12 months	Yes	No	No	No	1506
Initiation of cigarette use - young adults (percent)	Percent of young adults aged 18 to 25 years initiating cigarette use in the past 12 months	Yes	No	No	No	1510
Initiation of smokeless tobacco use- adolescents (percent)	Percent of children and adolescents aged 12 to 17 years initiating smokeless tobacco use in past 12 months	Yes	No	No	No	1507
Initiation of smokeless tobacco use- young adults (percent)	Percent of young adults aged 18 to 25 years initiating smokeless tobacco use in the past 12 months	Yes	No	No	No	1511
Initiation of tobacco use - adolescents (percent)	Percent of children and adolescents aged 12 to 17 years initiating tobacco use in past 12 months	Yes	No	No	No	1505
Initiation of tobacco use - young adults (percent)	Percent of young adults aged 18 to 25 years initiating tobacco products use in the past 12 months	Yes	No	No	No	1509
Injectable antidiabetic agents-ED visits for overdoses (per 10,000)	Rate of emergency department (ED) visits for overdoses from injectable antidiabetic agents	Yes	No	No	No	1204
Injuries due to falling of adults with balance and dizziness problems (percent)	Percent of adults with balance and dizziness problems who have been injured as a result of a fall for any reason in the past 12 months	Yes	No	No	No	770
Injury deaths (per 100,000)	Injury death rate	Yes	Yes	Yes	No	1071
Injury ED visits (per 100,000)	Rate of emergency department visits for nonfatal injuries	Yes	No	No	No	1073
Injury hospitalizations (per 100,000)	Hospitalization rate for nonfatal injuries	Yes	No	No	No	1072
Inpatient rehab Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who had covered inpatient rehabilitation services	Yes	Yes	No	Yes	207

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Inpatient rehab Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered inpatient rehabilitation facility services	Yes	Yes	No	Yes	208
Integration of Core Competencies in Public Health into curricula (percent)	Percent of Council on Education for Public Health (CEPH) accredited schools of public health, CEPH accredited academic programs, and schools of nursing (with a public health or community health component) that integrate Core Competencies for Public Health Professionals into curricula	Yes	No	No	No	1373
Intended pregnancies (percent)	Percent of pregnancies that are intended	Yes	No	No	No	779
Internet access (percent)	Percent of persons with access to the Internet	Yes	No	No	No	857
Interruption of activities of adults with balance or dizziness problems (percent)	Percent of adults with balance and dizziness problems in the past 12 months who were prevented from doing regular activities within the home or outside	Yes	No	No	No	767
Invasive colorectal cancer (per 100,000)	Invasive colorectal cancer incidence	Yes	No	No	No	509
Invasive pneumococcal infections among children (per 100,000)	Rate of new invasive pneumococcal infections among children under age 5 years	Yes	No	No	No	963
Invasive pneumococcal infections among elderly (per 100,000)	Rate of new invasive pneumococcal infections among adults aged 65 years and older	Yes	No	No	No	964
Invasive uterine cervical cancer (per 100,000)	Invasive uterine cervical cancer incidence	Yes	No	No	No	510
Involvement in health care decision making (percent)	Percent of persons who report that their health care providers always involved them in decisions about their health care as much as they wanted	Yes	No	No	No	866
IRF Medicare admissions (per 1,000 beneficiaries)	Rate of inpatient rehabilitation facility (IRF) admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2005
IRF Medicare days (per 1,000 beneficiaries)	Average number of inpatient rehabilitation facility (IRF) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2013

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Iron deficiency in children aged 1-2 years (percent)	Percent of young children aged 1 to 2 years who are deficient in iron	Yes	No	No	No	1209
Iron deficiency in children aged 3-4 years (percent)	Percent of young children aged 3 to 4 years who are deficient in iron	Yes	No	No	No	1210
Iron deficiency in females of childbearing age (percent)	Percent of females of childbearing age, aged 12 to 49 years, who are deficient in iron	Yes	No	No	No	1211
Iron deficiency in pregnant females (percent)	Iron deficiency among pregnant females	Yes	No	No	No	1212
Ischemic heart disease Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have ischemic heart disease	Yes	Yes	No	Yes	297
Ischemic heart disease Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have ischemic heart disease	Yes	Yes	No	Yes	298
Joint mobility among persons with hemophilia (percent)	Percent of persons with hemophilia who develop reduced joint mobility due to bleeding into joints	Yes	No	No	No	444
Joint pain among adults with arthritis (mean)	Mean level of joint pain among adults with doctor-diagnosed arthritis	Yes	No	No	No	418
Joint-related activities-Difficult to stoop, bend, kneel, adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-related activities-Stoop, bend, or kneel	Yes	No	No	No	433
Joint-related activities-Difficult to walk up 10 steps, adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-related activities-Walk up 10 steps without resting	Yes	No	No	No	432
Joint-related activities-Difficulty to grasp, adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-related activities-Use fingers to grasp or handle small objects	Yes	No	No	No	434
Joint-related activities-Difficulty to walk 3 blocks, adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-related activities-Walk a quarter of a mile-about 3 city blocks	Yes	No	No	No	431

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Joint-related symptoms--saw health care provider (percent)	Percent of adults with chronic joint symptoms who have seen a Health Care provider for their symptoms	Yes	No	No	No	425
Juvenile justice facilities screening for mental health problems (percent)	Percent of juvenile residential facilities that screen admissions for mental health problems	Yes	No	No	No	1111
Kidney transplantation within 3 years of terminal kidney failure (percent)	Percent of patients receiving a kidney transplant within 3 years of end-stage renal disease (ESRD)	Yes	No	No	No	473
Kindergarten vaccination record keeping, state participation (number)	Number of States collecting kindergarten vaccination coverage data according to CDC minimum standards	Yes	No	No	No	1026
Laboratory proficiency - chemical (percent)	Percent of LRN chemical laboratories that meet proficiency standards for chemical threat agents	Yes	No	No	No	1396
Laboratory test Medicare service events (per 1,000 beneficiaries)	Rate of laboratory test service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2024
Laboratory test Medicare users (number)	Count of Medicare fee-for-service beneficiaries 65+ years of age who were laboratory test users	Yes	Yes	No	Yes	2027
Laboratory test Medicare utilization (percent)	Percent of Medicare fee-for-service beneficiaries 65+ years of age who were laboratory test users	Yes	Yes	No	Yes	2028
Lactation care in birthing facilities (percent)	Percent of live births that occur in facilities that provide recommended care for lactating mothers and their babies	Yes	No	No	No	1186
Late preterm births (percent)	Percent of live births that are late preterm (34 to 36 weeks of gestation)	Yes	Yes	Yes	No	1138
Late-stage female breast cancer (per 100,000)	Late-stage female breast cancer incidence	Yes	No	No	No	511
Leisure-time physical activities among older adults (percent)	Percent of older adults with reduced physical or cognitive function who engage in light, moderate, or vigorous leisure-time physical activities	Yes	No	No	No	1258
Leisure-time physical activity-none (percent)	Percent of adults who engage in no leisure-time physical activity	Yes	No	No	No	1313

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Limited ability to work among adults with arthritis (percent)	Impact of doctor-diagnosed arthritis on employment in the working-aged population-Percent of adults with doctor-diagnosed arthritis who are limited in their ability to work for pay due to arthritis	Yes	No	No	No	424
Lipids under control--persons with diabetes (percent)	Lipid control among persons with diagnosed diabetes	Yes	No	No	No	561
Liquor stores (per 10,000 population)	Liquor store density rate	No	Yes	Yes	No	51
Live births (number)	Number of live births	Yes	Yes	Yes	No	10009
Local health departments with oral health components (percent)	Percent of local health departments that have oral health prevention or care programs	Yes	No	No	No	1288
Long term care hospital Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who had covered long-term care hospital services	Yes	Yes	No	Yes	200
Long term care hospital Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered long-term care hospital services	Yes	Yes	No	Yes	201
Long-Term Care Hospitals (number)	Number of long-term care hospitals participating in the Medicare program	No	Yes	No	Yes	10043
Low birth weight (percent)	Percent of live births that are low birth weight (LBW)	Yes	Yes	Yes	No	1135
Lower extremity amputations (per 1,000)	Rate of lower extremity amputations in persons with diagnosed diabetes	Yes	No	No	No	548
LTC hospital Medicare admissions (per 1,000 beneficiaries)	Rate of long term care (LTC) hospital admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2006
LTC hospital Medicare days (per 1,000 beneficiaries)	Average number of long term care (LTC) hospital days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2014
Lung cancer deaths (per 100,000)	Lung cancer death rate per 100,000	Yes	No	No	No	487
Lung cancer Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have lung cancer	Yes	Yes	No	Yes	303

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Lung cancer Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have lung cancer	Yes	Yes	No	Yes	304
Lung, trachea, and bronchus cancer deaths (per 100,000)	Lung, trachea, and bronchus cancer death rate per 100,000	Yes	Yes	Yes	No	88
Lyme disease (number)	Number of reported cases of Lyme disease	No	Yes	Yes	No	10024
M.D. training in counseling for health promotion and disease prevention (percent)	Percent of inclusion of core clinical prevention and population health content-Counseling for health promotion and disease prevention in health professions training-MD-granting medical schools	Yes	No	No	No	606
M.D. training in cultural diversity (percent)	Percent of inclusion of core clinical prevention and population health content-Cultural diversity in health professions training-MD-granting medical schools	Yes	No	No	No	607
M.D. training in environmental health (percent)	Percent of inclusion of core clinical prevention and population health content-Environmental health in health professions training-MD-granting medical schools	Yes	No	No	No	609
M.D. training in evaluation of health sciences literature. (percent)	Percent of inclusion of core clinical prevention and population health content-Evaluation of health sciences literature in health professions training-MD-granting medical schools	Yes	No	No	No	608
M.D. training in global health (percent)	Percent of inclusion of core clinical prevention and population health content-Global health in health professions training-MD-granting medical schools	Yes	No	No	No	611
M.D. training in public health systems (percent)	Percent of inclusion of core clinical prevention and population health content-Public health systems in health professions training-MD-granting medical schools	Yes	No	No	No	610
Major depressive episode (MDE) - adolescents (percent)	Percent of adolescents aged 12 to 17 years with Major Depressive Episode	Yes	No	No	No	1123
Major depressive episode (MDE) - adults (percent)	Percent of adults aged 18 years and older with Major Depressive Episode	Yes	No	No	No	1122

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Malaria in the U.S. (number)	Number of cases of malaria reported in the United States	Yes	No	No	No	844
Male aged 15 years and under who have never had sexual intercourse (percent)	Percent of male adolescents aged 15 years and under who have never had sexual intercourse	Yes	No	No	No	793
Male Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months in who are male	Yes	Yes	No	Yes	310
Male teens using a condom & hormonal or intrauterine contraception: first intercourse (percent)	Percent of sexually active males aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at first intercourse	Yes	No	No	No	799
Male teens using a condom & hormonal or intrauterine contraception: last intercourse (percent)	Percent of sexually active males aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at last intercourse	Yes	No	No	No	801
Males aged 15 to 17 years who have never had sexual intercourse (percent)	Percent of male adolescents aged 15 to 17 years who have never had sexual intercourse	Yes	No	No	No	791
Mammogram (percent)	Percent of women age 50+ that report a mammogram in the past 2 years	No	Yes	Yes	No	116
Mandatory ignition interlock laws (number)	Number of States with mandatory ignition interlock laws for first and repeat impaired driving offenders in the United States	Yes	No	No	No	1468
Maternal deaths (per 100,000 live births)	Rate of maternal death	Yes	No	No	No	1129
Maternal illness and complications due to pregnancy (percent)	Maternal illness and complications due to pregnancy (complications during hospitalized labor and delivery)	Yes	No	No	No	1130
Measles, total (number)	Number of reported cases of measles	No	Yes	Yes	No	10032
Measles, U.S.-acquired (number)	Number of measles cases (U.S.-acquired cases)	Yes	No	No	No	981
Median Household Income (dollars)	Small area estimates of median household Income	Yes	Yes	Yes	No	10003
Medicaid beneficiaries (number in thousands)	Number of Medicaid beneficiaries	Yes	Yes	No	No	10014

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Medicaid coverage for evidence based treatment of nicotine dependency-States (percent)	Comprehensive Medicaid insurance coverage of evidence-based treatment for nicotine dependency in States and the District of Columbia	Yes	No	No	No	1516
Medicaid managed care beneficiaries (percent)	Percent of Medicaid beneficiaries in managed care plans	Yes	Yes	No	No	10015
Medicaid payments per beneficiary (dollars)	Average payment per Medicaid beneficiary	Yes	Yes	No	No	10017
Medical evaluation of persons with chronic kidney disease (percent)	Percent of persons 65 years of age and over with chronic kidney disease who receive medical evaluation with serum creatinine, lipids, and microalbuminuria	Yes	No	No	No	477
Medical evaluation of persons with type 1 and type 2 diabetes and chronic kidney disease (percent)	Percent of persons 65 years of age and over with type 1 or type 2 diabetes and chronic kidney disease who receive medical evaluation with serum creatinine, microalbuminuria, HbA1c, lipids, and eye examinations	Yes	No	No	No	478
Medical treatment of persons with diabetes and chronic kidney disease (percent)	Percent of persons 65 years of age and over with Part D insurance with diabetes and chronic kidney disease who receive recommended medical treatment with angiotensin-converting enzyme (ACE) inhibitors or angiotensin II receptor blockers (ARBS)	Yes	No	No	No	479
Medicare beneficiaries 65+ who are also eligible for Medicaid (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months in who are also eligible for Medicaid	Yes	Yes	No	Yes	317
Medicare beneficiaries 65+ (number)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	307
Medicare managed care beneficiaries (percent)	Percent of Medicare beneficiaries enrolled in Medicare Advantage managed care plans	Yes	Yes	No	No	10020
Medication overdoses among children less than 5 years of age-ED visits (per 10,000)	Rate of emergency department (ED) visits for medication overdoses among children less than 5 years of age	Yes	No	No	No	1206
Melanoma cancer deaths (per 100,000)	Melanoma cancer death rate	Yes	No	No	No	493

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Meningococcal disease (per 100,000)	Meningococcal disease rate	Yes	No	No	No	968
Meningococcal vaccination among adolescents (percent)	Percent of adolescents who have 1 dose Meningococcal vaccine (MCV) by age 13 to 15 years	Yes	No	No	No	1012
Mental health treatment provided in primary care facilities (percent)	Percent of primary care facilities that provide mental health treatment onsite or by paid referral	Yes	No	No	No	1109
Mentally unhealthy days: adults (per person)	Average number of reported mentally unhealthy days per month among adults 18 years and over	No	Yes	Yes	No	9
Missed school days due to asthma (percent)	Percent of children aged 5 to 17 years with asthma who miss school days	Yes	No	No	No	1410
Missed work days due to asthma (percent)	Percent of adults aged 18 to 64 years with asthma who miss work days	Yes	No	No	No	1411
MMR vaccination (percent)	Percent of young children who have received universally recommended vaccines-1 dose measles-mumps-rubella (MMR) vaccine by age 19 to 35 months	Yes	No	No	No	996
MMR vaccination among kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-MMR vaccines (2 or more doses)	Yes	No	No	No	1004
Monitoring acute chemical poisoning (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: acute chemical poisoning	Yes	No	No	No	718
Monitoring arsenic poisoning (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: arsenic poisoning	Yes	No	No	No	716
Monitoring cadmium poisoning (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: cadmium poisoning	Yes	No	No	No	717

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Monitoring carbon monoxide poisoning (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: carbon monoxide poisoning	Yes	No	No	No	719
Monitoring lead poisoning (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: Lead poisoning	Yes	No	No	No	713
Monitoring mercury poisoning. (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: mercury poisoning	Yes	No	No	No	715
Monitoring pesticide poisoning (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that can be caused by exposure to environmental hazards: pesticide poisoning	Yes	No	No	No	714
Motor vehicle crash deaths (per 100,000,000 vehicle miles traveled)	Motor vehicle crash-related deaths death rate per 100 million vehicle miles traveled	Yes	No	No	No	1077
Motor vehicle crash deaths, unintentional (per 100,000)	Unintentional motor vehicle crash-related death rate per 100,000	Yes	Yes	Yes	No	1076
Motor vehicle deaths (per 100,000)	Unintentional motor vehicle related death rate	Yes	Yes	Yes	No	18
Motorcycle helmet use (percent)	Percent of motorcycle operators and passengers using helmets	Yes	No	No	No	1048
Mumps (number)	Number of reported cases of mumps	No	Yes	Yes	No	10031
Mumps, U.S.-acquired (number)	Number of mumps cases (U.S.-acquired cases)	Yes	No	No	No	982
Muscle-strengthening activity among adults (percent)	Percent of adults who perform muscle-strengthening activities on 2 or more days of the week	Yes	No	No	No	1321
Narrow-therapeutic-index medications-ED visits for overdoses (per 10,000)	Rate of emergency department (ED) visits for overdoses of narrow-therapeutic-index medications	Yes	No	No	No	1205
Neonatal deaths (per 1,000 live births)	Rate of neonatal deaths (within the first 28 days of life)	Yes	Yes	Yes	No	1159
Neonatal hearing screening (percent)	Percent of infants that receive screening for hearing loss no later than age 1 month	Yes	No	No	No	756

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
New cases of diabetes - 3-year average (per 1,000)	Rate of newly diagnosed diabetes among adults aged 18-84	Yes	No	No	No	555
New Hepatitis B cases in children (number)	Rate of new hepatitis B cases among persons aged 2 to 18 years	Yes	No	No	No	980
No exercise (percent)	Adults that report no leisure-time exercise in past month	No	Yes	Yes	No	120
No permanent tooth loss in older adults (percent)	Percent of older adults aged 65 to 74 years who have lost all of their natural teeth	Yes	No	No	No	1279
Non-fatal work-related injuries - Adolescents (per 100)	Rate of nonfatal work-related injuries among adolescent workers aged 15 to 19 years	Yes	No	No	No	1310
Non-fatal work-related injuries - with emergency departments (per 100)	Rate of injuries treated in emergency departments from work, or restricted work activity, as reported by employers	Yes	No	No	No	1309
Non-fatal work-related injuries - with medical treatment (per 100)	Rate of injuries in private sector industries resulting in medical treatment, lost time from work, or restricted work activity, as reported by employers	Yes	No	No	No	1308
Non-FDA-approved pain medications (number)	Number of non-FDA-approved pain medications	Yes	No	No	No	1198
Non-Hispanic White Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months in who are Non-Hispanic white	Yes	Yes	No	Yes	311
Nonfatal child maltreatment (per 1,000)	Nonfatal child maltreatment	Yes	No	No	No	1084
Nonfatal firearm-related injuries (per 100,000)	Nonfatal firearm-related injury rate	Yes	No	No	No	1043
Nonfatal intentional self-harm injuries (per 100,000)	Nonfatal intentional self-harm injury rate	Yes	No	No	No	1100
Nonfatal motor vehicle injuries (per 100,000)	Nonfatal motor vehicle crash-related injuries	Yes	No	No	No	1078
Nonfatal pedestrian injuries (per 100,000)	Rate of nonfatal pedestrian injuries on public roads	Yes	No	No	No	1046

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Nonfatal physical assault injuries (per 100,000)	Nonfatal physical assault injury rate	Yes	No	No	No	1099
Nonfatal poisonings (per 100,000)	Rate of nonfatal poisonings	Yes	No	No	No	1040
Nonfatal unintentional injuries (per 100,000)	Nonfatal unintentional injury rate	Yes	No	No	No	1075
Nurse practitioner training in counseling for health promotion and disease prevention (percent)	Percent of inclusion of core clinical prevention and population health content-Counseling for health promotion and disease prevention in health professions training-Nurse practitioner	Yes	No	No	No	624
Nurse practitioner training in cultural diversity (percent)	Percent of inclusion of core clinical prevention and population health content-Cultural diversity in health professions training-Nurse practitioner	Yes	No	No	No	625
Nurse practitioner training in environmental health (percent)	Percent of inclusion of core clinical prevention and population health content-Environmental health in health professions training-Nurse practitioner	Yes	No	No	No	627
Nurse practitioner training in evaluation of health sciences literature (percent)	Percent of inclusion of core clinical prevention and population health content-Evaluation of health sciences literature in health professions training-Nurse practitioner	Yes	No	No	No	626
Nurse practitioner training in global health (percent)	Percent of inclusion of core clinical prevention and population health content-Global health in health professions training-Nurse practitioner	Yes	No	No	No	629
Nurse practitioner training in public health systems (percent)	Percent of inclusion of core clinical prevention and population health content-Public health systems in health professions training-Nurse practitioner	Yes	No	No	No	628
Nursing home (Title 18/19) facilities (number)	Number of Medicare or Medicaid-certified Title 18/19 (nursing home) facilities	No	Yes	No	Yes	10053
Obesity in adolescents aged 12-19 years (percent)	Percent of children and adolescents aged 12 to 19 years who are considered obese	Yes	No	No	No	1215
Obesity in adults (percent)	Percent of adults who are obese	Yes	No	No	No	1208

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Obesity in children aged 2-5 years (percent)	Percent of children and adolescents aged 2 to 5 years who are considered obese	Yes	No	No	No	1213
Obesity in children aged 6-11 years (percent)	Percent of children and adolescents aged 6 to 11 years who are considered obese	Yes	No	No	No	1214
Obesity in children and adolescents aged 2-19 years (percent)	Percent of children and adolescents aged 2 to 19 years who are considered obese	Yes	No	No	No	1216
Obesity: adult (percent)	Percent of adults 18 years and over that report BMI \geq 30	No	Yes	Yes	No	15
Occupational eye injuries resulting in lost work days (per 10,000)	Rate of occupational eye injuries resulting in lost work days	Yes	No	No	No	1564
Occupational eye injuries treated in emergency departments (per 10,000)	Rate of occupational eye injuries treated in emergency departments	Yes	No	No	No	1565
Occupational skin diseases or disorders (per 10,000)	Rate of occupational skin diseases or disorders among full-time workers	Yes	No	No	No	1301
Older adults up to date on clinical preventive measures - Females (percent)	Percent of older adults who are up to date on a core set of clinical preventive measures-Females	Yes	No	No	No	1262
Older adults up to date on clinical preventive measures - Males (percent)	Percent of older adults who are up to date on a core set of clinical preventive measures-Males	Yes	No	No	No	1261
Oral and craniofacial State-based health surveillance system (number)	Number of States and the District of Columbia that have an oral and craniofacial health surveillance system	Yes	No	No	No	1268
Oral and pharyngeal cancer early detection (percent)	Percent of oral and pharyngeal cancers detected at the earliest stage	Yes	No	No	No	1264
Oral anticoagulants-ED visits for overdoses (per 10,000)	Rate of emergency department (ED) visits for overdoses from oral anticoagulants	Yes	No	No	No	1203
Oral health care system use (percent)	Percent of children, adolescents, and adults who used the oral health care system in the past year	Yes	No	No	No	1266
Organ Procurement Organizations (number)	Number of Organ Procurement Organizations participating in the Medicare program	No	Yes	No	Yes	10044

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Oropharyngeal cancer deaths (per 100,000)	Oropharyngeal cancer death rate	Yes	No	No	No	491
Osteoporosis among adults aged 50 years and older (percent)	Percent of adults aged 50 years and older with osteoporosis	Yes	No	No	No	427
Other hospital inpatient Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who had covered other hospital inpatient services	Yes	Yes	No	Yes	181
Other hospital inpatient Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered other hospital inpatient services	Yes	Yes	No	Yes	182
Other hospital IP Medicare admissions (per 1,000 beneficiaries)	Rate of other hospital inpatient (IP) admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2004
Other hospital IP Medicare days (per 1,000 beneficiaries)	Average number of other hospital inpatient (IP) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2012
Other or Unknown race Medicare beneficiaries 65+ (percent)	Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months in who are Other or Unknown race	Yes	Yes	No	Yes	316
Other Part B Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who used other Part B	Yes	Yes	No	Yes	275
Other Part B Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used other Part B	Yes	Yes	No	Yes	276
Other tests (Non-laboratory) Medicare service events (per 1,000 beneficiaries)	Rate of non-laboratory test service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2025
Other tests (non-laboratory) Medicare users (number)	Count of Medicare fee-for-service beneficiaries 65+ years of age who were non-laboratory test users	Yes	Yes	No	Yes	2029
Other tests (non-laboratory) Medicare utilization (percent)	Percent of Medicare fee-for-service beneficiaries 65+ years of age who were non-laboratory test users	Yes	Yes	No	Yes	2030
Otitis media visits in ambulatory care settings (per 1,000)	Rate of otitis media in children and adolescents	Yes	No	No	No	743

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Outbreak-associated infections associated with beef (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacter, Listeria, or Salmonella species associated with food commodity group: beef	Yes	No	No	No	839
Outbreak-associated infections associated with chicken (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacter, Listeria, or Salmonella species associated with food commodity group: poultry	Yes	No	No	No	843
Outbreak-associated infections associated with dairy (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacter, Listeria, or Salmonella species associated with food commodity group: dairy	Yes	No	No	No	840
Outbreak-associated infections associated with fruits and nuts (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacter, Listeria, or Salmonella species associated with food commodity group: fruits and nuts	Yes	No	No	No	841
Outbreak-associated infections associated with leafy vegetables (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacter, Listeria, or Salmonella species associated with food commodity group: leafy vegetables	Yes	No	No	No	842
Outpatient Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who used outpatient services	Yes	Yes	No	Yes	187
Outpatient Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used outpatient services	Yes	Yes	No	Yes	188
Outpatient physical therapy/speech pathology facilities (number)	Number of Outpatient physical therapy/speech pathology facilities participating in the Medicare program	No	Yes	No	Yes	10045
Overall cancer deaths (per 100,000)	Overall cancer death rate	Yes	Yes	Yes	No	486
Overexertion or repetitive motion (per 10,000)	Rate of injury and illness cases involving days away from work due to overexertion or repetitive motion	Yes	No	No	No	1297

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Ozone days (number)	Annual number of days with daily 8-hour maximum ozone concentration over the National Ambient Air Quality Standard	No	Yes	Yes	No	45
Pap test (percent)	Percent of women age 18+ that report Pap test in the last 3 years	No	Yes	Yes	No	115
Parents that read to child aged 0-5 years (percent)	Percent of parents who read to their young child	Yes	No	No	No	740
Part B drug Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who used Part B drugs	Yes	Yes	No	Yes	269
Part B drugs Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used Part B drugs	Yes	Yes	No	Yes	270
Particulate matter days (number)	Annual number of days with maximum 24-hour average PM 2.5 concentration over the National Ambient Air Quality Standard	No	Yes	Yes	No	44
PCV vaccination (percent)	Percent of young children who have received universally recommended vaccines-4 doses pneumococcal conjugate vaccine (PCV) by age 19 to 35 months	Yes	No	No	No	999
Pedal-cyclist deaths (per 100,000)	Death rate of pedal-cyclists on public roads	Yes	No	No	No	1103
Pedestrian deaths (per 100,000)	Rate of pedestrian deaths on public roads	Yes	No	No	No	1045
Pelvic inflammatory disease (percent)	Percent of females aged 15 to 44 years who have ever required treatment for pelvic inflammatory disease (PID)	Yes	No	No	No	1473
Penicillin-resistant pneumococcal infections among children (per 100,000)	Rate of invasive penicillin-resistant pneumococcal infections among children under age 5 years	Yes	No	No	No	965
Penicillin-resistant pneumococcal infections among elderly (per 100,000)	Rate of invasive penicillin-resistant pneumococcal infections among adults aged 65 years and older	Yes	No	No	No	966
Perinatal deaths (per 1,000 live births plus fetal deaths)	Rate of fetal and infant deaths during perinatal period (28 weeks of gestation to 7 days after birth)	Yes	No	No	No	1157

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Personal care limitations among adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who have difficulty in performing two or more personal care activities, thereby preserving independence	Yes	No	No	No	420
Persons receiving safe drinking water (percent)	Percent of persons served by community water systems who receive a supply of drinking water that meets the regulations of the Safe Drinking Water Act	Yes	No	No	No	659
Pertussis (number)	Number of reported cases of pertussis	No	Yes	Yes	No	10029
Pertussis among adolescents (number)	Number of pertussis cases among adolescents aged 11 to 18 years	Yes	No	No	No	984
Pertussis among infants (number)	Number of pertussis cases among children under age 1 year	Yes	No	No	No	983
Pesticide exposures resulting in visits to health care facility (number)	Number of pesticide exposures that result in visits to a health care facility	Yes	No	No	No	661
Physical activity policies in child care settings - number of minutes (number)	Number of States with licensing regulations for physical activity provided in child care-Require number of minutes of physical activity per day or by length of time in care	Yes	No	No	No	1345
Physical activity policies in child care settings - physical activity programs (number)	Number of States with licensing regulations for physical activity provided in child care-Require activity programs providing large muscle or gross motor activity, development, and/or equipment	Yes	No	No	No	1343
Physical activity policies in child care settings - vigorous or moderate physical activity (number)	Number of States with licensing regulations for physical activity provided in child care-Require children to engage in vigorous or moderate physical activity	Yes	No	No	No	1344
Physical assaults (per 1,000)	Rate of physical assaults among persons 12 years of age and over	Yes	No	No	No	1051
Physical education required daily in elementary schools (percent)	Percent of the Nation's public and private schools that require daily physical education for all students-Elementary schools	Yes	No	No	No	1314

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Physical education required daily in middle and junior high schools (percent)	Percent of the Nation's public and private schools that require daily physical education for all students-Middle and junior high schools	Yes	No	No	No	1315
Physical education required daily in senior high schools (percent)	Percent of the Nation's public and private schools that require daily physical education for all students-Senior high schools	Yes	No	No	No	1316
Physical fighting among adolescents (percent)	Percent of physical fighting among adolescents	Yes	Yes	No	No	1052
Physically or mentally unhealthy days (days)	Average number of reported physically or mentally unhealthy days per month	No	Yes	Yes	No	75
Physically unhealthy days: adults (per person)	Average number of reported physically unhealthy days per month among adults 18 years of age and over	No	Yes	Yes	No	7
Physician assistant training in counseling for health promotion and disease prevention (percent)	Percent of inclusion of core clinical prevention and population health content-Counseling for health promotion and disease prevention in health professions training-Physician assistant	Yes	No	No	No	630
Physician assistant training in cultural diversity (percent)	Percent of inclusion of core clinical prevention and population health content-Cultural diversity in health professions training-Physician assistant	Yes	No	No	No	631
Physician assistant training in environmental health (percent)	Percent of inclusion of core clinical prevention and population health content-Environmental health in health professions training-Physician assistant	Yes	No	No	No	633
Physician assistant training in evaluation of health sciences literature (percent)	Percent of inclusion of core clinical prevention and population health content-Evaluation of health sciences literature in health professions training-Physician assistant	Yes	No	No	No	632
Physician assistant training in global health (percent)	Percent of inclusion of core clinical prevention and population health content-Global health in health professions training-Physician assistant	Yes	No	No	No	635

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Physician assistant training in public health systems (percent)	Percent of inclusion of core clinical prevention and population health content-Public health systems in health professions training-Physician assistant	Yes	No	No	No	634
Physician counseling about physical activity (percent)	Percent of physician visits made by all child and adult patients that include counseling about exercise	Yes	No	No	No	1342
Physician counseling about physical activity - patients with chronic conditions (percent)	Percent of office visits made by patients with a diagnosis of cardiovascular disease, diabetes, or hyperlipidemia that include counseling or education related to exercise	Yes	No	No	No	1341
Physician E&M Medicare service events (per 1,000 beneficiaries)	Rate of physician evaluation and management service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2021
Physician E&M Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who used physician evaluation and management services	Yes	Yes	No	Yes	239
Physician E&M Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used physician evaluation and management services	Yes	Yes	No	Yes	240
Physician procedures Medicare service events (per 1,000 beneficiaries)	Rate of physician procedure service events among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2022
Physician procedures Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who used physician procedures services	Yes	Yes	No	Yes	245
Physician procedures Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used physician procedures services	Yes	Yes	No	Yes	246
Physician use delayed due to cost (percent)	Percent of adults 18+ who needed to see a doctor but could not because of cost	No	Yes	Yes	No	10019
Pneumococcal vaccination among elderly (percent)	Percent of adults who are vaccinated against pneumococcal disease-Noninstitutionalized adults aged 65 years and older	Yes	No	No	No	1018
Pneumococcal vaccination among high-risk adults (percent)	Percent of adults who are vaccinated against pneumococcal disease-Noninstitutionalized high-risk adults aged 18 to 64 years	Yes	No	No	No	1019

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Pneumococcal vaccination among nursing home residents (percent)	Percent of adults who are vaccinated against pneumococcal disease-Institutionalized adults (persons aged 18 years and older in long-term or nursing homes)	Yes	No	No	No	1020
Pneumococcal vaccination, adults 65+ (percent)	Percent of adults age 65+ that report ever having a pneumococcal vaccination	No	Yes	Yes	No	118
Pneumoconiosis deaths (number)	Annual number of pneumoconiosis deaths	Yes	No	No	No	1298
Pneumonia patient with blood cultures in emergency department before antibiotic administered (percent)	Hospital Compare: Pneumonia patient with blood cultures in emergency department before antibiotic administered	Yes	Yes	No	Yes	332
Pneumonia patients with appropriate initial antibiotic selection for community-acquired pneumonia in immunocompetent patients (percent)	Hospital Compare: Pneumonia patients with appropriate initial antibiotic selection for community-acquired pneumonia in immunocompetent patients	Yes	Yes	No	Yes	331
Pneumonia patients with influenza vaccination (percent)	Hospital Compare: Pneumonia patients with influenza vaccination	Yes	Yes	No	Yes	333
Pneumonia patients with initial antibiotic received within 6 hours of hospital arrival (percent)	Hospital Compare: Pneumonia patients with initial antibiotic received within 6 hours of hospital arrival	Yes	Yes	No	Yes	335
Pneumonia patients with pneumococcal vaccination (percent)	Hospital Compare: Pneumonia patients with pneumococcal vaccination	Yes	Yes	No	Yes	330
Pneumonia patients with smoking cessation counseling (percent)	Hospital Compare: Pneumonia patients with smoking cessation counseling	Yes	Yes	No	Yes	334
Poisoning deaths (per 100,000)	Poisoning death rate: All persons	Yes	Yes	Yes	No	1064
Poisoning deaths, 35-54 years (per 100,000)	Poisoning death rate: Persons aged 35 to 54 years	Yes	No	No	No	1065
Polio vaccination (percent)	Percent of young children who have received universally recommended vaccines-3 doses polio vaccine by age 19 to 35 months	Yes	No	No	No	997

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Polio vaccination among kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-Polio vaccine (3 or more doses)	Yes	No	No	No	1005
Polio, U.S.-acquired (number)	Number of acute paralytic poliomyelitis cases (U.S.-acquired cases)	Yes	No	No	No	985
Population density (per square mile)	Population density of United States geographic area per square mile	Yes	Yes	Yes	No	10001
Population distributions (percent)	Bridged-race population distributions according to 1977 OMB standard	Yes	Yes	Yes	No	10002
Population estimates (number)	Bridged-race population estimates according to 1977 OMB standard	Yes	Yes	Yes	No	10000
Portable X-Ray providers (number)	Number of portable X-Ray providers participating in the Medicare Program	No	Yes	No	Yes	10046
Positive communication with child aged 6-17 years (percent)	Percent of parents who use positive communication with their child	Yes	No	No	No	739
Post acute care Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who used post-acute care services	Yes	Yes	No	Yes	194
Post-acute care Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who used post-acute care services	Yes	Yes	No	Yes	193
Postneonatal deaths (per 1,000 live births)	Rate of postneonatal deaths (between 28 days and 1 year)	Yes	Yes	Yes	No	1160
Postoperative pulmonary embolism or deep vein thrombosis rate (per 100,000 beneficiaries)	Patient Safety Indicators: Postoperative pulmonary embolism or deep vein thrombosis rate (observed, among all surgical discharges based on secondary diagnosis field) among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2035

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Postoperative sepsis rate (per 100,000 beneficiaries)	Patient Safety Indicators: Postoperative sepsis rate (observed, among elective surgical discharges based on secondary diagnosis field) among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2036
PPS Medicare admissions (per 1,000 beneficiaries)	Rate of acute inpatient prospective payment system (PPS) admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2002
PPS Medicare days (per 1,000 beneficiaries)	Average number of acute inpatient prospective payment system (PPS) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2010
Pre-ESRD care from a nephrologist (percent)	Percent of chronic kidney disease patients receiving care from a nephrologist at least 12 months before the start of renal replacement therapy	Yes	No	No	No	468
Preconception abstinence from alcohol (percent)	Percent of women delivering a live birth who did not drink alcohol prior to pregnancy	Yes	No	No	No	1181
Preconception abstinence from smoking (percent)	Percent of women delivering a live birth who did not smoke prior to pregnancy	Yes	No	No	No	1180
Preconception intake of multivitamins/folic acid (percent)	Percent of women delivering a live birth who took multivitamins/folic acid prior to pregnancy	Yes	No	No	No	1179
Preemptive State tobacco control laws --advertising (number)	Number of States with State laws that preempt stronger local tobacco control laws-Preemption in advertising	Yes	No	No	No	1546
Preemptive State tobacco control laws --smoke-free indoor air (number)	Number of States with State laws that preempt stronger local tobacco control laws-Preemption on smoke-free indoor air	Yes	No	No	No	1545
Preemptive State tobacco control laws--youth access (number)	Number of States with State laws that preempt stronger local tobacco control laws-Preemption on youth access	Yes	No	No	No	1547
Preemptive transplantation in terminal kidney failure (percent)	Percent of patients who receive a preemptive transplant at the start of ESRD	Yes	No	No	No	474

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Pregnancies among females aged 15 to 17 years (per 1,000)	Pregnancy rate among adolescent females aged 15 to 17 years	Yes	No	No	No	788
Pregnancies among females aged 18 to 19 years (per 1,000)	Pregnancy rate among adolescent females aged 18 to 19 years	Yes	No	No	No	789
Pregnancies conceived within 18 months of a previous birth (percent)	Percent of pregnancies conceived within 18 months of a previous birth	Yes	No	No	No	784
Pregnancy despite use of reversible contraceptive method (percent)	Percent of females experiencing pregnancy despite use of a reversible contraceptive method	Yes	No	No	No	780
Prenatal abstinence from alcohol (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Alcohol	Yes	No	No	No	1142
Prenatal abstinence from binge drinking (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Binge drinking	Yes	No	No	No	1143
Prenatal abstinence from cigarettes (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Cigarette smoking	Yes	No	No	No	1144
Prenatal abstinence from illicit drugs (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Illicit drugs	Yes	No	No	No	1145
Prenatal care, early and adequate (percent)	Percent of pregnant women who receive early and adequate prenatal care	Yes	No	No	No	1132
Prenatal care, first trimester (percent)	Percent of pregnant women who receive prenatal care beginning in the first trimester of pregnancy	Yes	No	No	No	1131
Pressure ulcer-related hospitalizations among older adults (per 100,000)	Rate of pressure ulcer-related hospitalizations among older adults	Yes	No	No	No	1249
Pressure/decubitus ulcer rate (per 100,000 beneficiaries)	Patient Safety Indicators: Pressure/decubitus ulcer rate (observed, among medical and surgical discharges based on secondary diagnosis field) among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2032
Preterm births, total (percent)	Percent of live births that are preterm (< 37 weeks of gestation)	Yes	Yes	Yes	No	1137

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Primary and secondary syphilis among females (per 100,000)	Sustained domestic transmission of primary and secondary syphilis among females	Yes	No	No	No	1480
Primary and secondary syphilis among males (per 100,000)	Sustained domestic transmission of primary and secondary syphilis among males	Yes	No	No	No	1481
Primary care physicians who assess adult patients' body mass index (percent)	Percent of primary care physicians who regularly assess body mass index (BMI) in their adult patients	Yes	No	No	No	1234
Primary care physicians who assess child and adolescent patients' body mass index (percent)	Percent of primary care physicians who regularly assess body mass index (BMI) for age and gender in their child or adolescent patients	Yes	No	No	No	1235
Primary care provider rate (per 100,000)	Rate of primary care providers (PCP)	No	Yes	Yes	No	25
Professional development on behavior change-Health education teachers (percent)	Percent of required health education classes or courses with a teacher who has had professional development related to teaching personal and social skills for behavior change within the past 2 years	Yes	No	No	No	737
Proficiency in math-12th grade (percent)	Percent of 12th graders whose mathematics skills are at or above the proficient achievement level for 12th grade	Yes	No	No	No	401
Proficiency in math-4th grade (percent)	Percent of 4th graders whose mathematics skills are at or above the proficient achievement level for 4th grade	Yes	No	No	No	399
Proficiency in math-8th grade (percent)	Percent of 8th graders whose mathematics skills are at or above the proficient achievement level for 8th grade	Yes	No	No	No	400
Proficiency in reading-12th grade (percent)	Percent of 12th graders whose reading skills are at or above the proficient achievement level for 12th grade	Yes	No	No	No	398
Proficiency in reading-4th grade (percent)	Percent of 4th graders whose reading skills are at or above the proficient achievement level for 4th grade	Yes	No	No	No	396
Proficiency in reading-8th grade (percent)	Percent of 8th graders whose reading skills are at or above the proficient achievement level for 8th grade	Yes	No	No	No	397

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Prostate cancer Medicare beneficiaries (number)	Medicare beneficiaries age 65+ years with full FFS Medicare who have prostate cancer	Yes	Yes	No	Yes	305
Prostate cancer Medicare beneficiaries (percent)	Medicare beneficiaries age 65+ years with full FFS Medicare who have prostate cancer	Yes	Yes	No	Yes	306
Protective gear in intramural activities and clubs (percent)	Percent of public and private schools that require students to wear appropriate protective gear when engaged in school-sponsored Intramural activities or physical activity clubs	Yes	No	No	No	1083
Protective gear in physical education (percent)	Percent of public and private schools that require students to wear appropriate protective gear when engaged in school-sponsored physical education	Yes	No	No	No	1082
Psychiatric hospitals (number)	Number of Psychiatric hospitals participating in the Medicare program	No	Yes	No	Yes	10047
Public health agencies laboratory services: Disease prevention, control and surveillance (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Disease prevention, control, and surveillance	Yes	No	No	No	1349
Public health agencies laboratory services: Emergency response (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Emergency response	Yes	No	No	No	1356
Public health agencies laboratory services: Environmental health and protection (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Environmental health and protection	Yes	No	No	No	1352
Public health agencies laboratory services: Food safety (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Food safety	Yes	No	No	No	1353

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Public health agencies laboratory services: Integrated data management (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Integrated data management	Yes	No	No	No	1350
Public health agencies laboratory services: Laboratory improvement and regulation (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Laboratory improvement and regulation	Yes	No	No	No	1354
Public health agencies laboratory services: Partnerships and communication (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Partnerships and communication	Yes	No	No	No	1359
Public health agencies laboratory services: Policy development (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Policy development	Yes	No	No	No	1355
Public health agencies laboratory services: Public health-related research (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Public health-related research	Yes	No	No	No	1357
Public health agencies laboratory services: Reference and specialized testing (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Reference and specialized testing	Yes	No	No	No	1351
Public health agencies laboratory services: Training and education (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory services to support essential public health services-Training and education	Yes	No	No	No	1358
Public health majors (percent)	Percent of 4-year colleges and universities that offer public health or related majors	Yes	No	No	No	1385

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Public health minors (percent)	Percent of 4-year colleges and universities that offer public health or related minors	Yes	No	No	No	1386
Public health system assessment: Local public health systems (percent)	Percent of State and local public health jurisdictions that conduct a public health system assessment using national performance standards-Local public health systems	Yes	No	No	No	1376
Public health system assessment: State public health systems (percent)	Percent of State and local public health jurisdictions that conduct a public health system assessment using national performance standards-State public health systems	Yes	No	No	No	1375
Publicly funded family planning clinics offering emergency contraception (percent)	Percent of publicly funded family planning clinics that offer emergency contraception onsite	Yes	No	No	No	782
Publicly funded family planning clinics offering full range of contraception methods (percent)	Percent of publicly funded family planning clinics that offer the full range of FDA-approved methods of contraception onsite	Yes	No	No	No	781
Radon mitigation systems in homes (percent)	Percent of homes with an operating radon mitigation system for persons living in homes at risk for radon exposure	Yes	No	No	No	681
Radon-reducing new home construction (percent)	Percent of new single family homes (SFH) constructed with radon-reducing features, especially in high-radon-potential areas	Yes	No	No	No	682
Receipt of counseling about mammograms (percent)	Percent of women who were counseled by their providers about mammograms	Yes	No	No	No	494
Receipt of counseling about Pap tests (percent)	Percent of women who were counseled by their providers about Pap tests	Yes	No	No	No	495
Receipt of education on early signs/symptoms and responses to asthma episodes (percent)	Percent of persons with current asthma who receive education about appropriate response to an asthma episode, including recognizing early signs and symptoms or monitoring peak flow results	Yes	No	No	No	1415
Receipt of mental health services among homeless adults (percent)	Percent of homeless adults with mental health problems who receive mental health services	Yes	No	No	No	1107

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Receipt of oral health services at health centers (percent)	Percent of patients who receive oral health services at Federally Qualified Health Centers each year	Yes	No	No	No	1296
Receipt of proper-use instructions with asthma inhalers (percent)	Percent of persons with current asthma with prescribed inhalers who receive instruction on their use	Yes	No	No	No	1414
Receipt of written asthma plans from health care provider (percent)	Percent of persons with current asthma who receive written asthma management plans from their health care provider	Yes	No	No	No	1413
Recent hearing exams among adolescents (percent)	Percent of adolescents aged 12 to 19 years who have had a hearing examination in the past 5 years	Yes	No	No	No	750
Recent hearing exams among adults (percent)	Percent of adults aged 20 to 69 years who have had a hearing examination in the past 5 years	Yes	No	No	No	748
Recent hearing exams among older adults (percent)	Percent of adults aged 70 years and older who have had a hearing examination in the past 5 years	Yes	No	No	No	749
Recycling of municipal waste (percent)	Percent of municipal solid waste that is recycled or composted	Yes	No	No	No	662
Reducing fat/calories in diet--persons with pre-diabetes (percent)	Percent of persons at high risk for diabetes (with pre-diabetes) who report reducing the amount of fat or calories in their diet	Yes	No	No	No	564
Regularly scheduled recess - School districts (percent)	Percent of school districts that require regularly scheduled elementary school recess	Yes	No	No	No	1339
Regularly scheduled recess - States (number)	Number of States that require regularly scheduled elementary school recess	Yes	No	No	No	1338
Rehabilitation Hospitals (number)	Number of Rehabilitation hospitals participating in the Medicare program	No	Yes	No	Yes	10048
Religious Nonmedical Health Care Institutions (number)	Number of religious nonmedical health care institutions participating in the Medicare program	No	Yes	No	Yes	10049
Renal evaluation after acute kidney injury (percent)	Percent of hospital patients who incurred acute kidney injury and have followup renal evaluation in 6 months post discharge	Yes	No	No	No	482
Residential fire deaths (per 100,000)	Residential fire death rate	Yes	No	No	No	1050

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Risks posed by hazardous sites (number)	Number of hazardous sites that pose risks to human health and the environment	Yes	No	No	No	677
Rotavirus vaccination (percent)	Percent of young children who have received universally recommended vaccines-2 or more doses rotavirus vaccine (aged 19 to 35 months)	Yes	No	No	No	1002
Rubella, U.S.-acquired (number)	Number of rubella cases (U.S.-acquired cases)	Yes	No	No	No	986
Rural Health Clinics (number)	Number of rural health clinics participating in the Medicare program	No	Yes	No	Yes	10050
Safety belt use (percent)	Percent of persons observed using safety belts	Yes	No	No	No	1047
Salmonellosis (number)	Number of reported cases of salmonellosis	No	Yes	Yes	No	10026
Saturated fat consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake provided by saturated fat for persons aged 2 years and older	Yes	No	No	No	1221
School absenteeism -illness or injury (percent)	Percent of adolescents missing school 11 or more days due to illness or injury	Yes	No	No	No	403
School breakfast program (percent)	Percent of schools with a school breakfast program	Yes	No	No	No	417
School health education - Alcohol and drug use (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in alcohol and other drug use	Yes	No	No	No	571
School health education - All priority areas (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in All priority areas	Yes	No	No	No	566
School health education - growth and development (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Growth and development	Yes	No	No	No	654
School health education - hand washing (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Hand washing or hand hygiene	Yes	No	No	No	652

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
School health education - health screenings (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Importance of health screenings and checkups	Yes	No	No	No	658
School health education - Injury (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in unintentional injury	Yes	No	No	No	567
School health education - Nutrition (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in unhealthy dietary patterns	Yes	No	No	No	573
School health education - oral health (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Dental and oral health	Yes	No	No	No	653
School health education - Physical activity (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in inadequate physical activity	Yes	No	No	No	574
School health education - rest and sleep (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Benefits of rest and sleep	Yes	No	No	No	656
School health education - Suicide (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in suicide	Yes	No	No	No	569
School health education - sun safety (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Sun safety or skin cancer prevention	Yes	No	No	No	655
School health education - Tobacco use (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in tobacco use and addiction	Yes	No	No	No	570

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
School health education - Unintended pregnancy, HIV/AIDS, and STD (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in unintended pregnancy, HIV/AIDS, and STD infection	Yes	No	No	No	572
School health education - Violence (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health education to prevent health problems in violence	Yes	No	No	No	568
School health education - vision and hearing (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promote personal health and wellness-Ways to prevent vision and hearing loss	Yes	No	No	No	657
School health education -Concept comprehension (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)-Comprehending concepts related to health promotion and disease prevention (knowledge)	Yes	No	No	No	636
School nurse-to-student ratio - All schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered school nurse-to-student ratio of at least 1:750-All elementary, middle, and senior high schools	Yes	No	No	No	585
School nurse-to-student ratio - Elementary schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered school nurse-to-student ratio of at least 1:750-Elementary schools	Yes	No	No	No	588
School nurse-to-student ratio - High schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered school nurse-to-student ratio of at least 1:750-Senior high schools	Yes	No	No	No	586
School nurse-to-student ratio - Middle schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered school nurse-to-student ratio of at least 1:750-Middle schools	Yes	No	No	No	587

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
School work considered meaningful (percent)	Percent of adolescents who consider their school work to be meaningful and important	Yes	No	No	No	402
School-based centers with dental care programs (percent)	Percent of school-based health centers with an oral health component that includes dental care	Yes	No	No	No	1285
School-based centers with dental sealant programs (percent)	Percent of school-based health centers with an oral health component that includes dental sealants	Yes	No	No	No	1284
School-based centers with topical fluoride programs (percent)	Percent of school-based health centers with an oral health component that includes topical fluoride	Yes	No	No	No	1286
Schools - analyze influences on health (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Analyzing the influence of culture, media, technology, and other factors on health (skills)	Yes	No	No	No	639
Schools - health advocacy (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Advocating for personal, family, and community health (skills)	Yes	No	No	No	638
Schools - health information access (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Accessing valid information and health promoting products and services (skills)	Yes	No	No	No	637

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Schools - interpersonal communication to enhance health (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Using interpersonal communication skills to enhance health (skills)	Yes	No	No	No	642
Schools - practicing healthy behaviors (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Practicing health-enhancing behaviors and reducing health risks (skills)	Yes	No	No	No	640
Schools - skills to enhance health (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives that address the knowledge and skills articulated in the National Health Education Standards (high school, middle, elementary)- Using goal-setting and decision-making skills to enhance health (skills)	Yes	No	No	No	641
Schools located near highways (percent)	Number of new schools sited within 500 feet of an interstate or Federal or State highway	Yes	No	No	No	725
Schools that have a hazardous material plan (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Have a plan for how to use, label, store, and dispose of hazardous materials	Yes	No	No	No	685
Schools that have a pesticide area marking policy (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Exposure to pesticides by marking areas treated with pesticides	Yes	No	No	No	687

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Schools that have a pesticide information policy (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Exposure to pesticides by notifying staff, students, and families prior to application of the pesticide	Yes	No	No	No	688
Schools that have a policy to address mold problems (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Have a plan for how to address mold problems	Yes	No	No	No	684
Schools that have a spot use pesticide policy (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Exposure to pesticides by using spot treatments and baiting rather than widespread application of pesticide	Yes	No	No	No	686
Schools that have an air quality management program (percent)	Percent of the Nation's elementary, middle, and high schools that engage in practices that promote a healthy and safe physical school environment: Have an indoor air quality management program	Yes	No	No	No	683
Schools that inspect drinking water for bacteria (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Among schools with community water systems, test drinking water for bacteria	Yes	No	No	No	690
Schools that inspect drinking water for coliforms (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Among schools with community water systems, test drinking water for coliforms	Yes	No	No	No	691

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Schools that inspect drinking water for lead (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and engage in practices that promote a healthy and safe physical school environment: Test drinking water outlets for lead	Yes	No	No	No	689
Screen time - Adolescents (percent)	Percent of adolescents in grades 9 through 12 who view television, videos, or play video games for no more than 2 hours a day	Yes	Yes	No	No	1329
Screen time - Children 0-2 years (percent)	Percent of children aged 0 to 2 years who view no television or videos on an average weekday	Yes	No	No	No	1326
Screen time - Children 2-5 years (percent)	Percent of children aged 2 to 5 years who view television, videos, or play video games for no more than 2 hours a day	Yes	No	No	No	1327
Screen time - Children 6-14 years (percent)	Percent of children aged 6 to 14 years who view television, videos, or play video games for no more than 2 hours a day	Yes	No	No	No	1328
Screen time - Computer - Adolescent computer (percent)	Percent of adolescents in grades 9 through 12 who use a computer or play computer games outside of school (for nonschool work) for no more than 2 hours a day	Yes	Yes	No	No	1332
Screen time - Computer - Children 2-5 years (percent)	Percent of children aged 2 to 5 years who use a computer or play computer games outside of school (for nonschool work) for no more than 2 hours a day	Yes	No	No	No	1330
Screen time - Computer - Children 6-14 years (percent)	Percent of children aged 6 to 14 years who use a computer or play computer games outside of school (for nonschool work) for no more than 2 hours a day	Yes	No	No	No	1331
Seasonal flu vaccination among adolescents (percent)	Percent of children aged 13 to 17 years vaccinated annually against seasonal influenza	Yes	No	No	No	1032
Seasonal flu vaccination among adults (percent)	Percent of noninstitutionalized adults aged 18 to 64 years vaccinated annually against seasonal influenza	Yes	No	No	No	1033
Seasonal flu vaccination among children aged 2 to 4 years (percent)	Percent of children aged 2 to 4 years vaccinated annually against seasonal influenza	Yes	No	No	No	1030

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Seasonal flu vaccination among children aged 5 to 12 years. (percent)	Percent of children aged 5 to 12 years vaccinated annually against seasonal influenza	Yes	No	No	No	1031
Seasonal flu vaccination among health care workers (percent)	Percent of health care personnel vaccinated annually against seasonal influenza	Yes	No	No	No	1037
Seasonal flu vaccination among high-risk adults (percent)	Percent of noninstitutionalized high risk adults aged 18 to 64 years vaccinated annually against seasonal influenza	Yes	No	No	No	1034
Seasonal flu vaccination among infants (percent)	Percent of children aged 6 to 23 months vaccinated annually against seasonal influenza- (1 or 2 doses, depending on age-appropriateness and previous doses received)	Yes	No	No	No	1029
Seasonal flu vaccination among nursing home residents (percent)	Percent of institutionalized adults aged 18 years and older in long-term or nursing homes vaccinated annually against seasonal influenza	Yes	No	No	No	1036
Seasonal flu vaccination among older adults (percent)	Percent of noninstitutionalized adults aged 65 years and older vaccinated annually against seasonal influenza	Yes	No	No	No	1035
Seasonal flu vaccination among pregnant women (percent)	Percent of pregnant women vaccinated annually against seasonal influenza	Yes	No	No	No	1038
Self-blood-glucose-monitoring - Persons with diabetes - At least once daily (percent)	Percent of adults with diabetes who perform self-blood glucose-monitoring at least once daily	Yes	No	No	No	554
Serious psychological distress, adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who report serious psychological distress	Yes	No	No	No	435
Serious violent incidents in public schools (percent)	Percent of public schools with a serious violent incident	Yes	No	No	No	408
Sexually active females aged 15 to 19 years using a condom at first intercourse (percent)	Percent of sexually active females aged 15 to 19 years who use a condom at first intercourse	Yes	No	No	No	794

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Sexually active females aged 15 to 19 years using a condom at last intercourse (percent)	Percent of sexually active females aged 15 to 19 years who use a condom at last intercourse	Yes	No	No	No	796
Sexually active females who received reproductive health services (percent)	Percent of sexually active females aged 15 to 44 years who received reproductive health services	Yes	No	No	No	786
Sexually active males aged 15 to 19 years using a condom at first intercourse (percent)	Percent of sexually active males aged 15 to 19 years who use a condom at first intercourse	Yes	No	No	No	795
Sexually active males aged 15 to 19 years using a condom at last intercourse (percent)	Percent of sexually active males aged 15 to 19 years who use a condom at last intercourse	Yes	No	No	No	797
Sexually active males who received reproductive health services (percent)	Percent of sexually active males aged 15 to 44 years who received reproductive health services	Yes	No	No	No	787
Shigellosis (number)	Number of reported cases of shigellosis	No	Yes	Yes	No	10027
Shingles vaccination (percent)	Percent of adults who are vaccinated against zoster (shingles)	Yes	No	No	No	1027
Short stay hospitals (number)	Number of short stay hospitals participating in the Medicare program	No	Yes	No	Yes	10051
Sick leave for adults with balance or dizziness problems (percent)	Percent of adults who have missed days of work or school in the past 12 months because of balance and dizziness problems	Yes	No	No	No	768
Sigmoidoscopy/colonoscopy (percent)	Percent of adults age 50+ that report ever having had a sigmoidoscopy or colonoscopy	No	Yes	Yes	No	117
Single-parent households (percent)	Percent of households that are single-parent households	No	Yes	Yes	No	40
Skilled Nursing Home Title 18 Facilities (number)	Number of skilled nursing home Title 18 (Medicare only) facilities	No	Yes	No	Yes	10052
Smoke free homes --adults (percent)	Percent of adults living in smoke-free homes	Yes	No	No	No	1561

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Smoke-free indoor air laws - Private workplaces (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in private workplaces	Yes	No	No	No	1524
Smoke-free indoor air laws--Bars (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in bars	Yes	No	No	No	1527
Smoke-free indoor air laws--Commercial daycare centers (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in commercial daycare centers	Yes	No	No	No	1529
Smoke-free indoor air laws--home based daycare centers (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in home-based daycare centers	Yes	No	No	No	1530
Smoke-free indoor air laws--hotels and motels (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in hotels and motels	Yes	No	No	No	1532
Smoke-free indoor air laws--Multiunit housing (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in multiunit housing	Yes	No	No	No	1533
Smoke-free indoor air laws--Prisons/correctional facilities (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in prisons and correctional facilities	Yes	No	No	No	1535
Smoke-free indoor air laws--public transportation (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in public transportation	Yes	No	No	No	1531
Smoke-free indoor air laws--public workplaces (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in public workplaces	Yes	No	No	No	1525
Smoke-free indoor air laws--restaurants (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in restaurants	Yes	No	No	No	1526

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Smoke-free indoor air laws--Vehicles with children (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free indoor air that prohibit smoking in vehicles with children	Yes	No	No	No	1534
Smokeless tobacco use - Adults (percent)	Percent of adults aged 18 years and over using smokeless tobacco products	Yes	No	No	No	1499
Smoking cessation attempts by adolescent smokers (percent)	Percent of adolescent smokers in grades 9 through 12 who attempted smoking cessation in the past 12 months	Yes	Yes	No	No	1515
Smoking cessation attempts by adult smokers (percent)	Percent of adult smokers aged 18 years and over who attempted smoking cessation in the past 12 months	Yes	No	No	No	1513
Smoking cessation during pregnancy (percent)	Percent of females aged 18 to 49 years who ceased smoking during the first trimester of pregnancy	Yes	No	No	No	1491
Smoking cessation success (percent)	Percent of adults 18 years and older with smoking cessation success--6 months to 1 year ago	Yes	No	No	No	1555
Smoking: adults (percent)	Percent of adults 18 years and over that report currently smoking cigarettes	No	Yes	Yes	No	13
SNF Medicare admissions (per 1,000 beneficiaries)	Rate of skilled nursing facility (SNF) admissions among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2008
SNF Medicare days (per 1,000 beneficiaries)	Average number of skilled nursing facility (SNF) days among Medicare fee-for-service beneficiaries 65+ years of age	Yes	Yes	No	Yes	2016
SNF Medicare users (number)	Medicare fee-for-service beneficiaries 65+ years of age who had covered skilled nursing facility services	Yes	Yes	No	Yes	214
SNF Medicare utilization (percent)	Medicare fee-for-service beneficiaries 65+ years of age who had covered skilled nursing facility services	Yes	Yes	No	Yes	215
Social and emotional support, adults with disabilities (percent)	Percent of adults with disabilities who report sufficient social and emotional support	Yes	No	No	No	515
Social support (percent)	Percent of adults who report having friends or family members whom they talk with about their health	Yes	No	No	No	870

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Social-emotional support (percent)	Percent of adults 18 years and over that report receiving sufficient social-emotional support	No	Yes	Yes	No	39
Sodium consumption (milligrams)	Mean total daily sodium intake (in mg) by persons aged 2 years and older	Yes	No	No	No	1222
Solid fat and added sugar consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake provided by solid fats and added sugars for persons aged 2 years and older	Yes	No	No	No	1238
Solid fat consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake provided by solid fats for persons aged 2 years and older	Yes	No	No	No	1236
Source of ongoing care - 17 years and under (percent)	Percent of persons who have a specific source of ongoing care-children and youth aged 17 years and under	Yes	No	No	No	376
Source of ongoing care - 18 to 64 years (percent)	Percent of persons who have a specific source of ongoing care-adults aged 18 to 64 years	Yes	No	No	No	377
Source of ongoing care - 65 years and older (percent)	Percent of persons who have a specific source of care - adults aged 65 years and older	Yes	No	No	No	378
Source of ongoing care - all ages (percent)	Percent of persons who have a specific source of ongoing care-all ages	Yes	No	No	No	375
Special services for children with Autism spectrum disorders (percent)	Percent of children with an ASD enrolled in special services by 48 months of age	Yes	No	No	No	1190
Specialty health care for adults with moderate to severe balance or dizziness problems (percent)	Percent of adults with moderate to severe balance or dizziness problems who have seen or been referred to a Health Care specialist for evaluation or treatment	Yes	No	No	No	763
Specialty Health Care for adults with moderate to severe tinnitus (percent)	Percent of adults bothered by the onset of tinnitus in the past 5 years, for whom it is a moderate, big, or very big problem, who have seen or been referred to an audiologist or otolaryngologist (ENT physician)	Yes	No	No	No	760
Spina bifida births (per 100,000 live births)	Rate of spina bifida	Yes	No	No	No	1169
Spinal cord injury deaths (per 100,000)	Spinal cord injury death rate	Yes	No	No	No	1062

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Spinal cord injury hospitalizations (per 100,000)	Hospitalization rate for nonfatal traumatic spinal cord injuries	Yes	No	No	No	1063
Sports and recreation injuries (per 1,000)	Sports and recreation injury rate	Yes	No	No	No	1098
Spotted Fever Rickettsiosis (number)	Number of reported cases of spotted fever rickettsiosis	No	Yes	Yes	No	10028
State and local agencies with a dental professional directing their dental program (percent)	Percent of States (including the District of Columbia) and local health agencies that serve jurisdictions of 250,000 or more persons with a dental public health program directed by a dental professional with public health training	Yes	No	No	No	1291
State epidemiologists with formal training in epidemiology (percent)	Percent of Tribal, State, and local public health agencies that provide or assure comprehensive epidemiology services to support essential public health services-State epidemiologists with formal training in epidemiology	Yes	No	No	No	1360
State nutrition standards for child care (number)	Number of States and the District of Columbia with nutrition standards for foods and beverages provided to preschool-aged children in child care	Yes	No	No	No	1240
State vital event reporting: Certificates of birth (number)	Number of States that record vital events using the latest U.S. standard certificates and report-States using the standard certificate of birth	Yes	No	No	No	1382
State vital event reporting: Certificates of death (number)	Number of States that record vital events using the latest U.S. standard certificates and report - States using the standard certificate of death	Yes	No	No	No	1383
State vital event reporting: Report of fetal death (number)	Number of States that record vital events using the latest U.S. standard certificates and report - States using the standard report of fetal death	Yes	No	No	No	1384
State-level incentive policies for food retail (number)	Number of States and the District of Columbia that have State-level policies that incentivize food retail outlets to provide foods that are encouraged by the Dietary Guidelines	Yes	No	No	No	1239

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
States with "good" graduated driver licensing laws (number)	Number of States and the District of Columbia with "good" graduated driver licensing (GDL) laws	Yes	No	No	No	1058
States with asthma surveillance systems (number)	Numbers of States, Territories, and the District of Columbia with a comprehensive asthma surveillance system for tracking asthma cases, illness, and disability at the State level	Yes	No	No	No	1421
States with bicycle helmet laws (number)	Number of States and the District of Columbia with laws requiring bicycle helmets for bicycle riders	Yes	No	No	No	1049
States with emergency department routine E-code collection (percent)	Percent of States and the District of Columbia with statewide emergency department data systems that routinely collect external-cause-of injury codes for 90 percent or more of injury-related visits	Yes	No	No	No	1093
States with hospital discharge routine E-code collection (percent)	Percent of States and the District of Columbia with statewide hospital discharge data systems that routinely collect external-cause-of injury codes for 90 percent or more of injury-related discharges	Yes	No	No	No	1094
States with violent death surveillance systems (number)	Number of States that link data on violent deaths from death certificates, law enforcement, and coroner and medical examiner reports to inform prevention efforts at the State and local levels	Yes	No	No	No	1101
Statewide cancer registries (number)	Number of central, population-based registries from the 50 States and the District of Columbia that capture case information on at least 95 percent of the expected number of reportable cancers	Yes	No	No	No	497
Streptococcus pneumoniae, drug-resistant (number)	Number of reported cases of drug-resistant streptococcus pneumoniae	No	Yes	Yes	No	10025
Stroke deaths (per 100,000)	Stroke death rate	Yes	Yes	Yes	No	881
Student safety at school (percent)	Percent of adolescents whose parents consider them to be safe at school	Yes	No	No	No	407

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Sufficient sleep among 9th to 12th graders (percent)	Percent of students in grades 9 through 12 who get sufficient sleep	Yes	No	No	No	1471
Sufficient sleep among adults (percent)	Percent of adults who get sufficient sleep	Yes	No	No	No	1472
Suicide deaths (per 100,000)	Suicide death rate	Yes	Yes	Yes	No	1105
Surgery patients who received appropriate venous thromboembolism prophylaxis (percent)	Hospital Compare: Surgery patients who received appropriate venous thromboembolism prophylaxis	Yes	Yes	No	Yes	342
Surgery patients with appropriate hair removal (percent)	Hospital Compare: Surgery patients with appropriate hair removal	Yes	Yes	No	Yes	340
Surgery patients with appropriate prophylactic antibiotic selection (percent)	Hospital Compare: Surgery patients with appropriate prophylactic antibiotic selection	Yes	Yes	No	Yes	337
Surgery patients with prophylactic antibiotic received within one hour prior to surgery incision (percent)	Hospital Compare: Surgery patients with prophylactic antibiotic received within one hour prior to surgery incision	Yes	Yes	No	Yes	336
Surgery patients with prophylactic antibiotics discontinued within 24 hours after surgery end time (percent)	Hospital Compare: Surgery patients with prophylactic antibiotics discontinued within 24 hours after surgery end time	Yes	Yes	No	Yes	338
Surgery patients with recommended venous thromboembolism prophylaxis ordered (percent)	Hospital Compare: Surgery patients with recommended venous thromboembolism prophylaxis ordered	Yes	Yes	No	Yes	341
Surveillance programs for caregivers (States and DC) (number)	States and the District of Columbia with surveillance programs for caregivers	Yes	No	No	No	528
Survival after AIDS diagnosis (percent)	Percent of persons surviving more than 3 years after a diagnosis with AIDS	Yes	No	No	No	937
TB test confirmation timeliness (percent)	Average time for a laboratory to confirm and report tuberculosis cases	Yes	No	No	No	972
TB, Curative therapy (percent)	Percent of all tuberculosis patients who complete curative therapy within 12 months	Yes	No	No	No	970

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
TB, Treatment for latent (percent)	Percent of persons with latent tuberculosis infection who complete a course of treatment	Yes	No	No	No	971
Tdap vaccination among adolescents (percent)	Percent of adolescents who have 1 dose of tetanus-diphtheria-acellular pertussis (Tdap) booster vaccine by 13 to 15 years	Yes	No	No	No	1010
Time for recess (percent)	Percent of school districts that require or recommend elementary school recess for an appropriate period of time	Yes	No	No	No	1340
Timely fibrinolytic therapy-STEMI patients (percent)	Percent of eligible patients with heart attacks or strokes who receive timely artery-opening therapy as specified by current guidelines-Fibrinolytic therapy within 30 minutes of hospital arrival for patients with heart attacks	Yes	No	No	No	878
Timely PCI therapy-STEMI patients (percent)	Percent of eligible patients with heart attacks or strokes who receive timely artery-opening therapy as specified by current guidelines-Percutaneous intervention (PCI) within 90 minutes of hospital arrival for patients with heart attacks	Yes	No	No	No	879
Tobacco advertising and promotions: internet advertising exposure (percent)	Percent of adolescents and young adults grades 6 through 12 who are exposed to tobacco advertising and promotion-Internet advertising and promotion	Yes	No	No	No	1541
Tobacco advertising and promotions: magazine and newspaper advertising exposure (percent)	Percent of adolescents and young adults grades 6 through 12 who are exposed to tobacco advertising and promotion-Magazine and newspaper advertising and promotion	Yes	No	No	No	1542
Tobacco cessation counseling in hospital ambulatory care settings (percent)	Percent of tobacco users ages 12 and older who receive tobacco cessation counseling at hospital ambulatory care visits	Yes	No	No	No	1552
Tobacco cessation counseling in office-based ambulatory care settings (percent)	Percent of tobacco users ages 12 and older who receive tobacco cessation counseling at office based ambulatory care visits	Yes	No	No	No	1551
Tobacco free environments--Junior High Schools (percent)	Percent of Junior high schools with tobacco-free environments	Yes	No	No	No	1520

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Tobacco free environments--Middle High Schools (percent)	Percent of middle schools with tobacco-free environments	Yes	No	No	No	1521
Tobacco free environments--Senior High Schools (percent)	Percent of senior high schools with tobacco-free environments	Yes	No	No	No	1522
Tobacco screening--hospital ambulatory visits (percent)	Percent of persons aged 12 years and older who received tobacco screening in hospital ambulatory care settings	Yes	No	No	No	1558
Tobacco screening--office based ambulatory visits (percent)	Percent of persons aged 12 years and older who received tobacco screening in office-based ambulatory care settings	Yes	No	No	No	1557
Tobacco use by adolescents-Cigarettes (past month) (percent)	Percent of adolescent reporting use of cigarettes in the past month - Students (grades 9 through 12)	Yes	Yes	No	No	1502
Tobacco use by adolescents-Cigars (past month) (percent)	Percent of adolescent reporting use of cigars in the past month - Students (grades 9 through 12)	Yes	Yes	No	No	1504
Tobacco use by adolescents-Smokeless tobacco products (past month) (percent)	Percent of adolescent reporting use of smokeless tobacco products in the past month - Students (grades 9 through 12)	Yes	Yes	No	No	1503
Tobacco use by adolescents-Tobacco products (past month) (percent)	Percent of adolescent reporting use of tobacco products in the past month - Students (grades 9 through 12)	Yes	Yes	No	No	1501
Toxic chemicals (pounds)	Toxic chemicals released annually	No	No	Yes	No	105
Toxic pollutants (pounds)	Amount of toxic pollutants released into the environment	Yes	No	No	No	678
Transition planning for youth with special health care needs (percent)	Percent of youth with special health care needs whose health care provider has discussed transition planning from pediatric to adult Health Care	Yes	No	No	No	536
Trauma care access, by land mass (percent)	Percent of the land mass of the continental United States with access to trauma care	Yes	No	No	No	1097
Trauma care access, by population (percent)	Percent of the population residing within the continental United States with access to trauma care	Yes	No	No	No	1096
Trauma centers implementing alcohol Screening and Brief Interventions (number)	Number of Level I and Level II trauma centers and primary care settings that implement evidence-based alcohol Screening and Brief Intervention (SBI)	Yes	No	No	No	1460

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Traumatic brain injury deaths (per 100,000)	Traumatic brain injury death rate	Yes	No	No	No	1059
Traumatic brain injury ED visits (per 100,000)	Rate of emergency department visits for nonfatal traumatic brain injuries	Yes	No	No	No	1061
Traumatic brain injury hospitalizations (per 100,000)	Hospitalization rate for nonfatal traumatic brain injuries	Yes	No	No	No	1060
Treatment for children with mental health problems (percent)	Percent of children with mental health problems who receive treatment	Yes	No	No	No	1110
Treatment for co-occurring substance abuse and mental disorders (percent)	Percent of persons with co-occurring substance abuse and mental disorders who receive treatment for both disorders	Yes	No	No	No	1119
Treatment of adults with major depressive episode (percent)	Percent of adults aged 18 years and older with major depressive episode	Yes	No	No	No	1118
Treatment of adults with serious mental illness (percent)	Percent of adults aged 18 years and older with serious mental illness (SMI)	Yes	No	No	No	1117
Trying to lose weight--persons with pre-diabetes (percent)	Percent of persons at high risk for diabetes (with pre-diabetes) who report trying to lose weight	Yes	No	No	No	563
Tuberculosis (per 100,000)	Tuberculosis (TB) rate	Yes	No	No	No	969
Tuberculosis in foreign born persons (per 100,000)	Tuberculosis (TB) case rate for foreign-born persons living in the United States	Yes	No	No	No	845
Undergraduate nursing training in counseling for health promotion and disease prevention (percent)	Percent of inclusion of core clinical prevention and population health content-Counseling for health promotion and disease prevention in health professions training-Undergraduate nursing	Yes	No	No	No	618
Undergraduate nursing training in cultural diversity (percent)	Percent of inclusion of core clinical prevention and population health content-Cultural diversity in health professions training-Undergraduate nursing	Yes	No	No	No	619
Undergraduate nursing training in environmental health (percent)	Percent of inclusion of core clinical prevention and population health content-Environmental health in health professions training-Undergraduate nursing	Yes	No	No	No	621

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Undergraduate nursing training in evaluation of health sciences literature (percent)	Percent of inclusion of core clinical prevention and population health content-Evaluation of health sciences literature in health professions training-Undergraduate nursing	Yes	No	No	No	620
Undergraduate nursing training in global health (percent)	Percent of inclusion of core clinical prevention and population health content-Global health in health professions training-Undergraduate nursing	Yes	No	No	No	623
Undergraduate nursing training in public health systems (percent)	Percent of inclusion of core clinical prevention and population health content-Public health systems in health professions training-Undergraduate nursing	Yes	No	No	No	622
Unemployed (percent)	Percent of population age 16+ unemployed and looking for work	No	Yes	Yes	No	35
Unemployment among adults with arthritis (percent)	Impact of doctor-diagnosed arthritis on employment in the working-aged population-Unemployment rate among adults with doctor-diagnosed arthritis	Yes	No	No	No	423
Unemployment among adults with disabilities (percent)	Unemployment rates among adults aged 16-64 with disabilities	Yes	No	No	No	541
Unintentional and undetermined poisoning deaths (per 100,000)	Poisoning death rate of unintentional or undetermined intent among all persons	Yes	No	No	No	1066
Unintentional and undetermined poisoning deaths, 35-54 years (per 100,000)	Poisoning death rate of unintentional or undetermined intent among persons aged 35 to 54 years	Yes	No	No	No	1067
Unintentional injury deaths (per 100,000)	Unintentional injury death rate	Yes	Yes	Yes	No	1074
Unintentional Injury deaths: No MVA (per 100,000)	Unintentional injury death rate (excluding motor vehicle injuries)	Yes	Yes	Yes	No	79
Unintentional suffocation deaths (per 100,000)	Unintentional suffocation death rate: All persons	Yes	No	No	No	1068

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Unintentional suffocation deaths, 0-12 months (per 100,000)	Unintentional suffocation death rate: Infants 0 to 12 months	Yes	No	No	No	1069
Unintentional suffocation deaths, ages 65 years and older (per 100,000)	Unintentional suffocation death rate: Persons aged 65 years and older	Yes	No	No	No	1070
Untreated dental decay in adolescents (percent)	Percent of adolescents aged 13 to 15 years with untreated dental decay in their permanent teeth	Yes	No	No	No	1274
Untreated dental decay in adults (percent)	Percent of adults aged 35 to 44 years with untreated dental decay	Yes	No	No	No	1275
Untreated dental decay in children (percent)	Percent of children aged 6 to 9 years with untreated dental decay in their primary and permanent teeth	Yes	No	No	No	1273
Untreated dental decay in older adults (percent)	Percent of older adults aged 65 to 74 years with untreated coronal caries	Yes	No	No	No	1276
Untreated dental decay in the oldest adults (percent)	Percent of older adults aged 75 years and older with untreated root surface caries	Yes	No	No	No	1277
Untreated dental decay in young children (percent)	Percent of young children aged 3 to 5 years with untreated dental decay in their primary teeth	Yes	No	No	No	1272
Urinary tract infection admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Urinary tract infection admission rate among Medicare beneficiaries 65+ years of age with full fee-for-service coverage for 12 months	Yes	Yes	No	Yes	2044
Use of hearing protection devices among adolescents (per 1,000)	Rate of adolescents aged 12 to 19 years who have ever used hearing protection devices (earplugs, earmuffs) when exposed to loud sounds or noise	Yes	No	No	No	753
Use of hearing protection devices among adults (per 1,000)	Rate of adults aged 20 to 69 years who have ever used hearing protection devices (earplugs, earmuffs) when exposed to loud sounds or noise	Yes	No	No	No	752
Use of Internet to communicate with their provider (percent)	Percent of persons who use the Internet to communicate with their provider	Yes	No	No	No	868

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Use of Internet to keep track of personal health information (percent)	Percent of persons who use the Internet to keep track of personal health information, such as care received, test results, or upcoming medical appointments	Yes	No	No	No	867
Use of intervention therapy by adults with balance or dizziness problems (percent)	Percent of adults who have tried recommended methods for treating their balance or dizziness problem	Yes	No	No	No	765
Use of intervention therapy by adults with moderate to severe tinnitus (percent)	Percent of adults for whom tinnitus is a moderate to severe problem who have tried appropriate treatments	Yes	No	No	No	761
Use of mobile devices for internet access (percent)	Percent of persons who use mobile devices for internet use	Yes	No	No	No	859
Use of protective eyewear at home among adults (percent)	Percent of adults aged 18 years and over who use protective eyewear in recreational activities and hazardous situations around the home	Yes	No	No	No	1573
Use of protective eyewear at home among children (percent)	Percent of children and adolescents aged 6 to 17 years who use personal protective eyewear in recreational activities and hazardous situations around the home	Yes	No	No	No	1572
Usual primary care provider (percent)	Percent of persons with a usual primary care provider	Yes	No	No	No	372
Uterine cervix cancer deaths (per 100,000)	Uterine cervix cancer death rate	Yes	No	No	No	489
Vaccination coverage assessment of private health providers (percent)	Percent of private providers who have had vaccination coverage levels among children in their practice population measured within the past year	Yes	No	No	No	977
Vaccination coverage assessment of public health providers (percent)	Percent of public health providers who have had vaccination coverage levels among children in their practice population measured within the past year	Yes	No	No	No	976
Vaccination record keeping (percent)	Percent of children under age 6 years of age whose immunization records are in fully operational, population-based immunization information systems	Yes	No	No	No	1009

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Varicella among children (number)	Rate of varicella (chicken pox) cases among persons aged 17 years of age and under	Yes	No	No	No	987
Varicella vaccination (percent)	Percent of young children who have received universally recommended vaccines-1 dose varicella vaccine by age 19 to 35 months	Yes	No	No	No	998
Varicella vaccination among adolescents (percent)	Percent of adolescents who have 2 doses of varicella vaccine by age 13 to 15 years (excluding children who have had varicella)	Yes	No	No	No	1011
Varicella vaccination among kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-Varicella vaccine (2 or more doses)	Yes	No	No	No	1007
Vascular access for hemodialysis (percent)	Percent of adult hemodialysis patients who use arteriovenous fistulas as the primary mode of vascular access	Yes	No	No	No	469
Vegetable consumption (cup equivalents per 1,000 calories)	Mean daily intake of vegetables (in cup equivalents per 1,000 calories) by persons aged 2 years and older	Yes	No	No	No	1218
Vegetable consumption-dark green, orange, and legumes (cup equivalents per 1,000 calories)	Mean daily intake of dark green vegetables, orange vegetables, and legumes (in cup equivalents per 1,000 calories) by persons aged 2 years and older	Yes	No	No	No	1219
Vehicular crashes due to drowsy driving (per 100,000,000)	Rate of vehicular crashes per 100 million miles traveled that are due to drowsy driving	Yes	No	No	No	1470
Venous thromboembolism visits (per 10,000)	Rate of visits for venous thromboembolism (VTE)	Yes	No	No	No	441
Verification of newborn screening (number)	Number of States and the District of Columbia that verify through linkage with vital records that all newborns are screened shortly after birth for conditions mandated by their State-sponsored screening program	Yes	No	No	No	1173
Very low birth weight (percent)	Percent of live births that are very low birth weight (VLBW)	Yes	Yes	Yes	No	1136
Very low birth weight infants born at level III hospitals (percent)	Percent of very low birth weight (VLBW) infants born at level III hospitals or subspecialty perinatal centers	Yes	No	No	No	1195

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Very preterm births (percent)	Percent of live births that are very preterm (less than 32 weeks of gestation)	Yes	Yes	Yes	No	1140
Violent crime (per 100,000)	Violent crime rate	No	Yes	Yes	No	42
Vision screening for children (percent)	Percent of preschool children aged 5 years and under who receive vision screening	Yes	No	No	No	1562
Visitable features - homes and residential buildings (percent)	Percent of newly constructed or retrofitted U.S. homes and residential buildings that have visitable features	Yes	No	No	No	542
Visual impairment in children and adolescents (per 1,000)	Percent of children and adolescents aged 17 years and under with blindness and visual impairment	Yes	No	No	No	1563
Visual rehabilitation services use (per 1,000)	Rate of use of vision rehabilitation services by persons with visual impairment	Yes	No	No	No	1574
Waitlisting and/or transplantation (percent)	percent of dialysis patients wait-listed and/or receiving a deceased donor kidney transplant within 1 year of end-stage renal disease (ESRD) start (among patients under 70 years of age)	Yes	No	No	No	472
Waterborne disease outbreaks (number)	Number of waterborne disease outbreaks arising from water intended for drinking among persons served by community water systems	Yes	No	No	No	660
Weapon carrying by adolescents on school property (percent)	Percent of weapon carrying by adolescents on school property	Yes	Yes	No	No	1053
Welcome to Medicare benefit use (percent)	Percent of older adults who use the Welcome to Medicare benefit	Yes	No	No	No	1260
Whole grain consumption (ounce equivalents per 1,000 calories)	Mean daily intake of whole grains (in ounce equivalents per 1,000 calories) by persons aged 2 years and older	Yes	No	No	No	1220
Work-related assaults (per 10,000)	Rate of work-related assaults	Yes	No	No	No	1311
Work-related homicides (per 10,000)	Number of deaths due to work-related homicides	Yes	No	No	No	1299
Work-related injury deaths - Agriculture, forestry, fishing, and hunting (per 100,000)	Rate of work-related injury deaths - Agriculture, forestry, fishing, and hunting	Yes	No	No	No	1307

Short Description	Full Description	National	State	County	Hospital Referral Region	HIW ID
Work-related injury deaths - All industry (per 100,000)	Rate of work-related injury deaths - All industry	Yes	No	No	No	1303
Work-related injury deaths - Construction (per 100,000)	Rate of work-related injury deaths - Construction	Yes	No	No	No	1305
Work-related injury deaths - Mining (per 100,000)	Rate of work-related injury deaths - Mining	Yes	No	No	No	1304
Work-related injury deaths - Transportation (per 100,000)	Rate of work-related injury deaths - Transportation	Yes	No	No	No	1306
Work-related, noise-induced hearing loss (per 10,000 full-time workers)	Rate of work-related, noise-induced hearing loss	Yes	No	No	No	1302
Worksite lactation support programs (percent)	Percent of employers that have worksite lactation support programs	Yes	No	No	No	1184
Years of potential life lost before age 75 (per 100,000)	Rate per 100,000 of years of potential life lost before age 75	Yes	Yes	Yes	No	3
Young adult deaths aged 20-24 years (per 100,000)	Rates of adolescent and young adult deaths-Young adults aged 20 to 24 years	Yes	No	No	No	1128
Youth perpetration of violent crimes (per 100,000)	Rate of minor and young adult perpetration of violent crimes	Yes	No	No	No	413
Youth perpetration-serious property crimes (per 100,000)	Rate of minor and young adult perpetration of serious property crimes	Yes	No	No	No	414
Zero doses of vaccination (percent)	Percent of children in the United States who receive zero doses of recommended vaccines by age 19 to 35 months	Yes	No	No	No	1039