

Draft Environmental Assessment for a Marine Geophysical Survey of Portions of the Arctic Ocean, August-September, 2010

by

Beth Haley¹, Darren Ireland¹, and Jonathan R. Childs²

¹LGL ALASKA RESEARCH ASSOCIATES, INC.

Anchorage, AK 99518

²U.S. GEOLOGICAL SURVEY

Menlo Park, CA 94025

Technical assistance provided by:



Alaska Research Associates, Inc.

LGL Alaska Research Associates, Inc.
Anchorage, AK 99518

May 2010
LGL Document P1122-1

Suggested citation format:

Haley, B., and others, 2010. Draft Environmental Assessment for a marine geophysical survey of portions of the Arctic Ocean, August–September 2010;
U.S. Geological Survey, Menlo Park, CA

TABLE OF CONTENTS

| | Page |
|--|------------|
| TABLE OF CONTENTS | III |
| ABSTRACT | V |
| LIST OF ACRONYMS..... | VII |
| I. PURPOSE AND NEED | 1 |
| II. ALTERNATIVES INCLUDING PROPOSED ACTION | 3 |
| Proposed Action..... | 3 |
| (1) Project Objectives and Context | 3 |
| (2) Proposed Activities..... | 3 |
| (3) Mitigation Measures..... | 12 |
| Alternative Action: Conduct Survey during Alternative Time Period | 19 |
| No Action Alternative | 20 |
| III. AFFECTED ENVIRONMENT..... | 20 |
| Physical Environment | 20 |
| Biological Environment | 21 |
| Fish Resources | 21 |
| Seabirds | 22 |
| (1) Spectacled Eider | 23 |
| (2) Steller’s Eider | 23 |
| (3) Other Seabirds, Shorebirds, and Waterfowl..... | 24 |
| Marine Mammals | 25 |
| (1) Odontocetes | 25 |
| (2) Mysticetes..... | 30 |
| (3) Pinnipeds | 35 |
| (4) Carnivora..... | 40 |
| IV. ENVIRONMENTAL CONSEQUENCES OF PROPOSED ACTION | 42 |
| Direct Effects on Marine Mammals and their Significance | 42 |
| (1) Summary of Potential Effects of Airgun Sounds | 42 |
| (2) Possible Effects of Multibeam Echo Sounder Signals | 49 |
| (3) Possible Effects of Chirp Echo Sounder Signals..... | 51 |
| (4) Possible Effects of Chirp Sub-bottom Profiler | 52 |
| (5) Possible Effects of Helicopter Activities..... | 53 |
| (6) Possible Effects of Icebreaking Activities | 54 |
| (7) Mitigation Measures..... | 55 |
| (8) Numbers of Marine Mammals that May be “Taken by Harassment” | 56 |
| (9) Conclusions | 64 |

| | |
|--|------------|
| (10) Direct Effects on Fish, EFH, and Fisheries, and Their Significance..... | 65 |
| (11) Direct Effects on Seabirds and their Significance | 66 |
| (12) Indirect Effects to Marine Mammals and Their Significance | 67 |
| (13) Possible Effects on Subsistence Hunting and Fishing..... | 67 |
| (14) Cumulative Effects | 71 |
| (15) Unavoidable Impacts of Noise | 73 |
| (16) Coordination with Other Agencies and Processes..... | 74 |
| Alternative Action: Conduct Survey during Alternative Time Period | 75 |
| No Action Alternative | 76 |
| V. LIST OF PREPARERS..... | 77 |
| VI. LITERATURE CITED..... | 78 |
| APPENDIX A NOAA’S 2008 LETTER OF CATEGORICAL EXCLUSION FOR HEALY OPERATIONS IN INTERNATIONAL WATERS | 99 |
| APPENDIX B NOAA’S 2009 LETTER OF CATEGORICAL EXCLUSION FOR HEALY OPERATIONS IN INTERNATIONAL WATERS | 101 |
| APPENDIX C MARINE FISH OF THE BEAUFORT SEA AND ARCTIC OCEAN. FROM FISHBASE.ORG | 104 |
| APPENDIX D REVIEW OF THE EFFECTS OF AIRGUN SOUNDS ON MARINE MAMMALS..... | 112 |
| 1. Categories of Noise Effects | 112 |
| 2. Hearing Abilities of Marine Mammals..... | 112 |
| 2.1 Toothed Whales (Odontocetes)..... | 113 |
| 2.2 Baleen Whales (Mysticetes)..... | 113 |
| 2.3 Seals and Sea Lions (Pinnipeds) | 114 |
| 2.4 Manatees and Dugong (Sirenians) | 114 |
| 2.5 Sea Otter and Polar Bear | 115 |
| 3. Characteristics of Airgun Sounds | 115 |
| 4. Masking Effects of Airgun Sounds | 117 |
| 5. Disturbance by Seismic Surveys | 118 |
| 5.1 Baleen Whales..... | 120 |
| 5.2 Toothed Whales | 126 |
| 5.3 Pinnipeds..... | 131 |
| 5.4 Sirenians, Sea Otter and Polar Bear | 133 |
| 6. Hearing Impairment and Other Physical Effects of Seismic Surveys..... | 134 |
| 6.1 Temporary Threshold Shift (TTS) | 135 |
| 6.2 Permanent Threshold Shift (PTS) | 139 |
| 6.3 Strandings and Mortality..... | 141 |
| 6.4 Non-Auditory Physiological Effects | 143 |

7. Literature Cited.....143

APPENDIX E REVIEW OF THE EFFECTS OF AIRGUN SOUNDS ON FISH..... 159

1. Acoustic Capabilities.....159

2. Potential Effects on Fishes.....161

 2.1 Marine Fishes..... 161

 2.2 Freshwater Fishes..... 164

 2.3 Anadromous Fishes..... 164

3. Indirect Effects on Fisheries.....165

4. Literature Cited.....166

**APPENDIX F REVIEW OF THE EFFECTS OF AIRGUN SOUNDS ON MARINE
INVERTEBRATES 170**

1. Sound Production.....170

2. Sound Detection.....171

3. Potential Seismic Effects.....171

4. Literature Cited.....175

ABSTRACT

According to the United Nations Convention on the Law of the Sea (UNCLOS), individual nations' sovereign rights extend to 200 nautical miles (n.mi.) (370 km) offshore in an area called the continental shelf. These rights include jurisdiction over all resources in the water column and on and beneath the seabed. Article 76 of UNCLOS also establishes the criteria to determine areas beyond the 200 n.mi. (370 km) limit that could be defined as "extended continental shelf" where a nation could extend its sovereign rights over the seafloor and sub-seafloor¹. This jurisdiction provided in Article 76 includes resources on and below the seafloor but not in the water column. The United States has been acquiring data to determine the outer limits of its extended continental shelf (ECS) in the Arctic and has a vested interest in declaring and receiving international recognition of the reach of its extended continental shelf.

The U.S. collaborated with Canada in 2008 and 2009 on ECS studies in the Arctic Ocean. The U.S. Coast Guard (USCG) Cutter *Healy* worked with the Canadian Coast Guard ship *Louis S. St. Laurent* to map the continental shelf beyond 200 n.mi. (370 km) in the Arctic. Each icebreaking vessel contributed different capabilities in order to collect data needed by both nations more efficiently in order to save money, avoid redundancy and foster cooperation. Generally, the *Healy* collects bathymetric (seafloor topography) data and the *Louis S. St. Laurent* collects seismic reflection profile data. The vessels work in concert when ice conditions are heavy, with one vessel breaking ice for the ship collecting data. The Canadian Environmental Assessments for these projects are available on line at <http://www.ceaa.gc.ca/052/details-eng.cfm?pid=38185> (2008) and <http://www.ceaa.gc.ca/052/details-eng.cfm?pid=46518> (2009).

The U.S. Geological Survey (USGS) proposed a similar partnership again for 2010 in a limited area of U.S. waters during the period between ~6 to 12 August. The survey vessels will then proceed to international waters where surveying will proceed until ~3 September. The survey area will be bounded approximately by 145° to 158° W longitude and 71° to 84° N latitude in water depths ranging from ~2000-4000 m (Fig. 1). Ice conditions are expected to range from open water to 10/10 ice cover. The *Louis S. St. Laurent* will join accompanying vessel *Healy* in or near the survey area around 6 August to begin the survey.

As its energy source, the seismic system aboard *Louis S. St. Laurent* will employ a 3-airgun array that consisting of three Sercel G-airguns. Two guns will have a discharge volume of 500 in³ and the third a discharge volume of 150 in³ for a total array discharge volume of 1150 in³. The seismic survey will take place in water depths 2000–4000 m. This airgun array is identical to the system used in the 2008 and 2009 field programs by the Geological Survey of Canada.

The USGS requested that the National Marine Fisheries Service (NMFS) issue an Incidental Harassment Authorization (IHA) to authorize the incidental, i.e., not intentional, harassment of small numbers of cetaceans and seals should this occur during the seismic survey in US waters. USGS is also consulting with the U.S. Fish & Wildlife Service (USFWS) regarding concerns about disturbance to walrus and polar bears. Through informal consultation with the Office of Protected Resources with the National Oceanic and Atmospheric Administration (NOAA), USGS proposes that no ESA-listed marine species: bowhead, fin, humpback or sperm whale will be adversely affected by this project during the

¹ As used in UNCLOS, "continental shelf" refers to a legally defined region of the seafloor rather than a morphological shallow-water area adjacent to continents commonly used by geologists and hydrographers.

survey or transit to the survey area from Dutch Harbor. The information in this Environmental Assessment (EA) supports the IHA Application process, consultation with the USFWS, and provides information on marine species, some of which is also contained in the IHA Application to NMFS. Alternatives addressed in this EA consist of a similar program during a different time period along with issuance of an associated IHA, and the no action alternative, with no IHA and no seismic survey.

Several species of cetaceans and pinnipeds inhabit the Arctic Ocean. Few species that may be found in the survey area are listed as Endangered under the U.S. Endangered Species Act (ESA). The bowhead whale is the endangered species most likely to occur within the survey area. The polar bear, which was recently listed as Threatened under the ESA, may also occur in the survey area. The survey has been scheduled specifically to avoid the spring and fall bowhead whale migrations north of Barrow. Two additional species of special concern (birds) that might be encountered are the spectacled and Steller's eiders, which are listed as "threatened."

Potential impacts on the environment due to the seismic survey would be primarily a result of the operation of the airgun source. In addition to the airgun array, a Chirp pulse echo sounder will be operated on the *Louis S. St. Laurent*. The *Louis S. St. Laurent* will also tow a 3–5 kHz sub-bottom profiler while in open water and when not working with the *Healy*. The *Healy* will use a multibeam echo sounder, a sub-bottom profiler and a "piloting" echo sounder continuously when underway and during the seismic profiling. Acoustic Doppler current profilers may also be used on the *Healy*. The project will also involve vessel and helicopter traffic. Increased underwater noise from vessel traffic and use of geophysical equipment may result in avoidance behavior or other disturbance to some marine mammals and fish. An integral part of the planned survey is a monitoring and mitigation program to minimize impacts of the proposed activities on marine species and on fishing and subsistence activities, and to document the nature and extent of any effects. Injurious impacts to marine mammals have not been demonstrated to occur near airgun arrays, and the planned monitoring and mitigation measures would minimize the possibility of such effects should they occur.

Protection measures designed to mitigate the potential environmental impacts will include the following: a minimum of one dedicated protected species observer (PSO) maintaining a visual watch during all daylight airgun operations; two observers (when possible, otherwise a single observer) on watch 30 min before airgun operations start; and power downs or shut downs of the airgun array when mammals are detected in, or about to enter, designated safety radii. USGS and its collaborators are committed to apply these measures in order to minimize disturbance of marine mammals and to minimize the risk of injuries or other environmental impacts.

With the planned monitoring and mitigation measures, unavoidable impacts to each of the species of marine mammal that might be encountered are expected to be limited to short-term localized changes in behavior and distribution near the seismic vessel. At most, such effects may be interpreted as falling within the Marine Mammal Protection Act (MMPA) definition of "Level B Harassment". No long-term or significant effects are expected on individual marine mammals or marine mammal populations, or their habitats.

LIST OF ACRONYMS

| | |
|------------------|--|
| ~ | approximately |
| ACP | Arctic Coastal Plain |
| ADCP™ | Acoustic Doppler Current Profiler |
| ADFG | Alaska Department of Fish and Game |
| AEWC | Alaska Eskimo Whaling Commission |
| BASC | Barrow Arctic Science Consortium |
| BLM | Bureau of Land Management |
| CCGS | Canadian Coast Guard Ship |
| CI | Confidence Interval |
| CITES | Convention on International Trade in Endangered Species |
| CPA | Closest Point of Approach |
| CPUE | Catch per Unit Effort |
| dB re 1 μ Pa | decibels in relation to a reference pressure of 1 micropascal |
| DFO | Department of Fisheries and Oceans (Environment Canada) |
| EA | Environmental Assessment |
| EEZ | Exclusive Economic Zone |
| EFH | Essential Fish Habitat |
| ESA | (U.S.) Endangered Species Act |
| $f(0)$ | sighting probability density at zero perpendicular distance from survey track line |
| FMP | Fishery Management Plan |
| ft | feet |
| $g(0)$ | probability of seeing a group located directly on the survey trackline |
| G. gun | variant of an airgun, manufactured by Sodera, a French company owned by Sercel |
| GSC | Geological Survey of Canada (Natural Resources Canada) |
| h | hour |
| ICES | International Council for the Exploration of the Sea |
| IHA | Incidental Harassment Authorization (under MMPA) |
| in | inch |
| IUCN | International Union for the Conservation of Nature and Natural Resources |
| IWC | International Whaling Commission |
| kHz | kilohertz |
| kts | knots |
| kW | kilowatt |
| LT | Long ton = 1016 kg |
| L-DEO | Lamont-Doherty Earth Observatory of Columbia University |
| LME | Large Marine Ecosystem |
| m | meter |
| MCS | Multi-Channel Seismic |
| min | minute |
| PSO | Protected species observer |
| MMPA | (U.S.) Marine Mammal Protection Act |
| MMS | Minerals Management Service |
| ms | millisecond |
| MTTS | Masked Temporary Threshold Shift |

List of Acronyms

| | |
|---------|---|
| MW | Megawatt |
| n.mi. | nautical mile |
| NEPA | National Environmental Policy Act |
| NMFS | National Marine Fisheries Service |
| NMML | National Marine Mammal Laboratory |
| NOAA | National Oceanic and Atmospheric Administration |
| NPFMC | North Pacific Fisheries Management Council |
| NSB | North Slope Borough |
| NSB-DWM | North Slope Borough Department of Wildlife Management |
| NSF | National Science Foundation |
| OCS | Outer Continental Shelf |
| P.I. | Principal Investigator |
| pk | peak |
| psi | pounds per square inch |
| PTS | Permanent Threshold Shift |
| rms | root-mean-square |
| s | second |
| SE | Southeast |
| SEL | sound energy level |
| SIS | sea ice seismometer |
| SPL | sound pressure level |
| T | ton = 907.18 kg |
| TTS | Temporary Threshold Shift |
| U.K. | United Kingdom |
| UNEP | United Nations Environment Program |
| U.S. | United States of America |
| USCG | United States Coast Guard |
| USDI | United States Department of the Interior |
| USFWS | U.S. Fish and Wildlife Service |
| USN | U.S. Navy |
| USGS | United States Geological Survey |
| WCMC | World Conservation Monitoring Centre |

I. PURPOSE AND NEED

The U.S. Geological Survey (USGS) plans to conduct a seismic survey in the Arctic Ocean north of Alaska in cooperation with the Geological Survey of Canada (GSC). The geophysical survey will be conducted by the Canadian Coast Guard icebreaker, *Louis S. St. Laurent*, which will deploy a three-airgun array. The United States Coast Guard (USCG) Cutter *Healy*, a USCG icebreaker, will rendezvous with the *Louis S. St. Laurent* and accompany and break ice for the Canadian vessel if necessary as it conducts seismic operations. The *Healy* will further contribute by operating a multibeam echo sounder and 3.5 kHz Chirp sub-bottom profiler to map the sea floor. As currently scheduled, the survey will occur from ~6 August to 3 September 2010, though some variation is likely given the uncertainties in ice and other factors.

The purpose of this Environmental Assessment (EA) is to provide information needed to assess potential environmental impacts associated with use of a three-airgun array consisting of two 520-in³ Sercel G-airguns and one 150-in³ Sercel G. gun during the proposed cruise. The EA addresses potential impacts of the proposed seismic survey from the *Louis S. St. Laurent* on marine mammals including cetaceans and pinnipeds which are under the jurisdiction of the National Marine Fisheries Service (NMFS). Much of the information presented in the EA for cetacean and pinniped species is also included in USGS's permit application to NMFS. In addition the EA discusses other species not under NMFS jurisdiction including polar bear (*Ursus maritimus*), Pacific walrus (*Odobenus rosmarus*) and threatened eiders which are under the jurisdiction of the U.S. Fish and Wildlife Service (USFWS), and fisheries and subsistence harvesting in the Arctic Ocean.

The purpose of the proposed project is to survey potential areas of the "extended continental shelf" to which either Canada or the United States may legitimately lay claim. The United Nations Convention on the Law of the Sea (UNCLOS) established criteria within Article 76 to determine area beyond the 200 n.mi. (370 km) limit that could be defined as "extended continental shelf". Data collected that meet the criteria could extend a nation's jurisdiction from 200 n.mi. to as much as 350 n.mi. (648 km) or 100 n.mi. (185 km) beyond the 2500-m isobath, whichever is greater.

Several species of cetaceans and pinnipeds inhabit parts of the Arctic Ocean where this cruise will occur. A few species listed as endangered under the U.S. Endangered Species Act ESA may occur in portions of the survey area, most notably the bowhead whale (*Balaena mysticetus*), and (although very unlikely) the fin whale (*Balaenoptera physalus*). Through informal consultation with the Office of Protected Resources with the National Oceanic and Atmospheric Administration (NOAA), USGS proposes that no ESA-listed marine species: bowhead, fin, humpback or sperm whale will be adversely affected by this project during the survey or transit to the survey area from Dutch Harbor. The polar bear was recently listed as a "Threatened" species under the ESA (USFWS 2008). Other species of concern (birds) that might occur in the area close to Barrow are the spectacled (*Somateria fischeri*) and Steller's (*Polysticta stelleri*) eiders that are also listed as "Threatened".

Incidental Harassment Authorizations (IHAs) issued by the NMFS are often required prior to the start of offshore activities. IHAs authorize the "taking" (as defined under the Marine Mammal Protection Act) of small numbers of marine mammals incidental to the planned activities. To be eligible for an IHA, the proposed "taking" (with mitigation measures in place) must not cause serious physical injury or death of marine mammals, and must have negligible impacts on the species and stocks of marine mammals within the project area. The proposed project must "take by harassment" no more than small numbers of those species, and (where relevant) must not have an unmitigable adverse impact on the availability of the species or stocks for authorized subsistence uses. It is expected that all "takes" associated with the

I. Purpose and Need

proposed activities will be Level B takes involving temporary behavioral changes and that no Level A “takes” involving injury to marine mammals will occur.

IHAs or Letters of Authorization (LOAs) are also issued by the USFWS for species under its jurisdiction including Pacific walrus and polar bear. IHAs and LOAs issued by the USFWS may have compliance requirements similar to those of NMFS.

Protection measures designed to mitigate the potential environmental impacts are described in this EA as an integral part of the planned activities. With the mitigation measures in place, any impacts on marine mammals and other species of concern are expected to be limited to short-term, localized changes in behavior of small numbers of animals. No long-term or significant effects are expected on individual marine mammals or populations, on the subsistence harvest of marine mammals, on marine mammal habitat, or on the individuals and populations of other species.

II. ALTERNATIVES INCLUDING PROPOSED ACTION

Three alternatives are addressed: (1) the proposed seismic survey and issuance of an associated IHA, (2) a corresponding seismic survey program at an alternative time, along with issuance of an associated IHA, (3) the no-action alternative, with no IHA and no seismic survey.

Proposed Action

The project objectives, context, activities, and mitigation measures for the proposed project planned by USGS are described in the following subsections.

(1) Project Objectives and Context

The objective of the proposed study is to survey potential areas of the “extended continental shelf” to which the United States may legitimately have sovereign rights under Article 76 the United Nations Convention on the Law of the Sea (UNCLOS).

(2) Proposed Activities

(a) Location of the Activities

The survey area will be bounded approximately by 145° to 158° W longitude and 71° to 84° N latitude in water depths ranging from ~2000-4000 m (6562–13,123 ft; Fig. 1). Ice conditions are expected to range from open water to 10/10 (i.e. 100%) ice cover. The *Louis S. St. Laurent* will join accompanying vessel USCG cutter *Healy* in or near the survey area around 6 August to begin the survey.

(b) Description of the Activities

Two vessels will operate during the proposed geophysical survey although authorization for incidental takes of marine mammals is requested and is relevant only for activities by the *Louis S. St. Laurent* which will operate an airgun array. Another low-energy source, a Knudsen 320BR “Chirp” pulse echo sounder, will be operated at 12 kHz on the *Louis S. St. Laurent*. In addition, the *Louis S. St. Laurent* will tow a 3 - 5 kHz sub-bottom profiler while in open water when not working with the *Healy*. The *Healy* will use a multibeam echo sounder, (Kongsberg EM122), a sub-bottom profiler (Knudsen 3.5 kHz Chirp) and a “piloting” echo sounder (ODEC 1500) for marine safety that the bridge uses to monitor seafloor depths continuously when underway and during the seismic profiling. Acoustic Doppler current profilers (75-kHz and 150-kHz) may also be used on the *Healy*.

The *Louis S. St. Laurent* will access the survey area from Canada and rendezvous with the *Healy* vessel which will access the survey area from Dutch Harbor, AK. The *Louis S. St. Laurent* will deploy a relatively small airgun array comprised of three G-guns with a total volume of 1150 in³ and a multichannel hydrophone streamer ~300 m (984 ft) in length. In typical seismic operations, the *Louis S. St. Laurent* will follow the lead of the *Healy* which will operate ~1 to 2 n.mi. (1.9 to 3.7 km) ahead of the *Louis S. St. Laurent*. Sonobuoys deployed from the *Louis S. St. Laurent* will enhance reception of reflected seismic pulses from the airgun array. A helicopter will not be used for sonobuoy deployment but may be used for personnel transfer between vessels and ice reconnaissance. In ice conditions where seismic gear cannot be safely towed, the *Louis S. St. Laurent* may escort *Healy* to optimize bathymetric data collection. The survey will consist of eight transect lines that extend from ~108 km (~58 n.mi.) offshore of the Alaska coast to and beyond the 200 n.mi (370 km) limit of U.S. waters (Fig. 1). An additional 997 km (538 n.mi.) of seismic lines of interest to the U.S. will be collected in international waters beyond 200 n.mi. (370 km). As much as 1000 km (540 n.mi.) of multibeam bathymetric data will also be collected when seismic data are not being collected. These data will be collected from early August to early September (Table 1). Much of the proposed survey area will be >200 n.mi. (370 km)

II. Alternatives Including Proposed Action

offshore in water depths of ~2000 to 4000 m (6562–13,123 ft; Fig. 1). After completion of the survey the *Louis S-St. Laurent* will return to port in Canada, and the *Healy* will return to Barrow.




FIGURE 1. Proposed location of the USGS August–September 2010 geophysical survey area for priority lines of the U.S. This map does not show specific Canadian lines in international waters or in the Canadian exclusive economic zone.

TABLE 1. Proposed U.S. priority tracklines for USGS/ GSC 2010 Extended Continental Shelf Survey in the northern Beaufort Sea and Arctic Ocean.

| Location | End Point 1 | End Point 2 | km | Time (h) @ | |
|-------------------------------|----------------------|----------------------|------|------------|------------|
| | | | | n.mi. | 4 n.mi./hr |
| NS in central EEZ (south) | 71.22° N ; 145.17° W | 72.27° N ; 145.41° W | 118 | 64 | 16 |
| NS in central EEZ (north) | 72.27° N ; 145.41° W | 73.92° N ; 145.30° W | 183 | 100 | 25 |
| Central-western EEZ connector | 73.92° N ; 145.30° W | 71.84° N ; 151.82° W | 317 | 171 | 43 |
| NS in western EEZ | 71.84° N ; 151.82° W | 74.32° N ; 150.30° W | 281 | 152 | 39 |
| South Northwind Ridge | 74.32° N ; 150.30° W | 74.96° N ; 158.01° W | 239 | 129 | 32 |
| Northwind Ridge connector | 74.96° N ; 158.01° W | 76.30° N ; 155.88° W | 161 | 87 | 22 |
| Mid-Northwind Ridge | 76.30° N ; 155.88° W | 75.41° N ; 146.50° W | 274 | 148 | 37 |
| Northwind Ridge connector | 75.41° N ; 146.50° W | 76.57° N ; 146.82° W | 129 | 70 | 17 |
| Mid-Northwind Ridge | 76.57° N ; 146.82° W | 76.49° N ; 150.73° W | 102 | 55 | 14 |
| Totals | | | 1804 | 976 | 245 |

(c) Schedule

The proposed geophysical survey will be conducted for ~30 days from early August to early September 2010. The *Healy* will depart from Dutch Harbor in early August and the *Louis S. St. Laurent* will depart from Kugluktuk, Nunavut, Canada on approximately the same date. The two vessels will rendezvous to begin surveying in U.S. waters where survey activities will proceed from ~6-12 August. The two vessels will then move to international waters where survey activities will be completed by ~3 September. After completing the survey the *Louis S. St. Laurent* will return to Canadian waters and the *Healy* will return to Barrow.

(d) Description of Vessels Proposed for the 2010 Geophysical Survey***Louis S. St. Laurent***

The Canadian Coast Guard Ship (CCGS) *Louis S. St. Laurent* (Fig. 2) was built in 1969 by Canadian Vickers Ltd. in Montreal, Quebec, and underwent an extensive modernization in Halifax, Nova Scotia between 1988-1993.

The *Louis S. St. Laurent* is based at CCG Base Newfoundland in St. John's, Newfoundland. Current vessel activities involve summer voyages to the Canadian Arctic for sealifts to various coastal communities and scientific expeditions. A description of the *Louis S. St. Laurent* with vessel specifications is presented below and is available online at:

<http://www.ccg-gcc.gc.ca/eng/Fleet/Vessels?id=1111&info=5&subinfo>

Louis S. St. Laurent carries three protected species observers aboard following Department of Fisheries and Oceans (DFO) rules and regulations associated with Canadian permits for seismic work.



Figure 2. Photo of the Canadian Coast Guard vessel Louis S. St. Laurent available online at: **CCGS Louis S. St. Laurent Ship Characteristics**

| | | | |
|--------------------------|------------------------|------------------------------|---------------------|
| Length: | 119.8 m | Breadth: | 24.38 m |
| Draft: | 9.91 m | Freeboard: | 6.4 m |
| Hold 1: | 300 m ³ | Hatch Size 1 (l x w): | 3.5 m X 3 m |
| Hold 2: | 36 m ³ | Hatch Size 2 (l x w): | 3.5 m X 3 m |
| Main Deck Area: | 320 m ² | Boat Deck Area: | 216 m ² |
| Forcastle: | N/A | After Deck Area: | 120 m ² |
| Gross Tonnage: | 11345 grt | Net Tonnage: | 3403 nrt |
| Cruising Speed: | 16 kts | Max. Speed: | 20 kts |
| Cruising Range: | 23000 n.mi. | Endurance: | 205 days |
| Fuel Consumption: | 24 m ³ /day | Fuel Capacity: | 4800 m ³ |
| Fresh Water: | 200 m ³ | | |

| | | | |
|---------------------|-----------------------|-----|---------|
| Propulsion: | Diesel electric AC/DC | | |
| Description: | (5x) Krupp | Mak | 16M453C |
| | (3x) GE DC Motor | | |
| Power: | 20142 Kw | | |
| Propellers: | 3 - fixed pitch | | |

| | |
|------------------------|-------------------------------|
| Generators: | (2x) Krupp Mak 6M282 @ 1100kw |
| Emergency Gen.: | (1x) Caterpillar 3408 BDI |
| Bow: | Yes |
| Stern: | No |
| UPS: | No |

| | | | |
|--------------------------|--------------------|-----------------------|--------------------|
| Flight Deck Area: | 360 m ² | Hangar Area: | 132 m ² |
| Hangar Gear: | Yes | Fuel Capacity: | 40 m ³ |

Healy

The *Healy* (Fig. 3) is a USCG icebreaker, capable of traveling at 37 km/h or 20 knots (kts) through 1.4 m (4.6 ft) of ice and can operate at temperatures as low as -46°C (-50°F). A “Central Power Plant”, four Sultzer 12Z AU40S diesel generators, provides electric power for propulsion and ship’s services through a 60 Hz, 3-phase common bus distribution system. Propulsion power is provided by two electric AC Synchronous, 11.2 MW drive motors, fed from the common bus through a Cycloconverter system, that turns two fixed-pitch, four-bladed propellers. The *Healy* can accommodate a crew of 138 including space for ~50 scientists and is equipped with various types of echo sounding equipment.

The *Healy* is designed to conduct a wide range of research activities, with more than 4,200 square feet of scientific laboratory space, numerous electronic sensor systems, and oceanographic winches. The science community provided input on lab lay-outs and science capabilities during design and construction of the ship. The *Healy* is also a capable platform for supporting other potential missions in the polar regions, including logistics, search and rescue, ship escort, environmental protection, and enforcement of laws and treaties. The *Healy* will also serve as the platform from which vessel-based protected species observers will watch for marine mammals before and during airgun operations.



Figure 3. The U.S. Coast Guard cutter *Healy* is United States' newest and most technologically advanced polar icebreaker. A description with vessel specifications for the *Healy* is available online at: <http://www.uscg.mil/pacarea/cgcHealy/default.asp>

CGC *Healy* Ship Characteristics

| | |
|-------------------------|---------------------------------------|
| Length, Overall | 420'0" (128 meters) |
| Beam, Maximum | 82'0" (25 meters) |
| Draft, Full Load | 29'3" (8.9 meters) |
| Displacement, Full Load | 16,000 LT |
| Propulsion | Diesel Electric, AC/AC Cycloconvertor |
| Generating Plant | 4 Sultzer 12Z AU40S |
| Drive Motors | 2 AC Synchronous, 11.2 MW |
| Shaft Horsepower | 30,000 Max HP |
| Propellers | 2 Fixed Pitch, 4 Bladed |

| | |
|------------------------|---|
| Auxiliary Generator | EMD 16-645F7B, 2400kW |
| Fuel Capacity | 1,220,915 GAL (4,621,000 liters) |
| Cruising Speed | 12 knots @ 105 RPM |
| Max Speed | 17 knots @ 147 RPM |
| Icebreaking Capability | 4.5ft @ 3 knots (continuous) 8 ft (2.44 m) Backing and Ramming |
| Science Labs | Main, Bio-Chemical, Electronics, Meteorological, Photography |
| Accommodations | 19 Officer, 12 CPO, 54 Enlisted, 35 Scientists, 15 Surge, 2 Visitors |

(e) Airgun Description

The seismic source for the proposed geophysical survey will be comprised of three Sercel G-guns with a total volume of 1150 in³. The three-gun array will be comprised of two 500-in³ and one 150-in³ G-guns in a triangular configuration (Fig. 4). The single 150-in³ G-gun will be used if a power down is necessary for mitigation. The G-gun array will be towed behind the *Louis S. St. Laurent* at a depth of ~11 m (Fig. 5) along predetermined tracks in water depths ranging from ~2000-4000 m (6562–13,123 ft). One streamer ~300 m (984 ft) in length will be towed behind the airgun array. The distance from the source to the end of the multichannel hydrophone will be ~232 m (761 ft) which will be followed by anti-vibration and stretch sections.

A square wave trigger signal will be supplied to the firing system hardware by a FEI-Zyfer GPStarplus Clock model 565, based on GPS time (typically at ~14 to 20 sec intervals depending on water depths). Vessel speed will range from ~5.6 to 8.3 km/hr (3 to 4.5 kt) resulting in a shot interval ranging from ~39 to 56 m (128 to 184 ft). G-gun firing and synchronization will be controlled by a RealTime Systems LongShot fire controller, which will send a voltage to the gun solenoid to trigger firing with ~54.8 ms delay between trigger and fire point.

Pressurized air for the pneumatic G-guns will be supplied by two Hurricane compressors, model 6T-276-44SB/2500. These are air cooled, containerized compressor systems. Each compressor will be powered by a C13 Caterpillar engine which turns a rotary screw first stage compressor and a three stage piston compressor capable of developing a total air volume of 600 SCFM @ 2500 PSI. The seismic system will be operated at 1950 PSI. One compressor is used during typical operations with the other reserved for backup should the first compressor require maintenance.

Seismic acquisition will require a watchkeeper in the seismic lab, an airgun technician on watch or on call, and another in the compressor container. The seismic lab watchkeeper is responsible for data acquisition/recording, watching over-the-side equipment, gun firing and log keeping. A remote screen will permit monitoring of compressor pressures and alerts, as well as communication with the compressor watchkeeper. The compressor watchkeeper will be required to monitor the compressor for any emergency shut down and provide general maintenance that might be required during operations.

II. Alternatives Including Proposed Action




Figure 4. Configuration of three Sercel G-airguns during seismic operation from the *Louis S. St. Laurent*, 2010.




Figure 5. Geometric arrangement of the seismic source and streamer (source: Mosher et al. 2009).

(f) Low-energy sources

Along with the airgun operations, additional acoustic systems to be operated during the cruise will include a 12-kHz Chirp echo sounder and a 3–5 kHz sub-bottom profiler from the *Louis S. St. Laurent* and 12-kHz Kongsberg multibeam bathymetric echo sounder from the *Healy*. These sources will operate

throughout most of the cruise to map the bathymetry as necessary to meet the geophysical science objectives. During seismic operations, these sources will be deployed from the *Louis S. St. Laurent* and the *Healy* and will generally operate simultaneously with the airgun array deployed from the *Louis S. St. Laurent*. In 2008 and 2009 the *Louis S. St. Laurent* and the *Healy* surveyed together with a cooperative strategy similar to that proposed for 2010. The director of NOAA's Office of Ocean Exploration and Research deemed that the use of the *Healy's* multibeam echo sounder would not have significant impacts on marine mammals of a direct or cumulative nature. The U.S. portions of the projects were granted categorical exclusions from the need to prepare an Environmental Assessment (Appendices A and B). The Canadian Environmental Assessments for these projects are available on line at <http://www.ceaa.gc.ca/052/details-eng.cfm?pid=38185> (2008) and <http://www.ceaa.gc.ca/052/details-eng.cfm?pid=46518> (2009).

Chirp Echo Sounder (Knudsen 320BR)

Along with the airgun operations, an additional acoustic system to be operated on the *Louis S. St. Laurent* will include a 12-kHz Knudsen 320BR Chirp echo sounder. The Knudsen 320BR will provide information on depth and bottom profile. The Knudsen 320BR is a dual-frequency system with operating frequencies of 3.5 and 12 kHz, however, the unit will be functioning only at the higher 12 kHz frequency.

The calculated maximum source level (downward) of the Knudson 320BR is 215 dB re 1 μ Pa at 1 m. The pulse duration is typically 1.5 to 5 ms with a bandwidth of 3 kHz (FM sweep from 3 kHz to 6 kHz). The repetition rate is range dependent, but the maximum is a 1% duty cycle. Typical repetition rate is between 1/2 s (in shallow water) to 8 s in deep water.

A single 12 kHz transducer and one 3.5 kHz, low frequency (sub-bottom) transducer array, consisting of 16 elements in a 4 \times 4 array will be used for the Knudsen 320BR. The 12 kHz transducer (TC-12/34) emits a conical beam with a width of 30°.

Towed 3–5 kHz Sub-bottom Profiler

The 3–5 kHz chirp sub-bottom profiler will be towed by and operated from the *Louis S. St. Laurent* in open water when the *Louis S. St. Laurent* is not working in tandem with the *Healy*. The profiler provides information about sedimentary features and bottom topography. The chirp system has a maximum 7.2 kW transmit capacity into the towed array. The energy from the towed unit is directed downward by an array of eight transducers in a conical beamwidth of 80°. The interval between pulses will be no less than one pulse per second. Sub-bottom profilers of that frequency can produce sound levels of 200-230 dB re 1 μ Pa at 1 m (Richardson et al. 1995).

Multibeam Echo Sounder (Kongsberg EM122)

The Kongsberg EM 122 MBES operates at 10.5–13 (usually 12) kHz and is hull-mounted on the *Healy*. The transmitting beamwidth is 1° fore–aft and 150° athwartship. The maximum source level is 242 dB re 1 μ Pa \cdot m_{rms}. Each “ping” consists of eight (in water >1000 m deep) or four (in water <1000 m) successive fan-shaped transmissions, each ensonifying a sector that extends 1° fore–aft. Continuous-wave (CW) pulses increase in length from 2 to 15 ms in water depths up to 2600 m, and FM chirp pulses up to 100 ms long are used in water >2600 m. The successive transmissions span an overall cross-track angular extent of about 150°, with 2-ms gaps between the pulses for successive sectors.

Hydrographic Sub-bottom Profiler (Knudsen 320BR)

The Knudsen 320BR will provide information on sedimentary layering to depths between 20 and 70 m depending on bottom type and slope. The Knudsen 320BR is a dual-frequency system with operating frequencies of 3.5 and 12 kHz; only the low frequency will be used during this survey. At 3.5 kHz, the maximum output power into the transducer array, as wired on the *Healy* (where the array

impedance is approximately 125 ohms), is ~6000 watts (electrical), which results in a maximum source level of 221 dB re 1 μ Pa at 1 m downward. Pulse lengths range from 1.5 to 24 ms with a bandwidth of 3 kHz (FM sweep from 3 kHz to 6 kHz). The repetition rate is range dependent, but the maximum is a 1% duty cycle. Typical repetition rate is between 0.5 s (in shallow water) to 8 s in deep water.

The 3.5-kHz transducer array on the *Healy*, consisting of 16 (TR109) elements in a 4 \times 4 array, will be used for the Knudsen 320BR. At 3.5-kHz the sub-bottom profiler emits a downward conical beam with a width of approximately 26°.

Piloting Echo Sounder

The piloting echo sounder on the *Healy* is an Ocean Data Equipment Corporation (ODEC) Bathy-1500 which will provide information on water depth below the vessel. The ODEC system has a maximum 2-kW transmit capacity into the transducer and has two operating modes, single or interleaved dual frequency, with available frequencies of 12, 24, 33, 40, 100, and 200 kHz.

Acoustic Doppler Current Profiler (R D Instruments Ocean Surveyor 150 kHz)

The 150-kHz acoustic Doppler current profiler (ADCP™) has a minimum ping rate of 0.65 ms. There are four beam sectors and each beamwidth is 3°. The pointing angle for each beam is 30° from vertical with one each to port, starboard, forward, and aft. The four beams do not overlap. The 150-kHz ADCP's maximum depth range is 300 m.

Acoustic Doppler Current Profiler (R D Instruments Ocean Surveyor 75)

The Ocean Surveyor 75 is an ADCP operating at a frequency of 75 kHz, producing a ping every 1.4 s. The system is a four-beam phased array with a beam angle of 30°. Each beam has a width of 4° and there is no overlap. Maximum output power is 1 kW with a maximum depth range of 700 m.

(3) Mitigation Measures

Several species of marine mammals are known to occur in the proposed survey area. To minimize the likelihood that impacts will occur to marine mammal species and stocks, airgun operations will be conducted in accordance with all applicable U.S. Federal regulations and IHA requirements. USGS will coordinate all activities with the relevant U.S. Federal agencies, particularly the NMFS and the USFWS. The *Louis S. St. Laurent* will operate under provisions of a Canadian authorization from DFO based on that country's own environmental assessment and under the requirements and mitigation measures specified in an IHA which will be issued by NMFS and any stipulations in an IHA or LoA that may be issued by the USFWS for polar bears. Standard Canadian practices with respect to the mitigation of seismic sound (DFO 2008) will be followed when the *Louis S. St. Laurent*, a Canadian Coast Guard vessel, surveys international waters. Therefore, mitigation measures during the seismic survey will differ slightly according to the vessels' location. The following subsections provide more detailed information about the mitigation measures that are an integral part of the planned activities.

(a) Marine Mammal Monitoring

Vessel-based protected species observers (PSOs) will be stationed on both the *Louis S. St. Laurent* and the *Healy* during the proposed survey. The vessels will typically work together in tandem while making way through heavy ice with the *Healy* in lead breaking ice and collecting multibeam data. The *Louis S. St. Laurent* will follow collecting seismic reflection and refraction data. In light ice conditions, the vessels will separate to maximize data collection. "Real-time" communication between the two vessels regarding marine mammal detections will be available through VHF radio.

Three trained PSOs, knowledgeable about marine mammals of the Arctic, will be recruited through a Canadian Hunters and Trappers Committee to work on the *Louis S. St. Laurent*. These observers will

board the *Louis S. St. Laurent* in Kugluktuk, Nunavut, Canada. Three experienced PSOs and one Alaska Native observer will be aboard the *Healy* at the outset of the project. Before survey operations begin in U.S. waters, two of the PSOs on the *Healy* will transfer to the *Louis S. St. Laurent*. Thus, during operations in U.S. EEZ waters, a complement of five observers will work on the source vessel, the *Louis S. St. Laurent*, and two will be stationed on the *Healy*. When not surveying in U.S. waters, the distribution of PSOs will return to three on the *Louis S. St. Laurent* and four on the *Healy*.

In U.S. Waters PSOs on the *Louis S. St. Laurent* will monitor for marine mammals during all daylight airgun operations. At least one observer, and when practical two observers, will monitor for marine mammals from the *Louis S. St. Laurent* during ongoing daytime operations and nighttime start ups (when darkness is encountered). Use of two simultaneous observers will increase the proportion of animals present near the source vessel that are detected. PSOs will normally be on duty in shifts of no longer than 4 hours duration although more than one 4-hr shift may be worked per day with a maximum of 12 hr of daily watch time. Airgun operations will be shut down when marine mammals are observed within, or about to enter, designated safety radii (see below) where there may be a possibility of significant effects on hearing or other physical effects. PSOs on both the source vessel and the *Healy* will also watch for marine mammals within or near the safety radii for at least 30 min prior to the planned start of airgun operations after an extended shut down of the airgun array. When feasible, observations will also be made during periods without seismic operations (e.g., during transits). Environmental conditions will be recorded every half hour during PSO watch.

The PSOs aboard the *Healy* will also watch for marine mammals during daylight seismic activities conducted in both U.S. and international waters. They will maximize their time on watch but will not watch continuously because they will not have required mitigation duties and there will be only two PSOs aboard the *Healy*. The *Healy* PSOs will report sightings to the PSOs on the *Louis S. St. Laurent* to alert them of possible needs for mitigation.

The *Louis S. St. Laurent* crew will be instructed to assist in detecting marine mammals and implementing required mitigation (if practical). The crew will be given instruction on mitigation requirements and procedures for implementation of mitigation prior to the start of the seismic survey. Members of the *Healy* crew will be trained to monitor for marine mammals and asked to contact the *Healy* observers for sightings that occur while the PSOs are off-watch.

During seismic operations in international waters, PSOs aboard the *Louis S. St. Laurent* will conduct 8-hr watches. This schedule easily accommodates 24-hr/day monitoring by three PSOs which will be necessary during most of the survey when daylight will be continuous. *Healy* PSOs will limit watches to 4 hours in both U.S. and international waters.

The *Louis S. St. Laurent* and *Healy* are suitable platforms for marine mammal observations. When stationed on the flying bridge, eye level will be ~15.4 m (51 ft) above sea level on the *Louis S. St. Laurent* and ~24 m (78.7 ft) above sea level on the *Healy*. On both vessels the observer will have an unobstructed view around the entire vessel from the flying bridge. If surveying from the bridge of the *Louis S. St. Laurent* or the *Healy*, the observer's eye level will be 12.1 m (~40 ft) above sea level or 21.2 m (69 ft) above sea level, respectively. The PSO(s) will scan the area around the vessel systematically with laser range finding binoculars and with the unaided eye.

The survey will be conducted at high latitudes and continuous daylight will persist through much of the proposed survey area from mid- to late August. Day length will decrease to ~18 hr in the northern portion of the survey area by about early September. Laser range-finding binoculars will be available to assist with distance estimation; this equipment is useful in training observers to estimate distances visually, but is generally not useful in measuring distances to animals directly.

When mammals are detected within or about to enter the designated safety radius, the airgun(s) will be powered down or shut down immediately. The distinction between power downs and shut downs is described in section II(3)(c) below. Channels of communication between the PSOs and the airgun technicians will be established to assure prompt implementation of shutdowns when necessary as has been done in other recent seismic survey operations in the Arctic (e.g., Haley 2006). During power downs and shutdowns, PSOs will continue to maintain watch to determine when the animal(s) are outside the safety radius. Airgun operations will not resume until the animal is outside the safety radius. The animal will be considered to have cleared the safety radius if it is visually observed to have left the safety radius. Alternatively, in U.S. waters the safety zone will be considered clear if the animal has not been seen within the radius for 15 min for small odontocetes and pinnipeds or 30 min for mysticetes. Within international waters, the PSOs will apply a 30 minute period for all species.

All observations and airgun power downs or shut downs will be recorded in a standardized format. Data will be entered into a custom database using a notebook computer. The accuracy of the data entry will be verified by manual checking of the database. This will allow initial summaries of data to be prepared during and shortly after the field program, and will facilitate transfer of the data to statistical, graphical, or other programs for further processing and archiving.

Results from the vessel-based observations will provide:

1. The basis for real-time mitigation (airgun power down or shut down).
2. Information needed to estimate the number of marine mammals potentially taken by harassment, which must be reported to NMFS.
3. Data on the occurrence, distribution, and activities of marine mammals in the area where the geophysical survey is conducted.
4. Information to compare the distance and distribution of marine mammals relative to the source vessel at times with and without seismic activity.
5. Data on the behavior and movement patterns of marine mammals seen at times with and without seismic activity.

There is no plan to implement an acoustic monitoring program during the proposed seismic survey. Typically, marine mammal acoustic monitoring is conducted by listening to transmissions from a streamer or sonobuoy. Listening for marine mammal calls with a hydrophone streamer while surveying on an icebreaker would be unproductive because of masking caused by the high levels of ship noise and (when in the ice) icebreaking. Towing an additional streamer exclusively for acoustic monitoring presents the same problems and it is not practical to tow non-essential equipment through heavy ice.

During a *Healy* seismic survey across the Arctic Ocean in 2005, sonobuoys were periodically deployed by the geophysicists to relay seismic data. The use of sonobuoys for the survey provided an opportunity to monitor for marine mammal calls, and transmissions from sonobuoys were monitored by marine mammal observers (Haley and Ireland 2006). No marine mammal vocalizations however, were detected during a total 98 h (739 km) of monitoring. The use of sonobuoys to monitor marine mammal vocalizations would be particularly ineffective during surveys with lengthy transect lines as proposed for the current survey rather than a series of shorter, parallel lines that are often used during industry 3-D seismic surveys. Given these considerations, acoustic monitoring for marine mammals is not planned during the proposed survey.

Reports will be submitted to NMFS and USFWS within 90 days after completion of the cruise. The reports will describe the operations that were conducted in U.S. waters and the marine mammals that were detected near the operations. The reports will provide full documentation of methods, results, and

interpretation of all monitoring data. The 90-day report will summarize the dates and locations of seismic operations, and all marine mammal sightings (dates, times, locations, activities, associated seismic survey activities). The report will also include estimates of the amount and nature of potential “take” of marine mammals by harassment or in other ways. Analysis and reporting conventions will be consistent with those for other recent cruises in the Arctic and will add to the current database on the distribution and abundance of marine mammal species in the Arctic.

(b) Proposed Safety Radii

Under current NMFS guidelines (e.g., NMFS 2009b), “safety radii” for marine mammals around airgun arrays are customarily defined as the distances within which received pulsed sound levels are ≥ 180 dB (rms) for cetaceans and ≥ 190 dB re 1 μPa for pinnipeds. Those safety radii are based on an assumption that seismic pulses at lower received levels will not injure these mammals or impair their hearing abilities, but that higher received levels *might* have some such effects. Therefore, the safety zone applied in U.S. waters will be the modeled radii for the 180-dB rms and 190-dB rms received sound levels for cetaceans and pinnipeds, respectively.

Standard Canadian practices with respect to the mitigation of seismic sound (DFO 2008) will be followed when the *Louis S. St. Laurent* surveys international and Canadian waters. PSOs will establish a standard safety zone of 500 m or greater around the sound source that will be applied to all marine mammal species in international and Canadian waters.

The rms (root mean square) received sound pressure levels (SPLs) that are used as impact criteria for marine mammals are not directly comparable to the 0-to-peak or peak-to-peak values used to characterize source levels of airguns. The measurement units used to describe the airgun source, 0-to-peak or peak-to-peak dB, are always higher than the SPL rms dB units referred to in much of the biological literature. A measured received level of 160 dB rms in the far field would typically correspond to a peak measurement of about 170 to 172 dB, and to a peak-to-peak measurement of about 176 to 178 dB, *as measured for the same pulse received at the same location* (Greene 1997; McCauley et al. 1998, 2000). The precise difference between rms and 0-to-peak or peak-to-peak values for a given pulse depends on the frequency content and duration of the pulse, among other factors. However, the rms level is always lower than the 0-to-peak or peak-to-peak level for an airgun-type source. Thus the received sound levels in rms equivalents used by NMFS for potential injury (180 dB rms) or disturbance (160 dB rms) will occur at distances closer to the source than those reported for 0-to-peak values. The 0 to peak values described in the following paragraph have been adjusted by 10 dB to estimate the rms values.

A seismic calibration experiment was conducted for the 1150-in³ G-gun array during the 2009 program in the Arctic Ocean (Mosher et al. 2009; Roth and Schmidt 2010). A transmission loss model was then constructed assuming spherical (20LogR) spreading and using the source level estimate (235 dB re 1 μPa 0-peak; 225 dB re 1 μPa rms) from the measurements. The use of 20LogR spreading fit the data well out to ~1 km where variability in measured values increased (Fig. 6). Additionally, the Gundalf@ modeling package was used to model the airgun array and estimated a source level output of 236.7 dB 0-peak (226.7 dB rms). Using this slightly stronger source level estimate and 20LogR spreading the 180 and 190 dB rms radii were estimated to be 216 m and 68 m, respectively. As a conservative measure for the proposed safety radii, the sound-level radii indicated by the empirical data and source models have been increased to 500 m for the 180-dB isopleth and to 100 m for the 190-dB isopleth (Table 2). The actual 160 dB rms disturbance zone will be approximately 2157 m (1.34 mi), but USGS proposes using 2500 m (1.55 mi) as the radius for the 160 dB disturbance zone for the take estimates for this survey.

USGS has informed USFWS about the proposed survey and the potential incidental takes of walrus and polar bears in conjunction with the seismic activities proposed for this project. USFWS

granted permission for the survey with the agreement that USGS observe a ≥ 190 dB safety radius for polar bears and a ≥ 180 dB safety radius for walrus while conducting seismic operations within the U.S. EEZ, which is USFWS standard protocol in U.S. waters.




Figure 6. Measured peak sound pressure levels as a function of range for 1/3 and full octave bands. The blue line shows theoretical spherical spreading loss for a 235 dB marine source as a comparison (Roth and Schmidt 2010).

TABLE 2. Proposed sound level radii for the three-airgun array and mitigation airgun for the proposed USGS seismic survey.

| Seismic Source Volume | Estimated Distances for Received Levels (m) | | |
|--|---|------------|------------|
| | 190 dB rms | 180 dB rms | 160 dB rms |
| 150 in ³ mitigation gun | 30 | 75 | 750 |
| 1150 in ³ (three G-gun array) | 100 | 500 | 2500 |

(c) Mitigation during Operations

In addition to monitoring, mitigation measures that will be adopted within U.S. waters will include (1) speed or course alteration, provided that doing so will not compromise operational safety requirements, (2) power down or shut-down procedures, and (3) no start up of airgun operations unless the full safety zone is visible for at least 30 min. During foggy conditions or periods of darkness (which may be encountered in late August), the full safety radius may not be visible. In that case, the airguns could not start up after a full shut down until the entire safety radius is visible. If one or more airguns have been operational before nightfall, they can remain operational throughout the night, even though the entire safety radius may not be visible, in accordance with NMFS guidelines for mitigation.

Through informal consultation with the Office of Protected Resources, the USGS proposes that no ESA-listed marine species: bowhead, fin, or humpback whales will be adversely affected by this project. With that understanding, the USGS has included a precautionary measure for cetaceans for this survey. If a PSO observes a bowhead, fin, humpback or unidentified mysticete whale approaching or within the 160 dB received sound level radius (2.5 km) a power down will be implemented.

While in U.S., Canadian or international waters, PSOs will have the option of implementing the following mitigation procedures. The salient difference between mitigation in U.S. vs. international or Canadian waters is that a “power down” will not be an option if a marine mammal is sighted within or about to enter the 500 m or greater safety zone while following the Canadian statement of practice (DFO 2009b) outside of U.S. waters.

Speed or Course Alteration

If a marine mammal is detected outside the safety radius and, based on its position and relative motion, is likely to enter the safety radius, the vessel's speed and/or course may, when practical and safe, be changed in a manner that also minimizes the effect on the planned science objectives. The marine mammal activities and movements relative to the seismic vessel will be closely monitored to ensure that the marine mammal does not approach within the safety radius. If the mammal appears likely to enter the safety radius, further mitigative actions will be taken, i.e., either further course alterations or power down or shut down of the airgun(s).

Power-down Procedures

A power down involves decreasing the number of operating airguns resulting in a reduction of the radius of the safety zone. A power down may be implemented to reduce or eliminate the potential for marine mammal exposure to possibly harmful sound levels. A power down may also occur when the vessel is moving from one seismic line to another. During a power down, one airgun (or some other number of airguns less than the full airgun array) is operated. The continued operation of one airgun is intended to alert marine mammals to the presence of the seismic vessel in the area. In contrast, a shut down occurs when all airgun activity is suspended.

If a marine mammal is detected outside the safety radius but is likely to enter the safety radius, and if the vessel's speed and/or course cannot be changed to avoid having the mammal enter the safety radius, the airguns may (as an alternative to a complete shut down) be powered down before the mammal is within the safety radius. Likewise, if a mammal is already within the safety zone when first detected, the airguns will be powered down immediately if this is a reasonable alternative to a complete shut down. During a power down of the three G-gun array, only the 150-in³ G-gun will be operated. If a marine mammal is detected within or near the smaller safety radius around the 150-in³ G-gun, it will be shut down as well (see next subsection).

Following a power down, airgun activity will not resume until the marine mammal has cleared the safety zone. The animal will be considered to have cleared the safety zone if it

- is visually observed to have left the safety zone, or
- has not been seen within the zone for 15 min in the case of small odontocetes and pinnipeds, or
- has not been seen within the zone for 30 min in the case of mysticetes (large odontocetes do not occur within the study area).

A power down will not be available as a mitigation measure while the Canadian mitigation protocol with respect to seismic sound exposure levels is applied in international or Canadian waters (DFO 2009b). A safety zone of 500 m or greater will need to be maintained around the active array, even if the array has been powered down to a single airgun. Therefore, a complete shut down will be requested during seismic operations in international or Canadian waters if a marine mammal is observed within or approaching the 500 m or greater safety radius. However, a power down is an option that may be used for operational purposes, e.g., reducing sound during turns.

Shut-down Procedures

The operating airgun(s) will be shut down completely if a marine mammal approaches or enters the then-applicable safety radius and a power down is not practical. The operating airgun(s) will also be shut down completely if a marine mammal approaches or enters the estimated safety radius around the 150-in³ G-gun used during a power down.

Airgun activity will not resume until the marine mammal has cleared the safety radius. The same requirements described in the bullet points above following a power down will also apply for a shut down before airgun activity can resume. After a shut down in international or Canadian waters, however, all species of marine mammals will not be considered to have left the safety radius until the animal has not been seen within the zone for 30 minutes.

Ramp-up Procedures

A "ramp up" procedure will be followed when the airgun array begins operating after a specified-duration period without airgun operations. NMFS normally requires that the rate of ramp up be no more than 6 dB per 5 min period. The specified period depends on the speed of the source vessel and the size of the airgun array that is being used.

NMFS normally requires that, once ramp up commences, the rate of ramp up be no more than 6 dB per 5 min period. Ramp up will likely begin with a single airgun (the smallest airgun in the array). The precise ramp-up procedure has yet to be determined, but USGS intends to follow NMFS' guideline (or whatever guideline USFWS adopts) with a ramp up rate of no more than 6 dB per 5 min period when operating in U.S. waters. During the ramp-up, the safety zone for the full three G-gun array (or whatever smaller source might then be in use) will be maintained.

While operating within U.S. waters, if the complete 180 dB safety radius has not been visible for at least 30 min prior to the start of operations, ramp up will not commence unless at least one airgun has been operating during the interruption of seismic survey operations. This means that it will not be permissible to ramp up the airgun source from a complete shut down in thick fog or darkness (which may be encountered in late August) when the outer part of the 180 dB safety zone is not visible. If one airgun has operated during a power-down period, ramp up to full power will be permissible in poor visibility, on the assumption that marine mammals will be alerted to the approaching seismic vessel by the sounds from the single airgun and could move away if they choose. Ramp up of the airguns will not be initiated during the day if a marine mammal has been sighted within or near the applicable safety radii during the previous 15 min (for pinnipeds) or 30 min (for cetaceans), and has not been subsequently observed outside the applicable radius.

The Canadian statement of practices (DFO 2009b) will be observed during seismic activities in international or Canadian waters. The statement describes a ramp up as activating a single, preferably the smallest, airgun in the array and activating additional airguns gradually over a minimum period of 20 minutes. Ramp up will not be initiated in international waters unless no marine mammal species has been observed within the 500 m or greater safety zone for 30 minutes.

Alternative Action: Conduct Survey during Alternative Time Period

An alternative to issuing the IHA and conducting the survey during the period requested is to issue the IHA and conduct the survey during a different time period. The window of opportunity for an Arctic Ocean cruise however, is extremely narrow due to the dependence on ice conditions. Late summer is by far the most suitable time to conduct activities in the Arctic. The summer offers the least amount of pack ice and the most favorable weather conditions. Another consideration is the availability of the two icebreakers to work together. Delaying the cruise or conducting operations earlier than proposed could be impractical and could result in hazardous or unsafe operations due to ice and other weather related conditions.

A major scheduling consideration for the proposed seismic survey relates to the timing of the bowhead whale migration and associated subsistence hunt by Alaskan Natives. Spring whaling activities occur at Barrow from approximately early April to early June. Spring bowhead hunts do not occur at the other Beaufort Sea villages of Nuiqsut or Kaktovik. The proposed geophysical survey in August and September will not interfere with the spring bowhead hunt at Barrow.

Fall whaling activities begin at Kaktovik in late August or early September and the whaling season generally progresses on later dates at Nuiqsut and Barrow during the westward bowhead migration. The latest Beaufort Sea whaling occurs at Barrow from ~mid- September into October. Most of the proposed survey area is located >200 n.mi. (370 km) offshore, far to the north of traditional whaling areas along the Beaufort Sea coast. The geophysical survey activities will occur at a sufficient distance offshore that whaling activities at Kaktovik and Nuiqsut will not be affected. The *Healy* will return to the Barrow area in early September, generally before fall whaling at Barrow commences. The vessel will contact the whaling communities prior to approaching Barrow (or any other village) and coordinate their activities with those of whalers to eliminate any potential disturbance to ongoing whaling activities. Thus, the geophysical activities under the proposed schedule will have no impact on fall whaling by villagers along the Beaufort Sea coast.

The overall schedule for the *Louis S. St. Laurent* and the accompanying vessel has been established to accomplish this cruise and other objectives in a coordinated and optimized manner. The scientific personnel and specialized equipment to be deployed on the *Louis S. St. Laurent* and the accompanying

vessel are available for the planned period but not necessarily for other periods. Issuance of the IHA for a substantially different range of dates would require changes in scheduling of personnel and equipment which could result in cancellation of the 2010 cruise, given the probable inability to amend the schedules for all of the required project components.

Additionally, 2010 is the last year of a four-year program of data acquisition in the Canada Basin. Delaying the 2010 survey would result in a lost opportunity for the U.S. to meet data collection goals.

No Action Alternative

An alternative to conducting the proposed activities is the “No Action” alternative, i.e., the proposed geophysical survey will not be conducted. The “No Action” alternative would result in no disturbance to marine mammals attributable to the proposed activities, and no impacts of other types.

The proposed seismic survey would provide information valuable for determining the extent of area offshore of the coast of northern Alaska and beyond the 200 n.mi. (370 km) limit to which the United States may have legitimate, legal claims. The “No Action” alternative, through forcing cancellation of the planned seismic survey in the Arctic Ocean, would result in a loss of important scientific data and knowledge relevant to potential economic and strategic interests of the United States, and also seriously jeopardize the ability of the United States to substantiate the outer limits of its extended continental shelf in the Arctic.

III. AFFECTED ENVIRONMENT

Physical Environment

The Arctic Ocean is the smallest of the world’s oceans, covering 14,090,000 km². The Arctic region contains 12 of the world’s Large Marine Ecosystems (LME): West Greenland Shelf, East Greenland Shelf, Barents Sea, Norwegian Shelf, West Bering Sea, Chukchi Sea, Beaufort Sea, East Siberian Sea, Laptev Sea, Kara Sea, Hudson Bay, and Arctic Ocean (UN Atlas of the Oceans n.d.). Of these 12 LMEs, the proposed project will be active primarily within the Beaufort Sea and Arctic Ocean.

The Arctic Ocean LME lies between North America, Greenland and Asia beyond the Arctic Circle at a latitude of 66° N (UN Atlas of the Oceans n.d.). The oceanography and bathymetry of this region is complex. There are three main water layers in the Arctic Ocean: (1) relatively fresh, low salinity surface water (is this Pacific or Atlantic water?), (2) an intermediate layer that is composed of warmer, saltier Atlantic water, which enters north of Spitzbergen, and (3) cold, deep water which flows in across the submarine ridge between Spitzbergen and Greenland (Sverdrup et al. 1942; McLaughlin et al. 1996).

Surface water enters the Arctic Ocean mainly from the Pacific Ocean through the shallow Bering Strait and from the Atlantic Ocean through the eastern part of the Fram Strait. These source waters are modified by river runoff and meltwater in summer and by salt rejection during freezing in winter, resulting in a characteristic brackish surface layer (lower salinity) up to about 30–50 m (98–164 ft) in thickness. A smaller quantity of water is transported southward through the Barents and Kara seas and the Canadian Archipelago. Approximately 2% of the water entering the Arctic Ocean is fresh water, and precipitation in the region is ~10 times greater than loss by evaporation.

The core of the intermediate layer occurs at about 300 m and extends to a depth of about 400 m. Two water masses are evident within the bottom layer: (1) Eurasian Basin deep water, and (2) Canadian Basin deep water, separated by the Lomonosov Ridge (Woodgate et al. 2001). Warmer Atlantic water underlies the Arctic surface waters to a depth of about 900 m. As this water cools it becomes so dense that it slips below the surface layer as it enters the Arctic Basin. Cold bottom water extends beneath the Atlantic layer to the ocean floor.

Arctic surface waters are driven by wind and density differences and by a clockwise surface circulation pattern that reaches speeds of 15–40 cm per second. The deep boundary current in the Arctic Ocean appears to be characterized by weak mean flows and strong, isolated eddies (Aagaard 1989; Woodgate et al. 2001).

The Arctic is dominated by ice cover that opens significantly during summer only in the coastal seas to the north of Asia, Alaska, and northern Canada. Sea ice rarely forms in the open ocean below 60°N. Between 60°N and 75°N it is present seasonally. Above 75°N ice cover is present on a largely permanent basis. The Arctic has notable year-to-year variations in ice cover although an increasing trend in the retreat of the pack ice in recent years has been well documented (Stroeve et al. 2008).

When ice is present it suppresses wind stress and wind mixing and also reflects solar radiation, thereby lowering surface temperature and impeding evaporation. Wind and surface stresses keep the pack ice in constant motion, resulting in the formation of leads, polynyas, pressure ridges, shear zones, and other features.

The Beaufort Sea LME is a high-latitude marine region off the coast of northern Alaska and northwest Canada; it is dominated by an extreme arctic climate (UN Atlas of the Oceans n.d.). Most of the Beaufort Sea is ice-covered for the majority of the year, although there are major seasonal and annual variations. The Beaufort Gyral Stream forms a clockwise drift pattern. Leads can occur north of Barrow at any time of year, and in that area there are varying amounts of open water from late spring through autumn. During some years the southern edge of heavy pack ice can be 200 km or more off the coast of Barrow in August, but during other years the pack ice may extend south to the Barrow coast.

Biological Environment

The Arctic Ocean is classified as a low productivity ecosystem, a consequence of the extensive seasonal ice cover and extreme weather conditions. Arctic plankton show weak diurnal vertical migrations but pronounced seasonal ones. Arctic fauna is impoverished and consists mainly of organisms derived from the Atlantic Ocean. The biomass is low, often dominated by one of only a few species. Because of the extensive areas of sediments, arctic benthic fauna is mainly an infauna. Specialized endemic fish are not present in the Arctic. Marine mammals however, are relatively diverse.

The Beaufort Sea LME experiences highly variable seasonal productivity (UN Atlas of the Oceans n.d.). During winter there is limited light penetration due to low light conditions and the extent of sea-ice cover. Increasing daylight in the summer results in warmer temperatures, ice melt, and significantly higher productivity. The coastal region supports a wide diversity of organisms. The Beaufort coastal areas provide habitat for ducks, geese, swans, shorebirds, and marine birds. Many species of birds and fish rely on river deltas, estuaries, spits, lagoons, and islands in coastal waters for breeding habitat, food, shelter, and brood-rearing. Various waterbird and fish species depend on marine waters (mainly over the shallow waters of the continental shelf) for food and habitat during the summer.

Fish Resources

FishBase, a global information system on fishes available at fishbase.org, lists 102 marine fish species as being present in the Beaufort Sea (Appendix C). FishBase lists 123 species for the Arctic Ocean LME (Appendix C).

Fisheries

The majority of the fisheries in the Arctic Ocean and Beaufort Sea LMEs are of a subsistence nature and are conducted close to shore. There is no fishing activity along the planned geophysical survey route.

Twenty-one species of fish are harvested commercially in the Beaufort Sea, including arctic cisco (*Coregonus autumnalis*), broad whitefish (*C. nasus*), least cisco (*C. sardinella*), and Dolly Varden char (*Salvelinus malma*). Several species (including the Dolly Varden char) are anadromous and move seasonally between fresh water and underground springs in winter and salt water in summer. Figs. 7 and 8 present fisheries landings in the Beaufort Sea and Arctic Ocean, respectively.

These fish, however, remain in the coastal waters and it is unlikely that they will be farther offshore in the study area. These species have adapted to arctic conditions through complex migration patterns, late maturity and low recruitment rates.

Subsistence fishing occurs in the Barrow and Colville River delta areas but not in the proposed survey area. A small commercial fishery operates in the Colville River delta, >115 km southeast of the closest survey line. No large fisheries are operated in the Alaskan Beaufort Sea.

Essential Fish Habitat

The Magnuson-Stevens Fishery Conservation and Management Act (MSA; 16 U.S.C. §1801-1882) established Regional Fishery Management Councils and mandated that Fishery Management Plans (FMPs) be developed to manage exploited fish and invertebrate species responsibly in Federal waters of the U.S. In 1996 as the Sustainable Fisheries Act amended the MSA to require the description and identification of Essential Fish Habitat (EFH) and FMPs, adverse impacts on EFH, and actions to conserve and enhance EFH. Guidelines were developed by NMFS to assist fishery management councils in fulfilling the requirements set forth by the MSA.

The North Pacific Fisheries Management council (NPFMC) was tasked with preparation of a FMP for the Arctic Management Area which includes all marine waters in the U.S. Exclusive Economic Zone of the Chukchi and Beaufort seas from three nautical miles offshore of the Alaska coast to 200 n.mi (370 km) offshore. The FMP was approved by the Secretary of Commerce in August 2009 and governs commercial fishing for all stocks of fish including all finfish, shellfish, or other marine living resources, except commercial fishing for Pacific salmon and Pacific halibut. EFH established in the FMP includes all waters and substrate necessary to fish for spawning, breeding, feeding, or growth to maturity. Identification of EFH is based on the historical range of target species but may expand or contract based on a variety of factors including changes in environmental variables, population size, and predator/prey distribution. EFH may be specific to a specific life stage such as egg, larval, juvenile, etc. EFH is described for only one target species, arctic cod (*Boreogadus saida*), that is likely to occur in the proposed survey area (NPFMC 2009).

Seabirds

Two bird species of special concern may be encountered during transits off the coast of Alaska. Spectacled eiders (*Somateria fischeri*) travel west along the arctic coast after breeding across the Arctic Coastal Plain (ACP) of northern Alaska. Both marine and terrestrial (for males in particular) routes are used during migration (Troy 2003). Steller's eiders (*Polysticta stelleri*) also breed on the ACP and move to marine habitats after breeding (Fredrichsen 2001), but occur in much lower densities than spectacled eiders and would be less likely to be encountered by transiting vessels in the southern Beaufort Sea. Spectacled and Steller's eiders were listed as Threatened in the U.S. under the ESA in May 1993 and July 1997, respectively. The USFWS developed separate Recovery Plans for each species (USFWS 1996, 2002).

(1) Spectacled Eider

The spectacled eider is a medium-sized sea duck that breeds along coastal areas of western and northern Alaska and eastern Russia, and winters in the Bering Sea (Petersen et al. 2000). Three breeding populations have been described: one in the Yukon-Kuskokwim (Y-K) delta in western Alaska, a second on the North Slope of Alaska, and the third in northeastern Russia. Spectacled eider was listed as a Threatened species because of declines in the breeding population in the Y-K delta (Stehn et al. 1993; Ely et al. 1994). The North Slope spectacled eider population seems to be stable, at least since the initiation of aerial surveys of the ACP since 1992 (Larned et al. 2009).

Males leave the breeding grounds along the ACP around mid- to late June at the onset of incubation by female eiders. Males are followed by females whose nests fail, and finally by successful breeding females and young birds in August and September. Female spectacled eiders have been documented migrating west along the Alaska coast as far as 22 and 40 km offshore (TERA 1999). Large concentrations of spectacled eiders gather in Ledyard Bay in the eastern Chukchi Sea after the breeding season to feed and molt before moving to the Bering Sea wintering grounds. Ledyard Bay is located between Icy Cape and Cape Lisburne and was designated as the Ledyard Bay Critical Habitat Unit (LBCHU) by the USFWS in 2001.

The proposed 2010 geophysical activities will occur >200 n.mi. (370 km) offshore in the Arctic Ocean, well north of known spectacled eider migration routes. Small numbers of spectacled eiders could be encountered by the *Healy* during transit periods in the Chukchi Sea or southern Beaufort Sea. The *Louis S. St. Laurent* will enter and leave the project area from Canada which is far to the east of spectacled eider range.

Activities associated with the proposed geophysical survey are not likely to affect spectacled eiders or other marine birds. The primary concern relates to the potential for bird collisions with vessels which could result in injury or mortality. Spectacled eiders and other marine birds can easily avoid oncoming vessels and in general there is little potential for impacts to marine birds to result from the proposed activities. Few birds and no eiders are likely to occur in the proposed survey area and impacts to marine birds will likely be negligible during the survey period.

The *Louis S. St. Laurent* is not likely to encounter spectacled (or Steller's) eiders which would be extralimital along the vessel's proposed route. Spectacled eiders could be encountered by the accompanying vessel during periods of transit in the southern Beaufort Sea or the Chukchi Sea. The potential for bird collisions with vessels is greater during periods of darkness or poor visibility, and birds can sometimes be attracted to vessel lights. Any collisions of spectacled eiders should be reported to the USFWS.

The LBCHU is a high-use area for spectacled eiders from July through October. Many birds are flightless during portions of this period and may be energetically stressed and more susceptible to disturbance or displacement. Disturbance to eiders in LBCHU could result in reduced survival. MMS (2007) developed a stipulation to reduce or minimize disturbance from vessel traffic in LBCHU which is in effect from 1 July to 15 November during which time vessel traffic should avoid Ledyard Bay to the extent possible.

(2) Steller's Eider

Most Steller's eiders breed across coastal eastern Siberia and a small number breed on the ACP of Alaska, most conspicuously near Barrow. A smaller population also breeds in western Russia and winters in northern Europe (Fredrichson 2001). Steller's eiders were formerly common breeders in the Y-K delta, but numbers there declined drastically and Steller's eider is now apparently rare as a breeding species on

the Y-K delta (Kertell 1991; Flint and Herzog 1999). Steller's eider density on the ACP is low with the highest densities reported near Barrow; the largest population, located in eastern Russia, may number >128,000 birds (Hodges and Eldridge 2001).

Steller's eiders have been observed east of Barrow in the Prudhoe Bay area where they are considered rare (TERA 1997). Although Steller's eiders may breed in a relatively large area of the ACP as far east as the Prudhoe Bay area, densities are low. Steller's eiders apparently do not breed every year and breeding may be tied to the lemming cycle (Quakenbush et al. 2004).

After the breeding season Steller's eiders move to marine habitats and may use lagoon systems and coastal bays from Barrow to Cape Lisburne, the northeast Chukotka coast, and numerous locations in southwest Alaska (USFWS 2007). Few Steller's eiders would be likely to encounter transiting vessels in the southern Beaufort Sea and Steller's eiders would be unlikely to occur in the proposed survey area in the Arctic Ocean.

(3) Other Seabirds, Shorebirds, and Waterfowl

In addition to the two eider species described above, a portion of the project area is within the range of a number of other seabird, shorebird, and waterfowl species. Most of these species would be found mainly within 30 km of shore where no seismic activities will take place. Summer bird densities in offshore marine waters of the Beaufort Sea are considered to be lower than in other marine areas adjacent to Alaska (U.S. Army Corps of Engineers 1999). There is a general absence of diving seabirds in the offshore waters of the southern Beaufort Sea, with the exception of small numbers of thick-billed murres (*Uria lomvia*), horned puffins (*Fratercula corniculata*), loons (*Gavia* spp.) and black guillemots (*Cephus grylle*). A few species of surface-feeding birds also make use of offshore waters, including red and red-necked phalaropes (*Phalaropus fulicaria* and *P. lobatus*), pomarine, parasitic and long-tailed jaegers (*Stercorarius pomarinus*, *S. parasiticus*, and *S. longicaudus*), arctic tern (*Sterna paradisaea*), and glaucous gulls (*Larus hyperboreus*). Divoky (1979) reported a bird density during the open-water season in offshore waters deeper than 18 m (60 feet) of less than 10 birds/km².

Divoky (1983) conducted extensive boat-based surveys in the Beaufort Sea during early August through mid-September. The primary species observed during pelagic surveys were surface-feeding species including gulls, terns, phalaropes, and jaegers. Long-tailed ducks, loons, and migrant eiders as well as low densities of surface-feeding species were reported during nearshore surveys. Pelagic birds were feeding primarily on arctic cod while nearshore birds were feeding on epibenthic crustaceans and zooplankton.

Frame (1973) conducted seabird observations from an icebreaker in the Beaufort Sea during August 1969 and reported black-legged kittiwake (*Rissa tridactyla*) as the most abundant species, followed by Sabine's gull (*Xema sabini*). Pomarine and long-tailed jaegers were the other two most commonly observed species along with unidentified shorebirds.

Fisher and Larned (2004) conducted more recent aerial surveys of marine birds in 1999 and 2000 in areas to 100 km offshore of the Alaskan Beaufort Sea. Approximately 90% of birds observed were sea ducks, primarily long-tailed ducks (*Clangula hyemalis*), king eiders (*Somateria spectabilis*) and scoters (*Melanitta* spp.). Densities of most species decrease with distance offshore although king eiders densities were higher in deeper, offshore waters.

Harwood et al. (2005) recorded the distribution of birds during oceanographic studies through the Canadian Basin, Beaufort Sea, and Chukchi Sea from 16 August through 6 October 2002. Sixteen bird species and a total of 1213 individuals were recorded. The birds were found in greater density in areas where oceanographic features such as a shelf break, or an area of coastal upwelling, heightened productivity.

Marine Mammals

A total of nine cetacean species, five species of pinnipeds, and one ursid (polar bear) are known to or may occur in or near the proposed study area (Table 3). Three of these species, the bowhead, humpback, and fin whales, are listed as endangered under the ESA. Humpback and fin whales however, are unlikely to be encountered along the planned trackline.

The marine mammals that occur in the proposed survey area belong to three taxonomic groups: odontocetes (toothed cetaceans, such as beluga whale and narwhal whale), mysticetes (baleen whales), and carnivora (pinnipeds and polar bears). Cetaceans and pinnipeds (except walrus) are managed by the NMFS; Pacific walrus and polar bear are managed by the USFWS.

The marine mammal species most likely to be encountered during the seismic survey include two cetacean species (beluga whale and perhaps bowhead whale), two pinniped species (ringed and bearded seals), and polar bear. However, most species will occur in low numbers and encounters are likely to be most common within 100 km of shore where no seismic work is planned to occur. The marine mammal most likely to be encountered throughout the cruise is ringed seal. The most widely distributed marine mammals within the proposed survey area are expected to be the beluga whale, ringed seal, and polar bear.

Seven additional cetacean species— narwhal, killer whale, harbor porpoise, gray whale, Minke whale, fin whale, and humpback whale —could occur in the project area. Gray whale occurs regularly in continental shelf waters along the Chuckchi Sea coast in summer and to a lesser extent along the Beaufort Sea coast. Recent evidence from monitoring activities in the Chukchi and Beaufort seas during industry seismic surveys suggests that harbor porpoise and Minke whale, which have been considered uncommon or rare in the Chukchi and Beaufort seas, may be increasing in numbers in these areas (Funk et al. 2009). Small numbers of killer whales have also been recorded during these industry surveys, along with a few sightings of fin and humpback whales. The narwhal occurs in Canadian waters and occasionally in the Beaufort Sea, but is rare there and not expected to be encountered. Each of these species is uncommon or rare in the Chukchi and Beaufort seas, and relatively few if any encounters with these species are expected during the proposed geophysical program. No sightings of these species were recorded from the *Louis S. St. Laurent* during the 2009 survey in the Canada Basin (Mosher et al. 2009).

Additional pinniped species that could be encountered during the proposed geophysical survey include spotted and ribbon seals, and Pacific walrus. Spotted seals are more abundant in the Chukchi Sea and occur in small numbers in the Beaufort Sea. Ribbon seal is uncommon in the Chukchi Sea and there are few sightings in the Beaufort Sea. Pacific walrus is common in the Chukchi Sea but uncommon in the Beaufort Sea and not likely to occur in the deep waters of the proposed survey area. None of these species would likely be encountered during the proposed cruise other than perhaps during transit periods to or from the survey area.

Polar bears occur on the pack ice in low densities and may be encountered during the proposed geophysical survey. Small numbers of polar bears were recorded during recent seismic cruises in the Arctic (Haley 2006; Haley and Ireland 2006, Mosher et al. 2009).

(1) Odontocetes

(a) Beluga Whale (*Delphinapterus leucas*)

Beluga whale is the most likely cetacean species to occur in the proposed project area. Beluga whale is an arctic and subarctic species that includes several populations in Alaska and northern European waters. It has a circumpolar distribution in the Northern Hemisphere and occurs between 50° and 80°N

(Reeves et al. 2002). It is distributed in seasonally ice-covered seas and migrates to warmer coastal estuaries, bays, and rivers in summer for molting (Finley 1982).

In Alaska, beluga whales comprise five distinct stocks: Beaufort Sea, eastern Chukchi Sea, eastern Bering Sea, Bristol Bay, and Cook Inlet (O’Corry-Crowe et al. 1997). For the proposed project, only animals from the Beaufort Sea stock and eastern Chukchi Sea stock may be encountered. Some eastern Chukchi Sea animals enter the Beaufort Sea in late summer (Suydam et al. 2005a).

The **Beaufort Sea population** was estimated to contain 39,258 individuals as of 1992 (Angliss and Allen 2009). This estimate was based on the application of a sightability correction factor of 2× to the 1992 uncorrected census of 19,629 individuals made by Harwood et al. (1996). This estimate was obtained from a partial survey of the known range of the Beaufort Sea population and may be an underestimate of the true population size. This population is not considered by NMFS to be a strategic stock and is believed to be stable or increasing (DeMaster 1995).

Beluga whales of the Beaufort stock winter in the Bering Sea, summer in the eastern Beaufort Sea, and migrate in offshore waters of western and northern Alaska (Angliss and Allen 2009). The majority of belugas in the Beaufort stock migrate into the Beaufort Sea in April or May, although some whales may pass Point Barrow as early as late March and as late as July (Braham et al. 1984; Ljungblad et al. 1984; Richardson et al. 1995).

Much of the Beaufort Sea seasonal population enters the Mackenzie River estuary for a short period during July–August to molt their epidermis, but they spend most of the summer in offshore waters of the eastern Beaufort Sea, Amundsen Gulf and more northerly areas (Davis and Evans 1982; Harwood et al. 1996; Richard et al. 2001). Belugas are rarely seen in the central Alaskan Beaufort Sea during the early summer. During late summer and autumn, most belugas migrate westward far offshore near the pack ice (Frost et al. 1988; Hazard 1988; Clarke et al. 1993; Miller et al. 1999). Lyons et al. (2009) reported the highest beluga sighting rates during the first two weeks of September during aerial surveys in the Alaskan Beaufort Sea in 2007. Peak beluga sighting rates were reported in July in 2008 when these surveys were undertaken earlier in the year (Christie et al. 2009).

TABLE 3. The habitat, abundance, and conservation status of marine mammals inhabiting the proposed study area.

| Species | Habitat | Abundance | ESA ¹ | IUCN ² | CITES ³ |
|--|---|---|------------------|-------------------|--------------------|
| Odontocetes | | | | | |
| Beluga whale (<i>Delphinapterus leucas</i>) | Offshore, Coastal, Ice edges | 3710 ⁴ 39,258 ⁵ | Not listed | VU | II |
| Narwhal (<i>Monodon monoceros</i>) | Offshore, Ice edge | Rare ⁶ | Not listed | DD | II |
| Killer whale (<i>Orcinus orca</i>) | Widely distributed | Rare | Not listed | LR-cd | II |
| Harbor Porpoise (<i>Phocoena phocoena</i>) | Coastal, inland waters, shallow offshore waters | Common (Chukchi) Uncommon (Beaufort) | Not listed | VU | II |
| Mysticetes | | | | | |
| Bowhead whale (<i>Balaena mysticetus</i>) | Pack ice & coastal | 10,545 ⁷ | Endangered | LR-cd | I |
| Gray whale (<i>Eschrichtius robustus</i>) (eastern Pacific population) | Coastal, lagoons | 488 ⁸ 20,110 ⁹ | Not listed | LR-cd | I |
| Minke whale (<i>Balaenoptera acutorostrata</i>) | Shelf, coastal | Small numbers | Not listed | LR-cd | I |

| | | | | | |
|---|--|--|-----------------------|----|----|
| Fin whale (<i>Balaenoptera physalus</i>) | Slope, mostly pelagic | Rare (Chukchi) | Endangered | EN | I |
| Humpback whale (<i>Megaptera novaeangliae</i>) | Shelf, coastal | Rare | Endangered | - | - |
| Pinnipeds | | | | | |
| Bearded seal (<i>Erignathus barbatus</i>) | Pack ice, open water | 250,000-300,000 ¹⁰ | In review for listing | - | - |
| Spotted seal (<i>Phoca largha</i>) | Pack ice, open water, coastal haulouts | ~59,214 ¹¹ | Not listed in U.S. | - | - |
| Ringed seal (<i>Pusa hispida</i>) | Landfast & pack ice, open water | 18,000 ¹² ~208,000-252,000 ¹³ | In review for listing | - | - |
| Ribbon seal (<i>Histiophoca fasciata</i>) | Pack ice, open water | 90-100,000 ¹⁴ | Not listed | - | - |
| Pacific Walrus (<i>Odobenus rosmarus</i>) | Coastal, Pack ice, ice floes | ~200,000 to 246,000 ¹⁵ | In review for listing | - | II |
| Ursids | | | | | |
| Polar Bear (<i>Ursus maritimus</i>) | Pack ice | 4700 ¹⁶ | Threatened | | |

¹ Endangered Species Act.

² IUCN Red List of Threatened Species (2003). Codes for IUCN classifications: CR = Critically Endangered; EN = Endangered; VU = Vulnerable; LR = Lower Risk (-cd = Conservation Dependent; -nt = Near Threatened; -lc = Least Concern); DD = Data Deficient.

³ Convention on International Trade in Endangered Species of Wild Fauna and Flora (UNEP-WCMC 2004).

⁴ Eastern Chukchi Sea stock based on 1989-1991 surveys with a correction factor (Angliss and Allen 2009)

⁵ Beaufort Sea stock based on surveys in 1992 (Angliss and Allen 2009).

⁶ DFO (2004) states the population in Baffin Bay and the Canadian arctic archipelago is ~60,000; very few of these enter the Beaufort Sea.

⁷ Abundance of bowhead whales surveyed near Barrow, as of 2001 (George et al. 2004). Revised to 10,545 by Zeh and Punt (2005).

⁸ Southern Chukchi Sea and northern Bering Sea (Clark and Moore 2002).

⁹ Eastern North Pacific gray whale population (Rugh et al. 2008)

¹⁰ Based on earlier estimates, no current population estimate available (Angliss and Allen 2009)

¹¹ Alaska stock based on aerial surveys in 1992 (Angliss and Allen 2009).

¹² Beaufort Sea minimum estimate with no correction factor based on aerial surveys in 1996-1999 (Frost et al. 2002 in Angliss and Allen 2009).

¹³ Eastern Chukchi Sea population (Bengtson et al. 2005)

¹⁴ Bering Sea population (Bruns 1981 in Angliss and Allen 2009).

¹⁵ Pacific walrus population, 1975-1990 (Angliss and Allen 2009 and references therein).

¹⁶ Chukchi Sea and northern and southern Beaufort Sea populations combined (Aars et al. 2006; USFWS 2008).

Moore (2000) and Moore et al. (2000b) suggested that beluga whales select deeper slope water independent of ice cover. However, during the westward migration in late summer and autumn, small numbers of belugas are sometimes seen near the north coast of Alaska (e.g., Johnson 1979). Lyons et al. (2009) reported higher beluga sighting rates at locations >60 km offshore than at locations nearer shore during aerial surveys in the Alaskan Beaufort Sea in 2006 and 2007. No beluga whales were observed during seismic projects within latitudes of the proposed project – north of 71 °N – in 2005, 2006 or 2009 (Haley and Ireland 2006, Haley 2006, Mosher et al. 2009). Marine mammal observers did, however, record one sighting of more than two beluga whales within the southern-most latitude (71.37°N) of the proposed survey in 2008 (GSC unpubl. data, 2008). These animals were approximately 636 km east of the proposed project area on 23 August.

The main fall migration corridor of beluga whales is ~100+ km north of the coast. Satellite-linked telemetry data show that some belugas of this population migrate west considerably farther offshore, as far north as 76° to 78°N latitude (Richard et al. 1997, 2001).

The *eastern Chukchi Sea* population is estimated at 3710 animals (Angliss and Allen 2009). This estimate was based on surveys conducted in 1989–1991. Survey effort was concentrated on the 170 km long Kasegaluk Lagoon where belugas are known to occur during the open-water season. The actual number of beluga whales recorded during the surveys was much lower. Correction factors to account for animals that were underwater and for the proportion of newborns and yearlings that were not observed due to their small size and dark coloration were used to calculate the estimate. The calculation was considered to be a minimum population estimate for the eastern Chukchi stock because the surveys on which it was based did not include offshore areas where belugas are also likely to occur. This population is considered to be stable. It is assumed that beluga whales from the eastern Chukchi stock winter in Bering Sea (Angliss and Allen 2009).

Although beluga whales are known to congregate in Kasegaluk Lagoon during summer, evidence from a small number of satellite-tagged animals suggests that some of these whales may subsequently range into the Arctic Ocean north of the Beaufort Sea. Suydam et al. (2005a) put satellite tags on 23 beluga whales captured in Kasegaluk Lagoon in late June and early July 1998–2002. Five of these whales moved far into the Arctic Ocean and into the pack ice to 79–80°N. These and other whales moved to areas as far as 1,100 km offshore between Barrow and the Mackenzie River delta spending time in water with 90% ice coverage.

During aerial surveys in nearshore areas (i.e., ~37 km offshore) of the Chukchi Sea in 2006–2008, peak beluga sighting rates were recorded in July and the lowest monthly sighting rates were recorded in September (Thomas et al. 2009). Sighting rates were variable during other months but were lowest in August and September. Beluga whale sighting rates and number of individuals were generally highest in the band 25–35 km offshore. The largest single groups, however, were sighted at locations near shore in the band within 5 km of shore.

Beluga whales from the eastern Chukchi Sea stock are an important subsistence resource for residents of the village of Point Lay, adjacent to Kasegaluk Lagoon, and other villages in northwest Alaska. Each year, hunters from Point Lay drive belugas into the lagoon to a traditional hunting location. The belugas have been predictably sighted near the lagoon from late June through mid- to late July (Suydam et al. 2001). In 2007 approximately 70 belugas were also harvested at Kivalina located southeast of Point Hope.

Pod structure in beluga groups appears to be along matrilineal lines, with males forming separate aggregations. Small groups are often observed traveling or resting together. Belugas often migrate in groups of 100 to 600 animals (Braham and Krogman 1977). The relationships between whales within groups are not known, although hunters have reported that belugas form family groups with whales of different ages traveling together (Huntington 2000).

(b) Narwhal (*Monodon monoceros*)

Narwhals have a discontinuous arctic distribution (Hay and Mansfield 1989; Reeves et al. 2002). A large population inhabits Baffin Bay, West Greenland, and the eastern part of the Canadian Arctic archipelago, and much smaller numbers inhabit the Northeast Atlantic/East Greenland area. Narwhals are associated with sea ice. In the spring, as the ice breaks up, they follow the receding ice edge and enter deep sounds and fjords, where they remain during the summer and early fall (Reeves et al. 2002). As the ice reforms, narwhals move to offshore areas in the pack ice (Reeves et al. 2002), living in leads in the heavy pack ice throughout the winter. Most pods consist of 2–10 individuals but they may aggregate to

form larger herds of hundreds or even thousands of individuals (Jefferson et al. 1993). According to Hay (1985), segregation by age and sex is evident, with summering groups consisting of mature females with calves, immature and maturing males, and large mature males.

Population estimates for the narwhal are scarce, and the IUCN-World Conservation Union lists the species as Data Deficient (IUCN Red List of Threatened Species 2003). Innes et al. (2002) estimated a population size of 45,358 narwhals in the Canadian Arctic although little of the area was surveyed. There are scattered records of narwhal in Alaskan waters where the species is considered extralimital (Reeves et al. 2002). Narwhals were not recorded during survey projects within latitudes of the area of this proposed project in 2005, 2006, 2008 or 2009 (Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008; Mosher et al. 2009). Narwhals are not expected to be encountered during the proposed activity. If narwhals are observed during the survey, they would most likely be seen along the eastern portions of the proposed trackline where they would be considered extralimital.

(c) Killer Whale (*Orcinus orca*)

Killer whales are cosmopolitan and globally fairly abundant. The killer whale is very common in temperate waters, but also frequents the tropics and waters at high latitudes. Killer whales appear to prefer coastal areas, but are also known to occur in deep water (Dahlheim and Heyning 1999). The greatest abundance is thought to occur within 800 km of major continents (Mitchell 1975) and the highest densities occur in areas with abundant prey. Both resident and transient stocks have been described as well as an “offshore” ecotype. The resident and transient types are believed to differ in several aspects of morphology, ecology, and behavior including dorsal fin shape, saddle patch shape, pod size, home range size, diet, travel routes, dive duration, and social integrity of pods (Angliss and Allen 2009).

Killer whales are known to inhabit almost all coastal waters of Alaska, extending from southeast Alaska through the Aleutian Islands to the Bering and Chukchi seas (Angliss and Allen 2009). Killer whales probably do not occur regularly in the Beaufort Sea although sightings have been reported (Leatherwood et al. 1986; Lowry et al. 1987). George et al. (1994) reported that they and local hunters see a few killer whales at Point Barrow each year. Killer whales are more common southwest of Barrow in the southern Chukchi Sea and the Bering Sea. Based on photographic techniques, ~100 animals have been identified in the Bering Sea (ADFG 1994). Killer whales from either the North Pacific resident or transient stock could occur in the Chukchi Sea during the summer. The number of killer whales likely to occur in the Chukchi Sea during the proposed activity is unknown. Marine mammal observers (MMOs) onboard industry vessels in the Chukchi Sea recorded two killer whale sightings each in 2006 and 2008, and one sighting in 2007 (Haley et al. 2009). MMOs onboard industry vessels did not record any killer whale sighting in the Beaufort Sea in 2006-2008 (Savarese et al. 2009), and killer whales were not recorded during recent Arctic Ocean cruises (Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008; Mosher et al. 2009).

(d) Harbor Porpoise (*Phocoena phocoena*)

The harbor porpoise is a small odontocete that inhabits shallow, coastal waters—temperate, subarctic, and arctic—in the Northern Hemisphere (Read 1999). Harbor porpoises occur mainly in shelf areas where they can dive to depths of at least 220 m and stay submerged for more than 5 min (Harwood and Wilson 2001) feeding on small schooling fish (Read 1999). Harbor porpoises typically occur in small groups of only a few individuals and tend to avoid vessels (Richardson et al. 1995).

The subspecies *P. p. vomerina* ranges from the Chukchi Sea, Pribilof Islands, Unimak Island, and the south-eastern shore of Bristol Bay south to San Luis Obispo, California. Point Barrow, Alaska, is the approximate northeastern extent of their regular range (Suydam and George 1992), though there are extralimital records east to the mouth of the Mackenzie River in the Northwest Territories, Canada, and recent

sightings in the Beaufort Sea in the vicinity of Prudhoe Bay during aerial surveys in 2007 and 2008 (Lyons et al. 2009; LGL Limited, unpubl. data). MMOs onboard industry vessels reported one harbor porpoise sighting in the Beaufort Sea in 2006 but no sightings were recorded in 2007 or 2008 (Savarese et al. 2009). Monnett and Treacy (2005) did not report any harbor porpoise sightings during aerial surveys in the Beaufort Sea from 2002 through 2004.

Although separate harbor porpoise stocks for Alaska have not been identified, Alaskan harbor porpoises have been divided into three groups for management purposes. These groups include animals from southeast Alaska, Gulf of Alaska, and Bering Sea populations. Chukchi Sea harbor porpoises belong to the Bering Sea group which includes animals from Unimak Pass northward. Based on aerial surveys in 1999, the Bering Sea population was estimated at 48,215 animals, although this estimate is likely conservative because the surveyed area did not include known harbor porpoise range near the Pribilof Islands or waters north of Cape Newenham (~55°N; Angliss and Allen 2009). Suydam and George (1992) suggested that harbor porpoises occasionally occur in the Chukchi Sea and reported nine records of harbor porpoise in the Barrow area in 1985–1991.

More recent vessel-based surveys in the Chukchi Sea reported that harbor porpoise was one of the most abundant cetaceans during summer and fall in 2006–2008 (Haley et al. 2009; Ireland et al. 2009). Although these recent sightings suggest that harbor porpoise numbers may be increasing in the relatively shallow waters of the Chukchi Sea no recent information is available on their status in the deeper offshore waters of the proposed project area. Harbor porpoises were not recorded during arctic survey cruises in 2005, 2006, 2008 or 2009 (Haley and Ireland 2006; Haley 2006, GSC unpubl. data 2008, Mosher et al. 2009).

(2) *Mysticetes*

(a) *Bowhead Whale (Balaena mysticetus)*

The pre-exploitation population of bowhead whales in the Bering, Chukchi, and Beaufort seas is estimated to have been 10,400–23,000 whales. Commercial whaling activities may have reduced this population to perhaps 3000 animals (Woodby and Botkin 1993). Up to the early 1990s, the population size was believed to be increasing at a rate of about 3.2% per year (Zeh et al. 1996) despite annual subsistence harvests of 14–74 bowheads from 1973 to 1997 (Suydam et al. 1995). Allowing for an additional census in 2001, the latest estimates are based on an annual population growth rate of 3.4% (95% CI 1.7–5%) from 1978 to 2001 and a population size (in 2001) of ~10,470 animals (George et al. 2004, recently revised to 10,545 by Zeh and Punt 2005). Assuming a continuing annual population growth of 3.4%, the 2010 bowhead population may number around 14,247 animals. The large increases in population estimates that occurred from the late 1970s to the early 1990s were partly a result of actual population growth, but were also partly attributable to improved census techniques (Zeh et al. 1993). Although apparently recovering well, the BCB bowhead population is currently listed as endangered under the ESA and is classified as a *strategic stock* by NMFS and *depleted* under the MMPA (Angliss and Allen 2009).

Bowhead whales only occur at high latitudes in the northern hemisphere and have a disjunct circumpolar distribution (Reeves 1980). The bowhead is one of only three whale species that spend their entire lives in the Arctic. Bowhead whales are found in the western Arctic (Bering, Chukchi, and Beaufort seas), the Canadian Arctic and West Greenland (Baffin Bay, Davis Strait, and Hudson Bay), the Okhotsk Sea (eastern Russia), and the Northeast Atlantic from Spitzbergen westward to eastern Greenland. Four stocks are recognized for management purposes. The largest is the Western Arctic or Bering–Chukchi–Beaufort (BCB) stock, which includes whales that winter in the Bering Sea and migrate

through the Bering Strait, Chukchi Sea and Alaskan Beaufort Sea to the Canadian Beaufort Sea, where they feed during the summer. These whales migrate west through the Alaskan Beaufort Sea in the fall as they return to wintering areas in the Bering Sea. Satellite tracking data indicate that most bowhead whales continue migrating west past Barrow and through the Chukchi Sea to Russian waters before turning south toward the Bering Sea (Quakenbush 2007). Some bowhead may reach ~75°N latitude during the westward fall migration (Quakenbush 2009). Other researchers have also reported a westward movement of bowhead whales through the northern Chukchi Sea during fall migration (Moore et al. 1995; Mate et al. 2000).

The BCB stock of bowhead whales winter in the central and western Bering Sea and many of them summer in the Canadian Beaufort Sea (Moore and Reeves 1993). Spring migration through the Chukchi and the western Beaufort Sea occurs through offshore ice leads, generally from March through mid-June (Braham et al. 1984; Moore and Reeves 1993).

Some bowheads arrive in coastal areas of the eastern Canadian Beaufort Sea and Amundsen Gulf in late May and June, but most may remain among the offshore pack ice of the Beaufort Sea until mid-summer. After feeding primarily in the Canadian Beaufort Sea and Amundsen Gulf, bowheads migrate westward across the Beaufort Sea from late August through mid- or late October.

Bowhead activity in the Beaufort Sea in fall has been well studied in recent years. Fall migration into Alaskan waters is primarily during September and October. However, in recent years a small number of bowheads have been seen or heard offshore from the Prudhoe Bay region during the last week of August (Treacy 1993; LGL and Greeneridge 1996; Greene 1997; Greene et al. 1999; Blackwell et al. 2004, 2009; Greene et al. 2007). Consistent with this, Nuiqsut whalers have stated that the earliest arriving bowheads have apparently reached the Cross Island area earlier in recent years than formerly (T. Napageak, pers. comm.). In 2007 the MMS and the National Marine Mammal Laboratory (NMML) initiated the Bowhead Whale Feeding Ecology Study (BOWFEST) focusing on late summer oceanography and prey densities relative to bowhead distribution (Rugh 2009).

The Minerals Management Service (MMS) has conducted or funded late-summer/autumn aerial surveys for bowhead whales in the Alaskan Beaufort Sea since 1979 (e.g., Ljungblad et al. 1986, 1987; Moore et al. 1989; Treacy 1988–1998, 2000, 2002a,b; Monnett and Treacy 2005; Treacy et al. 2006). Bowheads tend to migrate west in deeper water (farther offshore) during years with higher-than-average ice coverage than in years with less ice (Moore 2000; Treacy et al. 2006). The migration corridor ranged from ~30 km offshore during light ice years to ~80 km offshore during heavy ice years (Treacy et al. 2006). In addition, the sighting rate tends to be lower in heavy ice years (Treacy 1997:67). During fall migration, most bowheads migrate west in water ranging from 15 to 200 m deep (Miller et al. 2002). Some individuals enter shallower water, particularly in light ice years, but very few whales are ever seen shoreward of the barrier islands in the Alaskan Beaufort Sea. Survey coverage far offshore in deep water is usually limited, and offshore movements may have been underestimated. However, the main migration corridor is over the continental shelf.

In autumn, westward-migrating bowhead whales typically reach the Kaktovik and Cross Island areas in early September, which is when the subsistence hunts for bowheads typically begin in those areas (Kaleak 1996; Long 1996; Galginaitis and Koski 2002; Galginaitis and Funk 2004, 2005; Koski et al. 2005). In recent years the hunts at those two locations have usually ended by mid- to late September.

Westbound bowheads typically reach the Barrow area in mid-September, and are in that area until late October (e.g., Brower 1996). Autumn bowhead whaling near Barrow normally begins in mid-September to early October, but may begin as early as August if whales are observed and ice conditions

are favorable (USDI/BLM 2005). Whaling near Barrow can continue into October, depending on the quota and conditions.

Over the years, local residents have reported small numbers of bowhead whales feeding off Barrow or in the pack ice off Barrow during the summer. Bowhead whales that are thought to be part of the Western Arctic stock may also occur in small numbers in the Bering and Chukchi seas during the summer (Rugh et al. 2003). Thomas et al. (2009) reported bowhead sightings during summer aerial surveys in nearshore areas of the Chukchi Sea from 2006-2008. All sightings were recorded in the northern portion of the study area north of 70°N latitude. Peak monthly bowhead sighting rates, however, were highest in October and November and lowest in July-September. A few bowhead whales were also recorded during vessel-based surveys in summer 2008 in the Chukchi Sea (LGL unpubl. data). Observers from the NMML reported 19 summer bowhead sightings in the Chukchi Sea during aerial surveys from 26 June through 26 July 2009 suggesting that some bowheads may summer in the Chukchi Sea (unpublished data available at http://www.afsc.noaa.gov/NMML/cetacean/bwasp/flights_COMIDA.php). Only one bowhead sighting was reported later in the year (22 August) during similar surveys in 2008. Sekiguchi et al. (2008) reported one sighting of an aggregation of ~30 bowheads during vessel-based operations about 130 km north of Cape Lisburne on 9 August 2007. Bowhead whales were not reported by vessel-based observers during arctic cruises in 2005, 2006, 2008 or 2009 (Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008; Mosher et al. 2009).

Most spring-migrating bowhead whales will likely pass through the Chukchi and Beaufort seas prior to the start of the proposed survey in August. However, a few whales that may remain in the Chukchi Sea or in the Barrow area during the summer could be encountered by transiting vessels. The potential for encounters with bowhead whales would be more likely during the westward fall migration in September. Much of the proposed survey area however, is in deep water well north of the known bowhead migration corridor and few if any bowheads are likely to be encountered during the survey activity.

(b) Gray Whale (Eschrichtius robustus)

Gray whales originally inhabited both the North Atlantic and North Pacific oceans. The Atlantic populations are believed to have become extinct by the early 1700s. There are two populations in the North Pacific. A relic population which survives in the Western Pacific summers near Sakhalin Island far from the proposed survey area. The larger eastern Pacific or California gray whale population recovered significantly from commercial whaling during its protection under the ESA until 1994 and numbered about 29,758 ±3122 in 1997 (Rugh et al. 2005). However, abundance estimates since 1997 indicate a consistent decline followed by the population stabilizing or gradually recovering. Rugh et al. (2005) estimated the population to be 18,178 ±1780 in winter 2001-2002. The population estimate increased during winter 2006-2007 to 20,110 ±1766 (Rugh et al. 2008). The eastern Pacific stock is not considered by NMFS to be endangered or to be a strategic stock.

Eastern Pacific gray whales calve in the protected waters along the west coast of Baja California and the east coast of the Gulf of California from January to April (Swartz and Jones 1981; Jones and Swartz 1984). At the end of the calving season, most of these gray whales migrate about 8000 km, generally along the west coast of North America, to the main summer feeding grounds in the northern Bering and Chukchi seas (Tomilin 1957; Rice and Wolman 1971; Braham 1984; Nerini 1984; Moore et al. 2003; Bluhm et al. 2007). Most gray whales begin a southward migration in November with breeding and conception occurring in early December (Rice and Wolman 1971).

Most summering gray whales have historically congregated in the northern Bering Sea, particularly off St. Lawrence Island in the Chirikov Basin (Moore et al. 2000a), and in the southern Chukchi Sea.

More recently, Moore et al. (2003) suggested that gray whale use of Chirikov Basin has decreased, likely as a result of the combined effects of changing currents resulting in altered secondary productivity dominated by lower quality food. Coyle et al (2007) noted that ampeliscid amphipod production in the Chirikov Basin had declined by 50% from the 1980s to 2002-3 and that as little as 3-6% of the current gray whale population could consume 10-20% of the ampeliscid amphipod annual production. These data support the hypotheses that changes in gray whale distribution may be caused by changes in food production and that gray whales may be approaching or have surpassed the carrying capacity of their summer feeding areas. Bluhm et al. (2007) noted high gray whale densities along ocean fronts and suggested that ocean fronts may play an important role in influencing prey densities in eastern North Pacific gray whale foraging areas. The northeastern-most of the recurring feeding areas is in the northeastern Chukchi Sea southwest of Barrow (Clarke et al. 1989).

Gray whales routinely feed in the Chukchi Sea during the summer. Moore et al. (2000b) reported that during the summer, gray whales in the Chukchi Sea were clustered along the shore primarily between Cape Lisburne and Point Barrow and were associated with shallow, coastal shoal habitat. In autumn, gray whales were clustered near shore at Point Hope and between Icy Cape and Point Barrow, as well as in offshore waters northwest of Point Barrow at Hanna Shoal and southwest of Point Hope. Thomas et al. (2009) reported that gray whale sighting rates and abundance were greater in the 0-5 km offshore band in 2006, and in the 25-30 km band in 2007 and 2008 during aerial surveys of the nearshore area of the eastern Chukchi Sea. They suggested that the difference in gray whale distribution in 2006 vs. 2007 and 2008 may have been due to differences in food availability and perhaps ice conditions.

Gray whales occur fairly often near Point Barrow, but historically only a small number of gray whales have been sighted in the Beaufort Sea east of Point Barrow. Hunters at Cross Island (near Prudhoe Bay) took a single gray whale in 1933 (Maher 1960). Only one gray whale was sighted in the central Alaskan Beaufort Sea during the extensive aerial survey programs funded by MMS and industry from 1979 to 1997. However, during September 1998, small numbers of gray whales were sighted on several occasions in the central Alaskan Beaufort Sea (Miller et al. 1999; Treacy 2000). More recently a single sighting of a gray whale was made on 1 August 2001 near the Northstar production island (Williams and Coltrane 2002). Several gray whale sightings were reported during both vessel-based and aerial surveys in the Beaufort Sea in 2006 and 2007 (Jankowski et al. 2008; Lyons et al. 2009) and during vessel-based surveys in 2008 (Savarese et al. 2009). Several single gray whales have been seen farther east in the Canadian Beaufort Sea (Rugh and Fraker 1981; LGL Ltd., unpubl. data), indicating that small numbers must travel through the Alaskan Beaufort during some summers. In recent years, ice conditions have become lighter near Barrow, and gray whales may have become more common there and perhaps in the Beaufort Sea. In the springs of 2003 and 2004, a few tens of gray whales were seen near Barrow by early-to-mid June (LGL Ltd and NSB-DWM, unpubl. data). In the spring of 2003 and 2004, a few tens of gray whales were seen near Barrow by early-to-mid June (LGL Ltd and NSB-DWM, unpubl. data). However, no gray whales were sighted during cruises in the Arctic Ocean north of Barrow in 2002, 2005, 2006, 2008 or 2009 (Harwood et al. 2005; Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008; Mosher et al. 2009).

Small numbers of gray whales could be encountered by survey vessels during transit periods. Gray whales occur in relatively shallow waters where they feed on benthic invertebrates and they are not likely to occur in the deeper water of the proposed survey area.

(c) Minke Whale (*Balaenoptera acutorostrata*)

Minke whales have a cosmopolitan distribution at ice-free latitudes (Stewart and Leatherwood 1985), and also occur in some marginal ice areas. Angliss and Allen (2009) recognize two minke whale

stocks in U.S. waters: (1) the Alaska stock, and (2) the California/Oregon/Washington stock. There is no abundance estimate for the Alaska stock. Provisional estimates of Minke whale abundance based on surveys in 1999 and 2000 are 810 and 1003 whales in the central-eastern and south-eastern Bering Sea, respectively (Moore et al. 2002). These estimates have not been corrected for animals that may have been submerged or otherwise missed during the surveys, and only a portion of the range of the Alaskan stock in the central eastern and southeastern Bering Sea was surveyed.

Minke whales range into the Chukchi Sea and a few sightings have been reported in the Beaufort Sea in recent years (Funk et al. 2009). The level of Minke whale use of the Chukchi Sea is unknown. Leatherwood et al. (1982, in Angliss and Allen 2009) indicated that Minke whales are not considered abundant in any part of their range, but that some individuals venture north of the Bering Strait in summer. Reiser et al. (2009) reported eight and five Minke whale sightings in 2006 and 2007, respectively, during vessel-based surveys in the Chukchi Sea, and Haley et al. (2009) reported 26 Minke whale sightings during similar vessel-based surveys in the Chukchi Sea in 2008. Savarese et al. (2009) reported two Minke whale sightings in the Beaufort Sea during vessel-based operations in 2006-2008. No Minke whale sightings were reported during arctic cruises in 2005, 2006, 2008 or 2009 (Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008; Mosher et al. 2009). Minke whales sometimes occur in areas with minimal ice cover and it is possible though unlikely that a few Minke whales could be encountered during the proposed survey activities.

(d) Fin Whale (*Balaenoptera physalus*)

Fin whale is listed as endangered under the ESA and by IUCN, is classified as a strategic stock by NMFS, and is a CITES Appendix I species (Table 3). Fin whales are widely distributed in all the world's oceans (Gambell 1985), but typically occur in temperate and polar latitudes and less frequently in the tropics (Reeves et al. 2002). Fin whales feed in northern latitudes during the summer where their prey includes plankton as well as schooling pelagic fish, such as herring, sandlance, and capelin (Jonsgård 1966a,b; Reeves et al. 2002). The North Pacific population summers from the Chukchi Sea in small numbers to California (Gambell 1985), but does not range into the Alaskan Beaufort Sea or waters of the northern Chukchi Sea. Reliable estimates of fin whale abundance in the Northeast Pacific are not available (Angliss and Allen 2009). Provisional estimates of fin whale abundance in the central-eastern and south-eastern Bering Sea are 3,368 and 683, respectively (Moore et al. 2002). Zerbini et al. (2006) reported numerous fin whale sightings from Kodiak Island to the central Aleutian Islands.

No estimates for fin whale abundance during the summer in the Chukchi Sea are available. Recently a fin whale was recorded in the southern Chukchi Sea during vessel-based surveys in 2006 (LGL unpublished data), and three fin whale sightings were recorded in the Chukchi Sea in 2008 (Haley et al. 2009). NMML observers also observed and photographed a fin whale off Pt. Lay in 2008 during the COMIDA aerial survey program. Fin whales were not recorded during vessel-based or aerial surveys in the Beaufort Sea in 2006-2008 (Savarese et al. 2009; Christie et al. 2009), and were not reported during arctic cruises in 2005, 2006, 2008 or 2009 (Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008; Mosher et al. 2009). Fin whale would be unlikely to occur in the proposed geophysical survey area.

(e) Humpback Whale (*Megaptera novaeangliae*)

Humpback whales are distributed in major oceans worldwide and their range in the North Pacific extends through the Bering Sea into the southern Chukchi Sea (Angliss and Allen 2009). In general, humpback whales spend winter in tropical and sub-tropical waters where breeding and calving occur, and migrate to higher latitudes for feeding during the summer.

Humpback whales were hunted extensively during the 20th century and worldwide populations may have been reduced to ~10% of their original numbers. The International Whaling Commission banned

commercial hunting of humpback whales in the Pacific Ocean in 1965 and humpbacks were listed as endangered under the ESA and depleted under the MMPA in 1973. Most humpback whale populations appear to be recovering well.

Humpbacks feed on euphausiids, copepods, and small schooling fish, notably herring, capelin, and sandlance (Reeves et al. 2002). As with other baleen whales, the food is trapped or filtered when large amounts of water taken into the mouth and the expanded throat area are forced out through the baleen plates. Individual humpback whales can often be identified by distinctive patterns on the tail flukes. They are frequently observed breaching or engaged in other surface activities. Adult male and female humpback whales average 14 and 15 m (46 and 49 ft) in length, respectively (Wynne 1997). Humpbacks have large, robust bodies and long pectoral flippers which may reach 1/3 of their body length. The dorsal fin is variable in shape and located well back toward the posterior 1/3 of the body on a hump which is particularly noticeable when the back is arched during a dive (Reeves et al. 2002).

Angliss and Allen (2009) reported that at least three humpback whale populations have been identified in the North Pacific. Two of these stocks may be relevant to the Chukchi Sea portion of the project area. The Central North Pacific stock winters in waters near Hawaii and migrates to British Columbia, Southeast Alaska, and Prince William Sound to Unimak Pass to feed during the summer. The Western North Pacific stock winters off the coast of Japan and probably migrates to the Bering Sea to feed during the summer. There may be some overlap between the Central and Western North Pacific stocks.

Humpback whale sightings in the Bering Sea have been recorded southwest of St. Lawrence Island, the southeastern Bering Sea, and north of the central Aleutian Islands (Moore et al. 2002; Angliss and Allen 2009). Recently there have been sightings of humpback whales in the Chukchi Sea and a single sighting in the Beaufort Sea (Green et al. 2007). Haley et al (2009) reported four humpback whales during vessel-based surveys in the Chukchi Sea in 2007 and two sightings in 2008. NMML observers recorded a humpback whale during aerial surveys in the Chukchi Sea in 2009. Green et al. (2007) reported and photographed a humpback whale cow/calf pair east of Barrow near Smith Bay in 2007. No humpback whales were reported during cruises in the Arctic Ocean in 2005, 2006, 2008 or 2009 (Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008; Mosher et al. 2009). Whether the recent humpback whale sightings in the Chukchi and Beaufort seas are related to climate changes in the Arctic in recent years is unknown. Humpback whales could occur in the Chukchi Sea and possibly in the Beaufort Sea but would be unlikely to occur in the deep offshore waters of the proposed survey area.

(3) Pinnipeds

(a) Pacific Walrus (*Odobenus rosmarus divergens*)

Walrus occur in moving pack ice over shallow waters of the circumpolar Arctic coast (King 1983). There are two recognized subspecies of walrus: the Pacific and Atlantic walrus (*O. r. divergens* and *O. r. rosmarus*, respectively.). Only the *divergens* subspecies could potentially occur within the proposed geophysical survey area.

Estimates of the pre-exploitation population of the Pacific walrus range from 200,000 to 400,000 animals (Angliss and Allen 2009 and references therein). Over the past 150 years, the population has been depleted by over-harvesting and then periodically allowed to recover (Fay et al. 1989). No current population estimate is available. The USFWS and the USGS are currently investigating new techniques, including remote sensing, for producing a more precise abundance estimate of the Pacific walrus population (Burn et al. 2006; Udevitz et al. 2008).

Pacific walrus range from the Bering Sea to the Chukchi Sea, occasionally moving into the East Siberian and Beaufort seas. Walrus are migratory, moving south with the advancing ice in autumn and north as the ice recedes in spring (Fay 1981). In the summer, most of the population of Pacific walrus moves to the Chukchi Sea, but several thousand aggregate in the Gulf of Anadyr and in Bristol Bay (Angliss and Allen 2009). Limited numbers of walrus inhabit the Beaufort Sea during the open water season, and they are considered extralimital east of Point Barrow (Sease and Chapman 1988). The northeast Chukchi Sea west of Barrow is the northeastern extent of the main summer range of the Pacific walrus, and only a few are seen farther east in the Beaufort Sea (e.g., Harwood et al. 2005; Savarese et al. 2009).

The estimated average annual walrus mortality due to subsistence harvest in Russia and the U.S. was 5789, which included animals wounded but not retrieved (Angliss and Allen 2009).

Walrus are most commonly found near the southern margins of the pack ice as opposed to deep in the pack where few open leads (polynyas) exist to afford access to the sea for foraging (Estes and Gilbert 1978; Gilbert 1989; Fay 1982). Walrus are not typically found in areas of >80% ice cover (Fay 1982). Ice serves as an important mobile platform providing walrus with a place to rest and nurse their young which is safe from predators and near feeding grounds.

This close relationship to the ice largely determines walrus distribution and the timing of their migrations. As the pack ice breaks up in the Bering Sea and recedes northward in May-June, a majority of subadults, females and calves migrate with it, either by swimming or resting on drifting ice sheets. Many males will choose to stay in the Bering Sea for the entire year, with concentrations near Saint Lawrence Island and further south in Bristol Bay. Two northward migration pathways are apparent, either toward the eastern Chukchi Sea near Barrow or northwestward toward Wrangel Island. By late June to early July, concentrations of walrus migrating northeastward spread along the Alaska coast concentrated within 200 km of the shore from Saint Lawrence Island to southwest of Barrow. In August, largely dependent on the retreat of the pack ice, walrus are found further offshore with principal concentrations northwest of Barrow. By October, a reverse migration occurs from the Chukchi Sea, with animals swimming ahead of the developing pack ice (Fay 1982).

Pacific walrus feed primarily on benthic invertebrates, occasionally fish and cephalopods, and more rarely, some adult males may prey on other pinnipeds (reviewed in Riedman 1990). Walrus typically feed in depths of 10–80 m (Vibe 1950; Fay 1982; Reeves et al. 2002). In a recent study in Bristol Bay, 98% of satellite locations of tagged walrus were in water depths of 60 m or less (Chadwick and Hills 2005). Though the deepest dive recorded for a walrus was 133 m, they are more likely to be found in depths of 80 m or less in coastal or continental shelf habitats, where they feed on clams and other marine mollusks (Fay 1982; Fay and Burns 1988; Reeves et al. 2002).

Recently global climate changes have apparently resulted in retreat of the pack ice beyond the shallow habitats of the Chukchi Sea into deeper waters of the Arctic Ocean during summer months. Water depths in the Arctic Ocean are too great to permit walrus feeding and many thousands of walrus hauled out to rest at terrestrial sites along the eastern Chukchi Sea coast in 2007 (Thomas et al. 2009). A similar situation occurred in 2009 when the pack ice retreated and walrus were forced to use terrestrial haulouts. In 2009 over 100 walrus, primarily smaller, young animals, died at haulouts apparently as a result of injuries sustained by stampeding adults. Similar mortality incidents were not reported in Alaska during 2007. Belikov et al. (1996) also reported similar use of terrestrial haulouts by walrus in years of excessive ice retreat in Russia. The Center for Biological Diversity petitioned the Secretary of Interior to list Pacific walrus as a threatened or endangered species under the ESA primarily as a result of potential impacts from global climate change and associated retreat of the pack ice (CBD 2008b).

The proposed geophysical survey will be conducted in the Arctic Ocean north of the Beaufort Sea in water depths that preclude walrus feeding, and walrus would be unlikely to occur in the vicinity of the proposed survey. Walrus could be encountered during transit periods in the Chukchi Sea but would not likely be encountered in the Beaufort Sea. Two sightings of seven total walrus were encountered between 71 and 74°N during the Healy's arctic survey in 2005 (Haley and Ireland 2006). However, these sightings occurred far to the west of the proposed 2010 survey area in water depths of <70 m.

(b) Bearded Seal (*Erignathus barbatus*)

Bearded seals are associated with sea ice and have a circumpolar distribution (Burns 1981). During the open-water period, bearded seals occur mainly in relatively shallow areas, because they are predominantly benthic feeders (Burns 1981). They prefer areas of water no deeper than 200 m (e.g., Harwood et al. 2005). No reliable estimate of bearded seal abundance is available for the Chukchi and Beaufort seas (Angliss and Allen 2009). The Alaska stock of bearded seals is not classified by NMFS as endangered or a strategic stock however there has recently been a petition to list this and other arctic seals due to the potential impact to seal habitats resulting from current warming trends (CBD 2008a). Bearded seal is currently under review as a possible candidate for listing.

In Alaskan waters, bearded seals occur over the continental shelves of the Bering, Chukchi, and Beaufort seas (Burns 1981). The Alaska stock of bearded seals may consist of about 300,000–450,000 individuals based on earlier accounts but no current population estimates are available (MMS 1996; Angliss and Allen 2009). Bengtson et al. (2005) reported bearded seal densities in the Chukchi Sea ranging from 0.07 to 0.14 seals/km² in 1999 and 2000, respectively. No population estimates could be calculated because these densities were not adjusted for haulout behavior. Bearded seals were more common in offshore pack ice with the exception of high bearded seal numbers observed near the shore south of the survey area near Kivalina. Haley et al. (2009) reported bearded seal densities up to 0.022 to 0.064 seals/km² in summer and fall, respectively during vessel-based surveys in the Chukchi Sea in 2006-2008. These densities were lower than those reported by Bengtson et al. (2005) but are not directly comparable because the latter densities were based on aerial surveys of seals at ice holes in the late May and early June.

In the Beaufort Sea Savarese et al. (2009) reported bearded seal densities up to 0.028 and 0.035 in the summer and fall, respectively during vessel-based surveys in 2006-2008. Haley and Ireland (2006) reported no sightings of bearded seals during an arctic cruise from the *Healy* in 2005 along ~361 km of monitored trackline within the latitudes of the proposed survey (71 -77 °N). Five bearded seal sightings were reported during the 2006 *Healy* cruise along 622 km of trackline within 71 -74 °N (Haley 2006).

Bearded seal is the largest of the northern phocids. Bearded seals have occasionally been reported to maintain breathing holes in sea ice and broken areas within the pack ice, particularly if the water depth is <200 m. Bearded seals apparently also feed on ice-associated organisms when they are present, and this allows a few bearded seals to live in areas considerably more than 200 m deep.

Seasonal movements of bearded seals are directly related to the advance and retreat of sea ice and to water depth (Kelly 1988). During winter, most bearded seals in Alaskan waters are found in the Bering Sea. In the Chukchi and Beaufort seas, favorable conditions are more limited, and consequently, bearded seals are less abundant there during winter. From mid-April to June as the ice recedes, some bearded seals that overwintered in the Bering Sea migrate northward through the Bering Strait. During the summer they are found near the widely fragmented margin of multi-year ice covering the continental shelf of the Chukchi Sea and in nearshore areas of the central and western Beaufort Sea. In the Beaufort Sea, bearded seals rarely use coastal haulouts.

In some areas, bearded seals are associated with the ice year-round; however, they usually move shoreward into open water areas when the pack ice retreats to areas with water depths greater than 200 m. During the summer, when the Bering Sea is ice-free, the most favorable bearded seal habitat is found in the central or northern Chukchi Sea along the margin of the pack ice. Suitable habitat is more limited in the Beaufort Sea where the continental shelf is narrower and the pack ice edge frequently occurs seaward of the shelf and over water too deep for benthic feeding. The preferred habitat in the western and central Beaufort Sea during the open-water period is the continental shelf seaward of the scour zone. WesternGeco conducted marine mammal monitoring during its open-water seismic program in the Alaskan Beaufort Sea from 1996 to 2001. Operations were conducted in nearshore waters, and of a total 454 seals that were identified to species while no guns were operating, 4.4% were bearded seals, 94.1% were ringed seals and 1.5% were spotted seals (Moulton and Lawson 2002). Haley and Ireland (2006) and Haley (2006) also reported much lower percentages of bearded compared to ringed seals during Healy cruises in the Arctic.

Small numbers of bearded seals would likely be encountered during the proposed geophysical survey. Bearded seals could also be encountered during transit periods in shallow areas closer to shore.

(c) Spotted Seal (*Phoca largha*)

Spotted seals (also known as largha seals) occur in the Beaufort, Chukchi, Bering and Okhotsk seas, and south to the northern Yellow Sea and western Sea of Japan (Shaughnessy and Fay 1977). They migrate south from the Chukchi Sea and through the Bering Sea in October (Lowry et al. 1998). Spotted seals overwinter in the Bering Sea and inhabit the southern margin of the ice during spring (Shaughnessy and Fay 1977).

In the Chukchi Sea, Kasegaluk Lagoon is an important area for spotted seals. Spotted seals haul out in the area from mid-July until freeze-up in late October or November. Frost and Lowry (1993) reported a maximum count of about 2200 spotted seals in the lagoon during aerial surveys. No spotted seals were recorded along the shore south of Pt. Lay. Based on satellite tracking data, Frost and Lowry (1993) reported that spotted seals at Kasegaluk Lagoon spent 94% of the time at sea. Extrapolating the count of hauled-out seals to account for seals at sea would suggest a Chukchi Sea population of about 36,000 animals.

An early estimate of the size of the world population of spotted seals was 370,000–420,000, and the size of the Bering Sea population, including animals in Russian waters, was estimated to be 200,000–250,000 animals (Bigg 1981). The total number of spotted seals in Alaskan waters is not known (Angliss and Allen 2009), but the estimate is most likely between several thousand and >50,000 (Rugh et al. 1997).

During the summer spotted seals are found in Alaska from Bristol Bay through western Alaska to the Chukchi and Beaufort seas. The ADF&G placed satellite transmitters on four spotted seals in Kasegaluk Lagoon and estimated that the proportion of seals hauled out was 6.8%. Based on an actual minimum count of 4145 hauled out seals, Angliss and Allen (2009) estimated the Alaskan population at 59,214 animals. The Alaska stock of spotted seals is not classified as endangered or as a strategic stock by NMFS (Hill and DeMaster 1998). CBD (2008a) recently petitioned to list the spotted seal under the ESA, however NMFS subsequently determined that the U.S. spotted seal populations did not warrant listing at this time (NMFS 2009).

During spring when pupping, breeding, and molting occur, spotted seals are found along the southern edge of the sea ice in the Okhotsk and Bering seas (Quakenbush 1988; Rugh et al. 1997). In late April and early May, adult spotted seals are often seen on the ice in female-pup or male-female pairs, or in male-female-pup triads. Subadults may be seen in larger groups of up to two hundred animals. During the summer, spotted seals are found primarily in the Bering and Chukchi seas, but some range into the

Beaufort Sea (Rugh et al. 1997; Lowry et al. 1998) from July until September. At this time of year, spotted seals haul out on land part of the time, but also spend extended periods at sea. Spotted seals are commonly seen in bays, lagoons and estuaries, but also range far offshore as far north as 69–72°N. Small numbers of spotted seals could occur near the southern portion of the proposed survey area, although in summer they are rarely seen on the pack ice except when the ice is very near shore. As the ice cover thickens with the onset of winter, spotted seals leave the northern portions of their range and move into the Bering Sea (Lowry et al. 1998).

Relatively low numbers of spotted seals are present in the Beaufort Sea. A small number of spotted seal haulouts are (or were) located in the central Beaufort Sea in the deltas of the Colville River and previously the Sagavanirktok River. Historically, these sites supported as many as 400–600 spotted seals, but in the 1990s <20 were seen at any one site (Johnson et al. 1999). A total of 12 spotted seals were positively identified near the source vessel during open-water seismic programs in the central Alaskan Beaufort Sea during the 6 years from 1996 to 2001 (Moulton and Lawson 2002). Numbers seen per year ranged from zero (in 1998 and 2000) to four (in 1999). More recently Green et al. (2007) reported 46 spotted seal sightings during barge operations between West Dock and Cape Simpson. Most sightings occurred from western Harrison Bay to Cape Simpson with only one sighting offshore of the Colville River delta. No spotted seals were recorded from the *Healy* during arctic cruises in 2005 or 2006 (Haley and Ireland 2006; Haley 2006). Spotted seals would be unlikely to occur in the proposed survey area in 2010 although some spotted seals could be encountered during transit periods.

(d) Ringed Seal (*Phoca hispida*)

Ringed seals have a circumpolar distribution and occur in all seas of the Arctic Ocean (King 1983). They are closely associated with ice, and in the summer they often occur along the receding ice edges or farther north in the pack ice. In the North Pacific, they occur in the southern Bering Sea and range south to the seas of Okhotsk and Japan. They are found throughout the Beaufort, Chukchi, and Bering seas (Angliss and Allen 2009).

During winter, ringed seals occupy landfast ice and offshore pack ice of the Bering, Chukchi and Beaufort seas. In winter and spring, the highest densities of ringed seals are found on stable shorefast ice. However, in some areas where there is limited fast ice but wide expanses of pack ice, including the Beaufort Sea, Chukchi Sea and Baffin Bay, total numbers of ringed seals on pack ice may exceed those on shorefast ice (Burns 1970; Stirling et al. 1982; Finley et al. 1983).

Ringed seals maintain breathing holes in the ice and occupy lairs in accumulated snow (Smith and Stirling 1975). They give birth in lairs from mid-March through April, nurse their pups in the lairs for 5–8 weeks, and mate in late April and May (Smith 1973; Hammill et al. 1991; Lydersen and Hammill 1993).

Ringed seals are year-round residents in the northern Chukchi and Beaufort seas and ringed seal is the most frequently encountered seal species in the area. No estimate for the size of the Alaska ringed seal stock is currently available (Angliss and Allen 2009). Past ringed seal population estimates in the Bering-Chukchi-Beaufort area ranged from 1–1.5 million (Frost 1985) to 3.3–3.6 million (Frost et al. 1988). Frost and Lowry (1981) estimated 80,000 ringed seals in the Beaufort Sea during summer and 40,000 during winter. More recent estimates based on extrapolation from aerial surveys and on predation estimates for polar bears (Amstrup 1995) suggest an Alaskan Beaufort Sea population of ~326,500 animals. During aerial surveys in 1999 and 2000, Bengtson et al. (2005) reported ringed seal densities 1.62 to 1.91 seals/km² in the eastern Chukchi Sea and estimated ringed seal abundance at >250,000 in the study area in 1999. The Alaska stock of ringed seals is not endangered, and is not classified as a strategic

stock by NMFS however there has recently been a petition to list this and other arctic seals due to the potential impact to seal habitats resulting from current warming trends (CBD 2008a).

Haley et al (2009) reported that ringed seal was the most abundant seal species during vessel-based surveys in the Chukchi Sea in 2006-2008 with densities up to 0.054 and 0.171 seals/km² in summer and fall, respectively. Savarese et al. (2009) also reported that ringed seal was the most abundant seal species in the Beaufort Sea during similar vessel-based surveys during the same period with densities up to 0.068 and 0.096 seals/km² in the summer and fall, respectively. Many unidentified seals during these surveys may have also been ringed seals and actual densities may have been higher.

Moulton et al. (2002) reported ringed seal densities (uncorrected) ranging from 0.43 to 0.63 seal per km² in water over 3 m in depth during spring aerial surveys in the central Alaskan Beaufort Sea. Densities were higher in nearshore than offshore locations. Ringed seal was the most frequently sighted seal identified to species from the *Healy* during arctic cruises in 2005 (3 sightings; Haley and Ireland 2006) and 2006 (10 sightings; Haley 2006). These sightings occurred over 361 km and 622 km of trackline, respectively within the latitudes of the proposed survey (71 -74 °N). Ringed seals likely would be encountered during the proposed geophysical survey.

(e) Ribbon Seal (*Histiophoca fasciata*)

Ribbon seals are found along the pack-ice margin in the southern Bering Sea during late winter and early spring and they move north as the pack ice recedes during late spring to early summer (Burns 1970; Burns et al. 1981). Little is known about their summer and fall distribution, but Kelly (1988) suggested that they move into the southern Chukchi Sea based on a review of sightings during the summer. During a recent satellite telemetry program sponsored by the National Marine Mammal Laboratory, a number of ribbon seals tagged in the Bering Sea in May had moved to the Chukchi Sea by July (NMML 2009). However, ribbon seals appeared to be relatively rare in the northern Chukchi Sea during recent vessel-based surveys in summer and fall of 2006-2009 with only three sightings among 1778 sightings of seals identified to species (Haley et al. 2009). Ribbon seals do not normally occur in the Beaufort Sea however three recent ribbon seal sightings were reported during vessel-based activities in the Beaufort Sea in 2007-2008 (Savarese et al 2009).

In response to a petition to list ribbon seal under the Endangered Species Act (CBD 2007), a recent announcement by NMFS indicated that listing of ribbon seal was not warranted at this time (NMFS 2008a). Ribbon seals were not reported during the arctic *Healy* cruises in 2005 and 2006, and would be unlikely to occur in the proposed survey area.

(4) Carnivora

(a) Polar Bear (*Ursus maritimus*)

Polar bears have a circumpolar distribution throughout the northern hemisphere (Amstrup et al. 1986) and occur in relatively low densities throughout most ice-covered areas (DeMaster and Stirling 1981). Polar bears are divided into 19 relatively distinct populations or management units although there may be overlap of some individuals among populations (Aars et al. 2006; USFWS 2008). Polar bears are common in the Chukchi and Beaufort Seas north of Alaska throughout the year, including the late summer period (Garner et al. 1990, Amstrup and Gardner 1994, Amstrup et al. 2000, Moulton and Williams 2003, Harwood et al. 2005). They also occur throughout the East Siberian, Laptev, and Kara Seas of Russia and the Barent's Sea of northern Europe. They are found in the northern part of the Greenland Sea, and are common in Baffin Bay, which separates Canada and Greenland, as well as through most of the Canadian Arctic Archipelago.

Current world population estimates for the polar bear range from ~20,000 to 30,000 animals (Derocher et al. 1998; Aars et al. 2006). Three polar bear populations are of concern for the proposed

geophysical survey. The Southern Beaufort Sea population with ~1500 bears ranges from the Baillie Islands, Canada, in the east to near Point Lay, Alaska, in the west. The Chukchi Sea population with ~2000 bears is found in much of the Chukchi Sea and the northern Bering Sea. The Northern Beaufort Sea population with ~1200 bears occurs in Canadian waters primarily north of the Southern Beaufort Sea and extending into Admunsen Gulf. USFWS (2008) designated the Northern Beaufort Sea population as stable, the Southern Beaufort Sea population as declining, and the Chukchi Sea population as data deficient. Data from tracking studies indicate wide-ranging movements of individual bears and overlap among polar bear populations (Garner et al. 1990; Amstrup 1995; Durner and Amstrup 1995).

Polar bear populations are protected under the Marine Mammal Protection Act of 1973, as well as by the International Agreement on the Conservation of Polar Bears, ratified in 1976. Countries participating in the latter treaty include Canada, Denmark, Norway, Russia (former USSR), and the USA. Article II of the agreement states, “Each contracting party...shall manage polar bear populations in accordance with sound conservation practices based on the best scientific data.”

USFWS (2008) listed polar bear as a threatened species under the U.S. ESA based on the expected continuation of declines in sea ice which is their principal habitat. No critical habitat for polar bears has as yet been officially defined, however USFWS (2009a) proposed designation of polar bear critical habitat to include sea ice over marine waters 300 m (984.2 ft) or less in depth that occur over the continental shelf. The deadline for final determination of the proposed critical habitat designation is 30 June 2010.

Polar bears usually forage in areas where there are high concentrations of ringed seal, which is their primary prey, and bearded seals (Larsen 1985; Stirling and McEwan 1975). This includes areas of land-fast ice, as well as moving pack ice. Polar bears are opportunistic feeders and feed on a variety of foods and carcasses including not only seals but also beluga whales, arctic cod, geese and their eggs, walrus, bowhead whales, and reindeer (Smith 1985; Jefferson et al. 1993; Smith and Hill 1996; Derocher et al. 2000).

Females give birth to 1 to 3 cubs at an average interval of every 3.6 years (Jefferson et al. 1993; Lentfer et al. 1980). Cubs remain with their mothers for 1.4 to 3.4 years (Derocher et al. 1993; Ramsay and Stirling 1988). Mating occurs from April to June followed by a delayed implantation during September to December. Females give birth usually the following December or January (Harington 1968; Jefferson et al. 1993). In general, females 6 years of age or older successfully wean more cubs than younger bears; however, females as young as 4 years old can produce offspring (Ramsay and Stirling 1988). An examination of reproductive rates of polar bears indicated that 5% of four-year-old females had cubs, whereas 50% of five year-old females had cubs (Ramsay and Stirling 1988). Females that were over 20 years had a very high rate of cub loss or did not successfully reproduce. The maximum reproductive age reported for Alaskan polar bears is 18 years (Amstrup and DeMaster 1988).

Polar bears typically range as far as 88°N (Ray 1971; Durner and Amstrup 1995) where the population thins dramatically. However, polar bears have been observed across the Arctic, including close to the North Pole (van Meurs and Splettstoesser 2003). Twenty-one sightings of 27 polar bears were made during the *Healy* cruise in 2005 (Haley and Ireland 2006). Most sightings were recorded between ~80 and 82°N latitude with one sighting at ~87°N. Proposed survey activities within US waters will occur between 71°N and 74°10'N where only one polar bear sightings of two individuals was recorded along ~2308 km of monitored trackline between 2005 and 2009 (Haley and Ireland 2006, Haley 2006, GSC unpubl. data 2008). Small numbers of polar bears will likely be encountered during the proposed geophysical survey.

IV. ENVIRONMENTAL CONSEQUENCES OF PROPOSED ACTION

Direct Effects on Marine Mammals and their Significance

The material in this section includes a summary of the anticipated effects (or lack thereof) on marine mammals of the medium-sized airgun source (three G-guns with a total discharge volume of 1150 in³) to be used during the proposed geophysical survey. A more detailed review of airgun effects on marine mammals appears in Appendix D. That appendix was recently updated and is similar to corresponding parts of previous EAs and associated IHA applications concerning seismic survey projects in the following areas: northern Gulf of Mexico; Hess Deep (eastern tropical Pacific); Norwegian Sea; Mid-Atlantic Ocean; Bermuda; SE Caribbean; southern Gulf of Mexico (Yucatan Peninsula); SE Alaska; Blanco Fracture Zone (northeast Pacific); off the Pacific coast of Central America; the Aleutian Islands, Alaska; and across the Arctic Ocean. The number of airguns used during recent industry exploratory activities in the Chukchi and Beaufort seas has ranged from 16 to 36, and array volume has ranged from 3147 to 3390 in³. Due to the size and configuration of the three-airgun, 1150 in³ source to be used in the present work, its effective size will be reduced compared to larger arrays used in the above projects, and anticipated impacts to marine mammals will likely also be somewhat reduced. This section also includes a discussion of the potential impacts of operations by bathymetric echo sounders and Chirp echo sounder.

Finally, this section includes estimates of the numbers of marine mammals that might be affected by the proposed geophysical survey in the Arctic Ocean in 2010. This section includes a description of the rationale for USGS's estimates of the potential numbers of harassment "takes" during the planned seismic survey.

(1) Summary of Potential Effects of Airgun Sounds

The effects of sounds from airguns might include one or more of the following: tolerance, masking of natural sounds, behavioral disturbance, and at least in theory, temporary or permanent hearing impairment, or non-auditory physical effects (Richardson et al. 1995). Given the moderate size of the sources planned for the proposed project, plus mitigation measures to be applied, it is unlikely that there would be any cases of temporary or especially permanent hearing impairment, or non-auditory physical effects. Also, behavioral disturbance could occur at longer distances than auditory effects.

Tolerance

Numerous studies have shown that pulsed sounds from airguns are often readily detectable in the water at distances of many kilometers. For a summary of the characteristics of airgun pulses, see Appendix D (3).

Numerous studies have shown that marine mammals at distances more than a few kilometers from operating seismic vessels often show no apparent response—see Appendix D (5). That is often true even in cases when the pulsed sounds must be readily audible to the animals based on measured received levels and the hearing sensitivity of that mammal group. Although various baleen whales, toothed whales, and (less frequently) pinnipeds have been shown to react behaviorally to airgun pulses under some conditions, at other times mammals of all three types have shown no overt reactions. In general, pinnipeds, small odontocetes, and sea otters seem to be more tolerant of exposure to airgun pulses than are baleen whales.

Masking

Masking effects of pulsed sounds (even from large arrays of airguns) on marine mammal calls and other natural sounds are expected to be limited, although there are very few specific data of relevance. Some whales are known to continue calling in the presence of seismic pulses. Their calls can be heard

between the seismic pulses (e.g., Richardson et al. 1986; McDonald et al. 1995; Greene et al. 1999; Nieukirk et al. 2004). Although there has been one report that sperm whales cease calling when exposed to pulses from a very distant seismic ship (Bowles et al. 1994), a more recent study reports that sperm whales off northern Norway continued calling in the presence of seismic pulses (Madsen et al. 2002). That has also been shown during recent work in the Gulf of Mexico (Tyack et al. 2003). Masking effects of seismic pulses are expected to be negligible in the case of the smaller odontocete cetaceans. Also, the sounds important to small odontocetes are predominantly at much higher frequencies than are airgun sounds. Masking effects, in general, are discussed further in Appendix D (4).

Disturbance Reactions

Disturbance includes a variety of effects, including subtle changes in behavior, more conspicuous changes in activities, and displacement. Based on NMFS (2001, p. 9293), we assume that simple exposure to sound, or brief reactions that do not disrupt behavioral patterns in a potentially significant manner, do not constitute harassment or “taking”. By potentially significant, we mean “in a manner that might have deleterious effects to the well-being of individual marine mammals or their populations”.

Reactions to sound, if any, depend on species, state of maturity, experience, current activity, reproductive state, time of day, and many other factors. If a marine mammal does react briefly to an underwater sound by changing its behavior or moving a small distance, the impacts of the change are unlikely to be significant to the individual, let alone the stock or the species as a whole. However, if a sound source displaces marine mammals from an important feeding or breeding area for a prolonged period, impacts on the animals could be significant. Given the many uncertainties in predicting the quantity and types of impacts of noise on marine mammals, it is common practice to estimate how many mammals were present within a particular distance of industrial activities, or exposed to a particular level of industrial sound. That likely overestimates the numbers of marine mammals that are affected in some biologically-important manner.

The sound criteria used to estimate how many marine mammals might be disturbed to some biologically-important degree by a seismic program are based on behavioral observations during studies of several species. However, information is lacking for many species. Detailed studies have been done on humpback, gray, and bowhead whales, and on ringed seals. Less detailed data are available for some other species of baleen whales, sperm whales, small toothed whales, and sea otters.

Baleen Whales—Baleen whales generally tend to avoid operating airguns, but avoidance radii are quite variable. Whales are often reported to show no overt reactions to pulses from large arrays of airguns at distances beyond a few kilometers, even though the airgun pulses remain well above ambient noise levels out to much longer distances. However, as reviewed in Appendix D (5), baleen whales exposed to strong noise pulses from airguns often react by deviating from their normal migration route and/or interrupting their feeding and moving away. In the case of the migrating gray and bowhead whales, the observed changes in behavior appeared to be of little or no biological consequence to the animals. They simply avoided the sound source by displacing their migration route to varying degrees, but within the natural boundaries of the migration corridors.

Studies of gray, bowhead, and humpback whales have determined that received levels of pulses in the 160–170 dB re 1 μ Pa rms range seem to cause obvious avoidance behavior in a substantial fraction of the animals exposed. In many areas, seismic pulses from large arrays of airguns diminish to those levels at distances ranging from 4.5 to 14.5 km from the source. A substantial proportion of the baleen whales within those distances may show avoidance or other strong disturbance reactions to the airgun array. Subtle behavioral changes sometimes become evident at somewhat lower received levels, and recent studies reviewed in Appendix D (5) have shown that some species of baleen whales, notably bowhead

and humpback whales, at times show strong avoidance at received levels lower than 160–170 dB re 1 μ Pa rms. Bowhead whales migrating west across the Alaskan Beaufort Sea in autumn, in particular, are unusually responsive, with substantial avoidance occurring out to distances of 20–30 km from a medium-sized airgun source (Miller et al. 1999; Richardson et al. 1999; see Appendix D [5]). However, more recent research on bowhead whales (Miller et al. 2005; Lyons et al. 2009; Christi et al. 2009) corroborates earlier evidence that, during the summer feeding season, bowheads are not as sensitive to seismic sources. In summer, bowheads typically begin to show avoidance reactions at a received level of about 160–170 dB re 1 μ Pa rms (Richardson et al. 1986; Ljungblad et al. 1988; Miller et al. 1999). The USGS project will be conducted primarily during fall migration at locations > 200 n.mi. offshore, well north of the known bowhead migration corridor. Recent evidence suggests that some bowheads feed during migration and feeding bowheads might be encountered in the central Alaskan Beaufort Sea during transit periods to and from Barrow (Lyons et al. 2009; Christi et al. 2009). The primary bowhead summer feeding grounds, however, are far to the east in the Canadian Beaufort Sea.

Malme et al. (1986, 1988) studied the responses of feeding eastern gray whales to pulses from a single 100 in³ airgun off St. Lawrence Island in the northern Bering Sea. They estimated, based on small sample sizes, that 50% of feeding gray whales ceased feeding at an average received pressure level of 173 dB re 1 μ Pa on an (approximate) rms basis, and that 10% of feeding whales interrupted feeding at received levels of 163 dB. Those findings were generally consistent with the results of experiments conducted on larger numbers of gray whales that were migrating along the California coast, and on observations of Western Pacific gray whales feeding off Sakhalin Island, Russia (Johnson 2002).

Data on short-term reactions (or lack of reactions) of cetaceans to impulsive noises do not necessarily provide information about long-term effects. It is not known whether impulsive noises affect reproductive rate or distribution and habitat use in subsequent days or years. However, gray whales continued to migrate annually along the west coast of North America despite intermittent seismic exploration and much ship traffic in that area for decades (Appendix A in Malme et al. 1984). Bowhead whales continued to travel to the eastern Beaufort Sea each summer despite seismic exploration in their summer and autumn range for many years (Richardson et al. 1987). Populations of both gray whales and bowhead whales grew substantially during this time. In any event, because the airguns are located on a moving ship, the brief exposures to sound pulses from the proposed airgun source are highly unlikely to result in prolonged effects.

Toothed Whales—Little systematic information is available about reactions of toothed whales to noise pulses. Few studies similar to the more extensive baleen whale/seismic pulse work summarized above and in Appendix D have been reported for toothed whales. However, systematic work on sperm whales is underway (Tyack et al. 2003), and there is an increasing amount of information about responses of various odontocetes to seismic surveys based on monitoring studies (e.g., Stone 2003; Smultea et al. 2004; Moulton and Miller 2005).

Seismic operators sometimes see dolphins and other small toothed whales near operating airgun arrays, but in general there seems to be a tendency for most delphinids to show some limited avoidance of seismic vessels operating large airgun systems. However, some dolphins seem to be attracted to the seismic vessel and floats, and some ride the bow wave of the seismic vessel even when large arrays of airguns are firing. Nonetheless, there have been indications that small toothed whales sometimes move away, or maintain a somewhat greater distance from the vessel, when a large array of airguns is operating than when it is silent (e.g., Goold 1996a,b,c; Calambokidis and Osmeck 1998; Stone 2003). Aerial surveys during seismic operations in the southeastern Beaufort Sea recorded much lower sighting rates of beluga whales within 10-20 km of an active seismic vessel. These results were consistent with the low number of

beluga sightings reported by observers aboard the seismic vessel, suggesting that some belugas might be avoiding the seismic operations at distances of 10- 20 km (Miller et al. 2005).

Similarly, captive bottlenose dolphins and (of some relevance in this project) beluga whales exhibit changes in behavior when exposed to strong pulsed sounds similar in duration to those typically used in seismic surveys (Finneran et al. 2000, 2002). However, the animals tolerated high received levels of sound (pk–pk level >200 dB re 1 μ Pa) before exhibiting aversive behaviors.

Odontocete reactions to large arrays of airguns are variable and, at least for small odontocetes, seem to be confined to a smaller radius than has been observed for mysticetes (Appendix B). A ≥ 170 dB disturbance criterion (rather than ≥ 160 dB) may be more appropriate for small odontocetes (and pinnipeds) which tend to be less responsive than other cetaceans. However, based on the limited existing evidence, belugas should not be grouped with delphinids in the “less responsive” category.

Pinnipeds—Pinnipeds are not likely to show a strong avoidance reaction to the medium-sized airgun source that will be used. Visual monitoring from seismic vessels has shown only slight (if any) avoidance of airguns by pinnipeds, and only slight (if any) changes in behavior—see Appendix D (5). Those studies show that pinnipeds frequently do not avoid the area within a few hundred meters of operating airgun arrays (e.g., Miller et al. 2005; Harris et al. 2001). However, initial telemetry work suggests that avoidance and other behavioral reactions to small airgun sources may at times be stronger than evident to date from visual studies of pinniped reactions to airguns (Thompson et al. 1998). Even if reactions of the species occurring in the proposed survey area are as strong as those evident in the telemetry study, reactions are expected to be confined to relatively small distances and durations, with no long-term effects on pinniped individuals or populations.

Polar Bears—Airgun effects on polar bears have not been studied. However, polar bears on the ice would be unaffected by underwater sound. Sound levels received by polar bears in the water would be attenuated because polar bears generally do not dive much below the surface. Received levels of airgun sounds are reduced near the surface because of the pressure release effect at the water’s surface (Greene and Richardson 1988; Richardson et al. 1995).

Hearing Impairment and Other Physical Effects

Temporary or permanent hearing impairment is a possibility when marine mammals are exposed to very strong sounds, but there has been no specific documentation of this for marine mammals exposed to sequences of airgun pulses. Current NMFS policy regarding exposure of marine mammals to high-level sounds is that cetaceans and pinnipeds should not be exposed to impulsive sounds ≥ 180 and 190 dB re 1 μ Pa (rms), respectively (NMFS 2000). These exposure levels have also been applied by the USFWS to walrus and polar bear, respectively. Those criteria have been used in defining the safety (shutdown) radii planned for the proposed seismic survey. However, those criteria were established before there were any data on the minimum received levels of sounds necessary to cause temporary auditory impairment in marine mammals. As discussed in Appendix D (6) and summarized here:

- the 180 dB criterion for cetaceans is probably quite precautionary, i.e., lower than necessary to avoid temporary threshold shift (TTS), let alone permanent auditory injury, at least for belugas and delphinids.
- the minimum sound level necessary to cause permanent hearing impairment is higher, by a variable and generally unknown amount, than the level that induces barely-detectable TTS.
- the level associated with the onset of TTS is often considered to be a level below which there is no danger of permanent damage.

NMFS is presently developing new noise exposure criteria for marine mammals that account for the now-available scientific data on TTS and other relevant factors in marine and terrestrial mammals (NMFS 2005; D. Wieting *in* <http://mmc.gov/sound/plenary2/pdf/plenary2summaryfinal.pdf>).

Several aspects of the planned monitoring and mitigation measures for this project are designed to detect marine mammals occurring near the airguns (and multi-beam bathymetric echo sounder), and to avoid exposing them to sound pulses that might, at least in theory, cause hearing impairment [see § II(3), MITIGATION MEASURES]. In addition, many cetaceans are likely to show some avoidance of the area with high received levels of airgun sound (see above). In those cases, the avoidance responses of the animals themselves will reduce or (most likely) avoid any possibility of hearing impairment.

Non-auditory physical effects might also occur in marine mammals exposed to strong underwater pulsed sound. Possible types of non-auditory physiological effects or injuries that theoretically might occur in mammals close to a strong sound source include stress, neurological effects, bubble formation, resonance effects, and other types of organ or tissue damage. It is possible that some marine mammal species (i.e., beaked whales) may be especially susceptible to injury and/or stranding when exposed to strong pulsed sounds. However, as discussed below, there is no definitive evidence that any of these effects occur even for marine mammals in close proximity to large arrays of airguns and beaked whales do not occur in the present study area. It is unlikely that any effects of these types would occur during the present project given the brief duration of exposure of any given mammal, and the planned monitoring and mitigation measures (see below). The following subsections discuss in somewhat more detail the possibilities of TTS, permanent threshold shift (PTS), and non-auditory physical effects.

Temporary Threshold Shift (TTS).—TTS is the mildest form of hearing impairment that can occur during exposure to a strong sound (Kryter 1985). While experiencing TTS, the hearing threshold rises and a sound must be stronger in order to be heard. TTS can last from minutes or hours to (in cases of strong TTS) days. For sound exposures at or somewhat above the TTS threshold, hearing sensitivity recovers rapidly after exposure to the noise ends. Only a few data on sound levels and durations necessary to elicit mild TTS have been obtained for marine mammals, and none of the published data concern TTS elicited by exposure to multiple pulses of sound.

For toothed whales exposed to single short pulses, the TTS threshold appears to be, to a first approximation, a function of the energy content of the pulse (Finneran et al. 2005, 2002). Given the available data, the received level of a single seismic pulse might need to be ~210 dB re 1 μ Pa rms (~221–226 dB pk–pk) in order to produce brief, mild TTS. Exposure to several seismic pulses at received levels near 200–205 dB (rms) might result in slight TTS in a small odontocete, assuming the TTS threshold is (to a first approximation) a function of the total received pulse energy. Seismic pulses with received levels of 200–205 dB or more are usually restricted to a radius of no more than 200 m around a seismic vessel operating a large array of airguns.

For baleen whales, there are no data, direct or indirect, on levels or properties of sound that are required to induce TTS. However, no cases of TTS are expected given the moderate size of the source, and the strong likelihood that baleen whales would avoid the approaching airguns (or vessel) before being exposed to levels high enough for there to be any possibility of TTS.

In pinnipeds, TTS thresholds associated with exposure to brief pulses (single or multiple) of underwater sound have not been measured. Initial evidence from prolonged exposures suggested that some pinnipeds may incur TTS at somewhat lower received levels than do small odontocetes exposed for similar durations (Kastak et al. 1999; Ketten et al. 2001; *cf.* Au et al. 2000). For harbor seal, which is closely related to the ringed seal, TTS onset apparently occurs at somewhat lower received energy levels than for odontocetes.

A marine mammal within a radius of ≤ 100 m (≤ 328 ft) around a typical large array of operating airguns might be exposed to a few seismic pulses with levels of ≥ 205 dB, and possibly more pulses if the mammal moved with the seismic vessel. The received sound levels will be reduced for the proposed three-gun array to be used during the current survey compared to the larger arrays thus reducing the potential for TTS for the proposed survey. (As noted above, most cetacean species tend to avoid operating airguns, although not all individuals do so.) However, several of the considerations that are relevant in assessing the impact of typical seismic surveys with airgun arrays are directly applicable here:

- “Ramping up” (soft start) is standard operational protocol during startup of airgun arrays in many jurisdictions. Ramping up involves starting the airguns in sequence, usually commencing with a single airgun and gradually adding additional airguns. This practice will be employed when the airgun array is operated during the proposed survey.
- It is unlikely that cetaceans would be exposed to airgun pulses at a sufficiently high level for a sufficiently long period to cause more than mild TTS, given the relative movement of the vessel and the marine mammal. For the current project the entire seismic survey will be in deep water where the radius of influence and duration of exposure to strong pulses is smaller.
- With a large array of airguns, TTS would be most likely to occur in odontocetes that bow-ride or otherwise linger near the airguns. However, no species that occur within the project area are expected to bow-ride.

NMFS (1995, 2000) concluded that cetaceans and pinnipeds should not be exposed to pulsed underwater noise at received levels exceeding, respectively, 180 and 190 dB re 1 μ Pa (rms). The predicted 180 and 190 dB distances for the airguns operated by USGS vary with water depth, however the proposed geophysical survey will be conducted entirely in deep water where sound levels are generally reduced compared to operations in shallower conditions. Furthermore, those sound levels are *not* considered to be the levels above which TTS might occur. Rather, they are the received levels above which, in the view of a panel of bioacoustics specialists convened by NMFS before TTS measurements for marine mammals started to become available, one could not be certain that there would be no injurious effects, auditory or otherwise, to marine mammals. As summarized above, data that are now available imply that TTS is unlikely to occur unless odontocetes are exposed to airgun pulses much stronger than 180 dB re 1 μ Pa rms and since no bow-riding species occur in the study area, it is unlikely such exposures will occur.

Permanent Threshold Shift (PTS).—When PTS occurs, there is physical damage to the sound receptors in the ear. In some cases, there can be total or partial deafness, whereas in other cases, the animal has an impaired ability to hear sounds in specific frequency ranges.

There is no specific evidence that exposure to pulses of airgun sound can cause PTS in any marine mammal, even with large arrays of airguns. However, given the possibility that mammals close to an airgun array might incur TTS, there has been further speculation about the possibility that some individuals occurring very close to airguns might incur PTS. Single or occasional occurrences of mild TTS are not indicative of permanent auditory damage in terrestrial mammals. Relationships between TTS and PTS thresholds have not been studied in marine mammals, but are assumed to be similar to those in humans and other terrestrial mammals. PTS might occur at a received sound level at least several decibels above that inducing mild TTS if the animal were exposed to the strong sound pulses with very rapid rise time—see Appendix D (6).

It is highly unlikely that marine mammals could receive sounds strong enough (and over a sufficient duration) to cause permanent hearing impairment during a project employing the medium-sized airgun sources planned here. For the proposed project, marine mammals are unlikely to be exposed to received levels of

seismic pulses strong enough to cause TTS. Marine mammals would probably need to be within 100-200 meters of the airguns and be exposed for some time period for TTS to occur. Given the higher level of sound necessary to cause PTS, it is even less likely that PTS could occur. In fact, even the levels immediately adjacent to the airgun may not be sufficient to induce PTS, especially because a mammal would not be exposed to more than one strong pulse unless it swam immediately alongside the airgun for a period longer than the inter-pulse interval. Baleen whales generally avoid the immediate area around operating seismic vessels. The planned monitoring and mitigation measures, including visual monitoring, power downs, and shut downs of the airguns when mammals are seen within the “safety radii”, will minimize the already-minimal probability of exposure of marine mammals to sounds strong enough to induce PTS.

Non-auditory Physiological Effects.—Non-auditory physiological effects or injuries that theoretically might occur in marine mammals exposed to strong underwater sound include stress, neurological effects, bubble formation, and other types of organ or tissue damage. However, studies examining such effects are very limited. If any such effects do occur, they probably would be limited to unusual situations when animals might be exposed at close range for unusually long periods. It is doubtful that any single marine mammal would be exposed to strong seismic sounds for a sufficient period of time that significant physiological stress would develop. That is especially so in the case of the proposed project where the airgun configuration is moderately sized, the ship is moving at speeds up to ~5 knots, and the tracklines will not “double back” through the same area.

Until recently, it was assumed that diving marine mammals are not subject to the bends or air embolism. This possibility was first explored at a workshop (Gentry [ed.] 2002) held to discuss whether the stranding of beaked whales in the Bahamas in 2000 (Balcomb and Claridge 2001; NOAA and USN 2001) might have been related to bubble formation in tissues caused by exposure to noise from naval sonar. However, the opinions were inconclusive. Jepson et al. (2003) first suggested a possible link between mid-frequency sonar activity and acute and chronic tissue damage that results from the formation *in vivo* of gas bubbles, based on the beaked whale stranding in the Canary Islands in 2002 during naval exercises. Fernández et al. (2005a) showed those beaked whales did indeed have gas bubble-associated lesions as well as fat embolisms. Fernández et al. (2005b) also found evidence of fat embolism in three beaked whales that stranded 100 km north of the Canaries in 2004 during naval exercises. Examinations of several other stranded species have also revealed evidence of gas and fat embolisms (e.g., Arbelo et al. 2005; Jepson et al. 2005a; Méndez et al. 2005). Most of the afflicted species were deep divers. There is speculation that gas and fat embolisms may occur if cetaceans ascend unusually quickly when exposed to aversive sounds, or if sound in the environment causes the destabilization of existing bubble nuclei (Potter 2004; Arbelo et al. 2005; Fernández et al. 2005a; Jepson et al. 2005b). Even if gas and fat embolisms can occur during exposure to mid-frequency sonar, there is no evidence that that type of effect occurs in response to airgun sounds. Also, most evidence for such effects has been related to beaked whales, which do not occur in the proposed study area.

In general, little is known about the potential for seismic survey sounds to cause auditory impairment or other physical effects in marine mammals. Available data suggest that such effects, if they occur at all, would be limited to short distances and probably to projects involving large arrays of airguns. However, the available data do not allow for meaningful quantitative predictions of the numbers (if any) of marine mammals that might be affected in those ways. Marine mammals that show behavioral avoidance of seismic vessels, including most baleen whales, some odontocetes (including belugas), and some pinnipeds, are especially unlikely to incur auditory impairment or other physical effects. Also, the planned monitoring and mitigation measures include shut downs of the airguns, which will reduce any such effects that might otherwise occur.

Strandings and Mortality

Marine mammals close to underwater detonations of high explosive can be killed or severely injured, and the auditory organs are especially susceptible to injury (Ketten et al. 1993; Ketten 1995). Airgun pulses are less energetic and have slower rise times, and there is no proof that they can cause serious injury, death, or stranding even in the case of large airgun arrays. However, the association of mass strandings of beaked whales with naval exercises and, in one case, an L-DEO seismic survey, has raised the possibility that beaked whales exposed to strong pulsed sounds may be especially susceptible to injury and/or behavioral reactions that can lead to stranding. Appendix D (6.3) provides additional details.

Seismic pulses and mid-frequency sonar pulses are quite different. Sounds produced by airgun arrays are broadband with most of the energy below 1 kHz. Typical military mid-frequency sonars operate at frequencies of 2–10 kHz, generally with a relatively narrow bandwidth at any one time. Thus, it is not appropriate to assume that there is a direct connection between the effects of military sonar and seismic surveys on marine mammals. However, evidence that sonar pulses can, in special circumstances, lead to physical damage and mortality (NOAA and USN 2001; Jepson et al. 2003; Fernández et al. 2005a), even if only indirectly, suggests that caution is warranted when dealing with exposure of marine mammals to any high-intensity pulsed sound.

In May 1996, 12 Cuvier's beaked whales stranded along the coasts of Kyparissiakos Gulf in the Mediterranean Sea. That stranding was subsequently linked to the use of low- and medium-frequency (250-3000 Hz) active sonar by a North Atlantic Treaty Organization (NATO) research vessel in the region (Frantzis 1998). In March 2000, a population of Cuvier's beaked whales being studied in the Bahamas disappeared after a U.S. Navy task force using mid-frequency tactical sonars passed through the area; some beaked whales stranded (Balcomb and Claridge 2001; NOAA and USN 2001).

In September 2002, a total of 14 beaked whales of various species stranded coincident with naval exercises in the Canary Islands (Martel n.d.; Jepson et al. 2003; Fernández et al. 2003). Also in September 2002, there was a stranding of two Cuvier's beaked whales in the Gulf of California, Mexico, when the L-DEO vessel *Maurice Ewing* was operating a 20-airgun, 8490 in³ array in the general area. The link between the stranding and the seismic surveys was inconclusive and not based on any physical evidence (Hogarth 2002; Yoder 2002). Nonetheless, that plus the incidents involving beaked whale strandings near naval exercises suggests a need for caution in conducting seismic surveys in areas occupied by beaked whales. However, no beaked whales are found within this project area and the planned monitoring and mitigation measures are expected to minimize any possibility for mortality of other species.

(2) Possible Effects of Multibeam Echo Sounder Signals

A Kongsberg 2112 multibeam 12 kHz echo sounder system will be operated from the *Healy* almost continuously during the planned geophysical survey. Details about the Kongsberg 2112 were provided in Section II. Sounds from the multibeam are very short pulses, depending on water depth. Most of the energy in the sound pulses emitted by the multibeam is at moderately high frequencies, centered at 12 kHz. The beam is narrow (~2°) in fore-aft extent and wide (~130°) in the cross-track extent. Any given mammal at depth near the trackline would be in the main beam for only a fraction of a second. Therefore, marine mammals that encounter the Kongsberg 2112 at close range are unlikely to be subjected to repeated pulses because of the narrow fore-aft width of the beam, and will receive only limited amounts of pulse energy because of the short pulses. Similarly, Kremser et al. (2005) noted that the probability of a cetacean swimming through the area of exposure when a multibeam echo sounder emits a pulse is small. The animal would have to pass the transducer at close range and be swimming at speeds similar to the vessel in order to be subjected to sound levels that could cause TTS. In 2008 and 2009 the *Louis S. St.*

Laurent and the *Healy* surveyed together with a cooperative strategy similar to that proposed for 2010. The director of NOAA's Office of Ocean Exploration and Research deemed that the use of the *Healy's* multibeam would not have significant impacts on marine mammals of a direct or cumulative nature. The U.S. portions of the projects were granted a categorical exclusion from the need to prepare an Environmental Assessment (Appendices A and B).

Navy sonars that have been linked to avoidance reactions and stranding of cetaceans (1) generally are more powerful than the Kongsberg 2112 echo sounder, (2) have longer pulse duration, and (3) are directed close to horizontally vs. downward for the Kongsberg 2112. The area of possible influence of the multibeam echo sounder is much smaller—a narrow band oriented in the cross-track direction below the source vessel. Marine mammals that encounter the multibeam echo sounder at close range are unlikely to be subjected to repeated pulses because of the narrow fore-aft width of the beam, and will receive only small amounts of pulse energy because of the short pulses. In assessing the possible impacts of a 15.5 kHz Atlas Hydrosweep multibeam echo sounder, Boebel et al. (2004) noted that the critical sound pressure level at which TTS may occur is 203.2 dB re 1 μ Pa (rms). The critical region included an area of 43 m in depth, 46 m wide athwartship, and 1 m fore-and-aft (Boebel et al. 2004). In the more distant parts of that (small) critical region, only slight TTS would be incurred.

Masking

Marine mammal communications will not be masked appreciably by the multibeam echo sounder signals given the low duty cycle of the echo sounder and the brief period when an individual mammal is likely to be within the echo sounder's beam. Furthermore, the 12 kHz multibeam will not overlap with the predominant frequencies in baleen whale calls, further reducing any potential for masking in that group.

Behavioral Responses

Behavioral reactions of free-ranging marine mammals to military and other sonars appear to vary by species and circumstance. Observed reactions have included silencing and dispersal by sperm whales (Watkins et al. 1985), increased vocalizations and no dispersal by pilot whales (Rendell and Gordon 1999), and the previously-mentioned beachings by beaked whales. Also, Navy personnel have described observations of dolphins bow-riding adjacent to bow-mounted mid-frequency sonars during sonar transmissions. During exposure to a 21–25 kHz whale-finding sonar with a source level of 215 dB re 1 μ Pa·m, gray whales showed slight avoidance (~200 m) behavior (Frankel 2005).

However, all of those observations are of limited relevance to the present situation. Pulse durations from the Navy sonars were much longer than those of the multibeam echo sounders to be used during the proposed study, and a given mammal would have received many pulses from the naval sonars. During USGS's operations, the individual pulses will be very short, and a given mammal would not receive many of the downward-directed pulses as the vessel passes by.

Captive bottlenose dolphins and a white whale exhibited changes in behavior when exposed to 1 s pulsed sounds at frequencies similar to those that will be emitted by the multibeam echo sounder to be used by USGS, and to shorter broadband pulsed signals. Behavioral changes typically involved what appeared to be deliberate attempts to avoid the sound exposure (Schlundt et al. 2000; Finneran et al. 2002; Finneran and Schlundt 2004). The relevance of those data to free-ranging odontocetes is uncertain, and in any case, the test sounds were quite different in either duration or bandwidth as compared with those from a multibeam echo sounder.

We are not aware of any data on the reactions of pinnipeds to echo sounder sounds at frequencies similar to those of the multibeam echo sounder (12 kHz). Based on observed pinniped responses to other

types of pulsed sounds, and the likely brevity of exposure to the multibeam echo sounder sounds, pinniped reactions to the echo sounder sounds are expected to be limited to startle or otherwise brief responses of no lasting consequence to the animals.

Polar bears would not occur below the *Healy* or elsewhere at sufficient depth to be in the main beam of the multibeam echo sounder, so would not be affected by the echo sounder sounds.

As noted earlier, NMFS (2001) has concluded that momentary behavioral reactions “do not rise to the level of taking”. Thus, brief exposure of cetaceans or pinnipeds to small numbers of signals from a multibeam multibeam echo sounder system would not result in a “take” by harassment.

Hearing Impairment and Other Physical Effects

Given recent stranding events that have been associated with the operation of naval sonar, there is concern that mid-frequency sonar sounds can cause serious impacts to marine mammals (see above). However, the multibeam echo sounder proposed for use by USGS is quite different from sonars used for navy operations. Pulse duration of the multibeam echo sounder is very short relative to the naval sonars. Also, at any given location, an individual cetacean or pinniped would be in the beam of the multibeam sonar for much less time given the generally downward orientation of the beam and its narrow fore-aft beamwidth. (Navy sonars often use near-horizontally-directed sound.) Those factors would all reduce the sound energy received from the bathymetric echo sounder relative to that from the sonars used by the Navy. Polar bears would not occur in the main beam of the echo sounder.

(3) Possible Effects of Chirp Echo Sounder Signals

A Knudsen 320BR Plus echo sounder will be operated from the *Louis S. St. Laurent* at nearly all times during the planned study. The Knudsen 320BR produces sound pulses with lengths of up to 24 ms every 0.5 to ~8 s, depending on water depth. The energy in the sound pulses emitted by this Chirp echo sounder is at moderately high frequency. The Knudsen 320BR can be operated with either a 3.5-kHz transducer, for sub-bottom profiling, or a 12-kHz transducer for sounding. The lower frequency (3.5 kHz) transducer is not installed and will not be used. The conical beamwidth for the 12-kHz transducer is 30° and is directed downward.

Source levels for the Knudsen 320BR operating at 12 kHz has been measured as a maximum of 215 dB re 1 μ Pa m. Received levels would diminish rapidly with increasing depth. Assuming circular spreading loss, received level directly below the transducer would diminish to 180 dB re 1 μ Pa at distances of about 56 m when operating at 12 kHz. The 180 dB distance in the horizontal direction (outside the downward-directed beam) would be substantially less. Kremser et al. (2005) noted that the probability of a cetacean swimming through the area of exposure when a bottom profiler emits a pulse is small, and if the animal was in the area, it would have to pass the transducer at close range in order to be subjected to sound levels that could cause TTS.

Navy sonars that have been linked to avoidance reactions and stranding of cetaceans (1) generally are more powerful than the Knudsen 320BR operating with the 12 kHz transducer, (2) have longer pulse duration, and (3) are directed close to horizontally vs. downward for the Knudsen 320BR. The area of possible influence of the Chirp echo sounder is much smaller—a narrow conical beam spreading downward from the vessel. Marine mammals that encounter the echo sounder at close range are unlikely to be subjected to repeated pulses because of the narrow width of the beam, and may receive only small amounts of pulse energy because of the short pulses.

Masking

Marine mammal communications will not be masked appreciably by the Chirp echo sounder signals given its relatively low duty cycle, directionality, and the brief period when an individual mammal is likely to be within its beam. Beluga whale is the only odontocete anticipated to be in the area of the proposed survey. Though belugas can hear sounds ranging from 1.2 to 120 kHz, their peak sensitivity is ~10-15 kHz, overlapping with the 12-kHz transducer signals (Fay 1988). Some level of masking could result for belugas whales in close proximity to the survey vessel during brief periods of exposure to the sound. However masking is unlikely to an issue for beluga whales because belugas are likely to avoid survey vessels. The 12-kHz frequency sonar signals will not overlap with the predominant low frequencies in baleen whale calls, thus reducing potential for masking in this group.

Behavioral Responses

Marine mammal behavioral reactions to other pulsed sound sources are discussed above, and responses to the Chirp echo sounder are likely to be similar to those for other pulsed sources if received at the same levels. When the 12 kHz transducer is in operation, the behavioral responses to the Knudsen 320BR are expected to be similar to those reactions to the Kongsberg bathymetric echo sounder system (as discussed above). NMFS (2001) has concluded that momentary behavioral reactions “do not rise to the level of taking”. Thus, brief exposure of cetaceans to small numbers of signals from the Chirp echo sounder would not result in a “take” by harassment.

Hearing Impairment and Other Physical Effects

When the 12-kHz transducer is operating, the source frequency is similar to that of the bathymetric echo sounder (as discussed above). As with the Kongsberg, the pulses are brief and concentrated in a downward beam. A marine mammal would be in the beam of the Chirp echo sounder only briefly, reducing its received sound energy. Thus, it is unlikely that the echo sounder produces pulse levels strong enough to cause hearing impairment or other physical injuries even in an animal that is (briefly) in a position near the source.

The Chirp echo sounder is usually operated simultaneously with other higher-power acoustic sources. Many marine mammals will move away in response to the approaching higher-power sources or the vessel itself before the mammals would be close enough for there to be any possibility of effects from the Chirp echo sounder (Appendix D). In the case of mammals that do not avoid the approaching vessel and its various sound sources, mitigation measures that would be applied to minimize effects of the higher-power sources [see § II (3)] would further reduce or eliminate any minor effects of the echo sounder.

(4) Possible Effects of Chirp Sub-bottom Profiler

A Knudsen 3260 sub-bottom profiler will be operated from the *Louis S. St. Laurent* in open water when the *Louis S. St. Laurent* is not working in tandem with the *Healy*. The Knudsen’s transducer will be towed behind the *Louis S. St. Laurent*. The chirp system has a maximum 7.2 kW transmit capacity into the towed array and generally operates at 3–5 kHz. The energy from the towed unit is directed downward by an array of eight transducers in a conical beamwidth of 80°. The interval between pulses will be no less than one pulse per second. Sub-bottom profilers of that frequency can produce sound levels of 200-230 dB re 1 µPa at 1 m (Richardson et al. 1995).

Masking

Marine mammal communications will not be masked appreciably by the sub-bottom profiler signals given its relatively low duty cycle, directionality, and the brief period when an individual mammal is likely to

be within its beam. In the case of most odontocetes, the 3–5 kHz chirp signals do not overlap with the predominant frequencies in their calls, which would avoid significant masking. Beluga whale is the only odontocete anticipated in the area of the proposed survey. Though belugas can hear sounds ranging from 1.2 to 120 kHz, their peak sensitivity is ~10–15 kHz, not overlapping with the 3–5 kHz signals (Fay 1988). The frequency of the low-energy chirp profiler signals does not overlap with the predominant low frequencies in baleen whale calls, further reducing potential for masking.

Behavioral Responses

Marine mammal behavioral reactions to other pulsed sound sources are discussed above, and responses to the sub-bottom profiler are likely to be similar to those for other pulsed sources if received at the same levels. However, the pulsed signals from the towed 3–5 kHz chirp sub-bottom profiler are weaker than those from the airgun array. Therefore, behavioral responses are not expected unless marine mammals are close to the source. NMFS (2001) has concluded that momentary behavioral reactions “do not rise to the level of taking”. Thus, brief exposure of cetaceans to small numbers of signals from the sub-bottom profiler would not result in a “take” by harassment.

Hearing Impairment and Other Physical Effects

The pulses from the chirp profiler are brief and directed downward. A marine mammal would be in the beam of the sub-bottom profiler only briefly, reducing its received sound energy. Thus, it is unlikely that the sub-bottom profiler produces pulse levels strong enough to cause hearing impairment or other physical injuries even in an animal that is (briefly) in a position near the source.

The sub-bottom profiler is usually operated simultaneously with other higher-power acoustic sources. Many marine mammals will move away in response to the approaching higher-power sources or the vessel itself before the mammals would be close enough for there to be any possibility of effects from the sub-bottom profiler (Appendix D). In the case of mammals that do not avoid the approaching vessel and its various sound sources, mitigation measures that would be applied to minimize effects of the higher-power sources would further reduce or eliminate any minor effects of the sub-bottom profiler.

(5) Possible Effects of Helicopter Activities

It is anticipated that a helicopter will be deployed daily, weather permitting, to conduct ice reconnaissance and spot bathymetry in open water at locations outside of the U.S. EEZ. The spot soundings will be recorded to maximize the area surveyed and the data will be collected off the ship’s survey lines. A 12-kHz transducer will be slung by the helicopter and placed in the water to a mark affixed to the tether. Data will then be logged to a laptop computer in the helicopter.

Levels and duration of sounds received by marine mammals underwater from a passing helicopter are a function of the type of helicopter, orientation of the helicopter, depth of the animal, and water depth. A Canadian Coast Guard helicopter, a Messerschmitt MBB BO105, will be providing air support for this project. Helicopter sounds are detectable underwater at greater distances when the receiver is in shallow rather than deep water. Generally, sound levels received underwater decrease as the altitude of the helicopter increases (Richardson et al. 1995). Helicopter sounds are audible for much greater distances in air than in water.

Cetaceans

The nature of sounds produced by helicopter activities above the surface of the water does not pose a direct threat to the hearing of marine mammals that are in the water; however minor and short-term behavioral responses of cetaceans to helicopters have been documented in several locations, including the

Beaufort Sea (Richardson et al. 1985a,b; Patenaude et al. 2002). Cetacean reactions to helicopters depend on several variables including the animal's behavioral state, activity, group size, habitat, and the helicopter flight pattern, among other variables (Richardson et al. 1995).

During spring migration in the Beaufort Sea, beluga whales reacted to helicopter noise more frequently and at greater distances than did bowhead whales (38% vs. 14% of observations, respectively; Patenaude et al. 2002). Most reaction occurred when the helicopter passed within 250 m lateral distance at altitudes ≤ 150 m. Neither species exhibited noticeable reactions to single passes at altitudes >150 m. Belugas within 250 m of stationary helicopters on the ice with the engine running showed the most overt reactions. Whales were observed to make only minor changes in direction in response to sounds produced by helicopters, so all reactions to helicopters were considered brief and minor. Cetacean reactions to helicopter disturbance are difficult to predict and may range from no reaction at all to minor changes in course or (infrequently) leaving the immediate area of the activity.

Pinnipeds

Few systematic studies of pinniped reactions to aircraft overflights have been completed. Documented reactions range from simply becoming alert and raising the head, to escape behavior such as hauled out animals rushing to the water. Ringed seals hauled out on the surface of the ice have shown behavioral responses to aircraft overflights with escape responses most probable at lateral distances <200 m and overhead distances ≤ 150 m (Born et al. 1999). Although specific details of altitude and horizontal distances are lacking from many largely anecdotal reports, escape reactions to a low flying helicopter (<150 m altitude) can be expected from all pinnipeds potentially encountered during the proposed operations. These responses would likely be relatively minor and brief in nature. Whether any response would occur when a helicopter is at the higher suggested operational altitudes (below) is difficult to predict and probably a function of several other variables including wind chill, relative wind chill, and time of day (Born et al. 1999).

As mentioned in the previous section, momentary behavioral reactions “do not rise to the level of taking” as defined by NMFS (NMFS 2001). In order to limit behavioral reactions of marine mammals during ice reconnaissance and spot bathymetry work, the helicopter will maintain a minimum altitude of 200 m (656 ft) above the sea ice except when taking off, landing or conducting spot bathymetry. Sea-ice landings are not planned at this time.

(6 Possible Effects of Icebreaking Activities)

Icebreakers produce more noise while breaking ice than ships of comparable size due primarily to the sounds of the propeller cavitation (Richardson et al. 1995). Multi-year ice, which is expected to be in the northern portion of the proposed survey area, is thicker than younger ice. Icebreakers typically ram into heavy ice until losing momentum, then back off to build momentum before ramming again. The highest noise levels usually occur while backing full astern in preparation to ram forward through the ice. Overall, the noise generated by an icebreaker pushing ice was 10-15 dB greater than the noise produced by the ship underway in open water (Richardson et al. 1995). In general, the Arctic Ocean is a noisy environment. Greening and Zakaruskas (1993) reported ambient sound levels of up to 180 dB/ $\mu\text{Pa}^2/\text{Hz}$ under multi-year pack ice in the central Arctic pack ice. Little information is available about the effect of the increased sound levels on marine mammals due to icebreaking.

Cetaceans

Few studies have been conducted to evaluate the potential interference of icebreaking noise with marine mammal vocalizations. Erbe and Farmer (1998) measured masked hearing thresholds of a captive beluga whale. They reported that the recording of a Canadian Coast Guard ship, *Henry Larsen*, ramming

ice in the Beaufort Sea, masked recordings of beluga vocalizations at a noise-to-signal pressure ratio of 18 dB. That occurred when the noise pressure level was eight times as high as the call pressure. Erbe and Farmer (2000) also predicted when icebreaker noise would affect beluga whales through software that combined a sound propagation model and beluga whale impact threshold models. They again used the data from the recording of the *Henry Larsen* in the Beaufort Sea and predicted that masking of beluga vocalizations could extend between 40 and 71 km near the surface.

Lesage et al. (1999) report that beluga whales changed their call type and call frequency when exposed to boat noise. It is possible that the whales were adapting to the ambient noise levels and were able to communicate despite the sound. Given the documented reaction of belugas to ships and icebreakers (see below) it is highly unlikely that beluga whales would remain in the proximity of vessels where their vocalizations would be masked.

Beluga whales have been documented swimming rapidly away from ships and icebreakers in the Canadian High Arctic when a ship approached to within 35-50 km, and they may travel up to 80 km from the vessel's track (Richardson et al. 1995). It is expected that belugas avoid icebreakers as soon as they detect the ships (Cosens and Dueck 1993). The reaction of beluga whales to ships vary greatly and some animals may become habituated to higher levels of ambient noise (Erbe and Farmer 2000).

Little information is available regarding the effects of icebreaking ships on baleen whales. Migrating bowhead whales appeared to avoid an area around a drillsite by >25 km where an icebreaker was working in the Beaufort Sea. There was intensive icebreaking daily in support of the drilling activities (Brewer et al. 1993). Migrating bowheads also avoided a nearby drillsite at the same time of year when little icebreaking was being conducted (LGL and Greeneridge 1987). It is unclear as to whether the drilling activities, icebreaking operations, or the ice itself might have been the cause for the whales' diversion.

Pinnipeds

Brueggeman et al. (1992) reported on the reactions of seals to an icebreaker during activities at two prospects in the Chukchi Sea. Reactions of seals to the icebreakers varied between the two prospects. Most (67%) seals did not react to the icebreaker at either prospect. Reaction at one prospect was greatest during icebreaking activity followed by general vessel activity (running/maneuvering/jogging) and was lowest while the vessel was at anchor or drifting. Frequency of reaction was greatest for animals within 0.23 km of the vessel and lowest for animals beyond 0.93 km. At the second prospect however, seal reaction was lowest during icebreaking activity with higher and similar levels of response during general (non-icebreaking) vessel operations and when the vessel was at anchor or drifting. The frequency of seal reaction generally declined with increasing distance from the vessel except during general vessel activity where it remained consistently high to about 0.46 km from the vessel before declining. Kanik et al. (1980 in Richardson et al. 1995) reported that most ringed seals and harp seals within 1-2 km from an icebreaker remained on ice but that seals closer to the icebreaker often dove into the water.

(7) Mitigation Measures

Several mitigation measures are built into the planned seismic survey as an integral part of the activities, as described in § II (3). Those measures include the following: one or two dedicated protected species observers on the source vessel maintaining a visual watch during all daylight airgun operations, two observers on the source vessel (when practical) for 30 min before and during the onset of activities, additional PSOs stationed on the *Healy* to assist with monitoring for mammals while the vessels work in tandem, power downs or shut downs when mammals are detected in or about to enter designated safety zones, no start ups of the airgun array unless the full safety radius is visible, and conducting the majority

of the survey before September to avoid migrating bowhead whales. Also, the seismic survey will be conducted in deep water, where sound propagation is less than in shallow water, and in the Arctic Ocean, where marine mammal densities are low.

Previous and subsequent analysis of potential impacts takes account of the planned mitigation measures. It would not be meaningful to analyze the effects of the planned activities without mitigation, as the mitigation (and associated monitoring) measures are an integral part of the activities.

(8) Numbers of Marine Mammals that May be “Taken by Harassment”

All anticipated takes would be “takes by harassment”, as described in Section I, involving temporary changes in marine mammal behavior. The mitigation measures to be applied will minimize the possibility of injurious takes. (However, as noted earlier and in Appendix D, there is no specific information demonstrating that injurious “takes” would occur even in the absence of the planned mitigation measures.) The sections below describe methods used to estimate “take by harassment” and present estimates of the numbers of marine mammals that might be affected during the proposed seismic survey in the Arctic Ocean. The estimates are based on data obtained during marine mammal surveys in and near the Arctic Ocean by Stirling et al. (1982), Kingsley (1986), Moore et al. (2000b), Haley and Ireland (2006), Haley (2006), GSC unpubl. data (2008), and Mosher et al. (2009), the Bowhead Whale Aerial Survey Program (BWASP), and on estimates of the sizes of the areas where effects could potentially occur. In some cases, these estimates were made from data collected from regions and habitats that differed from the proposed project area.

Detectability bias, quantified in part by $f(0)$, is associated with diminishing sightability with increasing lateral distance from the trackline. Availability bias ($g(0)$) refers to the fact that there is <100% probability of sighting an animal that is present along the survey trackline. Some sources of densities used below included these correction factors in their reported densities. In other cases the best available correction factors were applied to reported results when they had not been included in the reported data (e.g., Moore et al. 2000b). Adjustments to reported population or density estimates were made on a case by case basis to take into account differences between the source data and the general information on the distribution and abundance of the species in the project area.

Although several systematic surveys of marine mammals have been conducted in the southern Beaufort Sea, few data (systematic or otherwise) are available on the distribution and numbers of marine mammals in the northern Beaufort Sea or areas further offshore in the Arctic Ocean. The main sources of distributional and numerical data used in deriving the estimates are described in the next subsection. Both “maximum estimates” as well as “best estimates” of marine mammal densities (Table 4) and the numbers of marine mammals potentially exposed to underwater sound (Table 5) were calculated as described below. The best (or average) estimate is based on available distribution and abundance data and represents the most likely number of animals that may be encountered during the survey, assuming no avoidance of the airguns or vessel. The maximum estimate is either the highest estimate from applicable distribution and abundance data or the average estimate increased by a multiplier intended to produce a very conservative (over) estimate of the number of animals that may be present in the survey area. There is some uncertainty about how representative the available data are and the assumptions used below to estimate the potential “take by harassment”. However, the approach used here is accepted by NMFS as the best available at this time.

We have calculated exposures for marine mammals only within U.S. waters. After the *Louis S. St. Laurent* exits U.S. waters, the survey activities no longer fall under the jurisdiction of the United States or the MMPA.

The following estimates are based on a consideration of the number of marine mammals that might be disturbed appreciably over the ~806 line km (435 n.mi.) of seismic surveys within U.S. waters across the Arctic Ocean. An assumed total of 1007.5 km (544 n.mi.) of trackline includes a 25% allowance over and above the planned ~806 km to allow for turns, lines that might have to be repeated because of poor data quality, or for minor changes to the survey design.

The anticipated radii of influence of the lower energy sound sources including Chirp echo sounder (on the *Louis S. St. Laurent*) and bathymetric echo sounder (on the *Healy*) are less than that for the airgun configuration. It is assumed that during simultaneous operations of the airgun array and echo sounder, any marine mammals close enough to be affected by the echo sounder would already be affected by the airguns. However, whether or not the airguns are operating simultaneously with the echo sounder, marine mammals are expected to exhibit no more than short-term and inconsequential responses to the echo sounder given its characteristics (e.g., narrow downward-directed beam) and other considerations described in Section IV above. Similar minimal response levels are expected from marine mammals exposed to the *Healy's* bathymetric profiler. Such reactions are not considered to constitute "taking" as defined by NMFS (NMFS 2001). Therefore, no additional allowance is included for animals that might be exposed to sound sources other than the airguns.

Numbers of marine mammals that might be present and potentially disturbed are estimated below based on available data about mammal distribution and densities in the area and the area ensonified to sound levels ≥ 160 dB re 1 μ Pa (rms). "Take by harassment" is calculated by multiplying the expected densities of marine mammals likely to occur in the survey area by the area potentially ensonified to sound levels ≥ 160 dB re 1 μ Pa (rms).

(a) Marine Mammal Density Estimates

This section provides estimated densities of marine mammals that may occur in the survey area. Some surveys of marine mammals have been conducted near the southern end of the proposed project area, but few data on the species and abundance of marine mammals in the northern Beaufort Sea and the Arctic Ocean are available. No published densities of marine mammals are available for this region, although few marine mammals were encountered during vessel-based surveys through the general area in 2005, 2006, 2008 and 2009. A total of two polar bears, 36 seals, and a single beluga whale sighting(s) were recorded along ~2299 km of monitored trackline between 71°N and 74°N (Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008).

Given that the survey lines within U.S. waters extend from ~71° to 74°N latitude, it is likely that seismic operations will be conducted in both open-water and sea-ice conditions. Because densities of marine mammals often differ between open-water and pack-ice habitats, the likely extent of the pack ice and open water at the time of the survey was estimated to determine marine mammal densities for the two habitat types. Images of average monthly sea-ice concentration for August from 2005 through 2009 were available from the National Snow and Ice Data Center (NSIDC) and used to identify 74°N latitude as a reasonable ice-edge boundary applicable to the proposed survey period and location. Based on these satellite data, the majority of the survey in U.S. waters will be conducted in open water and unconsolidated pack ice in the southern latitudes of the survey area. This region will include the ice margin where the highest densities of cetaceans and pinnipeds are likely to be encountered. The proposed survey lines within U.S. waters reach ~74°10'N, and extend into the estimated ice-edge boundary for August 2010 by ~18.5 km (10 n.mi.). This comprises less than 4% of the total trackline within U.S. waters. We have divided the survey effort between the two habitat zones of open water and ice margin based on the 2005 – 2009 NSIDC satellite data described above and the planned location of the tracklines. NSIDC data from 2005-2009 suggest little ice will be present south of 74°N, although data from the 2009

cruise (Moser et al. 2009) show that inter-annual variability could result in a greater amount of ice being encountered than expected. As a conservative measure, we estimated that within U.S. waters 80% of the survey tracklines will occur in open water and 20% of the tracklines will occur within the ice margin.

During a 2009 survey, the *Louis S. St. Laurent* encountered mostly open water with occasional ice patch concentrations of 4/10 south of 75°N in mid-August (Mosher et al. 2009). This is roughly the region where all of the proposed survey lines within U.S. waters will occur in 2010. Because the proposed 2010 survey will start ~6 August, more ice may be present in the region. The NSIDC (2009) reported that more arctic sea ice cover in 2009 remained after the summer than in the record-setting low years of 2007 and 2008. We expect that sea ice density and extent in 2010 will be closer to the density and extent of sea ice in 2009 rather than the record-setting low years of 2007 and 2008. All animals observed during the 2009 survey (Mosher et al. 2009) were north of the proposed survey area, i.e., north of 74°N.

Cetaceans

Average and maximum densities for each cetacean species or species group reported to occur in U.S. waters of the Arctic Ocean within the proposed survey area are presented below. Densities were calculated based on the sightings and effort data from available survey reports. No cetaceans were observed during surveys near the proposed survey area in August/September 2005 (Haley and Ireland 2006), August 2006 (Haley 2006), August/September 2008 (GSC unpubl. data 2008), or August/September 2009 (Mosher et al. 2009).

Seasonal (summer and fall) differences in cetacean densities along the north coast of Alaska have been documented by Moore et al. (2000b). The proposed survey will be conducted in U.S. waters from ~6–12 August and is considered to occur during the summer season.

The summer *beluga* density (Table 4) was based on 41 sightings along 9022 km of on-transect effort that occurred over water >2000 m deep during the summer in the Beaufort Sea (Moore et al. 2000b). A mean group size of 2.8 (CV=1.0) derived from BWASP data of August beluga sightings in the Beaufort Sea in water depths >2000 m was used in the density calculation. An $f(0)$ value of 2.841 and a $g(0)$ value of 0.58 from Harwood et al. (1996) were also used in the density computation. The CV associated with group size was used to select an inflation factor of 2 to estimate the maximum density that may occur in the proposed survey area within U.S. waters. Most Moore et al. (2000b) sightings were south of the proposed seismic survey area. However, Moore et al. (2000b) found that beluga whales were associated with both light (1–10%) and heavy (70–100%) ice cover. Five of 23 beluga whales that Suydam et al. (2005a) tagged in Kasegaluk Lagoon (northeast Chukchi Sea) traveled to 79–80°N latitude into the pack ice and within the region of the proposed survey. These and other tagged whales moved into areas as far as 1100 km (594 n.mi.) offshore between Barrow and the Mackenzie River delta, spending time in water with 90% ice coverage. Therefore, we applied the observed density calculated from the Moore et al. (2000b) sightings as the average density for both “open-water” and “ice-margin” habitats. No beluga whales were sighted during recent surveys in the proposed survey area (Harwood et al. 2005; Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008; and Mosher et al. 2009), and the densities presented in Table 4 are likely higher than densities likely to be encountered.

By the time the survey begins in early August, most *bowhead* whales have typically traveled east of the proposed project area to summer in the eastern Beaufort Sea and Amundsen Gulf. Industry aerial surveys of the continental shelf near Camden Bay in 2008 recorded eastward migrating bowhead whales until 12 July (Lyons et al. 2009). No bowhead sightings were recorded again despite continued flights until 19 August. A summer bowhead whale density was derived from 9022 km of summer (July/August) aerial survey effort reported by Moore et al. (2000b) in the Alaskan Beaufort Sea during which six

sightings of bowhead whales were documented in water >2000 m deep. A mean group size for bowhead whale sightings in September in waters >2000 m deep was calculated to be 1.14 (CV=0.4) from BWASP data. An $f(0)$ value of 2 and a $g(0)$ value of 0.07, both from Thomas et al. (2002), were used to estimate a summer density for bowhead whales of 0.0109 whales/km². This density falls within the range of densities, i.e., 0.0099–0.0717 whales/km², reported by Lyons and Christie (2009) based on data from three July 2008 surveys.

Treacy et al. (2006) reported that in years of heavy ice conditions, bowhead whales occur farther offshore than in years of light to moderate ice. NSIDC (2009) reported that September 2009 had the third lowest sea ice extent since the start of their satellite records in 1979. The extent of sea ice at the end of the 2009 Arctic summer however, was greater than in 2007 or 2008. NSIDC does not expect 2010 to be a heavy ice year during which bowhead whales might occur farther offshore in the area of the proposed survey. During the lowest ice-cover year on record (2007), BWASP reported no bowhead whale sightings in locations far offshore with water depths ≥ 2000 m. Because few bowhead whales have been documented in the deep offshore waters of the proposed survey area, half of the bowhead whale density estimate from Moore et al. (2000b) was used as the average density (0.0054 whales/km²; Table 4). The CV of group size and standard errors reported in Thomas et al. (2002) for $f(0)$ and $g(0)$ correction factors suggest that an inflation factor of 2 applied to the average density is appropriate for estimating the maximum density. NSIDC did not forecast that 2010 would be a heavy ice year and we anticipate that bowheads will remain relatively close to shore, and in areas of light ice coverage. Therefore, we have applied the same density for bowheads to the open-water and ice-margin habitats. Bowhead whales were not sighted during recent surveys in the Arctic Ocean (Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008; Mosher et al. 2009) suggesting that the bowhead whale densities shown in Table 4 are likely higher actual densities in the proposed survey area.

For *other cetacean species* that may be encountered in the Beaufort Sea, densities are likely to be very low in the summer when the survey is scheduled. Fin and humpback whales are unlikely to occur in the Beaufort Sea. No gray whales were observed in the Beaufort Sea by Moore et al. (2000b) during summer aerial surveys in water >2000 m. Gray whales were not recorded in water >2000 m by the BWASP during August in 29 years of survey operation. Harbor porpoises are not expected to be present in large numbers in the Beaufort Sea during the fall although small numbers may be encountered during the summer. Neither gray whales nor harbour porpoises are likely to occur in the far-offshore waters of the proposed survey area. Narwhals are not expected to be encountered within the survey program area although a few individuals could be present if ice is nearby. Because these species occur so infrequently in the Beaufort Sea, little to no data are available for the calculation of densities. Minimal cetacean densities have therefore been assigned to these species for density calculation purpose and to allow for chance encounters (Table 4). Those densities include “0” for the average and 0.0001 individuals/km² for the maximum.

Pinnipeds

Extensive surveys of ringed and bearded seals have been conducted in the Beaufort Sea, but most surveys were conducted over the landfast ice during aerial surveys, and few seal surveys have occurred in open water or in the pack ice. Kingsley (1986) conducted *ringed seal* surveys of the offshore pack ice in the central and eastern Beaufort Sea during late spring (late June). These surveys provide the most relevant information on densities of ringed seals in ice-margin habitat of the Beaufort Sea. The density estimate in Kingsley (1986) was used as the average ringed seal density in ice-margin habitat of the proposed survey area (Table 4). The average density was multiplied by 4 to estimate maximum density, as was done for all seal species likely to occur within the survey area. Ringed seals are closely associated

with sea ice therefore the ice-margin densities were multiplied by a factor of 0.75 to estimate a summer open-water ringed-seal density for locations with water depth >2000 m.

Densities of *bearded seals* were estimated by multiplying the ringed seal densities by 0.051 based on the proportion of bearded seals to ringed seals reported in Stirling et al. (1982). Because bearded seals are associated with the pack ice edge and shallow water, their estimated summer ice-margin density was also multiplied by a factor of 0.75 for the open-water density estimate. Minimal values were used to estimate *spotted seal* and *Pacific walrus* densities because they are uncommon offshore in the Beaufort Sea and are not likely to be encountered.

Polar Bear

One polar bear sighting of two individuals was recorded along ~2308 km of monitored trackline between 71°N and 74°N (Haley and Ireland 2006; Haley 2006; GSC unpubl. data 2008) and all were hauled out on ice. This results in an average density of 0.0004 bears/ km² assuming all bears present within 1 km on either side of the vessel were observed. The maximum density in ice-margin habitat was assumed to be 4 times this value. The density of polar bears in open-water is expected to be much lower so minimal density estimates have been assumed (Table 4).

TABLE 4. Expected summer densities of marine mammals in U.S. waters offshore in the Beaufort Sea and Arctic Ocean. This area is expected to be mostly open water and may extend into the ice margin. Densities are corrected for *f*(0) and *g*(0) biases. Species listed as endangered are in italics.

| Species | Open Water | | Ice Margin | |
|-----------------------|--|--|--|--|
| | Average Density (# / km ²) | Maximum Density (# / km ²) | Average Density (# / km ²) | Maximum Density (# / km ²) |
| Odontocetes | | | | |
| Beluga | 0.0354 | 0.0709 | 0.0354 | 0.0709 |
| Narwhal | 0.0000 | 0.0001 | 0.0000 | 0.0002 |
| Delphinidae | | | | |
| Killer whale | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| Phocoenidae | | | | |
| Harbor porpoise | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| Mysticetes | | | | |
| <i>Bowhead whale</i> | 0.0061 | 0.0122 | 0.0061 | 0.0122 |
| Gray whale | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| Minke whale | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| <i>Fin whale</i> | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| <i>Humpback whale</i> | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| Pinnipeds | | | | |
| Bearded seal | 0.0096 | 0.0384 | 0.0128 | 0.0512 |
| Spotted seal | 0.0001 | 0.0004 | 0.0001 | 0.0004 |
| Ringed seal | 0.1883 | 0.7530 | 0.2510 | 1.0040 |
| Pacific Walrus | 0.0001 | 0.0004 | 0.0001 | 0.0004 |
| Ursidae | | | | |
| Polar Bear | 0.0001 | 0.0004 | 0.0001 | 0.0016 |

(b) Estimation of Area Ensonified to Sound Levels ≥ 160 dB rms

The area of water potentially exposed to sound levels ≥ 160 dB by the proposed operations was calculated by multiplying the planned trackline distance within U.S. waters by the cross-track distance of the sound propagation. Sound levels from the airgun array to be used for the proposed geophysical survey (comprised of two 500 in³ and one 150 in³ G-guns) were measured during a 2009 project in the Arctic Ocean (Mosher et al 2009; Roth and Schmidt 2010). The propagation experiment took place at 74°50.4'N; 156°34.31'W, in 3863 m of water. The location was near the northern end of the two proposed survey lines in U.S. waters. We expect the sound propagation by the airgun array in the planned 2010 survey will be similar to that measured in 2009 because of the similar water depths and relative locations of the test site and proposed survey area. The ≥ 160 dB rms sound level radius was estimated to be ~2500 m (1.3 n.mi.) based on modeling of the 0-peak energy of the airgun array (Roth and Schmidt 2010). The 0-peak values were corrected to rms by subtracting 10 dB.

Closely spaced survey lines and large cross-track distances of the ≥ 160 dB radii can result in repeated exposure of the same area of water. Excessive amounts of repeated exposure can lead to overestimation of the number of animals potentially exposed through double counting. The trackline for the proposed USGS survey in U.S. water, however, covers a large geographic area without adjacent tracklines and the potential for multiple or repeated exposure is unlikely to be a concern.

The USGS 2010 geophysical survey is planned to occur >108 km offshore along ~806 km (435 n.mi.) of survey lines in U.S. waters during the first half of August exposing a total of ~4109 km² of water to sound levels ≥ 160 dB rms. We included an additional 25% allowance over and above the planned tracklines within U.S. waters to allow for turns, lines that might have to be repeated because of poor data quality, or for minor changes to the survey design. The resulting estimate of 5136.5 m² was used to estimate the numbers of marine mammals exposed to underwater sound levels ≥ 160 dB rms.

(c) Potential Number of Marine Mammal “Exposures” to Sound Levels ≥ 160

Numbers of marine mammals that might be present and potentially disturbed are estimated below based on available data about mammal distribution and densities in two different habitats during the summer as described above. There is no evidence however, that exposure at received sound levels ≥ 160 dB would have significant effects on individual animals or that the subtle changes in behavior or movements would “rise to the level of taking” according to guidance by the NMFS (NMFS 2001). The number of individuals of each species potentially exposed to received levels ≥ 160 dB re 1 μ Pa (rms) was estimated by multiplying

- the anticipated area to be ensonified to levels ≥ 160 dB re 1 μ Pa (rms) in both open water and the ice margin, by
- the expected species density.

Some of the animals estimated to be exposed to sound levels ≥ 160 dB re 1 μ Pa, particularly migrating bowhead whales, might show avoidance reactions before actual exposure to this sound level (Appendix D). Thus, these calculations estimate the number of individuals potentially exposed to ≥ 160 dB rms that would occur if there were no avoidance of the area ensonified.

Based on the operational plans and marine mammal densities described above, the estimates of marine mammals potentially exposed to sound ≥ 160 dB in the proposed survey area are presented in Table 5. For the common species, the requested numbers are calculated as described above and based on the densities from the data reported in the studies mentioned above. For less common species, estimates

were set to minimal values to allow for chance encounters. Discussion of the number of potential exposures is summarized by species in the following subsections.

Cetaceans

Through consultation with the Office of Protected Resources (NOAA) USGS proposes that that no ESA-listed marine species: bowhead, fin, or humpback whale will be adversely affected by this project during the survey or transit to the survey area from Dutch Harbor. However, we estimated the number of takes of ESA species that might, although highly unlikely, be exposed to received sound levels of 160 dB.

Based on density estimates and the area ensonified, one endangered cetacean species (bowhead whale) is expected to be exposed to received sound levels ≥ 160 dB unless bowheads avoid the survey vessel before the received levels reach 160 dB. Migrating bowheads are likely to show an avoidance response, though many of the bowheads engaged in other activities, particularly feeding and socializing, may not. Our estimate of the number of bowhead whales potentially exposed to sound levels ≥ 160 dB in the portion of the survey area in U.S. waters is between 31 and 63 (Table 5). Other endangered cetacean species that may be encountered in the area, fin and humpback whales, are unlikely to be exposed given their minimal density in the area.

The only other cetacean species likely to occur in the proposed survey area is beluga whale. Average and maximum estimates of the number of beluga whales exposed to sound levels ≥ 160 dB rms are 182 and 364 respectively. Estimates for other cetacean species are minimal (Table 5).

TABLE 5. Estimates of the numbers of marine mammals potentially exposed to received sound levels ≥ 160 dB during USGS's proposed seismic program in U.S. waters in the northern Beaufort Sea and Arctic Ocean, ~6 - 12 August 2010. Species in italics are listed under the U.S. ESA as endangered.

| Species | Number of Exposures to Sound Levels ≥ 160 dB | | | | | |
|------------------------|---|-------------|------------|-------------|-------------|-------------|
| | Open Water | | Ice Margin | | Total | |
| | Average | Maximum | Average | Maximum | Average | Maximum |
| Odontocetes | | | | | | |
| Monodontidae | | | | | | |
| Beluga | 146 | 291 | 36 | 73 | 182 | 364 |
| Narwhal | 0 | 1 | 0 | 1 | 0 | 2 |
| Delphinidae | | | | | | |
| Killer whale | 0 | 0 | 0 | 0 | 0 | 1 |
| Phocoenidae | | | | | | |
| Harbor porpoise | 0 | 0 | 0 | 0 | 0 | 1 |
| Mysticetes | | | | | | |
| <i>Bowhead whale</i> | 25 | 50 | 6 | 13 | 31 | 63 |
| Gray whale | 0 | 0 | 0 | 0 | 0 | 1 |
| Minke whale | 0 | 0 | 0 | 0 | 0 | 0 |
| <i>Fin whale</i> | 0 | 0 | 0 | 0 | 0 | 0 |
| <i>Humpback whale</i> | 0 | 0 | 0 | 0 | 0 | 0 |
| Total Cetaceans | 171 | 344 | 43 | 87 | 213 | 430 |
| Pinnipeds | | | | | | |
| Bearded seal | 39 | 158 | 13 | 53 | 53 | 210 |
| Spotted seal | 0 | 2 | 0 | 0 | 1 | 2 |
| Ringed seal | 774 | 3094 | 258 | 1031 | 1031 | 4126 |
| Pacific walrus | 0 | 2 | 0 | 0 | 1 | 2 |
| Total Pinnipeds | 813 | 3254 | 271 | 1084 | 1085 | 4338 |
| Ursidea | | | | | | |
| Polar Bear | 0 | 2 | 0 | 2 | 1 | 3 |

Pinnipeds

Ringed seal is the most widespread and abundant pinniped in ice-covered arctic waters, and annual variation in abundance and distribution appears to be high. Ringed seals account for the vast majority of marine mammals expected to be encountered, and hence exposed to airgun sounds with received levels ≥ 160 dB re 1 μ Pa (rms) during the proposed marine survey. The average and maximum number of exposures of ringed seals to sound levels ≥ 160 dB rms were estimated to be 1031 and 4126 respectively.

Two additional pinniped species are likely to or may occur in the proposed project area. The average and maximum numbers of bearded seals exposed to sound levels ≥ 160 dB rms were estimated to be 53 and 210, respectively. Average and maximum numbers of spotted seals were estimated to be 1 and 2, respectively. Ribbon seal and Pacific Walrus are unlikely to be encountered in the survey area, but a chance encounter could occur.

Polar Bear

The average and maximum number of polar bears exposed to sound levels ≥ 160 dB rms were estimated to be 1 and 3, respectively. However, most polar bears are likely to be encountered when hauled-out on ice where they would not be exposed to sounds at the ≥ 160 dB rms level.

(9) Conclusions

The proposed survey in the Arctic Ocean will involve towing an airgun array that will introduce pulsed sounds into the ocean, along with simultaneous operation of a multibeam bathymetric echo sounder and Chirp echo sounder. Routine vessel operations, other than the proposed operations by the airguns, are conventionally assumed not to affect marine mammals sufficiently to constitute “taking”. No “taking” of marine mammals is expected in association with operations of the echo sounders given the considerations discussed in Section IV (2 and 3), i.e., sonar sounds are beamed downward, the beam is narrow, and the pulses are extremely short.

(a) Cetaceans

Bowhead whales are considered by NMFS to be disturbed after exposure to underwater sound levels ≥ 160 dB rms. The relatively small airgun array for the proposed geophysical survey limits the size of the 160 dB zone around the vessel and will result in few bowhead whale exposures to underwater sound levels sufficient to reach the disturbance criterion as defined by NMFS.

Odontocete reactions to seismic energy pulses are usually assumed to be limited to lesser distances from the airgun(s) than are those of mysticetes, probably in part because odontocete low-frequency hearing is assumed to be less sensitive than that of mysticetes. However, at least when in the Canadian Beaufort Sea in summer, belugas appear to be fairly responsive to seismic energy, with few being sighted within 10–20 km of seismic vessels during aerial surveys (Miller et al. 2005). Belugas will likely occur in small numbers in the project area within U.S. waters during the survey period. Most belugas will likely avoid the vicinity of the survey activities and few will likely be exposed to underwater sound levels ≥ 160 dB rms.

Taking into account the mitigation measures that are planned, effects on cetaceans are generally expected to be restricted to avoidance of a limited area around the survey operation and short-term changes in behavior, falling within the MMPA definition of “Level B harassment” (behavioral disturbance; no serious injury or mortality). Furthermore, the estimated numbers of animals potentially exposed to sound levels sufficient to cause appreciable disturbance are very low percentages of the population sizes in the Bering–Chukchi–Beaufort seas, as described below.

Based on the ≥ 160 dB disturbance criterion, the best (average) estimates of the numbers of cetacean exposures to sounds ≥ 160 dB re 1 μ Pa (rms) represent $<1\%$ of the populations of each species in the Chukchi Sea and adjacent waters. For species listed as endangered under the ESA, our estimates suggest it is unlikely that fin whales or humpback whales will be exposed to received levels ≥ 160 dB rms, but that ~ 31 bowheads may be exposed at this level. The latter is $<1\%$ of the Bering–Chukchi–Beaufort population of $>14,247$ assuming 3.4% annual population growth from the 2001 estimate of $>10,545$ animals (Zeh and Punt 2005).

Some monodontids may be exposed to sounds produced by the airgun arrays during the proposed survey, and the numbers potentially affected are small relative to the population sizes (Table 5). The best estimate of the number of belugas that might be exposed to ≥ 160 dB (182) represents $<1\%$ of their population.

Mitigation measures such as controlled vessel speed, dedicated protected species observers, non-pursuit, and shut downs or power downs when marine mammals are seen within defined distances from operating airguns will further reduce short-term reactions and minimize any effects on hearing sensitivity. In all cases, the effects are expected to be short-term, with no lasting biological consequence.

Subsistence issues are addressed below. USGS has adopted a spatial and temporal operational strategy that, when combined with its community outreach and engagement program, will provide effective protection to the bowhead migration and subsistence hunt.

(b) Pinnipeds

Several pinniped species may be encountered in the geophysical survey, but ringed seal is by far the most abundant marine mammal species in the survey area. The best (average) estimates of the numbers of individuals seals exposed to airgun sounds at received levels ≥ 160 dB re 1 μ Pa (rms) during the marine survey are as follows: ringed seals (1031), bearded seals (53), spotted seals (1), and Pacific walrus (1) representing $<1\%$ of the Bearing–Chukchi–Beaufort populations for each species. It is probable that only a small percentage of the pinnipeds exposed to sound level ≥ 160 dB would actually be disturbed. The short-term exposures of pinnipeds to airgun sounds are not expected to result in any long-term negative consequences for the individuals or their populations.

(c) Polar Bears

Effects on polar bears are anticipated to be minor at most. Small numbers of polar bears will likely be encountered during the proposed geophysical survey, however almost all would be on the ice and therefore unaffected by underwater sound from the airguns. For the few bears that may be in the water, levels of airgun sound would be attenuated because polar bears do not dive much below the surface. Received levels of airgun sound are reduced substantially near the surface relative to sound levels in deeper water due to pressure release effects.

(10) Direct Effects on Fish, EFH, and Fisheries, and Their Significance

(a) Effects on Fish and Invertebrates

One of the reasons for the adoption of airguns as the standard energy source for marine seismic surveys was that, unlike explosives, they do not result in any appreciable fish kill. However, the existing body of information relating to the impacts of seismic on marine fish and invertebrate species is limited. This information is reviewed in Appendix E for fish and Appendix F for invertebrates.

In water, acute injury and death of organisms exposed to seismic energy depends primarily on two features of the sound source: (1) the received peak pressure, and (2) the time required for the pressure to rise and decay (Hubbs and Rechnitzer 1952 *in* Wardle et al. 2001). Generally, the higher the received pressure and the less time it takes for the pressure to rise and decay, the greater the chance of acute pathological effects. Considering the peak pressure and rise/decay time characteristics of seismic airgun arrays used today, the pathological zone for fish and invertebrates would be expected to be within a few meters of the seismic source (Buchanan et al. 2004). For the proposed survey, any injurious effects on fish would be limited to very short distances (Appendix E).

The proposed Arctic Ocean seismic program for 2010 is predicted to have negligible to low physical effects on the various life stages of fish and invertebrates for its ~ 7 day duration and ~ 435 n.mi (806 km) extent. Therefore, physical effects of the proposed program would not adversely affect fish and invertebrates.

(b) EFH

The proposed survey off northern Alaska will occur in an area designated as Essential Fish Habitat (EFH) for Arctic cod (NPFMC 2009). The ~ 435 n.mi (806 km) of seismic survey that will be conducted in U.S. waters represents the maximum possible extent of potential EFH that would be ensonified during the project; the border of the U.S. EEZ defines the potential Arctic cod EFH boundary for Arctic cod. Effects on managed EFH species (Arctic cod) by the seismic operations assessed here would be

temporary and minor (see above). The main effect would be short-term disturbance that might lead to temporary and localized relocation of the EFH species or their food. The actual physical and chemical properties of the EFH will not be impacted.

(c) Fisheries

No active fishing is expected to be conducted within the study area during the time of the survey. Any on-going fisheries near the project area would be subsistence, and much closer to shore than the proposed survey.

(11) Direct Effects on Seabirds and their Significance

Investigations into the effects of airguns on seabirds are extremely limited. Stemp (1985) conducted opportunistic observations on the effects of seismic exploration on seabirds, and Lacroix et al. (2003) investigated the effect of seismic surveys on molting long-tailed ducks in the Beaufort Sea, Alaska. Stemp (1985) did not observe any effects of seismic testing, although he warned that his observations should not be extrapolated to areas with large concentrations of feeding or molting birds. In a more intensive and directed study, Lacroix et al. (2003) did not detect any effects of seismic exploration on molting long-tailed ducks in the inshore lagoon systems of Alaska's North Slope. Both aerial surveys and radio-tracking indicated that the proportion of ducks that stayed near their marking location from before to after seismic exploration was unaffected by nearby seismic survey activities. Seismic activity also did not appear to change the diving intensity of long-tailed ducks significantly. The predominant airgun source involved in the study by Lacroix et al. 2003 (Lawson 2002) was smaller in total volume than those planned for use here. However, it involved the same number of airguns (8), and number of airguns is the dominant influence on source level (Caldwell and Dragoset 2000). Consistent with that, the anticipated 180 and 190 dB radii in water >100 m deep during the planned Arctic Ocean survey are similar to those during the study of Lacroix et al. 2003 (*cf.* Lawson 2002) However, the anticipated 180 and 190 dB radii in shallow water (a small fraction of this survey) are considerably larger than those assumed in the Lacroix et al. (2003) study.

Birds might be affected slightly by seismic sounds from the proposed study, but the impacts are not expected to be significant to individual birds or their populations. The types of impacts that are possible are summarized below.

Localized, temporary displacement and disruption of feeding—Such displacements would be similar to those caused by other large vessels that passed through the area. Any adverse effects would be negligible.

Modified prey abundance—It is unlikely that prey species for birds will be affected by seismic activities to a degree that affects the foraging success of birds. If prey species exhibit avoidance of the ship, the avoidance is expected to be transitory and limited to a very small portion of a bird's foraging range.

Disturbance to breeding birds on island colonies—A vessel (seismic or otherwise) that approaches too close to a breeding colony could disturb adult birds from nests in response either to sonic or to visual stimuli. This is not applicable to the proposed Arctic Ocean survey, which will be in offshore waters away from any seabird colonies.

Egg and nestling mortality—Disturbance of adult birds from nests can lead to egg or nestling mortality *via* temperature stress or predation. There is no potential for this considering the distance that the seismic survey will occur from nesting colonies.

Chance injury or mortality—Many species of marine birds feed by diving to depths of several meters or more. Flocks of feeding birds consisting of hundreds or thousands of birds often occur in Alaskan waters. Also, some species of seabirds (particularly alcids) escape from boats by diving when the boat is close. It is possible that, during the course of normal feeding or escape behavior, some birds could be near enough to an airgun to be injured by a pulse. Although no specific information is available about the circumstances (if any) where this might occur, the negligible reactions of birds to airguns (see above) suggest that a bird would have to be very close to any airgun to receive a pulse with sufficient energy to cause injury, if that is possible at all.

Induced injury or mortality—By disorienting, injuring, or killing prey species, or by otherwise increasing the availability of prey species to marine birds, seismic activity could attract birds. Birds drawn too close to an airgun may be at risk of injury. However, available evidence from other seismic surveys has not shown a pattern of fish (or other prey) kills from airguns [see § IV(5)(a), above]. Thus, the potential that birds would be attracted and subsequently injured by the proposed seismic survey appears very low.

(12) Indirect Effects to Marine Mammals and Their Significance

The proposed airgun operations will not result in any permanent impact on habitats used by marine mammals, or to the food sources they use. The main impact issue associated with the proposed activities will be temporarily elevated noise levels and the associated direct effects on marine mammals, as discussed above.

During the seismic study only a small fraction of the available habitat would be strongly ensonified at any given time. Disturbance to fish species would be short-term and fish would return to their pre-disturbance behavior once the seismic activity ceased [§ IV(5)(a), above]. Thus, the proposed survey would have little, if any, impact on the abilities of marine mammals to feed in the area where seismic work is planned.

Some mysticetes, including bowhead whales, feed on concentrations of zooplankton. Although the main summering area for bowheads is in the Canadian Beaufort Sea, at least a few feeding bowhead whales may occur in offshore waters of the western Beaufort Sea in August and September during the proposed geophysical survey. A reaction by zooplankton to a seismic impulse would only be relevant to whales if it caused a concentration of zooplankton to scatter. Pressure changes of sufficient magnitude to cause that type of reaction would probably occur only very close to the source. Impacts on zooplankton behavior are predicted to be negligible, and that would translate into negligible impacts on feeding mysticetes.

(13) Possible Effects on Subsistence Hunting and Fishing

Subsistence hunting and fishing continue to be prominent in the household economies and social welfare of some Alaskan residents, particularly among those living in small, rural villages (Wolfe and Walker 1987). Subsistence remains the basis for Alaska Native culture and community. In rural Alaska, subsistence activities are often central to many aspects of human existence, including patterns of family life, artistic expression, and community religious and celebratory activities. Because of the importance of subsistence, the National Science Foundation offers guidelines for science coordination with native Alaskans at <http://www.arcus.org/guidelines/>.

(a) Subsistence Hunting for Marine Mammals

Marine mammals are legally hunted along the north coast of Alaska near Barrow, Nuiqsut, and Kaktovik by coastal Alaska Natives; species hunted include bowhead whales, beluga whales, ringed,

spotted, and bearded seals, walrus, and polar bears. In the Barrow area, bowhead whales provided ~69% of the total weight of marine mammals harvested from April 1987 to March 1990. During that time, ringed seals were harvested the most on a numerical basis (394 animals).

Bowhead whale hunting is the key activity in the subsistence economies of Barrow and two smaller communities to the east, Nuiqsut and Kaktovik. Whale harvests have a great influence on social relations by strengthening the sense of Inupiat culture and heritage in addition to reinforcing family and community ties.

An overall quota system for the hunting of bowhead whales was established by the International Whaling Commission in 1977; the quota is now regulated through an agreement between the National Marine Fisheries Service and the Alaska Eskimo Whaling Commission. The Alaska Eskimo Whaling Commission (AEWC) allots the number of bowhead whales that each whaling community may harvest annually (USDI/BLM 2005; NMFS 2008b).

The community of Barrow hunts bowhead whales in both the spring and fall during the whales' seasonal migrations along the coast. Often, the bulk of the Barrow bowhead harvest is taken during the spring hunt. However, with larger quotas in recent years, it is common for a substantial fraction of the annual Barrow quota to remain available for the fall hunt (Table 6). The communities of Nuiqsut and Kaktovik participate only in the fall bowhead harvest. The spring hunt at Barrow occurs after leads open due to the deterioration of pack ice; the spring hunt typically occurs from early April until the first week of June. The fall migration of bowhead whales that summer in the eastern Beaufort Sea typically begins in late August or September. The location of the fall subsistence hunt depends on ice conditions and (in some years) industrial activities that influence the bowheads movements as they move west (Brower 1996). In the fall, subsistence hunters use aluminum or fiberglass boats with outboards. Hunters prefer to take bowheads close to shore to avoid a long tow during which the meat can spoil, but Braund and Moorehead (1995) report that crews may (rarely) pursue whales as far as 80 km. The autumn hunt at Barrow usually begins in mid-September, and mainly occurs in the waters east and northeast of Point Barrow. The whales have usually left the Beaufort Sea by late October (Treacy 2002a,b).

The scheduling of this seismic survey has been discussed with representatives of those concerned with the subsistence bowhead hunt, most notably the Alaska Eskimo Whaling Commission and the Barrow Whaling Captains' Association. For this among other reasons, the project has been scheduled to commence in early August and terminate by mid-August in U.S. waters, before the start of the fall hunt at Barrow (or Nuiqsut or Kaktovik), to avoid possible conflict with whalers. In addition to scheduling the seismic operations prior to the start of the whaling season, the location of the surveys well offshore will further eliminate the potential for disturbance to the bowhead whale subsistence hunt.

Beluga whales are available to subsistence hunters at Barrow in the spring when pack-ice conditions deteriorate and leads open up. Belugas may remain in the area through June and sometimes into July and August in ice-free waters. Hunters usually wait until after the spring bowhead whale hunt is finished before turning their attention to hunting belugas. The average annual harvest of beluga whales taken by Barrow for 1962–1982 was five (MMS 1996). The Alaska Beluga Whale Committee recorded that 23 beluga whales had been harvested by Barrow hunters from 1987 to 2002, ranging from 0 in 1987, 1988 and 1995 to the high of 8 in 1997 (Fuller and George 1999; Alaska Beluga Whale Committee 2002 *in* USDI/BLM 2005). The proposed geophysical survey is unlikely to overlap with the beluga harvest, and the survey initiates well outside the area where impacts to beluga hunting by Barrow villagers could occur.

Table 6. Number of bowhead whale landing by year at Barrow, Cross Island (Nuiqsut), and Kaktovik, 1993-2008. Barrow numbers include the total number of whales landed for the year followed by the numbers landed during the fall hunt in parenthesis. Cross Isl. (Nuiqsut) and Kaktovik landings are in autumn.

| Year | Point Hope | Wainwright | Barrow | Cross Island | Kaktovik |
|------|------------|------------|---------|--------------|----------|
| 1993 | 2 | 5 | 23 (7) | 3 | 3 |
| 1994 | 5 | 4 | 16 (1) | 0 | 3 |
| 1995 | 1 | 5 | 19 (11) | 4 | 4 |
| 1996 | 3 | 3 | 24 (19) | 2 | 1 |
| 1997 | 4 | 3 | 30 (21) | 3 | 4 |
| 1998 | 3 | 3 | 25 (16) | 4 | 3 |
| 1999 | 2 | 5 | 24 (6) | 3 | 3 |
| 2000 | 3 | 5 | 18 (13) | 4 | 3 |
| 2001 | 4 | 6 | 27 (7) | 3 | 4 |
| 2002 | 0 | 1 | 22 (17) | 4 | 3 |
| 2003 | 4 | 5 | 16 (6) | 4 | 3 |
| 2004 | 3 | 4 | 21 (14) | 3 | 3 |
| 2005 | 7 | 4 | 29 (13) | 1 | 3 |
| 2006 | 0 | 2 | 22 (19) | 4 | 3 |
| 2007 | 3 | 4 | 20 (7) | 3 | 3 |
| 2008 | 2 | 2 | 21(12) | 4 | 3 |

Sources:USD/BLM and references therein; Burns et al. (1993); Koski et al. (2005); Suydam et al. 2004, 2005, 2006, 2007, 2008, 2009.

Ringed seals are hunted by villagers along the Alaskan north coast mainly from October through June. Hunting for these smaller mammals is concentrated during winter because bowhead whales, bearded seals and caribou are available through other seasons. Winter leads in the area off Pt. Barrow and along the barrier islands of Elson Lagoon to the east are used for hunting ringed seals. The average annual ringed seal harvest by the community of Barrow from the 1960s through much of the 1980s has been estimated as 394 (Table 7). Although ringed seals are available year-round, the seismic survey will not occur during the primary period when these seals are harvested. Also the seismic survey in offshore waters will not influence ringed seals in the nearshore areas where they are hunted.

Table 7. Average annual take of marine mammals other than bowhead whales harvested by the community of Barrow (as compiled by LGL Alaska Res. Assoc. 2004).

| Beluga Whales | Ringed Seals | Bearded Seals | Spotted Seals |
|---------------|--------------|---------------|---------------|
| 5 ** | 394 * | 174* | 1* |

* Average annual harvest for years 1987-90 (Braund et al. 1993).

** Average annual harvest for years 1962-82 (MMS 1996).

The **spotted seal** subsistence hunt peaked in July and August, at least in 1987 to 1990, but involves few animals. Spotted seals typically migrate south by October to overwinter in the Bering Sea. Admiralty Bay, <60 km to the east of Barrow, is a location where spotted seals are harvested. Spotted seals

are also occasionally hunted in the area off Pt. Barrow and along the barrier islands of Elson Lagoon to the east (USDI/BLM 2005). The average annual spotted seal harvest by the community of Barrow from 1987-1990 was one (Braund et al. 1993; Table 7). Spotted seals become less abundant at Nuiqsut and Kaktovik and few if any spotted seal are harvested at these villages. The seismic survey will commence at least 115 km offshore from the preferred nearshore harvest area of these seals.

Bearded seals, although not favored for their meat, are important to subsistence activities in Barrow because of their skins. Six to nine bearded seal hides are used by whalers to cover each of the skin-covered boats traditionally used for spring whaling. Because of their valuable hides and large size, bearded seals are specifically sought. Bearded seals are harvested during the summer months in the Beaufort Sea (USDI/BLM 2005). The animals inhabit the environment around the ice floes in the drifting ice pack, so hunting usually occurs from boats in the drift ice. Braund et al. (1993) estimated that 174 bearded seals were harvested annually at Barrow from 1987-1990 (Table 7). Braund et al. (1993) mapped the majority of bearded seal harvest sites from 1987 to 1990 as being within ~24 km of Point Barrow, well inshore of the proposed survey which is to start >115 km offshore and terminate >200 km offshore.

The USFWS has monitored the harvest of **polar bears** in Alaska using a mandatory marking, tagging, and reporting program implemented in 1988. Polar bears are harvested in winter and spring, but comprise a small percent of the annual Native subsistence harvest. Braund et al. (1993) reported that ~2% of the total edible pounds harvested by Barrow residents from 1987 to 1989 involved polar bears. The USFWS estimated that from 1996 to 2000 the average annual harvest of polar bears in Alaska was ~45 animals (Angliss and Allen 2009). It is not expected that the seismic survey will interfere with polar bear subsistence hunting due to the limited annual harvest documented by USFWS and the fact that the subsistence hunt typically takes place in the winter and spring, either well after or well before the scheduled survey (Angliss and Allen 2009; USFWS 2009b). **Walrus** are hunted primarily from June through mid-August to the west of Point Barrow and southwest to Peard Bay. Walrus rarely occur in the Beaufort Sea north and east of Barrow and become less abundant further east. The harvest effort peaks in July. The annual walrus harvest by Barrow residents ranged from 7 to 206 animals from 1990 to 2002 (Fuller and George 1999). It is possible, but unlikely, that accessibility to walrus during the subsistence hunt could be impaired during the *Healy's* transit north of Barrow to the starting point of the seismic survey. The area affected, however, would be in close proximity to the ship and disturbance would be of short duration as the vessel passed through the area. The majority of marine mammals are taken by hunters within ~33 km offshore (Fig. 9), and survey operations will not commence until the *Healy* rendezvous with the *Louis S. St. Laurent* significantly farther offshore (>115 km).

Helicopter operations will be far offshore where the seismic operations are occurring, and thus any reactions of marine mammals to the helicopter operations will have no effects on availability of marine mammals for subsistence. Furthermore, helicopter operations will be conducted in a manner that will minimize helicopter effects on marine mammals.

No survey operations are proposed in any areas used for subsistence purposes by Alaska Natives. The *Healy* may transit near some areas of subsistence use but disturbance would be temporary and unlikely to disrupt any subsistence hunting activities. The bowhead hunt near Barrow normally does not begin until mid-September, well after the *Healy* will pass the Barrow area prior to the start of the proposed surveys. Based on recent bowhead harvest dates (Suydam et al. 2005b, 2006, 2007, 2008, 2009), the *Healy* will likely complete the survey activities and transit through the Barrow area in mid-September prior to the start of the fall whaling season. The proposed geophysical survey activities, including transit periods, should have no effect of subsistence hunting activities for marine mammals.

(b) Subsistence Fishing

Subsistence fishing is conducted by Alaska Natives through the year, but most actively during the summer and fall months. Barrow residents often fish for camp food while hunting, so the range of subsistence fishing is widespread. Marine subsistence fishing occurs during the harvest of other subsistence resources in the summer. Fishing occurs in areas much closer to shore however, than the location of the proposed geophysical survey (MMS 1996) and subsistence fishing activity will not be affected by the proposed geophysical survey.

Seismic surveys can, at times, cause changes in the catchability of fish. Airgun operations are not planned to occur anywhere within 115 km of shore. However, in the highly unlikely event that subsistence fishing (or hunting) is occurring within 5 km (3 mi) of the survey operations, the airgun operations will be suspended until the *Louis S. St. Laurent* is >5 km away.

(c) Consultation with Local Barrow Community

USGS is working with the community of Barrow to identify and avoid areas of potential conflict. A representative of the project has met with North Slope Borough Department of Wildlife Management biologist Robert Suydam to discuss concerns of North Slope residents.

(14) Cumulative Effects

Cumulative effects refer to the impacts on the environment that result from a combination of past, existing, and imminent projects and human activities. Agents of cumulative effects can include multiple causes, multiple effects, effects of activities in more than one locale, and recurring events.

Human activities in the Alaskan Beaufort Sea and Arctic Ocean include whaling and sealing, commercial fishing, oil and gas development, and vessel traffic. These activities, when conducted separately or in combination with other activities, can affect marine mammals in the study area. Any cumulative effects caused by the addition of the seismic survey impacts on marine mammals will be extremely limited, especially considering the timeframe of the proposed activities and the location of the proposed survey area well offshore of the Alaska coast.

(a) Commercial Fishing

Commercial fisheries in the Alaskan Beaufort Sea are very limited. The Helmericks family operates an under-ice commercial gill net fishery during fall in the Colville River delta, well over 100 km southeast of the closest part of the present study area (Gallaway et al. 1983, 1989). The fishery typically operates from early October through the end of November. Fishing effort is concentrated in the Main (Kupigruak) and East Channels of the river near Anachilik Island. The three principal species targeted in the fishery are Arctic cisco, least cisco, and humpback whitefish. The timing of the proposed geophysical survey and its location well offshore will eliminate any conflicts with coastal fisheries.

The proposed survey will have a negligible impact on the marine mammals in the study area. The combination of USGS's activities with those of fisheries will not result in any detectable increment in impacts on marine mammals over and above the impacts from fisheries alone.

(b) Oil and Gas Development

Oil and gas development in the Alaskan Beaufort Sea and on the Arctic Coastal Plain has been considerable. USDI/MMS (2003) listed 17 offshore North Slope oil and gas discoveries and 46 onshore discoveries as of 1 July 2002.

Recent oil field developments include Alpine (onshore), which came on line in November 2000 and now produces ~90,000 barrels of oil per day; Northstar (offshore), which began production October 2001 and is currently producing ~22,477 barrels of oil per day; and the Pioneer Natural Resources

development at Oooguruk Drill Site in eastern Harrison Bay which began production in 2008. The Northstar production facility is the only one that is currently operating in the Beaufort Sea north (seaward) of the barrier islands. The offshore (but in a lagoon) Endicott field began production in 1987 and had produced 439 million barrels of oil through Feb 1995 (AOGCC 2005). The Niakuk, Pt. McIntyre, and Badami fields are located offshore, but production facilities are located onshore. The Alpine oil field is the westernmost of the oil field developments and is ~ 241 km southeast of Barrow. Two other developments which may come into production within the next several years include the BP Liberty development and the Eni Spy Island development.

The existing oil fields are serviced by land, air, and sea. Marine activities associated with the on-land oil developments in northern Alaska consist mainly of tug and barge traffic, mainly in nearshore waters along the north coast. Vessel traffic including barges and crew boats to Northstar Island have been ongoing during the open-water season, although much of the crew vessel traffic has been largely replaced by hovercraft and helicopter traffic, neither of which introduces much noise into the sea (Blackwell and Greene 2005). During the last several years barges and crew vessels have been used in support of activities at Pioneer's Oooruruk site, and in support of island construction by Eni at their Spy Island Drillsite located inside Spy Island in eastern Harrison Bay. Several supply vessels travel along the Beaufort Sea coast, transporting fuel and construction materials to communities and industrial centers. Two or three supply vessels routinely travel between Barrow and Kaktovik during the summer, with two additional vessels operating out of Prudhoe Bay.

Open-water industry seismic surveys were conducted in the Alaskan Chukchi and Beaufort seas each year since 2006 during the open-water season. BP and Eni also had smaller ocean bottom cable seismic survey programs in the general Prudhoe Bay area in 2008. Other seismic survey programs were conducted in the southern Alaskan Beaufort Sea from 1996–2001 (Richardson and Lawson 2002). These surveys occurred much closer to shore than the proposed USGS survey and may be ongoing in 2010. The timing of the potential industry surveys in 2010 is not precisely known but could overlap temporally but not spatially with the proposed *Louis S. St. Laurent* activities.

In addition to the potential for continued industry seismic exploration, offshore exploratory drilling may also occur in 2010 in the southern Beaufort Sea and the Chukchi Sea. Exploratory drilling activities in the Beaufort Sea would occur in nearshore locations and would not overlap spatially with the proposed USGS surveys.

Oil industry activities will likely be ongoing in the central part of Alaska's Beaufort Sea coast during the proposed seismic survey, but such activities are located >100 km south of the southernmost extent of the USGS surveys, with no spatial overlap. Noise generated by oil industry activities in the nearshore zone, such as at Northstar, generally is not detectable underwater more than a few km from facilities although vessel sound may be audible at greater distances (Blackwell and Greene 2006). Underwater sounds from vessels supporting oil industry activities are often detectable farther away. However, the proposed survey route will take the *Louis S. St. Laurent* well north of the central Alaskan Beaufort Sea coast and there will be no encounters with vessels servicing the oil fields. The activities associated with the proposed USGS survey may add incrementally to the overall cumulative impacts associated with various industry activities although any effects are expected to be negligible.

(c) Vessel Traffic

In heavily-traveled areas, shipping noise generally dominates ambient noise at frequencies from 20 to 300 Hz, although that is not the case in most of the Arctic (Richardson et al. 1995). Baleen whales are thought to be more sensitive to sound at those low frequencies than are toothed whales. There may be some avoidance by marine mammals of the two vessels operating in the proposed geophysical survey area. Bowhead whales,

in particular, often move away when vessels approach within several kilometers (Richardson and Malme 1993), and hunters at Barrow believe that vessel traffic near the coast southeast of Barrow can cause larger-scale displacement of bowheads. However, migrating bowheads are not expected to arrive in that area, or in the area where the *Louis S. St. Laurent* will operate, until after the *Louis S. St. Laurent* has completed the survey.

Responses of belugas to vessel traffic are highly variable (Richardson et al. 1995), and can extend to tens of kilometers in special circumstances (Finley et al. 1990). Belugas may also be tolerant of large vessels traveling in consistent directions but may flee from fast erratic movements from smaller boats (Richardson et al. 1995).

Aside from vessels supporting the oil industry (discussed in preceding subsection), vessel traffic in the proposed study area is limited. The majority of the other vessels will be within 20 km of the coast, and may include Native vessels used for fishing and hunting, cruise ships, icebreakers, Coast Guard vessels, and supply ships. Several supply vessels are also scheduled to visit the North Slope communities from Barrow to Kaktovik and on to Canada delivering fuel and construction equipment.

The addition of the proposed survey activities will not augment the impacts to marine mammals that occur due to routine vessel traffic in the area of the survey.

(d) Oil Spills

There is always the risk of an oil spill from a vessel. However, the *Healy* and *Louis S. St. Laurent* are well maintained and operated to high standards, with five engines and triple props. It is highly unlikely that these vessels will be the source of an oil spill of any significant size. Fuel capacities are relatively trivial when compared to the amount of oil produced from the offshore fields in the Beaufort Sea and the risk of a spill from either the *Healy* or the *Louis S. St. Laurent* is relatively low.

(e) Hunting

Marine mammals are legally hunted in Alaskan waters by coastal Alaska Natives. In the Alaskan Beaufort and Chukchi seas, bowhead whales, beluga whales, Pacific walruses, ringed, spotted, and bearded seals, and polar bears are hunted (see Section IV[8]). The hunting communities within the area of the proposed survey are Barrow, Nuiqsut, and Kaktovik in the Alaskan Beaufort Sea. The planned project (unlike subsistence hunting activities) will not result in directed or lethal takes of marine mammals. Also, the direct disturbance-related impacts of the project on individuals are anticipated to be short-term and inconsequential to the long-term well being of those individuals and their populations. Thus, the combined effects of the project and of subsistence hunting on marine mammal stocks are not expected to differ appreciably from those of subsistence hunting alone.

(f) Summary of Cumulative Impacts

For the majority of the proposed trackline, the *Healy* and *Louis S. St. Laurent* are unlikely to encounter any additional human activities, and thus the degree of cumulative impact will be minimal. Any such effects related to the cumulation of human activities near the start and end of the trackline will have no more than a negligible impact on the marine mammal populations encountered.

(15) Unavoidable Impacts of Noise

Unavoidable impacts to marine mammals, seabirds, or fish occurring in the proposed study area in the Arctic Ocean will be limited to short-term changes in behavior and local distribution. For cetaceans and pinnipeds, some of the changes in behavior may be sufficient to fall within the MMPA definition of “Level B Harassment” (behavioral disturbance; no serious injury or mortality). No long-term or significant impacts are expected on any individual marine mammals, seabirds, or on the populations to

which they belong. Effects on recruitment or survival are expected to be (at most) negligible. Also, any effects on accessibility of marine mammals for subsistence hunting and effects on commercial fishing are expected to be (at most) negligible.

(16) Coordination with Other Agencies and Processes

This EA has been prepared for and adopted by USGS to address issues relating to the request that an IHA be issued by NMFS to authorize “taking by harassment” (disturbance) of small numbers of cetaceans and pinnipeds during USGS’s planned seismic survey. USGS is the Federal funding agency for the geophysical survey work. Another important component has been to address potential impacts on polar bears, walruses, and seabirds, which are managed by USFWS.

USGS will coordinate the planned marine mammal monitoring program associated with the seismic survey in the Arctic Ocean with other parties that may have interest in this area and/or be conducting marine mammal studies in the same region during operations. No other marine mammal studies are expected to occur in the main (northern) parts of the survey area at the proposed time. However, other industry-funded seismic surveys may be occurring in the northeast Chukchi and/or western Beaufort Sea closer to shore, and those projects are likely to involve marine mammal monitoring. Further coordination of monitoring programs can occur during and after the planned Beaufort open-water peer review meeting in Anchorage in spring 2010.

The USGS has communicated with community authorities and residents of Barrow to foster understanding of the proposed survey. There are elements of the proposed survey, intrinsic to the project, that significantly limit the potential conflict with subsistence users. Operations will be conducted during early August before bowhead whale hunting typically occurs off Barrow and ~108 km offshore, farther offshore than traditional subsistence hunting grounds. USGS continues to work with the people of Barrow to identify and avoid areas of potential conflict.

- The USGS initiated contact with North Slope Borough (NSB) scientists and the chair of the AEWG in mid-December 2010 via an emailed description of the proposed survey that included components intended to minimize potential subsistence conflict.
- Invitations were extended on 31 December 2009 to members of the NSB, AEWG and North Slope Communities to attend a teleconference arranged for 11 January 2010. The teleconference served as a venue to promote understanding of the project and discuss shareholder concerns. Participants in the teleconference included Harry Brower, chair of the AEWG, and NSB wildlife biologist Dr. Robert Suydam.
- To further promote cooperation between the project researchers and the community, Dr. Deborah Hutchinson with USGS presented details of the proposed survey at a meeting of the AEWG in Barrow on 11 February 2010. Survey plans were explained to local hunters and whaling captains, including NSB Department of Wildlife Management biologists, Craig George and Robert Suydam. Dr. Hutchinson consulted with stakeholders about their concerns and discussed the aspects of the survey designed to mitigate impacts.
- Dr. Deborah Hutchinson of the USGS emailed a summary of the topics discussed during the teleconference and the AEWG meeting in Barrow to representatives of the NSB, AEWG and North Slope communities. These included:
 - Surveying within U.S. waters is scheduled early (~7-12 August) to avoid conflict with hunters

- The EA and IHA application will be distributed as early as possible to NSB and AEW C
- A community observer will be present aboard the *Healy* during the project
- Mitigation of the one crew transfer near Barrow in early September will be arranged – probably through Barrow Volunteer Search and Rescue
- Representatives of the USGS attended the Arctic Open-water Meeting in Anchorage, 22-24 March.
 - Dr. Deborah Hutchinson presented information regarding the proposed survey to the general assembly
 - Drs. Jonathan Childs (USGS) and Deborah Hutchinson met with stakeholders and agency representatives while at the meeting

Subsequent meetings with whaling captains, other community representatives, the AEW C, NSB, and any other interested parties will be held if necessary to coordinate the planned seismic survey operation with subsistence hunting activity. The USGS has informed the chairman of the AEW C, Harry Brower, Jr., of its survey plan and met with representatives during the open water meeting in Anchorage in March.

As noted above in § VIII, in the unlikely event that subsistence hunting or fishing is occurring within 5 km (3 mi) of the project vessel tracklines, or where potential impacts could occur, the airgun operations will be suspended until the vessel is >5 km away and otherwise not interfering with subsistence activities.

Pursuant to Council on Environmental Quality's regulations implementing NEPA (40 CFR 1501.1), the National Oceanic and Atmospheric Administration (NOAA), and US Coast Guard have been granted cooperating agency status in development of this EA. NOAA's National Marine Fisheries Service (NMFS) has jurisdiction by law to issue permits and authorizations under the Marine Mammal Protection Act (MMPA) and the Endangered Species Act (ESA). NOAA's Office of Oceanic and Atmospheric Research (OAR) is a partner agency for the project with the USGS and shares responsibility for funding the proposed action. It is the role of the NOAA NEPA Coordinator in the Office of Program Planning and Integration (PPI) to coordinate matters for NOAA on NEPA processes where multiple offices are involved; therefore, PPI is requesting cooperating agency status on behalf of both NMFS and OAR. The U.S. Coast Guard (Coast Guard) is a subject matter expert for topics related to navigation safety and maritime security for facilities and/or equipment located on, under, or adjacent to the navigable waters of the U.S. The Coast Guard seeks opportunity to contribute information on topics for which it has subject matter expertise.

Alternative Action: Conduct Survey during Alternative Time Period

The proposed project will take ~30 days and is expected to occur from approximately 6 August to 3 September 2010. An alternative to issuing the IHA for the period requested, and to conducting the project within that period, is to issue the IHA for another period, and to conduct the project during that alternative period. However, conducting the project at some other time of year outside the summer period could result in impracticalities related to ice conditions. In addition, the proposed period for the cruise is the period when the ships and all of the personnel and equipment essential to meet the overall project objectives are available. Postponing or changing the project period will delay this and potentially other scheduled projects during the rest of 2010.

Marine mammals are expected to be found throughout the proposed study area and throughout the time period during which the project may occur. Ringed seals, the most abundant marine mammal in the area of the survey, are year-round residents in Alaska (see § III, above), so altering the timing of the proposed project likely would result in no net benefits for that species. Bowhead and beluga whales are migratory, moving through the Beaufort Sea and possibly the proposed survey area in the spring and then again in the fall (see § III, above). The cruise has been timed to avoid the bowhead migration, and the main part of the beluga migration. Delay until later in the 2010 open-water period would move the *Healy* and *Louis S. St. Laurent* cruise closer to (or into) the main migration periods for those whale species. For other marine mammal species there are insufficient data to predict when their abundance may be highest.

Subsistence harvests of ringed seals, bearded seals, and bowhead whales occur near North Slope coastal villages, far south of the southern portion of the survey track. Marine mammal harvests take place year-round, but subsistence harvest peaks during the bowhead whale hunts in the spring and fall. The harvest is of great value to the Inupiat people, both culturally and as a food source. The survey has been scheduled to avoid the bowhead whale migration and subsistence harvest of bowheads.

The proposed 2010 survey will occur during the final year of a four-year data acquisition program. Postponement or delay of the 2010 survey would result in a lost opportunity for the U.S. to meet data collection goals critical to defining the ECS. Conducting surveys outside the US EEZ in previous years has served to ensure that there is an extended continental shelf beyond the US EEZ north of Alaska before surveying within the 200-nmi limit. After 2010, *Louis* will be moved to other parts of the Canadian continental margin for surveying; hence she will not be available for work in the Canada Basin.

No Action Alternative

An alternative to conducting the proposed activities is the “No Action” alternative, i.e., the geophysical survey is not conducted and no IHA is issued. If the research were not conducted, the “No Action” alternative would result in no disturbance to marine mammals attributable to the proposed seismic activities. Likewise, there would then be no possibility of effects on fisheries or on accessibility of marine mammals for subsistence hunting. Little reduction in impacts would occur however, if the project was not undertaken, given the expected negligible effects on marine mammals, seabirds, fish, subsistence hunting, and fisheries that are anticipated if the project goes ahead as planned. However, cancellation of this project would result in a loss of potentially important scientific data and knowledge that could affect the size of the offshore area under U.S. jurisdiction and the ability of the U.S. to develop the appropriate data for defining the outer limit of the ECS.

V. LIST OF PREPARERS

LGL Alaska Research Associates Inc. and LGL Ltd., environmental research associates

Robert Rodrigues, B.S., Anchorage, AK*

Beth Haley, B.A., Anchorage, AK *

Darren Ireland, M.Sc., Anchorage, AK*

U.S. Geological Survey

Deborah Hutchinson, PhD., Woods Hole, MA*

Jonathan R. Childs, M.Sc., Menlo Park, CA*

U.S. Coast Guard

National Oceanic and Atmospheric Administration

* Principal preparers of this specific document. Others listed above contributed to a lesser extent, or contributed substantially to previous related documents from which material has been excerpted.

VI. LITERATURE CITED

- Aagaard, K. 1989. A synthesis of the Arctic Ocean circulation. *Rapports et Proces-verbaux des Reunion Conseil International pour l'Exploration de la Mer* 188:11-22.
- Aars, J., N.J. Lunn, and A.E. Derocher. eds. 2006. Polar bears: proceedings of the 14th working meeting of the IUCN/SSC Polar Bear Specialist Group, 20-24 June, Seattle, Washington, USA. IUCN, Gland, Switzerland. 189 pp.
- ADFG (Alaska Department of Fish and Game). 1994. Orca: Wildlife Notebook Series. Alaska Dep. Fish & Game. Available at www.adfg.state.ak.us/pubs/notebook/marine/orca.php
- AOGCC (Alaska Oil and Gas Conservation Commission). 2005. Production Data Index. Available at www.state.ak.us/local/akpages/ADMIN/ogc/production/pindex.htm
- Amstrup, S.C. 1995. Movements, distribution, and population dynamics of polar bears in the Beaufort Sea. Ph.D. Dissertation. Univ. Alaska-Fairbanks, Fairbanks, AK. 299 p.
- Amstrup, S.C. and D.P. DeMaster. 1988. Polar bear (*Ursus maritimus*), p. 39-56 In: J.W. Lentfer, (ed.) Selected marine mammals of Alaska: Species Accounts with Research and Management Recommendations. Mar. Mamm. Comm., Washington, DC.
- Amstrup, S.C. and C. Gardner. 1994. Polar bear maternity denning in the Beaufort Sea. **J. Wildl. Manage.** 58(1):1-10.
- Amstrup, S.C., I. Stirling and J.W. Lentfer. 1986. Past and present status of polar bears in Alaska. **Wildl. Soc. Bull.** 14(3):241-254.
- Amstrup, S. C., G. M. Durner, I. Stirling, N. J. Lunn, and F. Messier. 2000. Movements and distribution of polar bears in the Beaufort Sea. **Can. J. Zool.** 78(6):948-966.
- Angliss, R.P. and K.L. Lodge. 2004. Alaska marine mammal stock assessments, 2003. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-144. 230 p.
- Angliss, R.P., and B.M. Allen. 2009. Alaska Marine Mammal Stock Assessments, 2008. NOAA Technical Memorandum NMFS-AFSC-193.
- Arbelo, M., M. Méndez, E. Sierra, P. Castro, J. Jaber, P. Calabuig, M. Carrillo and A. Fernández. 2005. Novel “gas embolic syndrome” in beaked whales resembling decompression sickness. Abstr. 16th Bien. Conf. Biol. Mar. Mamm., San Diego, CA, 12-16 Dec. 2005.
- Au, W.W.L., A.N. Popper and R.R. Fay. 2000. Hearing by Whales and Dolphins. Springer-Verlag, New York, NY. 458 p.
- Balcomb, K.C., III and D.E. Claridge. 2001. A mass stranding of cetaceans caused by naval sonar in the Bahamas. **Bahamas J. Sci.** 8(2):2-12.
- Barlow, J. 1999. Trackline detection probability for long-diving whales. p. 209-221 In: G.W. Garner, S.C. Amstrup, J.L. Laake, B.F.J. Manly, L.L. McDonald and D.G. Robertson (eds.), Marine mammal survey and assessment methods. A.A. Balkema, Rotterdam. 287 p.
- Belikov, S., A. Boltunov and Y. Gorbunov. 1996. Distribution and migration of polar bears, pacific walruses and gray whales depending on ice conditions in the Russian Arctic. *Proc. NIPS Symp. Polar Biol.* 9:263-274.
- Bengtson, J.L., L.M. Hiruki-Raring, M.A. Simpkins, and P.L. Boveng. 2005. Ringed and bearded seal densities in the eastern Chukchi Sea, 1999-2000. *Polar Biology* 28:833-845.
- Bigg, M.A. 1981. Harbour seal, *Phoca vitulina* and *P. largha*. p. 1-28 In: S.H. Ridgway and R.J. Harrison (eds.), Handbook of Marine Mammals, Vol. 2: Seals. Academic Press, New York, NY. 359 p.
- Blackwell, S.B., and C.R. Greene, Jr. 2005. Underwater and in-air sounds from a small hovercraft. **J. Acoust. Soc. Am.** 116(6):3646-3652.

- Blackwell, S.B., and C.R. Greene, Jr. 2006. Sounds from an oil production island in the Beaufort Sea in summer: Characteristics and contribution of vessels. **J. Acoust. Soc. Am.** 119(1):182-196.
- Blackwell, S.B., R.G. Norman, C.R. Greene Jr., M.W. McLennan, T.L. McDonald and W.J. Richardson. 2004. Acoustic monitoring of bowhead whale migration, autumn 2003. p. 71 to 744 *In*: Richardson, W.J. and M.T. Williams (eds.) 2004. Monitoring of industrial sounds, seals, and bowhead whales near BP's Northstar oil development, Alaskan Beaufort Sea, 1999-2003. [Dec. 2004 ed.] LGL Rep. TA4002. Rep. from LGL Ltd. (King City, Ont.), Greeneridge Sciences Inc. (Santa Barbara, CA) and WEST Inc. (Cheyenne, WY) for BP Explor. (Alaska) Inc., Anchorage, AK. 297 p. + Appendices A - N on CD-ROM.
- Blackwell, S.B., C.R. Greene, Jr., T.L. McDonald, M.W. McLennan, C.S. Nations, R.G. Norman, and A. Thode. 2009. Beaufort Sea acoustic monitoring program. Chapter 8 *In* Funk, D.W., R. Rodrigues, D.S. Ireland, and W.R. Koski (eds.). Joint monitoring program in the Chukchi and Beaufort seas, July–November 2007. LGL Alaska Report P971-2. Report from LGL Alaska Research Associates, Inc., Anchorage, Ak, LGL Ltd., environmental research associates, King City, Ont., JASCO Research, Victoria, B.C., and Greeneridge Sciences, Inc., Goleta, CA, for Shell Offshore, Inc., ConocoPhillips Alaska, Inc., and National Marine Fisheries Service, and U.S. Fish and Wildlife Service.
- Bluhm, B.A., K.O. Coyle, B. Konar and R. Highsmith. 2007. High gray whale relative abundances associated with an oceanographic front in the south-central Chukchi Sea. **Deep-sea Research II** 54:2919-2933.
- Boebel, O., H. Bornemann, M. Breitzke, E. Burkhardt, L. Kindermann, H. Klinck, J. Plotz, C. Ruholl and H.-W. Schenke. 2004. Risk Assessment of ATLAS HYDROSWEEP DS-2 Hydrographic Deep Sea Multi-beam Sweeping Survey Echo Sounder. Poster at the International Policy Workshop on Sound and Marine Mammals, Marine Mammal Commission and Joint Nature Conservation Committee, London, 2004. Available at www.mmc.gov/sound/internationalwrkshp/pdf/poster_03boebel.pdf
- Born, E.W., F.F. Riget, R. Dietz and D. Andriashek. 1999. Escape responses of hauled out ringed seals (*Phoca hispida*) to aircraft disturbance. **Polar Biol.** 21(3):171-178.
- Bowles, A.E., M. Smultea, B. Würsig, D.P. DeMaster and D. Palka. 1994. Relative abundance and behavior of marine mammals exposed to transmissions from the Heard Island Feasibility Test. **J. Acoust. Soc. Am.** 96(4):2469-2484.
- Braham, H.W. 1984. Distribution and migration of gray whales in Alaska. p. 249-266 *In*: M.L. Jones, S.L. Swartz and S. Leatherwood (eds.), The Gray Whale *Eschrichtius robustus*. Academic Press, Orlando, FL. 600 p.
- Braham, H.W. and B.D. Krogman. 1977. Population biology of the bowhead whale (*Balaena mysticetus*) and beluga (*Delphinapterus leucas*) whale in the Bering, Chukchi and Beaufort Seas. U.S. Dep. Comm., Seattle, WA.
- Braham, H.W., B.D. Krogman and G.M. Carroll. 1984. Bowhead and white whale migration, distribution, and abundance in the Bering, Chukchi, and Beaufort seas, 1975-78. NOAA Tech. Rep. NMFS SSRF-778. USDOC/NOAA/NMFS. 39 p. NTIS PB84-157908. 39 p.
- Braund, S.R. and E.L. Moorehead. 1995. Contemporary Alaska Eskimo bowhead whaling villages. p. 253-279 *In*: A.P. McCartney (ed.), Hunting the Largest Animals/Native Whaling in the Western Arctic and Subarctic. Studies in Whaling 3. Can. Circumpolar Inst., Univ. Alberta, Edmonton, Alb. 345 p.
- Braund, S.R., K. Brewster, L. Moorehead, T. Holmes and J. Kruse. 1993. North Slope subsistence study/Barrow 1987, 1988, 1989. OCS Study MMS 91-0086. Rep. from Stephen R. Braund & Assoc. and Inst. Social & Econ. Res., Univ. Alaska Anchorage. 466 p.
- Brewer, K.D., M.L. Gallagher, P.R. Regos, P.E. Isert and J.D. Hall. 1993. ARCO Alaska, Inc. Kuvlum #1 exploration prospect/Site specific monitoring program final report. Rep. from Coastal & Offshore Pacific Corp, Walnut Creek, CA, for ARCO Alaska Inc., Anchorage, AK. 80 p.
- Brower, H., Jr. 1996. Observations on locations at which bowhead whales have been taken during the fall subsistence hunt (1988 through 1995) by Eskimo hunters based in Barrow, Alaska. North Slope Borough Dep. Wildl. Manage., Barrow, AK. 8 p. Revised 19 Nov. 1996.

IV. Literature Cited

- Brueggeman, J.J., C.I. Malme, R.A. Grotefendt, D.P. Volsen, J.J. Burns, D.G. Chapman, D.K. Ljungblad and G.A. Green. 1990. Shell Western E & P Inc. 1989 Walrus Monitoring Program: The Klondike, Burger, and Popcorn Prospects in the Chukchi Sea. Report prepared by EBASCO Environmental for Shell Western E & P Inc. 157 p.
- Brueggeman, J.J., G.A. Green, R.A. Grotefendt, M.A. Smultea, D.P. Volsen, R.A. Rowlett, C.C. Swanson, C.I. Malme, R. Mlawskie and J.J. Burns. 1992. 1991 marine mammal monitoring program (walrus and polar bear) Crackerjack and Diamond prospects Chukchi Sea. Rep. from EBASCO Environmental for Shell Western E & P Inc. and Chevron U.S.A. Inc. Var. pag.
- Buchanan, R.A., J.R. Christian, V.D. Moulton, B. Mactavish and S. Dufault. 2004. 2004 Laurentian 2-D seismic survey environmental assessment. Rep. from LGL Ltd., St. John's, Nfld., and Canning & Pitt Associates, Inc., St. John's, Nfld., for ConocoPhillips Canada Resources Corp., Calgary, Alta. 274 p.
- Burn, D.M., M.A. Webber, and M.S. Udevitz. 2006. Application of airborne thermal imagery to surveys of Pacific walrus. *Wildlife society Bulletin* 34:51-58.
- Burns, J.J. 1970. Remarks on the distribution and natural history of pagophilic pinnipeds in the Bering and Chukchi Seas. *J. Mammal.* 51(3):445-454.
- Burns, J.J. 1981. Bearded seal *Erignathus barbatus* Erxleben, 1777. p. 145-170 *In* S.H. Ridgway and R.J. Harrison (eds.), *Handbook of Marine Mammals, Vol. 2 Seals*. Academic Press, New York.
- Burns, J.J., L.H. Shapiro, and F.H. Fay. 1981. Ice as marine mammal habitat in the Bering Sea. *In: The Eastern Bering Sea Shelf: Oceanography and Resources*, D.W. Hoome and J.A. Calder (eds.). Vo. II. Juneau, AK. OMPA, BLM.
- Burns, J.J., J.J. Montague and C.J. Cowles (eds.). 1993. The bowhead whale. *Spec. Publ. 2, Soc. Mar. Mamm.*, Lawrence, KS. 787 p.
- Calambokidis, J. and S.D. Osmek. 1998. Marine mammal research and mitigation in conjunction with air gun operation for the USGS SHIPS seismic surveys in 1998. Draft rep. from Cascadia Research, Olympia, WA, for U.S. Geol. Surv., Nat. Mar. Fish. Serv., and Minerals Manage. Serv.
- Caldwell, J. and W. Dragoset. 2000. A brief overview of seismic air-gun arrays. **The Leading Edge** 19(8):898-902.
- CBD. 2007. Petition to list the ribbon seal (*Histriophoca fasciata*) as a threatened or endangered species under the Endangered Species Act. Center for Biological Diversity, San Francisco, CA.
- CBD. 2008a. Petition to list three seal speices under the Endangered Species Act: ringed seal (*Pusa hispida*), bearded seal (*Erignathus barbatus*), an dspotted sea (*Phoca largha*). Center for Biological Diversity, San Francisco, CA.
- CBD. 2008b. Petition to list the Pacific walrus (*Odovenus rosmarus divergens*) as a threatened or endangered species under the Endangered Species Act. Center for Biological Diversity, San Francisco, CA.
- Chadwick, V.J. and S. Hills. 2005. Movements of walruses radio-tagged in Bristol Bay, Alaska. **Arctic** 58(2):192-202.
- Christi, K., C. Lyons, and W.R. Koski. 2009. Beaufort Sea aerial monitoring program. Chapter 7 *In* Funk, D.W., D.S. Ireland, R. Rodrigues, and W.R. Koski (eds.). 2009. Joint monitoring program in the Chukchi and Beaufort seas, July–November 2006-2008. LGL Alaska Report P1050-1. Report from LGL Alaska Research Associates, Inc., Anchorage, Ak, LGL Ltd., environmental research associates, King City, Ont., Greeneridge Sciences, Inc., Goleta, CA, and JASCO Research, Victoria, B.C., for Shell Offshore, Inc. and other Industry contributors, National Marine Fisheries Service, and U.S. Fish and Wildlife Service. 488 p. plus appendices.
- Clark, J.T. and S.E. Moore. 2002. A note on observations of gray whales in the southern Chukchi and northern Bering Seas, AuguSt. November, 1980-1989. **J. Cetac. Res. Manage.** 4(3):283-288.

- Clarke, J.T., S.E. Moore and D.K. Ljungblad. 1989. Observations on gray whale (*Eschrichtius robustus*) utilization patterns in the northeastern Chukchi Sea, July-October 1982-1987. **Can. J. Zool.** 67(11):2646-2654.
- Clarke, J.T., S.E. Moore and M.M. Johnson. 1993. Observations on beluga fall migration in the Alaskan Beaufort Sea, 198287, and northeastern Chukchi Sea, 198291. **Rep. Int. Whal. Comm.** 43:387-396.
- Cosens, S.E. and L.P. Dueck. 1993. Icebreaker noise in Lanaster Sound, N.W.T., Canada: Implications for marine mammal behaviour. **Mar. Mam. Sci** 9(3), 258-300.
- Coyle, K.O., B. Bluhm, B. Konar, A. Blanchard and R.C. Highsmith. 2007. Amphipod prey of gray whales in the northern Bering Sea: Comparison of biomass and distribution between the 1980s and 2002-3. **Deep-sea Research II** 54:2906-2918.
- Dahlheim, M.E. and J.E. Heyning. 1999. Killer whale *Orcinus orca* (Linnaeus, 1758). p. 281-322 *In*: S.H. Ridgway and R. Harrison (eds.), Handbook of Marine Mammals, Vol. 6: The Second Book of Dolphins and the Porpoises. Academic Press, San Diego, CA. 486 p.
- Davis, R.A. and C.R. Evans. 1982. Offshore distribution and numbers of white whales in the eastern Beaufort Sea and Amundsen Gulf, summer 1981. Rep. from LGL Ltd., Toronto, Ont., for Sohio Alaska Petrol. Co., Anchorage, AK, and Dome Petrol. Ltd., Calgary, Alb. (co-managers). 76 p.
- DeMaster, D.P. 1995. Minutes from the 4-5 and 11 January 1995 meeting of the Alaska Scientific Review Group. Anchorage, Alaska. 27 p. + app. Available upon request - D.P. DeMaster, Alaska Fisheries Science Center, 7600 Sand Point Way, NE, Seattle, WA 98115.
- DeMaster, D.P. and I. Stirling. 1981. *Ursus maritimus*. **Mamm. Species** 145. 7 p.
- Derocher, A.E., D. Andriashek and J.P.Y. Arnould. 1993. Aspects of milk composition and lactation in polar bears. **Can. J. Zool.** 71(3):561-567.
- Derocher, A.E., G.W. Garner, N.J. Lunn and Ø Wiig. 1998. Polar bears: Proceedings of the Twelfth Working Meeting of the IUCN/SSC Polar Bear Specialist Group, 3-7 February 1997, Oslo, Norway. Occasional Paper of the IUCN Species Survival Commission No. 19. IUCN, Gland.
- Derocher, A.E., Ø. Wiig and G. Bangjord. 2000. Predation of Svalbard reindeer by polar bears. **Polar Biol.** 23(10):675-678.
- DFO Canada. 2004. North Atlantic Right Whale. Fisheries and Oceans Canada. Available at http://www.mar.dfo-mpo.gc.ca/masaro/english/Species_Info/Right_Whale.html
- Divoky, G.J. 1979. Sea ice as a factor in seabird distribution and ecology in the Beaufort, Chukchi and Bering Seas. Pp. 9-18 in Conservation of marine birds of northern North America (J.C. Bartonek and D.N. Nettleship, eds.). U.S. Fish Wildl. Res. Rept. No. 11.
- Divoky, G.J. 1983. The pelagic and nearshore birds of the Alaskan Beaufort Sea. Outer Cont. Shelf Environ. Assess. Progr., Final Rep. Princ. Invest., NOAA, Anchorage, AK 23(1984):397-513. NTIS PB85-212595.
- Durner, G.M. and S.C. Amstrup. 1995. Movements of polar bear from north Alaska to northern Greenland. **Arctic** 48(4):338-341.
- Ely, C.R., C.P. Dau and C.A. Babcock. 1994. Decline in a population of spectacled eiders nesting on the Yukon-Kuskokwim delta, Alaska. **Northw. Natural.** 75:81-87.
- Erbe, C. and D.M. Farmer. 1998. Masked hearing thresholds of a beluga shwle (*Delphinapterus leucas*) in icebreaker noise. **Deep-Sea Research II** 45 (1998) 1373-1388.
- Erbe, C. and D.M. Farmer. 2000. Zones of impact around icebreakers affecting beluga whales in the Beaufort Sea. **J. Acoust. Soc. Am.** 108 (3), Pt. 1.
- Estes, J.A. and J.R. Gilbert. 1978. Evaluation of an aerial survey of Pacific walrus (*Odobenus rosmarus* divergens). **J. Fish. Res. Board Can.** 35:1130-1140.
- Fay, F.H. 1981. Walrus *Odobenus rosmarus* (Linnaeus, 1758). p. 1-23 *In*: S.H. Ridgway and R.J. Harrison (eds.), Handbook of Marine Mammals Vol. 1: The Walrus, Sea Lions, Fur Seals and Sea Otter. Academic Press, London. 235 p.

IV. Literature Cited

- Fay, F.H. 1982. Ecology and biology of the Pacific walrus, *Odobenus rosmarus divergens* Illiger. **N. Am. Fauna** 74:279 p.
- Fay, F.H. and Burns, J.J. 1988. Maximal feeding depth of walruses. **Arctic** 41(3):239-240.
- Fay, F.H., B.P. Kelly and J.L. Sease. 1989. Managing the exploitation of Pacific walrus: a tragedy of delayed response and poor communication. **Mar. Mamm. Sci.** 5(1):1-16.
- Fay, R.R. 1988. Hearing in vertebrates: A psychophysics databook. Hill-Fay Associates, Winnetka, IL. 621 p.
- Fernández, A., M. Arbelo, E. Degollada, M. André, A. Castro-Alonso, R. Jaber, V. Martín, P. Calabuig, P. Castro, P. Herraéz, F. Rodríguez and A. Espinosa de los Monteros. 2003. Pathological findings in beaked whales stranded massively in the Canary Islands (2002). p. 227-228 *In*: 17th Conf. Eur. Cetac. Soc., Las Palmas, March 2003/Conf. Guide & Abstr. European Cetacean Society.
- Fernández, A., J.F. Edwards, F. Rodriguez, A.E. de los Monteros, P. Herraéz, P. Castro, J.R. Jaber, V. Martin and M. Arbelo. 2005a. "Gas and fat embolic syndrome" involving a mass stranding of beaked whales (Family Ziphiidae) exposed to anthropogenic sonar signals. **Vet. Pathol.** 42(4):446-457.
- Fernández, A., M. Méndez, E. Sierra, A. Godinho, P. Herraéz, A.E. De los Monteros, F. Rodrigues and M. Arbelo. 2005b. New gas and fat embolic pathology in beaked whales stranded in the Canary Islands. Abstr. 16th Bien. Conf. Biol. Mar. Mamm., San Diego, CA, 12-16 Dec. 2005.
- Finley, K.J. 1982. The estuarine habitat of the beluga or white whale, *Delphinapterus leucas*. **Cetus** 4:4-5.
- Finley, K.J., G.W. Miller, R.A. Davis and W.R. Koski. 1983. A distinctive large breeding population of ringed seals (*Phoca hispida*) inhabiting the Baffin Bay pack ice. **Arctic** 36(2):162-173.
- Finley, K.J., G.W. Miller, R.A. Davis and C.R. Greene. 1990. Reactions of belugas, *Delphinapterus leucas*, and narwhals, *Monodon monoceros*, to ice-breaking ships in the Canadian high arctic. **Can. Bull. Fish. Aquatic Sci.** 224:97-117.
- Finneran, J.J. and C.E. Schlundt. 2004. Effects of intense pure tones on the behavior of trained odontocetes. TR 1913, SSC San Diego, San Diego, CA.
- Finneran, J.J., C.E. Schlundt, D.A. Carder, J.A. Clark, J.A. Young, J.B. Gaspin and S.H. Ridgway. 2000. Auditory and behavioral responses of bottlenose dolphins (*Tursiops truncatus*) and beluga whale (*Delphinapterus leucas*) to impulsive sounds resembling distant signatures of underwater explosions. **J. Acoust. Soc. Am.** 108(1):417-431.
- Finneran, J.J., C.E. Schlundt, R. Dear, D.A. Carder and S.H. Ridgway. 2002. Temporary shift in masked hearing thresholds in odontocetes after exposure to single underwater impulses from a seismic watergun. **J. Acoust. Soc. Am.** 111(6):2929-2940.
- Finneran, J.J., D.A. Carder, C.E. Schlundt and S.H. Ridgway. 2005. Temporary threshold shift in bottlenose dolphins (*Tursiops truncatus*) exposed to mid-frequency tones. **J. Acoust. Soc. Am.** 118(4):2696-2705.
- Fischer, J.B., and W.W. Larned. 2004. Summer distribution of marine birds in the western Beaufort Sea. **Arctic** 57(2):143-159.
- Flint, P.L. and M.P. Herzog. 1999. Breeding of Steller's eiders, *Polysticta stelleri*, on the Yukon-Kuskokwim delta, Alaska. **Can. Field-Nat.** 113(2):306-308.
- Frame, G.W. 1973. Occurrence of birds in the Beaufort Sea, summer 1969. **Auk** 90(3):552-563.
- Frankel, A.S. 2005. Gray whales hear and respond to a 21–25 kHz high-frequency whale-finding sonar. Abstr. 16th Bien. Conf. Biol. Mar. Mamm., San Diego, CA, 12-16 Dec. 2005.
- Frantzis, A. 1998. Does acoustic testing strand whales? **Nature** 392(6671):29.
- Fredrichson, L.H. 2001. Steller's Eider (*Polysticta stelleri*). *In*: A. Poole and F. Gill (eds.), The Birds of North America, No. 571. The Birds of North America, Inc., Philadelphia, PA.
- Frost, K.J. 1985. The ringed seal. Unpubl. Rep., Alaska Dep. Fish. and Game, Fairbanks, Alaska. 14 p.

- Frost, K.J. and L.F. Lowry. 1981. Foods and trophic relationships of cetaceans in the Bering Sea. p. 825-836 *In*: D.W. Hood and J.A. Calder (eds.) *The Eastern Bering Sea Shelf: Oceanography and Resources*, Vol. 2. Univ. Wash. Press, Seattle.
- Frost, K.J. and L.F. Lowry. 1993. Assessment of injury to harbor seals in Prince William Sound, Alaska, and adjacent areas following the *Exxon Valdez* oil spill. State-Federal Natural Resource Damage Assessment, Marine Mammals Study No. 5. 95 p.
- Frost, K.J., L.F. Lowry and J.J. Burns. 1988. Distribution, abundance, migration, harvest, and stock identity of belukha whales in the Beaufort Sea. p. 27-40 *In*: P.R. Becker (ed.), *Beaufort Sea (Sale 97) information update*. OCS Study MMS 86-0047. Nat. Oceanic & Atmos. Admin., Ocean Assess. Div., Anchorage, AK. 87 p.
- Frost, K.J., L.F. Lowry, G. Pendleton, and H.R. Nute. 2002. Monitoring distribution and abundance of ringed seals in northern Alaska. OCS Study MMS 2002-04.
- Fuller, A.S. and J.C. George. 1999. Evaluation of subsistence harvest data from the North Slope Borough 1993 census for eight North Slope villages for the calendar year 1992. North Slope Borough, Dep. Wildl. Manage., Barrow, AK.
- Funk, D.W., R. Rodrigues, D.S. Ireland, and W.R. Koski (eds.). 2009. Joint Monitoring Program in the Chukchi and Beaufort seas, July-November 2006-2008. LGL Alaska Report P1050-1. Report from LGL Alaska Research Associates, Inc., Anchorage, AK, LGL Ltd., environmental research associates, King City, Ont., Greeneridge Sciences, Inc., Santa Barbara, CA, and JASCO Research, Ltd., Victoria, BC, for Shell Offshore, Inc., Anchorage, AK, other Industry contributors, the National Marine Fisheries Service, Silver Springs, MD, and the U.S. Fish and Wildlife Service, Anchorage, AK. 488 p. plus appendices.
- Galginaitis, M. and D.W. Funk. 2004. Annual assessment of subsistence bowhead whaling near Cross Island, 2001 and 2002: ANIMIDA Task 4 final report. OCS Study MMS 2004-030. Rep. from Applied Sociocultural Res. and LGL Alaska Res. Assoc. Inc., Anchorage, AK, for U.S. Minerals Manage. Serv., Anchorage, AK. 55 p. + CD-ROM.
- Galginaitis, M. and D.W. Funk. 2005. Annual assessment of subsistence bowhead whaling near Cross Island, 2003: ANIMIDA Task 4 annual report. OCS Study MMS 2005-025. Rep. from Applied Sociocultural Research and LGL Alaska Res. Assoc. Inc., Anchorage, AK, for U.S. Minerals Manage. Serv., Anchorage, AK. 36 p. + Appendices.
- Galginaitis, M.S. and W.R. Koski. 2002. Kaktovikmiut whaling: historical harvest and local knowledge of whale feeding behavior. p. 2-1 to 2-30 (Chap. 2) *In*: W.J. Richardson and D.H. Thomson (eds.), *Bowhead whale feeding in the eastern Alaskan Beaufort Sea: update of scientific and traditional information*, vol. 1. OCS Study MMS 2002-012; LGL Rep. TA2196-7. Rep. from LGL Ltd., King City, Ont., for U.S. Minerals Manage. Serv., Anchorage, AK, and Herndon, VA. 420 p.
- Galloway, B.J., W.B. Griffiths, P.C. Craig, W.J. Gazey and J.W. Helmericks. 1983. An assessment of the Colville River Delta stock of Arctic Cisco - migrants from Canada? **Biol. Pap. Univ. Alaska** 21:4-23.
- Galloway, B.J., W.J. Gazey and L.L. Moulton. 1989. Population Trends for the Arctic Cisco (*Coregonus autumnalis*) in the Colville River of Alaska as reflected by the commercial fishery. **Biol. Pap. Univ. Alaska** 24:153-165.
- Gambell, R. 1985. Fin whale *Balaenoptera physalus* (Linnaeus, 1758). p. 171-192 *In*: S.H. Ridgway and R. Harrison (eds.), *Handbook of Marine Mammals*, Vol. 3: The Sirenians and Baleen Whales. Academic Press, London, U.K. 362 p.
- Garner, G.W., S.T. Knick and D.C. Douglas. 1990. Seasonal movements of adult female polar bears in the Bering and Chukchi Seas. **Int. Conf. Bear Res. Manage.** 8:219-226.
- Gentry, R. (ed.). 2002. Report of the workshop on acoustic resonance as a source of tissue trauma in cetaceans, Silver Spring, MD, April 2002. Nat. Mar. Fish. Serv. 19 p. Available at www.nmfs.noaa.gov/prot_res/PR2/Acoustics_Program/acoustics.html

IV. Literature Cited

- George, J.C., L.M. Philo, K. Hazard, D. Withrow, G.M. Carroll, and R. Suydam. 1994. Frequency of killer whale (*Orcinus orca*) attacks and ship collisions based on scarring on bowhead whales (*Balaena mysticetus*) of the Bering-Chukchi-Beaufort Seas stock. **Arctic** 47(3):247-255.
- George, J.C., J. Zeh, R. Suydam and C. Clark. 2004. Abundance and population trend (1978-2001) of Western Arctic bowhead whales surveyed near Barrow, Alaska. **Mar. Mamm. Sci.** 20(4):755-773.
- Gilbert, J.R. 1989. Aerial census of Pacific walruses in the Chukchi Sea, 1985. **Mar. Mamm. Sci.** 5(1):17-28.
- Goold, J.C. 1996a. Acoustic assessment of common dolphins off the west Wales coast, in conjunction with 16th round seismic surveying. Rep. from School of Ocean Sciences, Univ. Wales, Bangor, Wales, for Chevron UK Ltd, Repsol Explor. (UK) Ltd., and Aran Energy Explor. Ltd. 22 p.
- Goold, J.C. 1996b. Acoustic assessment of populations of common dolphin *Delphinus delphis* in conjunction with seismic surveying. **J. Mar. Biol. Assoc. U.K.** 76:811-820.
- Goold, J.C. 1996c. Acoustic cetacean monitoring off the west Wales coast. Rep. from School of Ocean Sciences, Univ. Wales, Bangor, Wales, for Chevron UK Ltd, Repsol Explor. (UK) Ltd, and Aran Energy Explor. Ltd. 20 p.
- Green, G.A., K. Hashagen, and D. Lee. 2007. Marine mammal monitoring program, FEX barging project, 2007. Report prepared by Tetra Tech EC, Inc., Bothell WA, for FEX L.P., Anchorage, AK.
- Greene, C.R., Jr. 1997. Physical acoustics measurements. (Chap. 3, 63 p.) *In*: W.J. Richardson (ed.), 1997. Northstar Marine Mammal Marine Monitoring Program, 1996. Marine mammal and acoustical monitoring of a seismic program in the Alaskan Beaufort Sea. Rep. TA2121-2. Rep. from LGL Ltd., King City, Ont., and Greeneridge Sciences Inc., Santa Barbara, CA, for BP Explor. (Alaska) Inc., Anchorage, AK, and U.S. Nat. Mar. Fish. Serv., Anchorage, AK, and Silver Spring, MD. 245 p.
- Greene, C.R., Jr. and W.R. Richardson. 1988. Characteristics of marine seismic survey sounds in the Beaufort Sea. **J. Acoust. Soc. Am.** 83(6):2246-2254.
- Greene, C.R., Jr., N.S. Altman and W.J. Richardson. 1999. Bowhead whale calls. p. 6-1 to 6-23 *In*: W.J. Richardson (ed.), Marine mammal and acoustical monitoring of Western Geophysical's open-water seismic program in the Alaskan Beaufort Sea, 1998. LGL Rep. TA2230-3. Rep. from LGL Ltd., King City, ON, and Greeneridge Sciences Inc., Santa Barbara, CA, for Western Geophysical, Houston, TX, and U.S. Nat. Mar. Fish. Serv., Anchorage, AK, and Silver Spring, MD. 390 p.
- Greene, C.R., Jr., R.G. Norman, S.B. Blackwell, and A. Thode. 2007. Acoustics research for studying bowhead migration, 2006. Chapter 10 *In* D.S. Ireland, D.W. Funk, R. Rodrigues, and W.R. Koski (eds.). Joint monitoring program in the Chukchi and Beaufort seas, July-November 2006. LGL Rep. P891-2. Prepared by LGL Alaska Research Associates, Inc., Anchorage, AK, and LGL Ltd., environmental research associates, King City, Ont., for Shell Offshore Inc., ConocoPhillips Alaska, Inc., GX Technology, the National Marine Fisheries Service, and the U.S. Fish and Wildlife Service.
- Greening, M.M., and P. Zakarauskas. 1993. Spatial and source level distributions of ice cracking in the Arctic Ocean. **J. Acoust. Soc. Am.** 95(2):783-790.
- Gundalf. 2009. Gundalf revision AIR6.1a, Date 2009-04-27, Epoch 2009-03-31, by Oakwood Computing Associates Limited Oakwood, 11 Carlton Road, New Malden, Surrey KT3 3AJ, Surrey KT3 3AJ, United Kingdom. Gundalf is a registered trademark of Oakwood Computing Associates Limited Oakwood.
- Haley, B. 2006. Marine mammal monitoring during University of Texas at Austin's marine geophysical survey of the western Canada Basin, Chukchi Borderland and Mendeleev Ridge, Arctic Ocean, July–August 2006. Report from LGL Alaska Research Associates, Inc., Anchorage AK, and LGL Ltd., King City, Ont., for the University of Texas at Austin, the Nat. Mar. Fish. Serv., Silver Springs, MD, and the U.S. Fish and Wildl. Serv., Anchorage, AK.
- Haley, B. and D. Ireland. 2006. Marine mammal monitoring during University of Alaska Fairbanks' marine geophysical survey across the Arctic Ocean, AuguSt. September 2005. LGL Rep. TA4122-3. Rep. from LGL Ltd., King City, Ont., for Univ. Alaska Fairbanks, Fairbanks, AK, and Nat. Mar. Fish. Serv., Silver Spring, MD. 80 p.

- Haley, B., J. Beland, D.S. Ireland, R. Rodrigues, and D.M. Savarese. 2009. Chapter 3 *In* Funk, D.W., D.S. Ireland, R. Rodrigues, and W.R. Koski (eds.). Joint monitoring program in the Chukchi and Beaufort seas, July–November 2006–2008. LGL Alaska Report P1050-1. Report from LGL Alaska Research Associates, Inc., Anchorage, Ak, LGL Ltd., environmental research associates, King City, Ont., Greeneridge Sciences, Inc., Goleta, CA, and JASCO Research, Victoria, B.C., for Shell Offshore, Inc. and other Industry contributors, National Marine Fisheries Service, and U.S. Fish and Wildlife Service. 488 p. plus appendices.
- Hammill, M.O., C. Lydersen, M. Ryg and T.G. Smith. 1991. Lactation in the ringed seal (*Phoca hispida*). **Can. J. Fish. Aquatic Sci.** 48(12):2471-2476.
- Harington, C.R. 1968. Denning habits of the polar bear (*Ursus maritimus*). **Can. Wildl. Serv. Rep. Ser.** 5:1-33.
- Harris, R.E., G.W. Miller and W.J. Richardson. 2001. Seal responses to airgun sounds during summer seismic surveys in the Alaskan Beaufort Sea. **Mar. Mamm. Sci.** 17(4):795-812.
- Harwood, J. and B. Wilson. 2001. The implications of developments on the Atlantic Frontier for marine mammals. **Cont. Shelf Res.** 21(8-10):1073-1093.
- Harwood, L., S. Innes, P. Norton and M. Kingsley. 1996. Distribution and abundance of beluga whales in the Mackenzie estuary, southeast Beaufort Sea, and the west Amundsen Gulf during late July 1992. **Can. J. Fish. Aquatic Sci.** 53(10):2262-2273.
- Harwood, L.A., F. McLaughlin, R.M. Allen, J. Illasiak Jr. and J. Alikamik. 2005. First ever marine mammal and bird observations in the deep Canada Basin and Beaufort/Chukchi seas: expeditions during 2002. **Polar Biol.** 28(3):250-253.
- Hay, K.A. 1985. The life history of the narwhal (*Monodon monoceros* L.) in the eastern Canadian Arctic. **Diss. Abst. Int. Pt. B Sci. Eng.** 45(10).
- Hay, K.A. and A.W. Mansfield. 1989. Narwhal - *Monodon monoceros* Linnaeus, 1758. p. 145-176 *In*: S.H. Ridgway and R. Harrison (eds.), Handbook of Marine Mammals, Vol. 4: River Dolphins and the Larger Toothed Whales. Academic Press, London, UK.
- Hazard, K. 1988. Beluga whale, *Delphinapterus leucas*. p. 195-235 *In*: J.W. Lentfer (ed.), Selected Marine Mammals of Alaska. Mar. Mamm. Comm., Washington, DC. 275 p. NTIS PB88-178462. 275 p.
- Hill, P.S. and D.P. DeMaster. 1998. Draft Alaska marine mammal stock assessments 1998. U.S. Nat. Mar. Fish. Serv., Nat. Mar. Mamm. Lab., Seattle, WA.
- Hodges, J.I. and W.D. Eldridge. 2001. Aerial surveys of eiders and other waterbirds on the eastern Arctic coast of Russia. **Wildfowl** 52:127-142.
- Hogarth, W.T. 2002. Declaration of William T. Hogarth in opposition to plaintiff's motion for temporary restraining order, 23 October 2002. Civ. No. 02-05065-JL. U.S. District Court, Northern District of California, San Francisco Div.
- Hubbs, C.L. and A.B. Rechnitzer. 1952. Report on experiments designed to determine effects of underwater explosions on fish life. **Calif. Fish & Game** 38(3):333-366.
- Huntington, H.P. 2000. Traditional knowledge of the ecology of belugas, *Delphinapterus leucas*, in Cook Inlet, Alaska. **Mar. Fish. Rev.** 62(3):134-140.
- Innes, S., M.P. Heide-Jørgensen, J. Laake, K. Laidre, H. Cleator and P. Richard. 2002. Surveys of belugas and narwhals in the Canadian high Arctic in 1996. **NAMMCO Sci. Publ.** 4:169-190.
- Ireland, D.S., R. Rodrigues, D. Funk, W. Koski, and D. Hannay (eds.). 2009. Marine mammal monitoring and mitigation during open-water seismic exploration by Shell Offshore Inc. in the Chukchi and Beaufort seas, July–October 2008: 90-day report. LGL Rep. P1049-1. Rep. from LGL Alaska Research Associates, Inc., Anchorage, AK, and JASCO Research Ltd., Victoria, BC, for Shell Offshore Inc., Anchorage, AK, the Nat. Mar. Fish. Serv., Silver Springs MD, and the U.S. Fish and Wildl. Serv., Anchorage, AK. 277 p + appendices.
- IUCN (The World Conservation Union). 2003. 2003 IUCN Red List of Threatened Species. <http://www.redlist.org>

- Jankowski, M., M. Fitzgerald, B. Haley, and H. Patterson. 2008. Beaufort sea vessel-based monitoring program. Chapter 6 *In* Funk, D.W., R. Rodrigues, D.S. Ireland, and W.R. Koski (eds.). Joint monitoring program in the Chukchi and Beaufort seas, July–November 2007. LGL Alaska Report P971-2. Report from LGL Alaska Research Associates, Inc., Anchorage, Ak, LGL Ltd., environmental research associates, King City, Ont., JASCO Research, Victoria, B.C., and Greeneridge Sciences, Inc., Goleta, CA, for Shell Offshore, Inc., ConocoPhillips Alaska, Inc., and National Marine Fisheries Service, and U.S. Fish and Wildlife Service.
- Jefferson, T.A., S. Leatherwood, and M.A. Webber. 1993. FAO Species Identification Guide. Marine Mammals of the World. UNEP/FAO, Rome.
- Jepson, P.D., M. Arbelo, R. Deaville, I.A.P. Patterson, P. Castro, J.R. Baker, E. Degollada, H.M. Ross, P. Herráez, A.M. Pocknell, F. Rodríguez, F.E. Howie, A. Espinosa, R.J. Reid, J.R. Jaber, V. Martin, A.A. Cunningham and A. Fernández. 2003. Gas-bubble lesions in stranded cetaceans. **Nature** 425(6958):575-576.
- Jepson, P.D., D.S. Houser, L.A. Crum, P.L. Tyack and A. Fernández. 2005a. Beaked whales, sonar and the “bubble hypothesis”. Abstr. 16th Bien. Conf. Biol. Mar. Mamm., San Diego, CA, 12-16 Dec. 2005.
- Jepson, P.D. R. Deaville, I.A.P. Patterson, A.M. Pocknell, H.M. Ross, J.R. Baker, F.E. Howie, R.J. Reid, A. Colloff and A.A. Cunningham. 2005b. Acute and chronic gas bubble lesions in cetaceans stranded in the United Kingdom. **Vet. Pathol.** 42(3):291-305.
- Johnson, C.B., B.E. Lawhead, J.R. Rose, M.D. Smith, A.A. Stickney and A.M. Wildman. 1999. Wildlife studies on the Colville River Delta, Alaska, 1998. Rep. from ABR, Inc., Fairbanks, AK, for ARCO Alaska, Inc., Anchorage, AK.
- Johnson, A., J. Burns, W. Dusenberry and R. Jones. 1982. Aerial survey of Pacific walrus, 1980. U.S. Fish and Wildlife Service, Anchorage, AK. Mimeo report. 32 p.
- Johnson, S.R. 1979. Fall observations of westward migrating white whales (*Delphinapterus leucas*) along the central Alaskan Beaufort Sea coast. **Arctic** 32(3):275-276.
- Johnson, S.R. 2002. Marine mammal mitigation and monitoring program for the 2001 Odoptu 3-D seismic survey, Sakhalin Island Russia: Executive summary. Rep. from LGL Ltd, Sidney, B.C., for Exxon Neftegas Ltd., Yuzhno-Sakhalinsk, Russia. 49 p. Also available as Working Paper SC/02/WGW/19, Int. Whal. Comm., Western Gray Whale Working Group Meeting, Ulsan, South Korea, 22-25 October 2002. 48 p.
- Jones, M.L. and S.L. Swartz. 1984. Demography and phenology of gray whales and evaluation of whale-watching activities in Laguna San Ignacio, Baja California Sur, Mexico. p. 309-374 *In*: M. L. Jones et al. (eds.), The Gray Whale *Eschrichtius robustus*. Academic Press, Orlando, FL. 600 p.
- Jonsgård, Å. 1966a. The distribution of Balaenopteridae in the North Atlantic Ocean. p. 114-124 *In*: K.S. Norris (ed.), Whales, Dolphins, and Porpoises. Univ. Calif. Press, Berkeley and Los Angeles, CA.
- Jonsgård, Å. 1966b. Biology of the North Atlantic fin whale *Balaenoptera physalus* (L.): taxonomy, distribution, migration and food. **Hvalrådets Skr.** 49:1-62.
- Kanik, B., M. Winsby and R. Tanasichuk. 1980. Observations of marine mammal and sea bird interaction with icebreaking activities in the High Arctic July 2-12, 1980. Rep. from Hatfield Consultants Ltd., West Vancouver, B.C., for Petro-Canada, Calgary, Alb. 53 p.
- Kastak, D., R.L. Schusterman, B.L. Southall and C.J. Reichmuth. 1999. Underwater temporary threshold shift induced by octave-band noise in three species of pinnipeds. **J. Acoust. Soc. Am.** 106(2):1142-1148.
- Kaleak, J. 1996. History of whaling by Kaktovik village. p. 69-71 *In*: Proc. 1995 Arctic Synthesis Meeting, Anchorage, AK, Oct. 1995. OCS Study MMS 95-0065. U.S. Minerals Manage. Serv., Anchorage, AK. 206 p. + Appendices.
- Kelly, B.P. 1988. Bearded seal, *Erignathus barbatus*. p. 77-94 *In*: J.W. Lentfer (ed.), Selected Marine Mammals of Alaska/Species Accounts with Research and Management Recommendations. Mar. Mamm. Comm., Washington, DC. 275 p.
- Kertell, K. 1991. Disappearance of the Steller’s eider from the Yukon-Kuskokwim delta, Alaska. **Arctic** 44(3):177-187.

- Ketten, D.R. 1995. Estimates of blast injury and acoustic trauma zones for marine mammals from underwater explosions. p. 391-407 *In*: R.A. Kastelein, J.A. Thomas and P.E. Nachtigall (eds.), *Sensory Systems of Aquatic Mammals*. De Spil Publ., Woerden, Netherlands. 588 p.
- Ketten, D.R., J. Lien and S. Todd. 1993. Blast injury in humpback whale ears: evidence and implications. **J. Acoust. Soc. Am.** 94(3, Pt. 2):1849-1850.
- Ketten, D.R., J. O'Malley, P.W.B. Moore, S. Ridgway and C. Merigo. 2001. Aging, injury, disease, and noise in marine mammal ears. **J. Acoust. Soc. Am.** 110(5, Pt. 2):2721.
- King, J.E. 1983. *Seals of the World*, 2nd ed. Cornell Univ. Press, Ithaca, NY. 240 p.
- Kingsley, M.C.S. 1986. Distribution and abundance of seals in the Beaufort Sea, Amundsen Gulf, and Prince Albert Sound, 1984. *Environ. Studies Revolving Funds Rep. No. 25*. 16 p.
- Koski, W.R. and R.A. Davis. 1994. Distribution and numbers of narwhals (*Monodon monoceros*) in Baffin Bay and Davis Strait. **Medd. Grønl., Biosci.** 39:15-40
- Koski, W.R., D.H. Thomson and W.J. Richardson. 1998. Descriptions of marine mammal populations. p. 1-182 + app. *In*: Point Mugu Sea Range Marine Mammal Technical Report. Rep. from LGL Ltd., King City, ON, for Naval Air Warfare Center, Weapons Div., Point Mugu, CA, and Southwest Div. Naval Facilities Eng. Command, San Diego, CA. 322 p.
- Koski, W.R., J.C. George, G. Sheffield and M.S. Galginitis. 2005. Subsistence harvests of bowhead whales (*Balaena mysticetus*) at Kaktovik, Alaska (1973-2000). **J. Cetac. Res. Manage.** 7(1):33-37.
- Kremser, U., P. Klemm and W.D. Kötz. 2005. Estimating the risk of temporary acoustic threshold shift, caused by hydroacoustic devices, in whales in the Southern Ocean. **Antarct. Sci.** 17(1):3-10.
- Kryter, K.D. 1985. *The Effects of Noise on Man*, 2nd ed. Academic Press, Orlando, FL. 688 p.
- Lacroix, D.L., R.B. Lancot, J.A. Reed and T. McDonald. 2003. Effect of underwater seismic surveys on molting male Long-tailed Ducks in the Beaufort Sea, Alaska. **Can. J. Zool.** 81(11):1862-1875.
- Larned, W., R. Stehn and R. Platte. 2009. Eider breeding population survey, Arctic Coastal Plain, Alaska 2008. USFWS, Migratory Bird Management, Waterfowl Management, Soldatna and Anchorage, AK.
- Lawson, J.W. 2002. Seismic program described, 2001. p. 2-1 to 2-22 *In*: W.J. Richardson and J.W. Lawson (eds.), *Marine mammal monitoring of WesternGeco's open-water seismic program in the Alaskan Beaufort Sea, 2001*. LGL Rep. 2564-4. Rep. from LGL Ltd., King City, ON, for WesternGeco LLC, Anchorage, AK, BP Explor. (Alaska) Inc., Anchorage, AK, and Nat. Mar. Fish. Serv., Anchorage, AK, and Silver Spring, MD. 95 p.
- Larsen, T. 1985. Polar bear denning and cub production in Svalbard, Norway. **J. Wildl. Manage.** 49(2):320-326.
- Leatherwood, S., R.R. Reeves, W.F. Perrin, and W.E. Evans. 1982. Whales, dolphins, and porpoises of the eastern North Pacific and adjacent Arctic waters: A guide to their identification. U.S. Dept. of Commerce, NOAA Tech. Rep., NMFS Circular 444.
- Leatherwood, S., A.E. Bowles, and R. Reeves. 1986. Aerial surveys of marine mammals in the southeastern Bering Sea. U.S. Department of Commerce, NOAA, OCSEAP Final Report 42:147-490.
- Lentfer, W.J., R.J. Hensel, J.R. Gilbert and F.E. Sorensen. 1980. Population characteristics of Alaskan polar bears. **Int. Conf. Bear Res. Manage.** 3:109-115.
- Lesage, V. C. Barrette, M.C.S. Kingsley, and B. Sjare. 1999. The effect of vessel noise on the vocal behavior of belugas in the St. Lawrence River Estuary, Canada. *Mar. Mam Sci.* 15(1) 65-84.
- LGL and Greeneridge. 1987. Responses of bowhead whales to an offshore dirlling operation in the Alaskan Beaufort Sea, autumn 1986. Rep. from LGL Ltd., King City, Ont., and Greeneridge Sciences Inc., Santa Barbara, CA for Shell Western E & P Inc., Anchorae, AK. 371 p.
- LGL and Greeneridge. 1996. Northstar Marine Mammal Monitoring Program, 1995: Baseline surveys and retrospective analyses of marine mammal and ambient noise data from the Central Alaskan Beaufort Sea. Rep.

IV. Literature Cited

- from LGL Ltd., King City, Ont., and Greeneridge Sciences Inc., Santa Barbara, CA, for BP Explor. (Alaska) Inc., Anchorage, AK. 104 p.
- Ljungblad, D.K., S.E. Moore and D.R. Van Schoik. 1984. Aerial surveys of endangered whales in the Beaufort, eastern Chukchi, and northern Bering Seas, 1983: with a five year review, 1979-1983. NOSC Tech Rep. 955. Rep. from Naval Ocean Systems Center, San Diego, CA for U.S. Minerals Manage. Serv., Anchorage, AK. 356 p. NTIS AD-A146 373/6.
- Ljungblad, D.K., S.E. Moore and D.R. Van Schoik. 1986. Seasonal patterns of distribution, abundance, migration and behavior of the Western Arctic stock of bowhead whales, *Balaena mysticetus* in Alaskan seas. **Rep. Int. Whal. Comm., Spec. Iss.** 8:177:205.
- Ljungblad, D.K., S.E. Moore, J.T. Clarke and J.C. Bennett. 1987. Distribution, abundance, behavior and bioacoustics of endangered whales in the Alaskan Beaufort and eastern Chukchi Seas, 1979-86. NOSC Tech. Rep. 1177; OCS Study MMS 87-0039. Rep. from Naval Ocean Systems Center, San Diego, CA, for U.S. Minerals Manage. Serv., Anchorage, AK. 391 p. NTIS PB88-116470.
- Ljungblad, D.K., B. Würsig, S.L. Swartz and J.M. Keene. 1988. Observations on the behavioral responses of bowhead whales (*Balaena mysticetus*) to active geophysical vessels in the Alaskan Beaufort Sea. **Arctic** 41(3):183-194.
- Long, F., Jr. 1996. History of subsistence whaling by Nuiqsut. p. 73-76 *In: Proc. 1995 Arctic Synthesis Meeting*, Anchorage, AK, Oct. 1995. OCS Study MMS 95-0065. U.S. Minerals Manage. Serv., Anchorage, AK. 206 p. + Appendices.
- Lowry, L.F., R.R. Nelson, and K.J. Frost. 1987. Observations of killer whales, (*Orcinus orca*) in western Alaska: Sightings, strandings and predation on other marine mammals. **Canadian Field-Naturalist** 101:6-12.
- Lowry, L.F., K.J. Frost, R. Davis, D.P. DeMaster and R.S. Suydam. 1998. Movements and behavior of satellite-tagged spotted seals (*Phoca largha*) in the Bering and Chukchi Seas. **Polar Biol.** 19(4):221-230.
- Lyderson, C. and M.O. Hammill. 1993. Diving in ringed seal (*Phoca hispida*) pups during the nursing period. **Can. J. Zool.** 71(5):991-996.
- Lyons, C., W. Koski, and D. Ireland. 2009. Chapter 7 *In* Ireland, D.S., D.W. Funk, R. Rodrigues, and W.R. Koski (eds.). 2009. Joint monitoring program in the Chukchi and Beaufort seas, July–November 2007. LGL Alaska Report P971-2. Report from LGL Alaska Research Associates, Inc., Anchorage, Ak, LGL Ltd., environmental research associates, King City, Ont., JASCO Research, Victoria, B.C., and Greeneridge Sciences, Inc., Goleta, CA, for Shell Offshore, Inc., ConocoPhillips Alaska, Inc., the National Marine Fisheries Service, silver Springs, MD, and U.S. Fish and Wildlife Service, Anchorage, AK. 445 p. plus appendices.
- Maher, W.J. 1960. Recent records of the California gray whale (*Eschrichtius glaucus*) along the north coast of Alaska. **Arctic** 13(4):257-265.
- Madsen, P.T., B. Møhl, B.K. Nielsen and M. Wahlberg. 2002. Male sperm whale behavior during exposures to distant seismic survey pulses. **Aquat. Mamm.** 28(3):231-240.
- Malme, C.I., P.R. Miles, C.W. Clark, P. Tyack and J.E. Bird. 1984. Investigations of the potential effects of underwater noise from petroleum industry activities on migrating gray whale behavior/Phase II: January 1984 migration. BBN Rep. 5586. Rep. from Bolt Beranek & Newman Inc., Cambridge, MA, for U.S. Minerals Manage. Serv., Anchorage, AK. NTIS PB86-218377.
- Malme, C.I., B. Würsig, J.E. Bird and P. Tyack. 1986. Behavioral responses of gray whales to industrial noise: feeding observations and predictive modeling. **Outer Cont. Shelf Environ. Assess. Progr., Final Rep. Princ. Invest., NOAA**, Anchorage, AK 56(1988):393-600. BBN Rep. 6265. 600 p. OCS Study MMS 88-0048; NTIS PB88-249008.
- Malme, C.I., B. Würsig, J.E. Bird and P. Tyack. 1988. Observations of feeding gray whale responses to controlled industrial noise exposure. p. 55-73 *In: W.M. Sackinger, M.O. Jeffries, J.L. Imm and S.D. Treacy (eds.), Port*

- and Ocean Engineering under Arctic Conditions, Vol. II. Geophysical Inst., Univ. Alaska, Fairbanks, AK. 111 p.
- Martel, V.M. n.d. Summary of the report on the atypical mass stranding of beaked whales in the Canary Islands in September 2002 during naval exercises. Society for the Study of Cetaceans in the Canary Islands.
- Mate, B.R., G.K. Krutzikowski, and M.H. Winsor. 2000. Satellite-monitored movements of radio-tagged bowhead whales in the Beaufort and Chukchi seas during the late-summer feeding season and fall migration. **Can. J. Zool.** 78:1168-1181.
- McCauley, R.D., M.N. Jenner, C. Jenner, K.A. McCabe and J. Murdoch. 1998. The response of humpback whales (*Megaptera novangliae*) to offshore seismic survey noise: preliminary results of observations about a working seismic vessel and experimental exposures. **APPEA (Austral. Petrol. Product. Explor. Assoc.) Journal** 38:692-707.
- McCauley, R.D., J. Fewtrell, A.J. Duncan, C. Jenner, M.N. Jenner, J.D. Penrose, R.I.T. Prince, A. Adhitya, J. Murdoch and K. McCabe. 2000. Marine seismic surveys: Analysis of airgun signals and effects of air gun exposure on humpback whales, sea turtles, fishes and squid. Rep. from Centre for Marine Science and Technology, Curtin Univ., Perth, W.A., for Austral. Petrol. Prod. Assoc., Sydney, N.S.W. 188 p.
- McDonald, M.A., J.A. Hildebrand and S.C. Webb. 1995. Blue and fin whales observed on a seafloor array in the Northeast Pacific. **J. Acoust. Soc. Am.** 98(2, Pt.1):712-721.
- McLaughlin, F.A., E.C. Carmack, R.W. Macdonald and J.K.B. Bishop. 1996. Physical and chemical properties across the Atlantic/Pacific water mass front in the southern Canadian Basin. **J. Geophys. Res.** 101, No. C1: 1183-1198. [Abstract]
- Méndez, M., M. Arbelo, E. Sierra, A. Godinho, M.J. Caballero, J. Jaber, P. Herráez and A. Fernández. 2005. Lung fat embolism in cetaceans stranded in Canary Islands. Abstr. 16th Bien. Conf. Biol. Mar. Mamm., San Diego, CA, 12-16 Dec. 2005.
- Miller, G.W., R.E. Elliott, W.R. Koski, V.D. Moulton and W.J. Richardson. 1999. Whales. p. 5-1 to 5-109 *In*: W.J. Richardson (ed.), Marine mammal and acoustical monitoring of Western Geophysical's open-water seismic program in the Alaskan Beaufort Sea, 1998. LGL Rep. TA2230-3. Rep. from LGL Ltd., King City, Ont., and Greeneridge Sciences Inc., Santa Barbara, CA, for Western Geophysical, Houston, TX and U.S. Nat. Mar. Fish. Serv., Anchorage, AK, and Silver Spring, MD. 390 p.
- Miller, G.W., R.E. Elliot, T.A. Thomas, V.D. Moulton, and W.R. Koski. 2002. Distribution and numbers of bowhead whales in the eastern Alaskan Beaufort Sea during late summer and autumn, 1979-2000. Chapter 9 *In* Richardson, W.J. and D.H. Thomson (eds). 2002. Bowhead whale feeding in the eastern Alaskan Beaufort Sea: update of scientific and traditional information. OCS Study MMS 2002-012; LGL Rep. TA2196-7. Rep. from LGL Ltd., King City, Ont., for U.S. Minerals Manage. Serv., Anchorage, AK, and Herndon, VA. xlv + 697 p. 2 vol. NTIS PB2004-101568. Available from www.mms.gov/alaska/ref/AKPUBS.HTM#2002.
- Miller, G.W., V.D. Moulton, R.A. Davis, M. Holst, P. Millman, A. MacGillivray and D. Hannay. 2005. Monitoring seismic effects on marine mammals—southeastern Beaufort Sea, 2001-2002. p. 511-542 *In*: S.L. Armsworthy, P.J. Cranford, and K. Lee (eds.), Offshore Oil and Gas Environmental Effects Monitoring/Approaches and Technologies. Battelle Press, Columbus, OH.
- Mitchell, E.D. 1975. Report on the meeting on small cetaceans, Montreal, April 1-11, 1974. **J. Fish. Res. Board Can.** 32:914-91.
- MMS. 1996. Beaufort Sea Planning Area oil and gas lease sale 144/Final Environmental Impact Statement. OCS EIS/EA MMS 96-0012. U.S. Minerals Manage. Serv., Alaska OCS Reg., Anchorage, AK. Two Vol. Var. pag.
- MMS. 2007. Seismic surveys in the Beaufort and Chukchi seas, Alaska. OCS EIS/EA MMS 2007-001.
- Monnett, C. and S.D. Treacy. 2005. Aerial surveys of endangered whales in the Beaufort Sea, fall 2002-2004. OCS Study MMS 2005-037. Minerals Manage. Serv., Anchorage, AK. xii + 153 p.

- Moore, S.E. 2000. Variability in cetacean distribution and habitat selection in the Alaskan Arctic, autumn 1982-91. **Arctic** 53(4):448-460
- Moore, S.E. and R.R. Reeves. 1993. Distribution and movement. p. 313-386 *In*: J.J. Burns, J.J. Montague and C.J. Cowles (eds.), *The Bowhead Whale*. Spec. Publ. 2. Soc. Mar. Mammal., Lawrence, KS. 787 p.
- Moore, S.E., and DeMaster, D.P. 1997. Cetacean habitats in the Alaskan Arctic. **J. Northw. Atl. Fish. Sci.** 22:55-69
- Moore, S.E., J.T. Clarke and D.K. Ljungblad. 1989. Bowhead whale (*Balaena mysticetus*) spatial and temporal distribution in the central Beaufort Sea during late summer and early fall 1979-86. **Rep. Int. Whal. Comm.** 39:283-290.
- Moore, S.E., J.C. George, K.O. Coyle, and T.J. Weingartner. 1995. Bowhead whales along the Chukotka coast in autumn. **Arctic** 48(2):155-160.
- Moore, S.E., J.M. Waite, L.L. Mazzuca and R.C. Hobbs. 2000a. Mysticete whale abundance and observations of prey associations on the central Bering Sea shelf. **J. Cetac. Res. Manage.** 2(3): 227-234.
- Moore, S.E., D.P. DeMaster and P.K. Dayton. 2000b. Cetacean habitat selection in the Alaskan Arctic during summer and autumn. **Arctic** 53(4):432-447.
- Moore, S.E., J.M. Waite, N.A. Friday, and T. Honkalehto. 2002. Distribution and comparative estimates of cetacean abundance on the central and southeastern Bering Sea shelf with observations on bathymetric and prey associations. **Progr. Oceanogr.** 55:249-262.
- Moore, S.E., J.M. Grebmeier and J.R. Davies. 2003. Gray whale distribution relative to forage habitat in the northern Bering Sea: current conditions and retrospective summary. **Can. J. Zool.** 81(4):734-742.
- Mosher, D.C., J.W. Shimeld, and D.R. Hutchinson. 2009. 2009 Canada Basin seismic reflection and refraction survey, western Arctic Ocean: CCGS Louis S. St. Laurent expedition report. Geological Survey of Canada, Ottawa, Ontario.
- Moulton, V.D. and J.W. Lawson. 2002. Seals, 2001. p. 3-1 to 3-46 *In*: W.J. Richardson and J.W. Lawson (eds.), *Marine mammal monitoring of WesternGeco's open-water seismic program in the Alaskan Beaufort Sea, 2001*. LGL Rep. TA2564-4. Rep. from LGL Ltd., King City, Ont., for WesternGeco LLC, Anchorage, AK; BP Explor. (Alaska) Inc., Anchorage, AK; and Nat. Mar. Fish. Serv., Anchorage, AK, and Silver Spring, MD. 95 p.
- Moulton, V.D. and M.T. Williams. 2003. Incidental sightings of polar bears during monitoring activities for BP's Northstar oil development, Alaskan Beaufort Sea, 2002. Rep. from LGL Ltd. for BP Explor. (Alaska) Inc. and USFWS Office of Mar. Mamm. Manage, Anchorage, AK.
- Moulton, V.D. and G.W. Miller. 2005. Marine mammal monitoring of a seismic survey on the Scotian Slope, 2003. p. 29-40 in K. Lee, H. Bain and G.V. Hurley, eds. 2005. *Acoustic Monitoring and Marine Mammal Surveys in the Gully and Outer Scotian Shelf before and during Active Seismic Programs*. **Environmental Studies Research Funds Report** No. 151. 154 p.
- Moulton, V.D., W.J. Richardson, T.L. McDonald, R.E. Elliot, and M.T. Williams. 2002. Factors influencing local abundance and haulout behaviour of ringed seals (*Phoca hispida*) on landfast ice of the Alaskan Beaufort Sea. **Can. J. Zool.** 80:1900-1917.
- Nerini, M. 1984. A review of gray whale feeding ecology. p. 423-450 *In* M.L. Jones, S.L. Swartz and S. Leatherwood (eds.), *The Gray Whale, Eschrichtius robustus*. Academic Press, Inc. Orlando, FL. 600 p.
- Nieukirk, S.L., K.M. Stafford, D.K. Mellinger, R.P. Dziak and C.G. Fox. 2004. Low-frequency whale and seismic airgun sounds recorded in the mid-Atlantic Ocean. **J. Acoust. Soc. Am.** 115(4):1832-1843.
- NMFS. 1995. Small takes of marine mammals incidental to specified activities; offshore seismic-activities in southern California. **Fed. Regist.** 60(200, 17 Oct.):53753-53760.
- NMFS. 2000. Small takes of marine mammals incidental to specified activities; marine seismic-reflection data collection in southern California/Notice of receipt of application. **Fed. Regist.** 65(60, 28 Mar.):16374-16379.

- NMFS. 2001. Small takes of marine mammals incidental to specified activities; oil and gas exploration drilling activities in the Beaufort Sea/Notice of issuance of an incidental harassment authorization. **Fed. Regist.** 66(26, 7 Feb.):9291-9298.
- NMFS. 2005. Endangered fish and wildlife; Notice of Intent to prepare an Environmental Impact Statement. **Fed. Regist.** 70(7, 11 Jan.):1871-1875.
- NMFS. 2008a. Endangered and threatened wildlife; notice of 12-month finding on a petition to list the ribbon seal as a Threatened or Endangered species. **Fed. Regist.** 73(250, 30 Dec.):79822-79828.
- NMFS. 2008b. Final Environmental Impact Statement for issuing annual quotas to the Alaska Exkimo Whaling Commission for a subsistence hunt on bowhead whales for the years 2008 through 2012. NOAA/NMFS.
- NMFS. 2009. Endangered and threatened wildlife and plants; proposed threatened and not warranted status for distinct population segments of the spotted seal. **Fed. Regist.** 74(201, 20 Oct.):53,683-53,696.
- NMML. 2009. Polar Ecosystems Project. Data available online at:
<http://www.afsc.noaa.gov?Quarterly/amj2007/divruptsNMML4.htm>.
- NOAA and USN. 2001. Joint interim report: Bahamas marine mammal stranding event of 14–16 March 2000. U.S. Dep. Commer., Nat. Oceanic Atmos. Admin., Nat. Mar. Fish. Serv., Sec. Navy, Assis. Sec. Navy, Installations and Envir. 61 p.
- NPFMC. 2009. Fishery Management Plan for Fish Resources of the Arctic Management Area. Prepared by the North Pacific Fishery Management Council, Anchorage, AK.
- O'Corry-Crowe, G.M., R.S. Suydam, A. Rosenberg, K.J. Frost and A.E. Dizon. 1997. Phylogeography, population structure and dispersal patterns of the beluga whale *Delphinapterus leucas* in the western Nearctic revealed by mitochondrial DNA. **Molec. Ecol.** 6(10):955-970.
- Patenaude, N.J., W.J. Richardson, M.A. Smultea, W.R. Koski, and G.W. Miller. 2002. Aircraft sound and disturbance to bowhead and beluga whales during spring migration in the Alaskan Beaufort Sea. **Marine Mammal Science** 18(2):309–335.
- Petersen, M.R., J.B. Grand and C.P. Dau. 2000. Spectacled Eider (*Somateria fischeri*). In: A. Poole and F. Gill (eds.), The Birds of North America, No. 547. The Birds of North America, Inc., Philadelphia, PA.
- Potter, J.R. 2004. A possible mechanism for acoustic triggering of decompression sickness symptoms in deep-diving marine mammals. Paper presented to the 2004 IEEE International Symposium on Underwater Technology, Taipei, Taiwan, 19-23 April 2004. Available at http://www.zifios.com/documentos-oficiales/documentos/Singapore_John_R_Potter_UT04.pdf.
- Quakenbush, L.T. 1988. Spotted seal, *Phoca largha*. p. 107-124 In: J.W. Lentfer (ed.), Selected Marine Mammals of Alaska/Species Accounts with Research and Management Recommendations. Marine Mammal Commis., Washington, DC. 275 p.
- Quakenbush, L.T. 2007. Preliminary satellite telemetry results for Bering-Chukchi-Beaufort bowhead whales. *Rep. Int. Whal. Comm.* SC/59/BRG12.
- Quakenbush, L.T. 2009. Summary of maps of fall movements of bowhead whales in the Chukchi Sea. Alaska Dept. of Fish and Game, online at http://wildlife.alaska.gov/management/mm/bow_move_Chukchi_sea.pdf.
- Quakenbush, L., R. Suydam, T. Obritschkewitsch, and M. Deering. 2004. Breeding biology of Steller's eiders (*Polysticta stelleri*) near Barrow, Alaska, 1991-99. **Arctic** 57(2):166-182.
- Ramsay, M.A. and I. Stirling. 1988. Reproductive biology and ecology of female polar bears (*Ursus maritimus*). **J. Zool.** 214:601-634.
- Ray, C.E. 1971. Polar bear and mammoth on the Pribilof Islands. **Arctic** 24(1):9-19.
- Read, A.J. 1999. Harbour porpoise *Phocoena phocoena* (Linnaeus, 1758). p. 323-355 In: S.H. Ridgway and R. Harrison (eds.), Handbook of Marine Mammals Vol. 6: The Second Book of Dolphins and the Porpoises. Academic Press, San Diego, CA. 486 p.
- Reeves, R.R. 1980. Spitsbergen bowhead stock: a short review. **Mar. Fish. Rev.** 42(9/10):65-69.

IV. Literature Cited

- Reeves, R.R., B.S. Stewart, P.J. Clapham and J.A. Powell. 2002. Guide to Marine Mammals of the World. Chanticleer Press, New York, NY.
- Reiser, C.R., B. Haley, D. Savarese, and D. Ireland. 2009. Chapter 3 In Funk, D.W., R. Rodrigues, D.S. Ireland, and W.R. Koski (eds.). Joint monitoring program in the Chukchi and Beaufort seas, July–November 2007. LGL Alaska Report P971-2. Report from LGL Alaska Research Associates, Inc., Anchorage, Ak, LGL Ltd., environmental research associates, King City, Ont., JASCO Research, Victoria, B.C., and Greeneridge Sciences, Inc., Goleta, CA, for Shell Offshore, Inc., ConocoPhillips Alaska, Inc., and National Marine Fisheries Service, and U.S. Fish and Wildlife Service.
- Rendell, L.E. and J.C.D. Gordon. 1999. Vocal response of long-finned pilot whales (*Globicephala melas*) to military sonar in the Ligurian Sea. **Mar. Mamm. Sci.** 15(1):198-204.
- Rice, D.W. and A.A. Wolman. 1971. The life history and ecology of the gray whale (*Eschrichtius robustus*). **Am. Soc. Mamm. Spec. Publ.** 3. 142 p.
- Richard, P.R., A.R. Martin and J.R. Orr. 1997. Study of summer and fall movements and dive behaviour of Beaufort Sea belugas, using satellite telemetry: 1992-1995. ESRF Rep. 134. Environ. Stud. Res. Funds, Calgary, Alb. 38 p.
- Richard, P.R., A.R. Martin and J.R. Orr. 2001. Summer and autumn movements of belugas of the eastern Beaufort Sea stock. **Arctic** 54(3):223-236.
- Richardson, W.J. and C.I. Malme. 1993. Man-made noise and behavioral responses. p. 631-700 In: J.J. Burns, J.J. Montague and C.J. Cowles (eds.), The bowhead whale. Spec. Publ. 2. Soc. Mar. Mammal., Lawrence, KS. 787 p.
- Richardson, W.J., and J.W. Lawson (eds.). 2002. Marine mammal monitoring of WesternGeco's open-water seismic program in the Alaskan Beaufort Sea, 2001. LGL Rep. TA2564-4. Rep. from LGL Ltd., King City, Ont. for WesternGeco LLC, Anchorage, AK, BP Exploration (Alaska) Inc., Anchorage, AK, and Nat. Mar. Fish. Serv., Anchorage, AK and Silver Springs, MD.
- Richardson, W.J. and D.H. Thomson (eds). 2002. Bowhead whale feeding in the eastern Alaskan Beaufort Sea: update of scientific and traditional information. OCS Study MMS 2002-012; LGL Rep. TA2196-7. Rep. from LGL Ltd., King City, Ont., for U.S. Minerals Manage. Serv., Anchorage, AK, and Herndon, VA. xlv + 697 p. 2 vol. NTIS PB2004-101568. Available from www.mms.gov/alaska/ref/AKPUBS.HTM#2002.
- Richardson, W.J., M.A. Fraker, B. Würsig E and R.S. Wells. 1985a. Behaviour of bowhead whales, *Balaena mysticetus*, summering in the Beaufort Sea: reactions to industrial activities. **Biol. Conserv.** 32(3):185-230.
- Richardson, W. J., Wells, R.S. and B. Wursig. 1985b. Disturbance responses of bowheads, 1980-84. p. 89-196 In: W.J. Richardson (ed.), Behavior, Disturbance Responses and Distribution of Bowhead Whales *Balaena mysticetus* in the Eastern Beaufort Sea, 1980-84. OCS Study MMS 85-0034. Rep. from LGL Ecol. Res. Assoc. Inc., Bryan, TX, for U.S. Minerals, Manage. Serv., Reston, VA. 306 p. NTIS PB87-124376.
- Richardson, W.J., B. Würsig and C.R. Greene. 1986. Reactions of bowhead whales, *Balaena mysticetus*, to seismic exploration in the Canadian Beaufort Sea. **J. Acoust. Soc. Am.** 79(4):1117-1128.
- Richardson, W.J., R.A. Davis, C.R. Evans, D.K. Ljungblad and P. Norton. 1987. Summer distribution of bowhead whales, *Balaena mysticetus*, relative to oil industry activities in the Canadian Beaufort Sea, 1980-84. **Arctic** 40(2):93-104.
- Richardson, W.J., C.R. Greene, Jr., C.I. Malme and D.H. Thomson. 1995. Marine Mammals and Noise. Academic Press, San Diego. 576 p.
- Richardson, W.J., G.W. Miller and C.R. Greene Jr. 1999. Displacement of migrating bowhead whales by sounds from seismic surveys in shallow waters of the Beaufort Sea. **J. Acoust. Soc. Am.** 106(4, Pt. 2):2281.
- Riedman, M. 1990. The Pinnipeds: Seals, Sea Lions, and Walruses. Univ. Calif. Press, Berkeley and Los Angeles, CA. 439 p.

- Roth, E.H., and V. Schmidt. 2010. U.S. Geological Survey coastal and marine geology report on cooperative agreement G09AC00352: Analysis of acoustic sound pressure levels generated by research icebreakers and marine seismic sources in the deep-water, Arctic Ocean. Report prepared by the Marine Physical Laboratory of the Scripps Institution of Oceanography, University of California, San Deigo, La Holla, CA,
- Rugh, D. (ed.) 2009. Bowhead Whale Feeding Ecology Study (BOWFEST) in the Western Beaufort Sea; 2008 Annual Report. MMS-4500000120. Produced through the National Marine Mammal Laboratory, Alaska Fisheries Science Center, NMFS, NOAA, 7600 Sand Point Way, NE Seattle, WA 98115-6349.
- Rugh, D.J., and M.A. Fraker. 1981. Gray whale (*Eschrichtius robustus*) sightings in eastern Beaufort Sea. **Arctic** 34(2):186-187.
- Rugh, D.J., K.E.W. Shelden and D.E. Withrow. 1997. Spotted seals, *Phoca largha*, in Alaska. **Mar. Fish. Rev.** 59(1):1-18.
- Rugh, D.J., D. DeMaster, A. Rooney, J. Breiwick, K. Sheldon and S. Moore. 2003. A review of bowhead whale (*Balaena mysticetus*) stock identity. **J. Cetacean Res. Manage.** 5(3):267-279.
- Rugh, D.J., R.C. Hobbs, J.A. Lerczak and J.M. Breiwick. 2005. Estimates of abundance of the eastern North Pacific stock of gray whales (*Eschrichtius robustus*) 1997-2002. **J. Cetac. Res. Manage.** 7(1):1-12.
- Rugh, D., J. Breiwick, M. Muto, R. Hobbs, K. Shelden, C. D'Vincent, I.M. Laursen, S. Reif, S. Maher, and S. Nilson. 2008. Report of the 2006-2007 census of the eastern North Pacific stock of gray whales. Alaska Fisheries Science Center Report 2008-03.
- Savarese, D.M., C.R. Reiser, D.S. Ireland, and R. Rodrigues. 2009. Beaufort Sea vessel-based monitoring program. Chapter 6 *In* Funk, D.W., D.S. Ireland, R. Rodrigues, and W.R. Koski (eds.). 2009. Joint monitoring program in the Chukchi and Beaufort seas, July–November 2006-2008. LGL Alaska Report P1050-1. Report from LGL Alaska Research Associates, Inc., Anchorage, Ak, LGL Ltd., environmental research associates, King City, Ont., Greeneridge Sciences, Inc., Goleta, CA, and JASCO Research, Victoria, B.C., for Shell Offshore, Inc. and other Industry contributors, National Marine Fisheries Service, and U.S. Fish and Wildlife Service. 488 p. plus appendices.
- Schlundt, C.E., J.J. Finneran, D.A. Carder and S.H. Ridgway. 2000. Temporary shift in masking hearing thresholds of bottlenose dolphins, *Tursiops truncatus*, and white whales, *Delphinapterus leucas*, after exposure to intense tones. **J. Acoust. Soc. Am.** 107(6):3496-3508.
- Sease, J.L. and D.G. Chapman. 1988. Pacific walrus (*Odobenus rosmarus divergens*). p. 17-38 *In* J.W. Lentfer (eds) Selected Marine Mammals of Alaska: Species Accounts with Research and Management Recommendations. Mar. Mamm. Comm., Washington, D.C. NTIS PB88-178462.
- Sekiguchi, K., T. Uyama, and K. Yamshiro. 2008. Cetacean sighting survey during T/S Oshoro Maru Bering and Chukchi Sea cruise, 1 July–28 August 2007.
- Shaughnessy, P.D. and F.H. Fay. 1977. A review of the taxonomy and nomenclature of North Pacific harbor seals. **J. Zool. (Lond.)** 182:385-419.
- Smith, A.E. and M.R.J. Hill. 1996. Polar bear, *Ursus maritimus*, depredation of Canada Goose, *Branta canadensis*, nests. **Can. Field-Nat.** 110(2):339-340.
- Smith, T.G. 1973. Population dynamics of the ringed seal in the Canadian eastern arctic. **Fish. Res. Board Can. Bull.** 181. 55 p.
- Smith, T.G. 1985. Polar Bears, *Ursus maritimus*, as predators of Belugas *Delphinapterus leucas*. **Can. Field-Nat.** 99(1):71-75.
- Smith, T.G. and I. Stirling. 1975. The breeding habitat of the ringed seal (*Phoca hispida*). the birth lair and associated structures. **Can. J. Zool.** 53(9):1297-1305.
- Smultea, M.A., M. Holst, W.R. Koski and S. Stoltz. 2004. Marine mammal monitoring during Lamont-Doherty Earth Observatory's seismic program in the Southeast Caribbean Sea and adjacent Atlantic Ocean, April-June 2004. LGL Rep. TA2822-26. Rep. from LGL Ltd., King City, ON, for Lamont-Doherty Earth Observatory of Columbia Univ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD. 106 p.

IV. Literature Cited

- Stehn, R.A., C.P. Dau, B. Conant, and W.I. Butler, Jr. 1993. Decline of spectacled eiders nesting in western Alaska. **Arctic** 46(3):264-277.
- Stemp, R. 1985. Observations on the effects of seismic exploration on seabirds. p. 217-233 *In*: G.D. Greene, F.R. Engelhardt and R.J. Paterson (eds.), Proc. workshop on effects of explosives use in the marine environment, January 29 to 31, 1985, Halifax. Tech. Rep. 5. Energy, Mines and Resources Canada and Indian and Northern Affairs Canada, Canada Oil and Gas Lands Administration, Environ. Prot. Branch, Ottawa, Ont.
- Stewart, B.S. and S. Leatherwood. 1985. Minke whale *Balaenoptera acutorostrata* Lacépède, 1804. p. 91-136 *In*: S.H. Ridgway and R. Harrison (eds.), Handbook of Marine Mammals, Vol. 3.: The Sirenians and Baleen Whales. Academic Press, London, U.K. 362 p.
- Stirling, I. and E.H. McEwan. 1975. The caloric value of whole ringed seals (*Phoca hispida*) in relation to polar bear (*Ursus maritimus*) ecology and hunting behavior. **Can. J. Zool.** 53(8):1021-1027.
- Stirling, I., M. Kingsley and W. Calvert. 1982. The distribution and abundance of seals in the eastern Beaufort Sea, 1974-79. **Can. Wildl. Serv. Occas. Pap.** 47:25 p.
- Stone, C.J. 2003. The effects of seismic activity on marine mammals in UK waters 1998-2000. JNCC Report 323. Joint Nature Conservation Committee, Aberdeen, Scotland. 43 p.
- Stroeve, J., M. Serreze, S. Drobot, S. Gearheard, M. Holland, J. Maslanik, W. Meier, and T. Scambos. 2008. Arctic sea ice extent plummets in 2007. EOS, **Transactions, American Geophysical Union** 89(2):13-14.
- Suydam, R.S. and J.C. George. 1992. Recent sightings of harbor porpoises, *Phocoena phocoena*, near Point Barrow, Alaska. **Can. Field Nat.** 106(4): 489-492.
- Suydam, R.S., R.P. Angliss, J.C. George, S.R. Braund and D.P. DeMaster. 1995. Revised data on the subsistence harvest of bowhead whales (*Balaena mysticetus*) by Alaska eskimos, 1973-1993. **Rep. Int. Whal. Comm.** 45:335-338.
- Suydam, R.S., L.F. Lowry, K.J. Frost, G.M. O'Corry-Crowe and D. Pikok Jr. 2001. Satellite tracking of eastern Chukchi Sea beluga whales into the Arctic Ocean. **Arctic** 54(3):237-243.
- Suydam, R.S., L.F. Lowry, and K.T. Frost. 2005a. Distribution and movements of beluga whales from the eastern Chukchi Sea Stock during summer and early autumn. Report prepared the North Slope Borough Department of Wildlife Management, Barrow, Alaska.
- Suydam, R.S., J.C. George, C. Hanns, and G. Sheffield. 2005b. Subsistence harvest of bowhead whales (*Balaena mysticetus*) by Alaskan Eskimos during 2004. SC/57/BRG15.
- Suydam, R.S., J.C. George, C. Hanns, and G. Sheffield. 2006. Subsistence harvest of bowhead whales (*Balaena mysticetus*) by Alaskan Eskimos during 2005. SC/58/BRG21.
- Suydam, R.S., J.C. George, C. Rosa, B. Person, C. Hanns, G. Sheffield, and J. Bacon. 2007. Subsistence harvest of bowhead whales (*Balaena mysticetus*) by Alaskan Eskimos during 2006. SC/59/BRG4.
- Suydam, R.S., J.C. George, C. Rosa, B. Person, C. Hanns, G. Sheffield, and J. Bacon. 2008. Subsistence harvest of bowhead whales (*Balaena mysticetus*) by Alaskan Eskimos during 2007. SC/60/BRG10.
- Suydam, R.S., J.C. George, C. Rosa, B. Person, C. Hanns, G. Sheffield, and J. Bacon. 2009. Subsistence harvest of bowhead whales (*Balaena mysticetus*) by Alaskan Eskimos during 2008. SC/61/BRG6.
- Sverdrup, H.U., M.W. Johnson and R.H. Fleming. 1942. The Oceans. Prentice-Hall, Inc., Englewood Cliffs, NJ. 1087 p.
- Swartz, S.L. and M.L. Jones. 1981. Demographic studies and habitat assessment of gray whales, *Eschrichtius robustus*, in Laguna San Ignacio, Baja California, Mexico. U.S. Mar. Mamm. Comm. Rep. MMC-78/03. 34 p. NTIS PB-289737.
- TERA. 1997. Distribution and abundance of spectacled eiders in the vicinity of Prudhoe Bay, Alaska: 1997 status report. Unpublished report prepared by Troy Environmental Research Associates, Anchorage, AK.
- TERA. 1999. Spectacled eiders in the Beaufort Sea: distribution and timing of use. Unpublished report prepared by Troy Ecological Research Associates, Anchorage, AK.

- Thomas, T.A., W.R. Koski and W.J. Richardson. 2002. Correction factors to calculate bowhead whale numbers from aerial surveys of the Beaufort Sea. p. 15-1 to 15-28 (Chap. 15) *In*: W.J. Richardson and D.H. Thomson (eds.), Bowhead whale feeding in the eastern Alaskan Beaufort Sea: update of scientific and traditional information, vol. 1. OCS Study MMS 2002-012; LGL Rep. TA2196-7. Rep. from LGL Ltd., King City, Ont., for U.S. Minerals Manage. Serv., Anchorage, AK, and Herndon, VA. 420 p.
- Thomas, T., W.R. Koski, and D.S. Ireland. 2009. Chukchi Sea nearshore aerial surveys. Chapter 4 *In* Funk, D.W., D.S. Ireland, R. Rodrigues, and W.R. Koski (eds.). Joint monitoring program in the Chukchi and Beaufort seas, July–November 2006–2008. LGL Alaska Report P1050-1. Report from LGL Alaska Research Associates, Inc., Anchorage, Ak, LGL Ltd., environmental research associates, King City, Ont., Greeneridge Sciences, Inc., Goleta, CA, and JASCO Research, Victoria, B.C., for Shell Offshore, Inc. and other Industry contributors, National Marine Fisheries Service, and U.S. Fish and Wildlife Service. 488 p. plus appendices.
- Thompson, D., M. Sjöberg, E.B. Bryant, P. Lovell and A. Bjørge. 1998. Behavioural and physiological responses of harbour (*Phoca vitulina*) and grey (*Halichoerus grypus*) seals to seismic surveys. p. 134 *In*: World Marine Mammal Science Conf. Abstract volume, Monaco. 160 p.
- Tolstoy, M., J. Diebold, S. Webb, D. Bohnenstiehl and E. Chapp. 2004a. Acoustic calibration measurements. Chapter 3 *In*: W.J. Richardson (ed.), Marine mammal and acoustic monitoring during Lamont-Doherty Earth Observatory's acoustic calibration study in the northern Gulf of Mexico, 2003. Revised ed. Rep. from LGL Ltd., King City, ON, for Lamont-Doherty Earth Observ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD. [Advance copy of updated Chapter 3. Available at: http://www.nmfs.noaa.gov/pr/readingrm/mmpa_small_take/gom_90d_report_final.pdf
- Tolstoy, M., J.B. Diebold, S.C. Webb, D.R. Bohnenstiehl, E. Chapp, R.C. Holmes and M. Rawson. 2004b. Broadband calibration of R/V *Ewing* seismic sources. **Geophys. Res. Lett.** 31:L14310.
- Tomilin, A.G. 1957. Mammals of the U.S.S.R. and adjacent countries. Vol, 9: Cetaceans. Israel Progr. Sci. Transl. (1967), Jerusalem. 717 p. NTIS TT 65-50086.
- Troy, D. 2003. Molt migration of spectacled eiders in the Beaufort Sea region. Unpublished report prepared by Troy Ecological Research Associates, Anchorage, AK.
- Treacy, S.D. 1988. Aerial surveys of endangered whales in the Beaufort Sea, fall 1987. OCS Study MMS 88-0030. U.S. Minerals Manage. Serv., Anchorage, AK. 142 p. NTIS PB89-168785.
- Treacy, S.D. 1989. Aerial surveys of endangered whales in the Beaufort Sea, fall 1988. OCS Study MMS 89-0033. U.S. Minerals Manage. Serv., Anchorage, AK. 102 p. NTIS PB90-161464.
- Treacy, S.D. 1990. Aerial surveys of endangered whales in the Beaufort Sea, fall 1989. OCS Study MMS 90-0047. U.S. Minerals Manage. Serv., Anchorage, AK. 105 p. NTIS PB91-235218.
- Treacy, S.D. 1991. Aerial surveys of endangered whales in the Beaufort Sea, fall 1990. OCS Study MMS 91-0055. U.S. Minerals Manage. Serv., Anchorage, AK. 108 p. NTIS PB92-176106.
- Treacy, S.D. 1992. Aerial surveys of endangered whales in the Beaufort Sea, fall 1991. OCS Study MMS 92-0017. U.S. Minerals Manage. Serv., Anchorage, AK. 93 p.
- Treacy, S.D. 1993. Aerial surveys of endangered whales in the Beaufort Sea, fall 1992. OCS Study MMS 93-0023. U.S. Minerals Manage. Serv., Anchorage, AK. 136 p.
- Treacy, S.D. 1994. Aerial surveys of endangered whales in the Beaufort Sea, fall 1993. OCS Study MMS 94-0032. U.S. Minerals Manage. Serv., Anchorage, AK. 133 p.
- Treacy, S.D. 1995. Aerial surveys of endangered whales in the Beaufort Sea, fall 1994. OCS Study MMS 95-0033. U.S. Minerals Manage. Serv., Anchorage, AK. 116 p.
- Treacy, S.D. 1996. Aerial surveys of endangered whales in the Beaufort Sea, fall 1995. OCS Study MMS 96-0006. U.S. Minerals Manage. Serv., Anchorage, AK. 121 p. NTIS PB97-115752
- Treacy, S.D. 1997. Aerial surveys of endangered whales in the Beaufort Sea, fall 1996. OCS Study MMS 97-0016. U.S. Minerals Manage. Serv., Anchorage, AK. 115 p. NTIS PB97-194690

IV. Literature Cited

- Treacy, S.D. 1998. Aerial surveys of endangered whales in the Beaufort Sea, fall 1997. OCS Study MMS 98-0059. U.S. Minerals Manage. Serv., Anchorage, AK. 143 p. Published 1999.
- Treacy, S.D. 2000. Aerial surveys of endangered whales in the Beaufort Sea, fall 1998-1999. OCS Study MMS 2000-066. U.S. Minerals Manage. Serv., Anchorage, AK. 135 p.
- Treacy, S.D. 2002a. Aerial surveys of endangered whales in the Beaufort Sea, fall 2000. OCS Study MMS 2002-014. U.S. Minerals Manage. Serv., Anchorage, AK. 111 p.
- Treacy, S.D. 2002b. Aerial surveys of endangered whales in the Beaufort Sea, fall 2001. OCS Study MMS 2002-061. U.S. Minerals Manage. Serv., Anchorage, AK. 117 p.
- Treacy, S.D., J.S. Gleason and C.J. Cowles. 2006. Offshore distances of bowhead whales (*Balaena mysticetus*) observed during fall in the Beaufort Sea, 1982-2000: an alternative interpretation. *Arctic* 59(1):83-90.
- Troy, D.M. 2003. Molt migration of spectacled eiders in the Beaufort Sea region. Unpublished report prepared by Troy Ecological Research Associates, Anchorage, AK, for BP Exploration (Alaska) Inc., Anchorage, AK. 17p.
- Tyack, P., M. Johnson and P. Miller. 2003. Tracking responses of sperm whales to experimental exposures of airguns. p. 115-120 *In*: A.E. Jochens and D.C. Biggs (eds.), Sperm whale seismic study in the Gulf of Mexico/Annual Report: Year 1. OCS Study MMS 2003-069. Rep. from Texas A&M Univ., College Station, TX, for U.S. Minerals Manage. Serv., Gulf of Mexico OCS Reg., New Orleans, LA.
- Tynan, C.T., and D.P. DeMaster. 1997. Observations and predictions of Arctic climate change: Potential effects on marine mammals. *Arctic* 50(4):308-322.
- Udevitz, M.S., D.M. Burn, and M.A. Webber. 2008. Estimation of walrus populations on sea ice with infrared imagery and aerial photograph[hy. *Marine Mammal Science* 24(1):57-70.
- UN Atlas of the Oceans. n. d. United Nations Atlas of the Oceans. Available at <http://www.oceansatlas.org/index.jsp>
- UNEP-WCMC. 2004. UNEP-WCMC species database: CITES-listed species. Available at <http://www.unep-wcmc.org/index.html?http://sea.unep-wcmc.org/isdb/CITES/Taxonomy/tax-gs-search1.cfm?displaylanguage=eng&source=animals~main>
- U.S. Army Corps of Engineers (USACE). 1999. Final Environmental Impact Statement, Beaufort Sea Oil and Gas Development/Northstar Project. Prepared by U.S. Army Corps of Engineers, Alaska.
- USDI/MMS (U.S. Department of the Interior/Minerals Management Service). 1996. Beaufort Sea Planning Area Oil and Gas Lease Sale 144 Final Environmental Impact Statement.
- USDI/BLM (U.S. Department of the Interior/Bureau of Land Management). 2003. Northwest National Petroleum Reserve – Alaska; Final Amended Integrated Activity Plan/Environmental Impact Statement.
- USDI/BLM (U.S. Department of the Interior/Bureau of Land Management). 2005. Northwest National Petroleum Reserve – Alaska; Final Amended Integrated Activity Plan/Environmental Impact Statement.
- USDI/MMS (U.S. Department of the Interior/Minerals Management Service). 2003. Beaufort Sea Planning Area Oil and gas lease sales 186, 195, and 202 – Final Environmental Impact Statement.
- U.S. Fish and Wildlife Service (USFWS). 1996. Spectacled eider recovery plan. Anchorage, AK. 157 p.
- U.S. Fish and Wildlife Service (USFWS). 2002. Steller's Eider Recovery Plan. Fairbanks, AK. 27 p.
- USFWS. 2000a. Pacific walrus (*Odobenus rosmarus divergens*): Alaska Stock. p. 185-190 *In*: R.C. Ferrero, D.P. DeMaster, P.S. Hill, M.M. Muto, and A.L. Lopez (eds.) Alaska Marine Mammal Stock Assessments, 2000. NOAA Tech. Memo. NMFS-AFSC-119. U.S. Dep. Comm. NOAA, NMFS, Alaska Fisheries Science Center.
- USFWS. 2000b. Polar Bear: Alaska Chukchi/Bering Seas. p. 175-179 *In*: R.C. Ferrero, D.P. DeMaster, P.S. Hill, M.M. Muto, and A.L. Lopez (eds.) Alaska Marine Mammal Stock Assessments, 2000. NOAA Tech. Memo. NMFS-AFSC-119. U.S. Dep. Comm. NOAA, NMFS, Alaska Fisheries Science Center.
- USFWS. 2000c. Polar bear: Alaska southern Beaufort Sea. p. 180-184 *In*: R.C. Ferrero, D.P. DeMaster, P.S. Hill, M.M. Muto, and A.L. Lopez (eds.) Alaska Marine Mammal Stock Assessments, 2000. NOAA Tech. Memo. NMFS-AFSC-119. U.S. Dep. Comm. NOAA, NMFS, Alaska Fisheries Science Center.

- USFWS. 2006. Draft Study Plan for Estimating the Size of the Pacific Walrus Population. Marine Mammals Management, U.S. Fish and Wildlife Service, Alaska Science Center, U.S. Geological Survey, GiproRybFlot, Research and Engineering Institute for the Development and Operation of Fisheries, ChukotTINRO, Pacific Research Institute of Fisheries and Oceanography.
- USFWS. 2007. Biological Opinion for Chukchi Sea Planning Area Oil and Gas Lease Sale 193 and associated seismic surveys and exploratory drilling. Consultation with the Minerals Management Service–Alaska OCS Region, Anchorage, AK.
- USFWS. 2008. Endangered and Threatened Wildlife and Plants; determination of Threatened status for the polar bear (*Ursus maritimus*) throughout its range; Final Rule. **Fed. Regist.** 73(95, 15 May):28,212-28,302.
- USFWS. 2009a. Endangered and Threatened Wildlife and Plants; designation of critical habitat for the polar bear (*Ursus maritimus*) in the United States. **Fed. Regist.** 74(208, 29 Oct):56058-56086.
- USFWS. 2009b. Polar bear harvest management in Alaska. Fact Sheet published by U.S. Fish & Wildlife Service available at <http://alaska.fws.gov/fisheries/mmm/polarbear/facts.htm>.
- van Meurs, R. and J.F. Splettstoesser. 2003. Letter to the editor–Farthest North Polar Bear. **Arctic** 56(3):309.
- Vibe, C. 1950. The marine mammals and the marine fauna in the Thule District (northwest Greenland) with observations on ice conditions in 1939-41 **Medd. Grønl.** 150:1-115.
- Wardle, C.S., T.J. Carter, G.G. Urquhart, A.D.F. Johnstone, A.M. Ziolkowski, G. Hampson and D. Mackie. 2001. Effects of seismic air guns on marine fish. **Cont. Shelf Res.** 21(8-10):1005-1027.
- Watkins, W.A., K.E. Moore and P. Tyack. 1985. Sperm whale acoustic behaviors in the southeast Caribbean. **Cetology** 49:1-15.
- Williams, M.T. and J.A. Coltrane (eds.). 2002. Marine mammal and acoustical monitoring of the Alaska Gas Producers Pipeline Team's open water pipeline route survey and shallow hazards program in the Alaskan Beaufort Sea, 2001. LGL Rep. P643. Rep. from LGL Alaska Res. Assoc. Inc., Anchorage, AK, for BP Explor. (Alaska) Inc., ExxonMobil Production, Phillips Alaska Inc., and Nat. Mar. Fish. Serv. 103 p.
- Wilson, D. E. 1976. Cranial variation in polar bears. *Int. Conf. Bear Res. Manage.* **IUCN Publ. New Series** 40: 447-453.
- Wolfe, R. and R. Walker. 1987. Subsistence economies in Alaska: Productivity, geography and development impacts. **Arctic Anthr.** 24(2):56-81.
- Woodby, D.A. and D.B. Botkin. 1993. Stock sizes prior to commercial whaling. p. 387-407 *In: J.J. Burns, J.J. Montague and C.J. Cowles (eds.), The Bowhead Whale. Spec. Publ. 2. Soc. Mar. Mamm., Lawrence, KS.* 787 p.
- Woodgate, R.A., K. Aagard, R.D. Muench, J. Gunn, G. Bork, B. Rudels, A.T. Roach and U. Schauer. 2001. The Arctic Ocean boundary current along the Eurasian slope and the adjacent Lomonosov Ridge: water mass properties, transports and transformations from moored instruments. **Deep-Sea Res.**
- Wynn, K. 1997. Guide to Marine Mammals of Alaska. Alaska Sea Grant College Program, University of Alaska, Fairbanks.
- Yoder, J.A. 2002. Declaration of James A. Yoder in opposition to plaintiff's motion for temporary restraining order, 28 October 2002. Civ. No. 02-05065-JL. U.S. District Court, Northern District of California, San Francisco Division.
- Zeh, J.E. and A.E. Punt. 2005. Updated 1978-2001 abundance estimates and their correlations for the Bering-Chukchi-Beaufort Seas stock of bowhead whales. **J. Cetac. Res. Manage.** 7(2):169-175.
- Zeh, J.E., C.W. Clark, J.C. George, D. Withrow, G.M. Carroll and W.R. Koski. 1993. Current population size and dynamics. p. 409-489 *In: J.J. Burns, J.J. Montague and C.J. Cowles (eds.), The Bowhead Whale. Spec. Publ. 2. Soc. Mar. Mamm., Lawrence, KS.* 787 p.
- Zeh, J.E., A.E. Raftery, and A.A. Schaffner. 1996. Revised estimates of bowhead population size and rate of increase. **Rep. Int. Whal. Comm.** 46:670.

IV. Literature Cited

Zerbini, A.N., J.M. Waite, J.L. Laake, and P.R. Wade. 2006. Abundance, trends and distribution of baleen whales off western Alaska and the central Aleutian Islands. **Deep-sea Research**, Part 1. 53(11):1772-1790.

Marine mammal distribution and abundance in the western Canadian high Arctic has generally been restricted to regions within 100 to 200 km from the shore (LGL 2005). The survey area is more than 200nm from the shore. In the survey area, primarily the Canadian Basin, the marine mammals that could be potentially encountered belong to three taxonomic groups: odontocetes (toothed whales), mysticetes (baleen whales) and carnivora (seals and polar bears). Marine mammals that could be encountered include four (4) cetacean species (beluga whale, bowhead whale, narwhal and killer whale), five (5) piniped species (walrus, bearded seal, ringed seal, hooded seal and harp seal), and the polar bear. However, it must be noted that most of these species will: 1), occur in low numbers in the study area, 2) are not likely to be encountered in survey area and 3) most require regularly spaced breathing holes or open water to survive. In the *Healy* Expedition survey region where active multibeam data collection will occur, the extreme thickness of the pack-ice precludes the permanent establishment and continual maintenance of breathing holes for most if not all of the fully aquatic marine mammals. In addition, it is unlikely that polar bears will be frequently encountered due to the lack of their prime food resource, seals, which are not able to establish and maintain regular breathing holes in the thick pack-ice of the study area and consequently will restrict their distribution to nearshore open waters with less severe ice conditions. [Ref: *Draft Environmental Assessment for Canadian Polar Margin Seismic Reflection Survey in Waters Offshore of the Western Canadian Arctic Islands in Support of the Law of the Sea, Sept 6-Oct 1, 2008; prepared by Triton for the Canada Dept of Natural Resources, May 2008.*]

In addition, there has been a review of a number of Environmental Impacts Statements from previous expeditions on the *Healy* as well as the Environmental Assessment conducted by Canada for this expedition. While it is clear that there are environmental impacts from the use of seismic air guns in collecting seismic data, there is no evidence of potential environmental impacts from the use of multi-beam echo sounders. Consultation with the NMFS Office of Protected Resources confirmed that this is the current view of their office as well. However, OPR suggested that NOAA take this opportunity to collect data on the behavior of any marine mammals encountered for use in future expeditions. In addition, NOAA had an analysis done by one of its university partners in regard to the SeaBeam 2112 multi-beam echo sounder that is currently installed in the *Healy*. See Attachment II. In sum, the survey or mapping activities are not expected to have significant impacts of a direct or cumulative nature. NOAA's own hydrographic fleet routinely operates the sonar frequencies that will be used on this project. Knowledgeable experts who are aware of the sensitivities of the marine environment will conduct the at-sea portions of these projects. In particular, the plan is that anywhere where they are most likely to see a marine mammal they will operate the echo sounder at levels below the thresholds of concern identified in the documents reviewed for this decision.

CATEGORICAL EXCLUSION

This project would not result in any changes to the human environment. As defined in Sections 5.05 and 6.03.c.3(a) of NAO 216-6, these are research projects of limited size or magnitude or with only short term effects on the environment and for which any cumulative effects are negligible. As such, this project is categorically excluded from the need to prepare an Environmental Assessment.

APPENDIX B
NOAA's 2009 LETTER OF CATEGORICAL EXCLUSION FOR *HEALY*
OPERATIONS IN INTERNATIONAL WATERS



UNITED STATES DEPARTMENT OF COMMERCE
OCEANIC & ATMOSPHERIC RESEARCH
Office of Ocean Exploration and Research
1315 East West Hwy, 10th Floor, R/OE
Silver Spring, MD 20910

June 29, 2009

MEMORANDUM FOR: The Record

FROM: CAPT Harris B. Halverson, NOAA
Acting Director
Office of Ocean Exploration and Research

SUBJECT: Categorical Exclusion for Extended Continental Shelf (ECS)
Expedition to Collect Bathymetric & Gravity Data, August-
September 2009

NAO 216-6, Environmental Review Procedures, requires all proposed U.S. Federal actions to be reviewed with respect to consequences on the natural and human environment. This memorandum addresses the NAO's applicability to the U.S. role in the survey activities described below.

PROJECT DESCRIPTION

On August 7, 2009, NOAA's Office of Ocean Exploration and Research (OER) is embarking on an international expedition in cooperation with Canada and the U.S. Geological Survey to collect data on the seaward limits of the U.S. and Canadian extended continental shelves in portions of the western Arctic Ocean north of Barrow, AK and offshore of the western Canadian Arctic Islands. All the activities will be conducted in the high seas/international waters beyond the 200 nm EEZ limit of the United States. The U.S. Coast Guard Cutter *Healy* will be the lead vessel with the Canadian Coast Guard Ship *Louis S. St. Laurent* following. The *Healy* will be breaking up the ice for the *Louis* and collecting multi-beam data. The Canadian vessel, *Louis*, will be collecting seismic data. In addition to collecting data required for a submission under Article 76 of the Law of the Sea Convention, the project will involve the conduct of multi-disciplinary

ocean mapping and exploration activities designed to increase knowledge of marine resources in the Arctic. No on-ice operations are planned.

For more details see attached science plan (Attachment I).

PROJECT EFFECTS

In the survey area, the marine mammals that potentially could be encountered include four (4) cetacean species (beluga whale, bowhead whale, narwhal and killer whale), five (5) pinniped species (walrus, bearded seal, ringed seal, hooded seal and harp seal), and polar bears. The extreme thickness of the area's pack-ice is anticipated to preclude the permanent establishment and continual maintenance of breathing holes for most, if not all, of the above fully aquatic marine mammals. Hence, most of these species, including the polar bears that prey upon the fully aquatic marine mammals: 1) are not likely to be encountered in the survey area, and/or 2) will occur in low numbers.

Nonetheless, sounds produced in conjunction with this collection of ECS data, specifically those produced by seismic equipment, have the potential to affect the behavior of marine mammals that inhabit and/or migrate through the area. In preparation for the seismic data collection, Natural Resources Canada conducted an environmental assessment and subsequently the *Louis* received an authorization/permit from the Canadian Department of Fisheries and Oceans for their use of multichannel seismic equipment. As a condition of the permit, the operators of the *Louis* intend to alter vessel speed/course as necessary to avoid marine mammal interactions, provided it will not compromise operational safety requirements. Also in compliance with the permit issued under applicable Canadian law, *Louis* is using a safety radius/zone of 1 km such that when marine mammals are detected within 1 km of *Louis*, the seismic airgun will be silenced. Canada will have three protected species observers (PSOs) on board the *Louis* pursuant to the permit issued under Canadian law. The Canadian ship will also have members of the Hunters and Trappers Community present to monitor and mitigate, as necessary, any potential effects on marine mammals of interest to subsistence use villages.

While it is clear there is potential for environmental impact from the use of seismic air guns to collect ECS data, there is no evidence of similar potential impact from the use of the *Healy's* multi-beam echo sounders or ice-breaking capacities. Consultation with the NMFS Office of Protected Resources (OPR) confirmed that this is the current view of their office as well. In addition, NOAA Office of Coast Survey had one of its university partners conduct an analysis of marine mammal exposure to noise from the *Healy* SeaBeam 2112 multi-beam echo sounder. In sum, the SeaBeam survey or mapping activities are not expected to have significant impacts of a direct or cumulative nature. See Attachment II for more details. Even so, the U.S is sensitive to the fact that little information exists on the effects of sound on marine mammals in the Arctic and is viewing this expedition as an opportunity to collect information that will inform future ECS endeavors. Onboard *Healy*, there will be one PSO and one Alaska Native community participant; each will serve as additional lookouts for the observers on the Canadian vessel. The PSO and the community observer on *Healy* will collect marine mammal sighting and behavioral data during operations. This information, including the approximate location and distance from

the *Healy*, will be shared in real-time with the protected species observers on *Louis*, who have the authority to implement measures to mitigate further harm/disturbance to marine mammals during these field exercises. The data will also be collected to contribute to the existing body of data relating to marine mammal noise exposure and to inform future research endeavors.

CATEGORICAL EXCLUSION

The U.S. portion of this project would not result in any sustained changes to the human or natural environment. As defined in Sections 5.05 and 6.03.c.3(a) of NAO 216-6, it is a research effort of limited size or magnitude or with only short term effects on the environment and for which any cumulative effects are negligible. As such, it is categorically excluded from the need to prepare an Environmental Assessment.

APPENDIX C
MARINE FISH OF THE BEAUFORT SEA AND ARCTIC OCEAN.
FROM FISHBASE.ORG

| Species | Family | Common Name | Habitat | Length (Total Length; cm) | Trophic level | Status | Region |
|---------------------------------|-----------------|------------------------|---------------|---------------------------|---------------|--------------|---------------------------|
| <i>Agonus cataphractus</i> | Agonidae | Hooknose | demersal | 21 TL | 3.4 | native | Arctic Ocean |
| <i>Acantholumpenus mackayi</i> | Stichaeidae | Pighead prickleback | demersal | 86 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Amblyraja radiata</i> | Rajidae | Thorny skate | demersal | 100 TL | 4 | native | Beaufort Sea/Arctic Ocean |
| <i>Ammodytes dubius</i> | Ammodytidae | Northern sand lance | demersal | 25 TL | 3.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Ammodytes hexapterus</i> | Ammodytidae | Pacific sand lance | benthopelagic | 27 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Anarhichas denticulatus</i> | Anarhichadidae | Northern wolfish | benthopelagic | 180 TL | 3.8 | native | Beaufort Sea/Arctic Ocean |
| <i>Anarrhichthys ocellatus</i> | Anarhichadidae | Wolf eel | demersal | 240 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Anisarchus medius</i> | Stichaeidae | Stout eel blenny | demersal | 18 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Arctogadus borisovi</i> | Gadidae | East Siberian cod | demersal | 56 TL | 3.8 | questionable | Beaufort Sea/Arctic Ocean |
| <i>Arctogadus glacialis</i> | Gadidae | Arctic cod | bathypelagic | 33 TL | 3.7 | questionable | Beaufort Sea/Arctic Ocean |
| <i>Argyropelecus hemigymnus</i> | Sternoptychidae | Half-naked hatchetfish | bathypelagic | 5 TL | 3.3 | native | Arctic Ocean |
| <i>Arctediellus scaber</i> | Cottidae | Hamecon | demersal | 9 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Arctediellus uncinatus</i> | Cottidae | Arctic hookear sculpin | demersal | 10 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Aspidophoroides bartoni</i> | Agonidae | Aleutian alligatorfish | demersal | 22 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |

| Species | Family | Common Name | Habitat | Length (Total Length; cm) | Trophic level | Status | Region |
|--|-----------------|---------------------|---------------|---------------------------|---------------|--------|---------------------------|
| <i>Atheresthes stomias</i> | Pleuronectidae | Arrowtooth flounder | demersal | 84 TL | 4.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Bathymaster signatus</i> | Bathymasteridae | Searcher | demersal | 38 TL | 3.6 | native | Beaufort Sea/Arctic Ocean |
| <i>Boreogadus saida</i> | Gadidae | Polar cod | demersal | 40 TL | 3.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Careproctus reinhardti</i> | Liparidae | Sea tadpole | bathydemersal | 30 TL | 3.5 | native | Arctic Ocean |
| <i>Clupea pallasii pallasii</i> | Clupeidae | Pacific herring | pelagic | 46 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Coregonus autumnalis autumnalis</i> | Salmonidae | Arctic cisco | pelagic | 64 TL | 3.6 | native | Beaufort Sea/Arctic Ocean |
| <i>Coregonus laurettae</i> | Salmonidae | Bering cisco | pelagic | 54 TL | 3.8 | native | Beaufort Sea/Arctic Ocean |
| <i>Coregonus muksun</i> | Salmonidae | Muksun | benthopelagic | 64 TL | 3.3 | native | Arctic Ocean |
| <i>Coregonus nasus</i> | Salmonidae | Broad whitefish | demersal | 71 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Coregonus peled</i> | Salmonidae | Peled | demersal | 50 TL | 3 | native | Arctic Ocean |
| <i>Coregonus pidschian</i> | Salmonidae | Humpback whitefish | demersal | 46 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Coregonus sardinella</i> | Salmonidae | Common whitefish | pelagic | 47 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Cottunculus microps</i> | Psychrolutidae | Polar sculpin | bathydemersal | 37 TL | 3.4 | native | Arctic Ocean |
| <i>Cottunculus sadko</i> | Psychrolutidae | Fathead | bathydemersal | 19 TL | 3.3 | native | Arctic Ocean |
| <i>Cyclopteropsis jordani</i> | Cyclopteridae | Smooth lumpfish | demersal | 8 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Cyclopteropsis mc alpini</i> | Cyclopteridae | Arctic lumpsucker | demersal | 8 TL | 3.4 | native | Arctic Ocean |
| <i>Dipturus lintea</i> | Rajidae | Sailray | bathydemersal | 123 TL | 3.5 | native | Arctic Ocean |

| Species | Family | Common Name | Habitat | Length (Total Length; cm) | Trophic level | Status | Region |
|----------------------------------|---------------|---------------------------|---------------|---------------------------|---------------|--------|---------------------------|
| <i>Eleginus gracilis</i> | Gadidae | Saffron cod | demersal | 55 TL | 4.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Eleginus nawaga</i> | Gadidae | Navaga | demersal | 42 TL | 4.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Eumesogrammus praecisus</i> | Stichaeidae | Fourline snakeblenny | benthopelagic | 22 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Eumicrotremus andriashevi</i> | Cyclopteridae | Pimpled lumpsucker | demersal | 6 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Eumicrotremus derjugini</i> | Cyclopteridae | Leatherfin lumpsucker | demersal | 13 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Eumicrotremus orbis</i> | Cyclopteridae | Pacific spiny lumpsucker | demersal | 13 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Eumicrotremus spinosus</i> | Cyclopteridae | Atlantic spiny lumpsucker | demersal | 13 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Gadus ogac</i> | Gadidae | Greenland cod | demersal | 77 TL | 3.6 | native | Beaufort Sea/Arctic Ocean |
| <i>Gymnelus andersoni</i> | Zoarcidae | Eelpout | bathydemersal | 14 TL | 3.3 | native | Arctic Ocean |
| <i>Gymnelus hemifasciatus</i> | Zoarcidae | Bigeye unernak | demersal | 13 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Gymnelus viridis</i> | Zoarcidae | Fish doctor | demersal | 56 TL | 3.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Gymnocanthus pistilliger</i> | Cottidae | Threaded sculpin | demersal | 23 TL | 3.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Gymnocanthus tricuspis</i> | Cottidae | Arctic staghorn sculpin | demersal | 30 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Hemilepidotus papilio</i> | Cottidae | Butterfly sculpin | demersal | 37 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Hemilepidotus zapus</i> | Cottidae | Longfin Irish lord | demersal | 13 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |

| Species | Family | Common Name | Habitat | Length (Total Length; cm) | Trophic level | Status | Region |
|----------------------------------|----------------|------------------------|---------------|---------------------------|---------------|--------|---------------------------|
| <i>Hexagrammos stelleri</i> | Hexagrammidae | Whitespotted greenling | demersal | 48 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Hippoglossoides robustus</i> | Pleuronectidae | Bering flounder | demersal | 37 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Hippoglossus stenolepis</i> | Pleuronectidae | Pacific halibut | demersal | 267 TL | 4.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Icelus bicornis</i> | Cottidae | Twohorn sculpin | demersal | 20 TL | 3.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Icelus spatula</i> | Cottidae | Spatulate sculpin | demersal | 14 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Leptagonus decagonus</i> | Agonidae | Atlantic poacher | demersal | 21 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Leptoclinus maculatus</i> | Stichaeidae | Daubed shanny | demersal | 20 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Lethenteron camtschaticum</i> | Petromyzonidae | Arctic lamprey | demersal | 62 TL | 4.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Limanda aspera</i> | Pleuronectidae | Yellowfin sole | demersal | 47 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Liopsetta glacialis</i> | Pleuronectidae | Arctic flounder | demersal | 35 TL | 3.6 | native | Beaufort Sea/Arctic Ocean |
| <i>Liparis bristolensis</i> | Liparidae | Snailfish | demersal | 20 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Liparis fabricii</i> | Liparidae | Gelatinous snailfish | bathydemersal | 20 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Liparis gibbus</i> | Liparidae | Variiegated snailfish | demersal | 52 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Liparis tunicatus</i> | Liparidae | Kelp snailfish | demersal | 16 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Lumpenus fabricii</i> | Stichaeidae | Slender eelblenny | benthopelagic | 36 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |

| Species | Family | Common Name | Habitat | Length (Total Length; cm) | Trophic level | Status | Region |
|----------------------------------|-----------|--------------------|---------------|---------------------------|---------------|---------|---------------------------|
| <i>Lycenchelys kolthoffi</i> | Zoarcidae | Eelpout | bathydemersal | 29 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycenchelys muraena</i> | Zoarcidae | Eelpout | bathydemersal | 28 TL | 3.5 | native | Arctic Ocean |
| <i>Lycodes eudipleurostictus</i> | Zoarcidae | Doubleline eelpout | demersal | 55 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes frigidus</i> | Zoarcidae | Eelpout | bathydemersal | 69 TL | 3.8 | native | Arctic Ocean |
| <i>Lycodes jugoricus</i> | Zoarcidae | Shulupaoluk | demersal | 26 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes luetkenii</i> | Zoarcidae | Eelpout | bathydemersal | 44 TL | 3.4 | native | Arctic Ocean |
| <i>Lycodes mcallisteri</i> | Zoarcidae | Eelpout | bathydemersal | 46 TL | 3.4 | endemic | Arctic Ocean |
| <i>Lycodes mucosus</i> | Zoarcidae | Saddled eelpout | demersal | 25 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes palearis</i> | Zoarcidae | Wattled eelpout | bathydemersal | 51 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes pallidus</i> | Zoarcidae | Pale eelpout | demersal | 26 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes polaris</i> | Zoarcidae | Canadian eelpout | demersal | 25 TL | 3.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes raridens</i> | Zoarcidae | Eelpout | demersal | 31 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes reticulatus</i> | Zoarcidae | Arctic eelpout | bathydemersal | 36 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes rossi</i> | Zoarcidae | Threespot eelpout | demersal | 31 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes sagittarius</i> | Zoarcidae | Archer eelpout | bathydemersal | 34 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes seminudus</i> | Zoarcidae | Longear eelpout | bathydemersal | 52 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |

| Species | Family | Common Name | Habitat | Length (Total Length; cm) | Trophic level | Status | Region |
|-----------------------------------|---------------|---------------------|-----------------|---------------------------|---------------|--------------|---------------------------|
| <i>Lycodes squamiventer</i> | Zoarcidae | Scalebelly eelpout | bathydemersal | 26 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes turneri</i> | Zoarcidae | Polar eelpout | demersal | 25 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Lycodes vahlii</i> | Zoarcidae | Vahl's eelpout | bathydemersal | 52 TL | 3.4 | native | Arctic Ocean |
| <i>Magnisudis atlantica</i> | Paralepididae | Duckbill baracudina | pelagic | 69 TL | 4.1 | native | Arctic Ocean |
| <i>Mallotus villosus</i> | Osmeridae | Capelin | pelagic | 26 TL | 3.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Megalocottus platycephalus</i> | Cottidae | Belligerent sculpin | demersal | 42 TL | 4.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Myoxocephalus jaok</i> | Cottidae | Plain sculpin | demersal | 46 TL | 4.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Myoxocephalus scorpioides</i> | Cottidae | Arctic sculpin | demersal | 22 TL | 3.3 | questionable | Beaufort Sea/Arctic Ocean |
| <i>Myoxocephalus scorpius</i> | Cottidae | Shorthorn sculpin | demersal | 90 TL | 3.9 | native | Beaufort Sea/Arctic Ocean |
| <i>Myoxocephalus stelleri</i> | Cottidae | Steller's sculpin | reef-associated | 49 TL | 3.9 | native | Beaufort Sea/Arctic Ocean |
| <i>Myoxocephalus verrucosus</i> | Cottidae | Sculpin | demersal | 44 TL | 3.8 | native | Beaufort Sea/Arctic Ocean |
| <i>Myxine limosa</i> | Myxinidae | Hagfish | demersal | 51 TL | 3.4 | native | Arctic Ocean |
| <i>Ocella dodecaedron</i> | Agonidae | Bering poacher | demersal | 27 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Oncorhynchus gorbuscha</i> | Salmonidae | Pink salmon | demersal | 76 TL | 4.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Oncorhynchus keta</i> | Salmonidae | Chum salmon | benthopelagic | 111 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Oncorhynchus kisutch</i> | Salmonidae | Coho salmon | demersal | 108 TL | 4.2 | native | Beaufort Sea/Arctic Ocean |

| Species | Family | Common Name | Habitat | Length (Total Length; cm) | Trophic level | Status | Region |
|--|-----------------|-----------------------|---------------|---------------------------|---------------|--------|---------------------------|
| <i>Oncorhynchus mykiss</i> | Salmonidae | Rainbow trout | benthopelagic | 120 TL | 4.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Oncorhynchus nerka</i> | Salmonidae | Sockeye salmon | pelagic | 84 TL | 3.7 | native | Beaufort Sea/Arctic Ocean |
| <i>Oncorhynchus tshawytscha</i> | Salmonidae | Chinook salmon | benthopelagic | 150 TL | 4.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Osmerus mordax dentex</i> | Osmeridae | Arctic rainbow smelt | pelagic | 33 TL | 4.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Petromyzon marinus</i> | Petromyzontidae | Sea lamprey | demersal | 120 TL | 4.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Pholis fasciata</i> | Pholidae | Banded gunnel | demersal | 30 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Pholis gunnellus</i> | Pholidae | Rock gunnel | demersal | 31 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Platichthys flesus</i> | Pleuronectidae | Flounder | demersal | 60 TL | 3.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Platichthys stellatus</i> | Pleuronectidae | Starry flounder | demersal | 91 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Pleuronectes platessa</i> | Pleuronectidae | European plaice | demersal | 122 TL | 3.3 | native | Arctic Ocean |
| <i>Pleuronectes quadrituberculatus</i> | Pleuronectidae | Alaska plaice | demersal | 74 TL | 3.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Podothecus accipenserinus</i> | Agonidae | Sturgeon poacher | demersal | 31 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Pollachius virens</i> | Gadidae | Pollock | demersal | 130 TL | 4.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Pungitius pungitius</i> | Gasterosteidae | Ninespine stickleback | benthopelagic | 9 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Reinhardtius hippoglossoides</i> | Pleuronectidae | Greenland halibut | benthopelagic | 120 TL | 4.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Salangichthys microdon</i> | Salangidae | Japanese icefish | demersal | 12 TL | 3.7 | native | Arctic Ocean |

| Species | Family | Common Name | Habitat | Length (Total Length; cm) | Trophic level | Status | Region |
|--------------------------------------|-------------|-----------------------|---------------|---------------------------|---------------|--------|---------------------------|
| <i>Salmo salar</i> | Salmonidae | Atlantic salmon | benthopelagic | 150 TL | 4.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Salvelinus alpinus</i> | Salmonidae | Charr | benthopelagic | 107 TL | 4.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Salvelinus malma malma</i> | Salmonidae | Dolly varden | benthopelagic | 127 TL | 4.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Somniosus microcephalus</i> | Dalatiidae | Greenland shark | benthopelagic | 730 TL | 4.2 | native | Beaufort Sea/Arctic Ocean |
| <i>Somniosus pacificus</i> | Dalatiidae | Pacific sleeper shark | benthopelagic | 440 TL | 4.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Stichaeus punctatus punctatus</i> | Stichaeidae | Arctic shanny | demersal | 22 TL | 3.1 | native | Beaufort Sea/Arctic Ocean |
| <i>Theragra chalcogramma</i> | Gadidae | Alaska pollock | benthopelagic | 91 TL | 3.5 | native | Beaufort Sea/Arctic Ocean |
| <i>Triglops nybelini</i> | Cottidae | Bigeye sculpin | demersal | 17 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Triglops pingelii</i> | Cottidae | Ribbed sculpin | demersal | 25 TL | 3.4 | native | Beaufort Sea/Arctic Ocean |
| <i>Triglopsis quadricornis</i> | Cottidae | Fourhorn sculpin | demersal | 60 TL | 3.7 | native | Beaufort Sea/Arctic Ocean |
| <i>Ulcina olrikii</i> | Agonidae | Arctic alligatorfish | demersal | 9 TL | 3.3 | native | Beaufort Sea/Arctic Ocean |
| <i>Zeus faber</i> | Zeidae | John dory | benthopelagic | 90 TL | 4.5 | native | Arctic Ocean |

APPENDIX D

REVIEW OF THE EFFECTS OF AIRGUN SOUNDS ON MARINE MAMMALS²

The following subsections review relevant information concerning the potential effects of airguns on marine mammals. Because this review is intended to be of general usefulness, it includes references to types of marine mammals that will not be found in some specific regions.

1. Categories of Noise Effects

The effects of noise on marine mammals are highly variable, and can be categorized as follows (adapted from Richardson et al. 1995):

1. The noise may be too weak to be heard at the location of the animal, i.e., lower than the prevailing ambient noise level, the hearing threshold of the animal at relevant frequencies, or both;
2. The noise may be audible but not strong enough to elicit any overt behavioral response, i.e., the mammal may tolerate it, either without or with some deleterious effects (e.g., masking, stress);
3. The noise may elicit behavioral reactions of variable conspicuousness and variable relevance to the well being of the animal; these can range from subtle effects on respiration or other behaviors (detectable only by statistical analysis) to active avoidance reactions;
4. Upon repeated exposure, animals may exhibit diminishing responsiveness (habituation), or disturbance effects may persist; the latter is most likely with sounds that are highly variable in characteristics, unpredictable in occurrence, and associated with situations that the animal perceives as a threat;
5. Any man-made noise that is strong enough to be heard has the potential to reduce (mask) the ability of marine mammals to hear natural sounds at similar frequencies, including calls from conspecifics, echolocation sounds of odontocetes, and environmental sounds such as surf noise or (at high latitudes) ice noise. However, intermittent airgun or sonar pulses could cause strong masking for only a small proportion of the time, given the short duration of these pulses relative to the inter-pulse intervals;
6. Very strong sounds have the potential to cause temporary or permanent reduction in hearing sensitivity, or other physical or physiological effects. Received sound levels must far exceed the animal's hearing threshold for any temporary threshold shift to occur. Received levels must be even higher for a risk of permanent hearing impairment.

2. Hearing Abilities of Marine Mammals

The hearing abilities of marine mammals are functions of the following (Richardson et al. 1995; Au et al. 2000):

1. Absolute hearing threshold at the frequency in question (the level of sound barely audible in the absence of ambient noise). The "best frequency" is the frequency with the lowest absolute threshold.
2. Critical ratio (the signal-to-noise ratio required to detect a sound at a specific frequency in the presence of background noise around that frequency).

² By **W. John Richardson** and **Valerie D. Moulton**, with subsequent updates (to November 2009) by WJR and VDM plus **Patrick Abgrall**, **William E. Cross**, **Meike Holst**, and **Mari A. Smultea**, all of LGL Ltd., environmental research associates

3. The ability to determine sound direction at the frequencies under consideration.
4. The ability to discriminate among sounds of different frequencies and intensities.

Marine mammals rely heavily on the use of underwater sounds to communicate and to gain information about their surroundings. Experiments and monitoring studies also show that they hear and may react to many man-made sounds including sounds made during seismic exploration (Richardson et al. 1995; Gordon et al. 2004; Nowacek et al. 2007; Tyack 2008).

2.1 Toothed Whales (*Odontocetes*)

Hearing abilities of some toothed whales (odontocetes) have been studied in detail (reviewed in Chapter 8 of Richardson et al. [1995] and in Au et al. [2000]). Hearing sensitivity of several species has been determined as a function of frequency. The small to moderate-sized toothed whales whose hearing has been studied have relatively poor hearing sensitivity at frequencies below 1 kHz, but extremely good sensitivity at, and above, several kHz. There are very few data on the absolute hearing thresholds of most of the larger, deep-diving toothed whales, such as the sperm and beaked whales. However, Cook et al. (2006) found that a stranded juvenile Gervais' beaked whale showed evoked potentials from 5 kHz up to 80 kHz (the entire frequency range that was tested), with best sensitivity at 40–80 kHz. An adult Gervais' beaked whale had a similar upper cutoff frequency (80–90 kHz; Finneran et al. 2009).

Most of the odontocete species have been classified as belonging to the “mid-frequency” (MF) hearing group, and the MF odontocetes (collectively) have functional hearing from about 150 Hz to 160 kHz (Southall et al. 2007). However, individual species may not have quite so broad a functional frequency range. Very strong sounds at frequencies slightly outside the functional range may also be detectable. The remaining odontocetes—the porpoises, river dolphins, and members of the genera *Cephalorhynchus* and *Kogia*—are distinguished as the “high frequency” (HF) hearing group. They have functional hearing from about 200 Hz to 180 kHz (Southall et al. 2007).

Airguns produce a small proportion of their sound at mid- and high-frequencies, although at progressively lower levels with increasing frequency. In general, most of the energy in the sound pulses emitted by airgun arrays is at low frequencies; strongest spectrum levels are below 200 Hz, with considerably lower spectrum levels above 1000 Hz, and smaller amounts of energy emitted up to ~150 kHz (Goold and Fish 1998; Sodal 1999; Goold and Coates 2006; Potter et al. 2007).

Despite the relatively poor sensitivity of small odontocetes at the low frequencies that contribute most of the energy in pulses of sound from airgun arrays, airgun sounds are sufficiently strong, and contain sufficient mid- and high-frequency energy, that their received levels sometimes remain above the hearing thresholds of odontocetes at distances out to several tens of kilometers (Richardson and Würsig 1997). There is no evidence that most small odontocetes react to airgun pulses at such long distances. However, beluga whales do seem quite responsive at intermediate distances (10–20 km) where sound levels are well above the ambient noise level (see below).

In summary, even though odontocete hearing is relatively insensitive to the predominant low frequencies produced by airguns, sounds from airgun arrays are audible to odontocetes, sometimes to distances of 10s of kilometers.

2.2 Baleen Whales (*Mysticetes*)

The hearing abilities of baleen whales (mysticetes) have not been studied directly. Behavioral and anatomical evidence indicates that they hear well at frequencies below 1 kHz (Richardson et al. 1995; Ketten 2000). Frankel (2005) noted that gray whales reacted to a 21–25 kHz whale-finding sonar. Some baleen whales react to pinger sounds up to 28 kHz, but not to pingers or sonars emitting sounds at 36 kHz

or above (Watkins 1986). In addition, baleen whales produce sounds at frequencies up to 8 kHz and, for humpbacks, with components to >24 kHz (Au et al. 2006). The anatomy of the baleen whale inner ear seems to be well adapted for detection of low-frequency sounds (Ketten 1991, 1992, 1994, 2000; Parks et al. 2007b). Although humpbacks and minke whales (Berta et al. 2009) may have some auditory sensitivity to frequencies above 22 kHz, for baleen whales as a group, the functional hearing range is thought to be about 7 Hz to 22 kHz and they are said to constitute the “low-frequency” (LF) hearing group (Southall et al. 2007). The absolute sound levels that they can detect below 1 kHz are probably limited by increasing levels of natural ambient noise at decreasing frequencies (Clark and Ellison 2004). Ambient noise levels are higher at low frequencies than at mid frequencies. At frequencies below 1 kHz, natural ambient levels tend to increase with decreasing frequency.

The hearing systems of baleen whales are undoubtedly more sensitive to low-frequency sounds than are the ears of the small toothed whales that have been studied directly. Thus, baleen whales are likely to hear airgun pulses farther away than can small toothed whales and, at closer distances, airgun sounds may seem more prominent to baleen than to toothed whales. However, baleen whales have commonly been seen well within the distances where seismic (or other source) sounds would be detectable and often show no overt reaction to those sounds. Behavioral responses by baleen whales to seismic pulses have been documented, but received levels of pulsed sounds necessary to elicit behavioral reactions are typically well above the minimum levels that the whales are assumed to detect (see below).

2.3 Seals and Sea Lions (Pinnipeds)

Underwater audiograms have been obtained using behavioral methods for three species of phocinid seals, two species of monachid seals, two species of otariids, and the walrus (reviewed in Richardson et al. 1995: 211ff; Kastak and Schusterman 1998, 1999; Kastelein et al. 2002, 2009). The functional hearing range for pinnipeds in water is considered to extend from 75 Hz to 75 kHz (Southall et al. 2007), although some individual species—especially the eared seals—do not have that broad an auditory range (Richardson et al. 1995). In comparison with odontocetes, pinnipeds tend to have lower best frequencies, lower high-frequency cutoffs, better auditory sensitivity at low frequencies, and poorer sensitivity at the best frequency.

At least some of the phocid seals have better sensitivity at low frequencies (≤ 1 kHz) than do odontocetes. Below 30–50 kHz, the hearing thresholds of most species tested are essentially flat down to ~ 1 kHz, and range between 60 and 85 dB re 1 μ Pa. Measurements for harbor seals indicate that, below 1 kHz, their thresholds under quiet background conditions deteriorate gradually with decreasing frequency to ~ 75 dB re 1 μ Pa at 125 Hz (Kastelein et al. 2009).

For the otariid (eared) seals, the high frequency cutoff is lower than for phocinids, and sensitivity at low frequencies (e.g., 100 Hz) is poorer than for seals (harbor seal).

2.4 Manatees and Dugong (Sirenians)

The West Indian manatee can apparently detect sounds and low-frequency vibrations from 15 Hz to 46 kHz, based on a study involving behavioral testing methods (Gerstein et al. 1999, 2004). A more recent study found that, in one Florida manatee, auditory sensitivity extended up to 90.5 kHz (Bauer et al. 2009). Thus, manatees may hear, or at least detect, sounds in the low-frequency range where most seismic energy is released. It is possible that they are able to feel these low-frequency sounds using vibrotactile receptors or because of resonance in body cavities or bone conduction.

Based on measurements of evoked potentials, manatee hearing is apparently best around 1–1.5 kHz (Bullock et al. 1982). However, behavioral tests suggest that best sensitivities are at 6–20 kHz (Gerstein

et al. 1999) or 8–32 kHz (Bauer et al. 2009). The ability to detect high frequencies may be an adaptation to shallow water, where the propagation of low frequency sound is limited (Gerstein et al. 1999, 2004).

2.5 Sea Otter and Polar Bear

No data are available on the hearing abilities of sea otters (Ketten 1998), although the in-air vocalizations of sea otters have most of their energy concentrated at 3–5 kHz (McShane et al. 1995; Thomson and Richardson 1995). Sea otter vocalizations are considered to be most suitable for short-range communication among individuals (McShane et al. 1995). However, Ghoual et al. (2009) noted that the in-air “screams” of sea otters are loud signals (source level of 93–118 dB re 20 μPa_{pk}) that may be used over larger distances; screams have a frequency of maximum energy ranging from 2 to 8 kHz. In-air audiograms for two river otters indicate that this related species has its best hearing sensitivity at the relatively high frequency of 16 kHz, with some sensitivity from about 460 Hz to 33 kHz (Gunn 1988). However, these data apply to a different species of otter, and to in-air rather than underwater hearing.

Data on the specific hearing capabilities of polar bears are limited. A recent study of the in-air hearing of polar bears applied the auditory evoked potential method while tone pips were played to anesthetized bears (Nachtigall et al. 2007). Hearing was tested in $\frac{1}{2}$ octave steps from 1 to 22.5 kHz, and best hearing sensitivity was found between 11.2 and 22.5 kHz. Although low-frequency hearing was not studied, the data suggested that medium- and some high-frequency sounds may be audible to polar bears. However, polar bears’ usual behavior (e.g., remaining on the ice, at the water surface, or on land) reduces or avoids exposure to underwater sounds.

3. Characteristics of Airgun Sounds

Airguns function by venting high-pressure air into the water. The pressure signature of an individual airgun consists of a sharp rise and then fall in pressure, followed by several positive and negative pressure excursions caused by oscillation of the resulting air bubble. The sizes, arrangement, and firing times of the individual airguns in an array are designed and synchronized to suppress the pressure oscillations subsequent to the first cycle. The resulting downward-directed pulse has a duration of only 10–20 ms, with only one strong positive and one strong negative peak pressure (Caldwell and Dragoset 2000). Most energy emitted from airguns is at relatively low frequencies. For example, typical high-energy airgun arrays emit most energy at 10–120 Hz. However, the pulses contain significant energy up to 500–1000 Hz and some energy at higher frequencies (Goold and Fish 1998; Potter et al. 2007). Studies in the Gulf of Mexico have shown that the horizontally-propagating sound can contain significant energy above the frequencies that airgun arrays are designed to emit (DeRuiter et al. 2006; Madsen et al. 2006; Tyack et al. 2006a). Energy at frequencies up to 150 kHz was found in tests of single 60-in³ and 250-in³ airguns (Goold and Coates 2006). Nonetheless, the predominant energy is at low frequencies.

The pulsed sounds associated with seismic exploration have higher peak levels than other industrial sounds (except those from explosions) to which whales and other marine mammals are routinely exposed. The nominal source levels of the 2- to 36-airgun arrays used by Lamont-Doherty Earth Observatory (L-DEO) from the R/V *Maurice Ewing* (now retired) and R/V *Marcus G. Langseth* (36 airguns) are 236–265 dB re 1 $\mu\text{Pa}_{\text{p-p}}$. These are the nominal source levels applicable to downward propagation. The effective source levels for horizontal propagation are lower than those for downward propagation when the source consists of numerous airguns spaced apart from one another. Explosions are the only man-made sources with effective source levels as high as (or higher than) a large array of airguns. However, high-power sonars can have source pressure levels as high as a small array of airguns, and signal duration can be longer for a sonar than for an airgun array, making the source energy levels of some sonars more comparable to those of airgun arrays.

Several important mitigating factors need to be kept in mind. (1) Airgun arrays produce intermittent sounds, involving emission of a strong sound pulse for a small fraction of a second followed by several seconds of near silence. In contrast, some other sources produce sounds with lower peak levels, but their sounds are continuous or discontinuous but continuing for longer durations than seismic pulses. (2) Airgun arrays are designed to transmit strong sounds downward through the seafloor, and the amount of sound transmitted in near-horizontal directions is considerably reduced. Nonetheless, they also emit sounds that travel horizontally toward non-target areas. (3) An airgun array is a distributed source, not a point source. The nominal source level is an estimate of the sound that would be measured from a theoretical point source emitting the same total energy as the airgun array. That figure is useful in calculating the expected received levels in the far field, i.e., at moderate and long distances, but not in the near field. Because the airgun array is not a single point source, there is no one location within the near field (or anywhere else) where the received level is as high as the nominal source level.

The strengths of airgun pulses can be measured in different ways, and it is important to know which method is being used when interpreting quoted source or received levels. Geophysicists usually quote peak-to-peak (p-p) levels, in bar-meters or (less often) dB re 1 $\mu\text{Pa} \cdot \text{m}$. The peak (= zero-to-peak, or 0-p) level for the same pulse is typically ~ 6 dB less. In the biological literature, levels of received airgun pulses are often described based on the “average” or “root-mean-square” (rms) level, where the average is calculated over the duration of the pulse. The rms value for a given airgun pulse is typically ~ 10 dB lower than the peak level, and 16 dB lower than the peak-to-peak value (Greene 1997; McCauley et al. 1998, 2000a). A fourth measure that is increasingly used is the energy, or Sound Exposure Level (SEL), in dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$. Because the pulses, even when stretched by propagation effects (see below), are usually < 1 s in duration, the numerical value of the energy is usually lower than the rms pressure level. However, the units are different.³ Because the level of a given pulse will differ substantially depending on which of these measures is being applied, it is important to be aware which measure is in use when interpreting any quoted pulse level. In the past, the U.S. National Marine Fisheries Service (NMFS) has commonly referred to rms levels when discussing levels of pulsed sounds that might “harass” marine mammals.

Seismic sound pulses received at any given point will arrive via a direct path, indirect paths that include reflection from the sea surface and bottom, and often indirect paths including segments through the bottom sediments. Sounds propagating via indirect paths travel longer distances and often arrive later than sounds arriving via a direct path. (However, sound traveling in the bottom may travel faster than that in the water, and thus may, in some situations, arrive slightly earlier than the direct arrival despite traveling a greater distance.) These variations in travel time have the effect of lengthening the duration of the received pulse, or may cause two or more received pulses from a single emitted pulse. Near the source, the predominant part of a seismic pulse is ~ 10 – 20 ms in duration. In comparison, the pulse duration as received at long horizontal distances can be much greater. For example, for one airgun array operating in

³ The rms value for a given airgun array pulse, as measured at a horizontal distance on the order of 0.1 km to 1–10 km in the units dB re 1 μPa , usually averages 10–15 dB higher than the SEL value for the same pulse measured in dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$ (e.g., Greene 1997). However, there is considerable variation, and the difference tends to be larger close to the airgun array, and less at long distances (Blackwell et al. 2007; MacGillivray and Hannay 2007a,b). In some cases, generally at longer distances, pulses are “stretched” by propagation effects to the extent that the rms and SEL values (in the respective units mentioned above) become very similar (e.g., MacGillivray and Hannay 2007a,b).

the Beaufort Sea, pulse duration was ~300 ms at a distance of 8 km, 500 ms at 20 km, and 850 ms at 73 km (Greene and Richardson 1988).

The rms level for a given pulse (when measured over the duration of that pulse) depends on the extent to which propagation effects have “stretched” the duration of the pulse by the time it reaches the receiver (e.g., Madsen 2005). As a result, the rms values for various received pulses are not perfectly correlated with the SEL (energy) values for the same pulses. There is increasing evidence that biological effects are more directly related to the received energy (e.g., to SEL) than to the rms values averaged over pulse duration (Southall et al. 2007).

Another important aspect of sound propagation is that received levels of low-frequency underwater sounds diminish close to the surface because of pressure-release and interference phenomena that occur at and near the surface (Urlick 1983; Richardson et al. 1995; Potter et al. 2007). Paired measurements of received airgun sounds at depths of 3 vs. 9 or 18 m have shown that received levels are typically several decibels lower at 3 m (Greene and Richardson 1988). For a mammal whose auditory organs are within 0.5 or 1 m of the surface, the received level of the predominant low-frequency components of the airgun pulses would be further reduced. In deep water, the received levels at deep depths can be considerably higher than those at relatively shallow (e.g., 18 m) depths and the same horizontal distance from the airguns (Tolstoy et al. 2004a,b).

Pulses of underwater sound from open-water seismic exploration are often detected 50–100 km from the source location, even during operations in nearshore waters (Greene and Richardson 1988; Burgess and Greene 1999). At those distances, the received levels are usually low, <120 dB re 1 μ Pa on an approximate rms basis. However, faint seismic pulses are sometimes detectable at even greater ranges (e.g., Bowles et al. 1994; Fox et al. 2002). In fact, low-frequency airgun signals sometimes can be detected thousands of kilometers from their source. For example, sound from seismic surveys conducted offshore of Nova Scotia, the coast of western Africa, and northeast of Brazil were reported as a dominant feature of the underwater noise field recorded along the mid-Atlantic ridge (Nieukirk et al. 2004).

4. Masking Effects of Airgun Sounds

Masking is the obscuring of sounds of interest by interfering sounds, generally at similar frequencies (Richardson et al. 1995). Introduced underwater sound will, through masking, reduce the effective communication distance of a marine mammal species if the frequency of the source is close to that used as a signal by the marine mammal, and if the anthropogenic sound is present for a significant fraction of the time (Richardson et al. 1995). If little or no overlap occurs between the introduced sound and the frequencies used by the species, communication is not expected to be disrupted. Also, if the introduced sound is present only infrequently, communication is not expected to be disrupted much if at all. The duty cycle of airguns is low; the airgun sounds are pulsed, with relatively quiet periods between pulses. In most situations, strong airgun sound will only be received for a brief period (<1 s), with these sound pulses being separated by at least several seconds of relative silence, and longer in the case of deep-penetration surveys or refraction surveys. A single airgun array might cause appreciable masking in only one situation: When propagation conditions are such that sound from each airgun pulse reverberates strongly and persists for much or all of the interval up to the next airgun pulse (e.g., Simard et al. 2005; Clark and Gagnon 2006). Situations with prolonged strong reverberation are infrequent, in our experience. However, it is common for reverberation to cause some lesser degree of elevation of the background level between airgun pulses (e.g., Guerra et al. 2009), and this weaker reverberation presumably reduces the detection range of calls and other natural sounds to some degree.

Although masking effects of pulsed sounds on marine mammal calls and other natural sounds are expected to be limited, there are few specific studies on this. Some whales continue calling in the presence of seismic pulses and whale calls often can be heard between the seismic pulses (e.g., Richardson et al. 1986; McDonald et al. 1995; Greene et al. 1999a,b; Nieu Kirk et al. 2004; Smultea et al. 2004; Holst et al. 2005a,b, 2006; Dunn et al. 2009). However, there is one recent summary report indicating that calling fin whales distributed in one part of the North Atlantic went silent for an extended period starting soon after the onset of a seismic survey in the area (Clark and Gagnon 2006). It is not clear from that preliminary paper whether the whales ceased calling because of masking, or whether this was a behavioral response not directly involving masking. Also, bowhead whales in the Beaufort Sea may decrease their call rates in response to seismic operations, although movement out of the area might also have contributed to the lower call detection rate (Blackwell et al. 2009a,b). In contrast, Di Iorio and Clark (2009) found evidence of *increased* calling by blue whales during operations by a lower-energy seismic source—a sparker.

Among the odontocetes, there has been one report that sperm whales ceased calling when exposed to pulses from a very distant seismic ship (Bowles et al. 1994). However, more recent studies of sperm whales found that they continued calling in the presence of seismic pulses (Madsen et al. 2002; Tyack et al. 2003; Smultea et al. 2004; Holst et al. 2006; Jochens et al. 2008). Madsen et al. (2006) noted that airgun sounds would not be expected to mask sperm whale calls given the intermittent nature of airgun pulses. Dolphins and porpoises are also commonly heard calling while airguns are operating (Gordon et al. 2004; Smultea et al. 2004; Holst et al. 2005a,b; Potter et al. 2007). Masking effects of seismic pulses are expected to be negligible in the case of the smaller odontocetes, given the intermittent nature of seismic pulses plus the fact that sounds important to them are predominantly at much higher frequencies than are the dominant components of airgun sounds.

Pinnipeds, sirenians and sea otters have best hearing sensitivity and/or produce most of their sounds at frequencies higher than the dominant components of airgun sound, but there is some overlap in the frequencies of the airgun pulses and the calls. However, the intermittent nature of airgun pulses presumably reduces the potential for masking.

A few cetaceans are known to increase the source levels of their calls in the presence of elevated sound levels, shift their peak frequencies in response to strong sound signals, or otherwise modify their vocal behavior in response to increased noise (Dahlheim 1987; Au 1993; reviewed in Richardson et al. 1995:233ff, 364ff; Lesage et al. 1999; Terhune 1999; Nieu Kirk et al. 2005; Scheifele et al. 2005; Parks et al. 2007a, 2009; Di Iorio and Clark 2009; Hanser et al. 2009). It is not known how often these types of responses occur upon exposure to airgun sounds. However, blue whales in the St. Lawrence Estuary significantly increased their call rates during sparker operations (Di Iorio and Clark 2009). The sparker, used to obtain seismic reflection data, emitted frequencies of 30–450 Hz with a relatively low source level of 193 dB re 1 $\mu\text{Pa}_{\text{pk-pk}}$. If cetaceans exposed to airgun sounds sometimes respond by changing their vocal behavior, this adaptation, along with directional hearing and preadaptation to tolerate some masking by natural sounds (Richardson et al. 1995), would all reduce the importance of masking by seismic pulses.

5. Disturbance by Seismic Surveys

Disturbance includes a variety of effects, including subtle to conspicuous changes in behavior, movement, and displacement. In the terminology of the 1994 amendments to the U.S. Marine Mammal Protection Act (MMPA), seismic noise could cause “Level B” harassment of certain marine mammals. Level B harassment is defined as “...disruption of behavioral patterns, including, but not limited to, migration, breathing, nursing, breeding, feeding, or sheltering.”

There has been debate regarding how substantial a change in behavior or mammal activity is required before the animal should be deemed to be “taken by Level B harassment”. NMFS has stated that

“...a simple change in a marine mammal’s actions does not always rise to the level of disruption of its behavioral patterns. ... If the only reaction to the [human] activity on the part of the marine mammal is within the normal repertoire of actions that are required to carry out that behavioral pattern, NMFS considers [the human] activity not to have caused a disruption of the behavioral pattern, provided the animal’s reaction is not otherwise significant enough to be considered disruptive due to length or severity. Therefore, for example, a short-term change in breathing rates or a somewhat shortened or lengthened dive sequence that are within the animal’s normal range and that do not have any biological significance (i.e., do not disrupt the animal’s overall behavioral pattern of breathing under the circumstances), do not rise to a level requiring a small take authorization.” (NMFS 2001, p. 9293).

Based on this guidance from NMFS, and on NRC (2005), simple exposure to sound, or brief reactions that do not disrupt behavioral patterns in a potentially significant manner, do not constitute harassment or “taking”. In this analysis, we interpret “potentially significant” to mean in a manner that might have deleterious effects on the well-being of individual marine mammals or their populations.

Even with this guidance, there are difficulties in defining what marine mammals should be counted as “taken by harassment”. Available detailed data on reactions of marine mammals to airgun sounds (and other anthropogenic sounds) are limited to relatively few species and situations (see Richardson et al. 1995; Gordon et al. 2004; Nowacek et al. 2007; Southall et al. 2007). Behavioral reactions of marine mammals to sound are difficult to predict in the absence of site- and context-specific data. Reactions to sound, if any, depend on species, state of maturity, experience, current activity, reproductive state, time of day, and many other factors (Richardson et al. 1995; Wartzok et al. 2004; Southall et al. 2007; Weilgart 2007). If a marine mammal reacts to an underwater sound by changing its behavior or moving a small distance, the impacts of the change are unlikely to be significant to the individual, let alone the stock or population. However, if a sound source displaces marine mammals from an important feeding or breeding area for a prolonged period, impacts on individuals and populations could be significant (e.g., Lusseau and Bejder 2007; Weilgart 2007). Also, various authors have noted that some marine mammals that show no obvious avoidance or behavioral changes may still be adversely affected by noise (Brodie 1981; Richardson et al. 1995:317ff; Romano et al. 2004; Weilgart 2007; Wright et al. 2009). For example, some research suggests that animals in poor condition or in an already stressed state may not react as strongly to human disturbance as would more robust animals (e.g., Beale and Monaghan 2004).

Studies of the effects of seismic surveys have focused almost exclusively on the effects on individual species or related groups of species, with little scientific or regulatory attention being given to broader community-level issues. Parente et al. (2007) suggested that the diversity of cetaceans near the Brazil coast was reduced during years with seismic surveys. However, a preliminary account of a more recent analysis suggests that the trend did not persist when additional years were considered (Britto and Silva Barreto 2009).

Given the many uncertainties in predicting the quantity and types of impacts of sound on marine mammals, it is common practice to estimate how many mammals would be present within a particular distance of human activities and/or exposed to a particular level of anthropogenic sound. In most cases, this approach likely overestimates the numbers of marine mammals that would be affected in some biologically important manner. One of the reasons for this is that the selected distances/isopleths are based on limited studies indicating that some animals exhibited short-term reactions at this distance or

sound level, whereas the calculation assumes that all animals exposed to this level would react in a biologically significant manner.

The definitions of “taking” in the U.S. MMPA, and its applicability to various activities, were slightly altered in November 2003 for military and federal scientific research activities. Also, NMFS is proposing to replace current Level A and B harassment criteria with guidelines based on exposure characteristics that are specific to particular groups of mammal species and to particular sound types (NMFS 2005). Recently, a committee of specialists on noise impact issues has proposed new science-based impact criteria (Southall et al. 2007). Thus, for projects subject to U.S. jurisdiction, changes in procedures may be required in the near future.

The sound criteria used to estimate how many marine mammals might be disturbed to some biologically significant degree by seismic survey activities are primarily based on behavioral observations of a few species. Detailed studies have been done on humpback, gray, bowhead, and sperm whales, and on ringed seals. Less detailed data are available for some other species of baleen whales and small toothed whales, but for many species there are no data on responses to marine seismic surveys.

5.1 Baleen Whales

Baleen whales generally tend to avoid operating airguns, but avoidance radii are quite variable among species, locations, whale activities, oceanographic conditions affecting sound propagation, etc. (reviewed in Richardson et al. 1995; Gordon et al. 2004). Whales are often reported to show no overt reactions to pulses from large arrays of airguns at distances beyond a few kilometers, even though the airgun pulses remain well above ambient noise levels out to much longer distances. However, baleen whales exposed to strong sound pulses from airguns often react by deviating from their normal migration route and/or interrupting their feeding and moving away. Some of the major studies and reviews on this topic are Malme et al. (1984, 1985, 1988); Richardson et al. (1986, 1995, 1999); Ljungblad et al. (1988); Richardson and Malme (1993); McCauley et al. (1998, 2000a,b); Miller et al. (1999, 2005); Gordon et al. (2004); Moulton and Miller (2005); Stone and Tasker (2006); Johnson et al. (2007); Nowacek et al. (2007) and Weir (2008a). Although baleen whales often show only slight overt responses to operating airgun arrays (Stone and Tasker 2006; Weir 2008a), strong avoidance reactions by several species of mysticetes have been observed at ranges up to 6–8 km and occasionally as far as 20–30 km from the source vessel when large arrays of airguns were used. Experiments with a single airgun showed that bowhead, humpback and gray whales all showed localized avoidance to a single airgun of 20–100 in³ (Malme et al. 1984, 1985, 1986, 1988; Richardson et al. 1986; McCauley et al. 1998, 2000a,b).

Studies of gray, bowhead, and humpback whales have shown that seismic pulses with received levels of 160–170 dB re 1 $\mu\text{Pa}_{\text{rms}}$ seem to cause obvious avoidance behavior in a substantial portion of the animals exposed (Richardson et al. 1995). In many areas, seismic pulses from large arrays of airguns diminish to those levels at distances ranging from 4–15 km from the source. More recent studies have shown that some species of baleen whales (bowheads and humpbacks in particular) at times show strong avoidance at received levels lower than 160–170 dB re 1 $\mu\text{Pa}_{\text{rms}}$. The largest avoidance radii involved migrating bowhead whales, which avoided an operating seismic vessel by 20–30 km (Miller et al. 1999; Richardson et al. 1999). In the cases of migrating bowhead (and gray) whales, the observed changes in behavior appeared to be of little or no biological consequence to the animals—they simply avoided the sound source by displacing their migration route to varying degrees, but within the natural boundaries of the migration corridors (Malme et al. 1984; Malme and Miles 1985; Richardson et al. 1995). Feeding bowhead whales, in contrast to migrating whales, show much smaller avoidance distances (Miller et al.

2005; Harris et al. 2007), presumably because moving away from a food concentration has greater cost to the whales than does a course deviation during migration.

The following subsections provide more details on the documented responses of particular species and groups of baleen whales to marine seismic operations.

Humpback Whales.—Responses of humpback whales to seismic surveys have been studied during migration, on the summer feeding grounds, and on Angolan winter breeding grounds; there has also been discussion of effects on the Brazilian wintering grounds. McCauley et al. (1998, 2000a) studied the responses of migrating humpback whales off Western Australia to a full-scale seismic survey with a 16-airgun 2678-in³ array, and to a single 20 in³ airgun with a (horizontal) source level of 227 dB re 1 $\mu\text{Pa} \cdot \text{m}_{\text{p-p}}$. They found that the overall distribution of humpbacks migrating through their study area was unaffected by the full-scale seismic program, although localized displacement varied with pod composition, behavior, and received sound levels. Observations were made from the seismic vessel, from which the maximum viewing distance was listed as 14 km. Avoidance reactions (course and speed changes) began at 4–5 km for traveling pods, with the closest point of approach (CPA) being 3–4 km at an estimated received level of 157–164 dB re 1 $\mu\text{Pa}_{\text{rms}}$ (McCauley et al. 1998, 2000a). A greater stand-off range of 7–12 km was observed for more sensitive resting pods (cow-calf pairs; McCauley et al. 1998, 2000a). The mean received level for initial avoidance of an approaching airgun was 140 dB re 1 $\mu\text{Pa}_{\text{rms}}$ for humpback pods containing females, and at the mean CPA distance the received level was 143 dB re 1 $\mu\text{Pa}_{\text{rms}}$. One startle response was reported at 112 dB re 1 $\mu\text{Pa}_{\text{rms}}$. The initial avoidance response generally occurred at distances of 5–8 km from the airgun array and 2 km from the single airgun. However, some individual humpback whales, especially males, approached within distances of 100–400 m, where the maximum received level was 179 dB re 1 $\mu\text{Pa}_{\text{rms}}$. The McCauley et al. (1998, 2000a,b) studies show evidence of greater avoidance of seismic airgun sounds by pods with females than by other pods during humpback migration off Western Australia.

Humpback whales on their summer feeding grounds in southeast Alaska did not exhibit persistent avoidance when exposed to seismic pulses from a 1.64-L (100 in³) airgun (Malme et al. 1985). Some humpbacks seemed “startled” at received levels of 150–169 dB re 1 μPa . Malme et al. (1985) concluded that there was no clear evidence of avoidance, despite the possibility of subtle effects, at received levels up to 172 re 1 μPa on an approximate rms basis.

Among wintering humpback whales off Angola ($n = 52$ useable groups), there were no significant differences in encounter rates (sightings/hr) when a 24-airgun array (3147 in³ or 5085 in³) was operating vs. silent (Weir 2008a). There was also no significant difference in the mean CPA (closest observed point of approach) distance of the humpback sightings when airguns were on vs. off (3050 m vs. 2700 m, respectively).

It has been suggested that South Atlantic humpback whales wintering off Brazil may be displaced or even strand upon exposure to seismic surveys (Engel et al. 2004). The evidence for this was circumstantial and subject to alternative explanations (IAGC 2004). Also, the evidence was not consistent with subsequent results from the same area of Brazil (Parente et al. 2006), or with direct studies of humpbacks exposed to seismic surveys in other areas and seasons (see above). After allowance for data from subsequent years, there was “no observable direct correlation” between strandings and seismic surveys (IWC 2007, p. 236).

Bowhead Whales.—Responsiveness of bowhead whales to seismic surveys can be quite variable depending on their activity (feeding vs. migrating). Bowhead whales on their summer feeding grounds in the Canadian Beaufort Sea showed no obvious reactions to pulses from seismic vessels at distances of 6–99 km and received sound levels of 107–158 dB on an approximate rms basis (Richardson et al. 1986);

their general activities were indistinguishable from those of a control group. However, subtle but statistically significant changes in surfacing–respiration–dive cycles were evident upon statistical analysis. Bowheads usually did show strong avoidance responses when seismic vessels approached within a few kilometers (~3–7 km) and when received levels of airgun sounds were 152–178 dB (Richardson et al. 1986, 1995; Ljungblad et al. 1988; Miller et al. 2005). They also moved away when a single airgun fired nearby (Richardson et al. 1986; Ljungblad et al. 1988). In one case, bowheads engaged in near-bottom feeding began to turn away from a 30-airgun array with a source level of 248 dB re 1 $\mu\text{Pa} \cdot \text{m}$ at a distance of 7.5 km, and swam away when it came within ~2 km; some whales continued feeding until the vessel was 3 km away (Richardson et al. 1986). This work and subsequent summer studies in the same region by Miller et al. (2005) and Harris et al. (2007) showed that many feeding bowhead whales tend to tolerate higher sound levels than migrating bowhead whales (see below) before showing an overt change in behavior. On the summer feeding grounds, bowhead whales are often seen from the operating seismic ship, though average sighting distances tend to be larger when the airguns are operating. Similarly, preliminary analyses of recent data from the Alaskan Beaufort Sea indicate that bowheads feeding there during late summer and autumn also did not display large-scale distributional changes in relation to seismic operations (Christie et al. 2009; Koski et al. 2009). However, some individual bowheads apparently begin to react at distances a few kilometers away, beyond the distance at which observers on the ship can sight bowheads (Richardson et al. 1986; Citta et al. 2007). The feeding whales may be affected by the sounds, but the need to feed may reduce the tendency to move away until the airguns are within a few kilometers.

Migrating bowhead whales in the Alaskan Beaufort Sea seem more responsive to noise pulses from a distant seismic vessel than are summering bowheads. Bowhead whales migrating west across the Alaskan Beaufort Sea in autumn are unusually responsive, with substantial avoidance occurring out to distances of 20–30 km from a medium-sized airgun source at received sound levels of around 120–130 dB re 1 $\mu\text{Pa}_{\text{rms}}$ (Miller et al. 1999; Richardson et al. 1999; see also Manly et al. 2007). Those results came from 1996–98, when a partially-controlled study of the effect of Ocean Bottom Cable (OBC) seismic surveys on westward-migrating bowheads was conducted in late summer and autumn in the Alaskan Beaufort Sea. At times when the airguns were not active, many bowheads moved into the area close to the inactive seismic vessel. Avoidance of the area of seismic operations did not persist beyond 12–24 h after seismic shooting stopped. Preliminary analysis of recent data on traveling bowheads in the Alaskan Beaufort Sea also showed a stronger tendency to avoid operating airguns than was evident for feeding bowheads (Christie et al. 2009; Koski et al. 2009).

Bowhead whale calls detected in the presence and absence of airgun sounds have been studied extensively in the Beaufort Sea. Early work on the summering grounds in the Canadian Beaufort Sea showed that bowheads continue to produce calls of the usual types when exposed to airgun sounds, although numbers of calls detected may be somewhat lower in the presence of airgun pulses (Richardson et al. 1986). Studies during autumn in the Alaskan Beaufort Sea, one in 1996–1998 and another in 2007–2008, have shown that numbers of calls detected are significantly lower in the presence than in the absence of airgun pulses (Greene et al. 1999a,b; Blackwell et al. 2009a,b; Koski et al. 2009; see also Nations et al. 2009). This decrease could have resulted from movement of the whales away from the area of the seismic survey or a reduction in calling behavior, or a combination of the two. However, concurrent aerial surveys showed that there was strong avoidance of the operating airguns during the 1996–98 study, when most of the whales appeared to be migrating (Miller et al. 1999; Richardson et al. 1999). In contrast, aerial surveys during the 2007–08 study showed less consistent avoidance by the bowheads, many of which appeared to be feeding (Christie et al. 2009; Koski et al. 2009). The reduction in call detection rates during periods of airgun operation may have been more dependent on actual avoidance

during the 1996–98 study and more dependent on reduced calling behavior during the 2007–08 study, but further analysis of the recent data is ongoing.

There are no data on reactions of bowhead whales to seismic surveys in winter or spring.

Gray Whales.—Malme et al. (1986, 1988) studied the responses of feeding eastern gray whales to pulses from a single 100-in³ airgun off St. Lawrence Island in the northern Bering Sea. They estimated, based on small sample sizes, that 50% of feeding gray whales stopped feeding at an average received pressure level of 173 dB re 1 μ Pa on an (approximate) rms basis, and that 10% of feeding whales interrupted feeding at received levels of 163 dB re 1 μ Pa_{rms}. Malme et al. (1986) estimated that an average pressure level of 173 dB occurred at a range of 2.6–2.8 km from an airgun array with a source level of 250 dB re 1 μ Pa_{peak} in the northern Bering Sea. These findings were generally consistent with the results of studies conducted on larger numbers of gray whales migrating off California (Malme et al. 1984; Malme and Miles 1985) and western Pacific gray whales feeding off Sakhalin, Russia (Würsig et al. 1999; Gailey et al. 2007; Johnson et al. 2007; Yazvenko et al. 2007a,b), along with a few data on gray whales off British Columbia (Bain and Williams 2006).

Malme and Miles (1985) concluded that, during migration off California, gray whales showed changes in swimming pattern with received levels of ~160 dB re 1 μ Pa and higher, on an approximate rms basis. The 50% probability of avoidance was estimated to occur at a CPA distance of 2.5 km from a 4000-in³ airgun array operating off central California. This would occur at an average received sound level of ~170 dB re 1 μ Pa_{rms}. Some slight behavioral changes were noted when approaching gray whales reached the distances where received sound levels were 140 to 160 dB re 1 μ Pa_{rms}, but these whales generally continued to approach (at a slight angle) until they passed the sound source at distances where received levels averaged ~170 dB re 1 μ Pa_{rms} (Malme et al. 1984; Malme and Miles 1985).

There was no indication that western gray whales exposed to seismic noise were displaced from their overall feeding grounds near Sakhalin Island during seismic programs in 1997 (Würsig et al. 1999) and in 2001 (Johnson et al. 2007; Meier et al. 2007; Yazvenko et al. 2007a). However, there were indications of subtle behavioral effects among whales that remained in the areas exposed to airgun sounds (Würsig et al. 1999; Gailey et al. 2007; Weller et al. 2006a). Also, there was evidence of localized redistribution of some individuals within the nearshore feeding ground so as to avoid close approaches by the seismic vessel (Weller et al. 2002, 2006b; Yazvenko et al. 2007a). Despite the evidence of subtle changes in some quantitative measures of behavior and local redistribution of some individuals, there was no apparent change in the frequency of feeding, as evident from mud plumes visible at the surface (Yazvenko et al. 2007b). The 2001 seismic program involved an unusually comprehensive combination of real-time monitoring and mitigation measures designed to avoid exposing western gray whales to received levels of sound above about 163 dB re 1 μ Pa_{rms} (Johnson et al. 2007). The lack of strong avoidance or other strong responses was presumably in part a result of the mitigation measures. Effects probably would have been more significant without such intensive mitigation efforts.

Gray whales in British Columbia exposed to seismic survey sound levels up to ~170 dB re 1 μ Pa did not appear to be strongly disturbed (Bain and Williams 2006). The few whales that were observed moved away from the airguns but toward deeper water where sound levels were said to be higher due to propagation effects (Bain and Williams 2006).

Rorquals.—Blue, sei, fin, and minke whales (all of which are members of the genus *Balaenoptera*) often have been seen in areas ensonified by airgun pulses (Stone 2003; MacLean and Haley 2004; Stone and Tasker 2006), and calls from blue and fin whales have been localized in areas with airgun operations (e.g., McDonald et al. 1995; Dunn and Hernandez 2009). Sightings by observers on seismic vessels during 110 large-source seismic surveys off the U.K. from 1997 to 2000 suggest that, during times of good

sightability, sighting rates for mysticetes (mainly fin and sei whales) were similar when large arrays of airguns were shooting vs. silent (Stone 2003; Stone and Tasker 2006). However, these whales tended to exhibit localized avoidance, remaining significantly further (on average) from the airgun array during seismic operations compared with non-seismic periods ($P = 0.0057$; Stone and Tasker 2006). The average CPA distances for baleen whales sighted when large airgun arrays were operating vs. silent were about 1.6 vs. 1.0 km. Baleen whales, as a group, were more often oriented away from the vessel while a large airgun array was shooting compared with periods of no shooting ($P < 0.05$; Stone and Tasker 2006). In addition, fin/sei whales were less likely to remain submerged during periods of seismic shooting (Stone 2003).

In a study off Nova Scotia, Moulton and Miller (2005) found little difference in sighting rates (after accounting for water depth) and initial average sighting distances of baleen whales when airguns were operating (mean = 1324 m) vs. silent (mean = 1303 m). However, there were indications that these whales were more likely to be moving away when seen during airgun operations. Baleen whales at the average sighting distance during airgun operations would have been exposed to sound levels (via direct path) of about 169 dB re 1 $\mu\text{Pa}_{\text{rms}}$ (Moulton and Miller 2005). Similarly, ship-based monitoring studies of blue, fin, sei and minke whales offshore of Newfoundland (Orphan Basin and Laurentian Sub-basin) found no more than small differences in sighting rates and swim directions during seismic vs. non-seismic periods (Moulton et al. 2005, 2006a,b). Analyses of CPA data yielded variable results.⁴ The authors of the Newfoundland reports concluded that, based on observations from the seismic vessel, some mysticetes exhibited localized avoidance of seismic operations (Moulton et al. 2005, 2006a).

Minke whales have occasionally been observed to approach active airgun arrays where received sound levels were estimated to be near 170–180 dB re 1 μPa (McLean and Haley 2004).

Discussion and Conclusions.—Baleen whales generally tend to avoid operating airguns, but avoidance radii are quite variable. Whales are often reported to show no overt reactions to airgun pulses at distances beyond a few kilometers, even though the airgun pulses remain well above ambient noise levels out to much longer distances. However, studies done since the late 1990s of migrating humpback and migrating bowhead whales show reactions, including avoidance, that sometimes extend to greater distances than documented earlier. Avoidance distances often exceed the distances at which boat-based observers can see whales, so observations from the source vessel can be biased. Observations over broader areas may be needed to determine the range of potential effects of some large-source seismic surveys where effects on cetaceans may extend to considerable distances (Richardson et al. 1999; Bain and Williams 2006; Moore and Angliss 2006). Longer-range observations, when required, can sometimes be obtained via systematic aerial surveys or aircraft-based observations of behavior (e.g., Richardson et al. 1986, 1999; Miller et al. 1999, 2005; Yazvenko et al. 2007a,b) or by use of observers on one or more support vessels operating in coordination with the seismic vessel (e.g., Smultea et al. 2004; Johnson et al. 2007). However, the presence of other vessels near the source vessel can, at least at times, reduce sightability of cetaceans from the source vessel (Beland et al. 2009), thus complicating interpretation of sighting data.

⁴ The CPA of baleen whales sighted from the seismic vessels was, on average, significantly closer during non-seismic periods vs. seismic periods in 2004 in the Orphan Basin (means 1526 m vs. 2316 m, respectively; Moulton et al. 2005). In contrast, mean distances without vs. with seismic did not differ significantly in 2005 in either the Orphan Basin (means 973 m vs. 832 m, respectively; Moulton et al. 2006a) or in the Laurentian Sub-basin (means 1928 m vs. 1650 m, respectively; Moulton et al. 2006b). In both 2005 studies, mean distances were greater (though not significantly so) *without* seismic.

Some baleen whales show considerable tolerance of seismic pulses. However, when the pulses are strong enough, avoidance or other behavioral changes become evident. Because the responses become less obvious with diminishing received sound level, it has been difficult to determine the maximum distance (or minimum received sound level) at which reactions to seismic become evident and, hence, how many whales are affected.

Studies of gray, bowhead, and humpback whales have determined that received levels of pulses in the 160–170 dB re 1 $\mu\text{Pa}_{\text{rms}}$ range seem to cause obvious avoidance behavior in a substantial fraction of the animals exposed. In many areas, seismic pulses diminish to these levels at distances ranging from 4 to 15 km from the source. A substantial proportion of the baleen whales within such distances may show avoidance or other strong disturbance reactions to the operating airgun array. However, in other situations, various mysticetes tolerate exposure to full-scale airgun arrays operating at even closer distances, with only localized avoidance and minor changes in activities. At the other extreme, in migrating bowhead whales, avoidance often extends to considerably larger distances (20–30 km) and lower received sound levels (120–130 dB re 1 $\mu\text{Pa}_{\text{rms}}$). Also, even in cases where there is no conspicuous avoidance or change in activity upon exposure to sound pulses from distant seismic operations, there are sometimes subtle changes in behavior (e.g., surfacing–respiration–dive cycles) that are only evident through detailed statistical analysis (e.g., Richardson et al. 1986; Gailey et al. 2007).

Mitigation measures for seismic surveys, especially nighttime seismic surveys, typically assume that many marine mammals (at least baleen whales) tend to avoid approaching airguns, or the seismic vessel itself, before being exposed to levels high enough for there to be any possibility of injury. This assumes that the ramp-up (soft-start) procedure is used when commencing airgun operations, to give whales near the vessel the opportunity to move away before they are exposed to sound levels that might be strong enough to elicit TTS. As noted above, single-airgun experiments with three species of baleen whales show that those species typically do tend to move away when a single airgun starts firing nearby, which simulates the onset of a ramp up. The three species that showed avoidance when exposed to the onset of pulses from a single airgun were *gray whales* (Malme et al. 1984, 1986, 1988); *bowhead whales* (Richardson et al. 1986; Ljungblad et al. 1988); and *humpback whales* (Malme et al. 1985; McCauley et al. 1998, 2000a,b). Since startup of a single airgun is equivalent to the start of a ramp-up (=soft start), this strongly suggests that many baleen whales will begin to move away during the initial stages of a ramp-up.

Data on short-term reactions by cetaceans to impulsive noises are not necessarily indicative of long-term or biologically significant effects. It is not known whether impulsive sounds affect reproductive rate or distribution and habitat use in subsequent days or years. However, gray whales have continued to migrate annually along the west coast of North America despite intermittent seismic exploration (and much ship traffic) in that area for decades (Appendix A in Malme et al. 1984; Richardson et al. 1995), and there has been a substantial increase in the population over recent decades (Angliss and Outlaw 2008). The western Pacific gray whale population did not seem affected by a seismic survey in its feeding ground during a prior year (Johnson et al. 2007). Similarly, bowhead whales have continued to travel to the eastern Beaufort Sea each summer despite seismic exploration in their summer and autumn range for many years (Richardson et al. 1987), and their numbers have increased notably (Angliss and Outlaw 2008). Bowheads also have been observed over periods of days or weeks in areas ensonified repeatedly by seismic pulses (Richardson et al. 1987; Harris et al. 2007). However, it is generally not known whether the same individual bowheads were involved in these repeated observations (within and between years) in strongly ensonified areas. In any event, in the absence of some unusual circumstances, the history of coexistence between seismic surveys and baleen whales suggests that brief exposures to sound pulses from any single seismic survey are unlikely to result in prolonged effects.

5.2 Toothed Whales

Little systematic information is available about reactions of toothed whales to noise pulses. Few studies similar to the more extensive baleen whale/seismic pulse work summarized above have been reported for toothed whales. However, there are recent systematic data on sperm whales (e.g., Gordon et al. 2006; Madsen et al. 2006; Winsor and Mate 2006; Jochens et al. 2008; Miller et al. 2009). There is also an increasing amount of information about responses of various odontocetes to seismic surveys based on monitoring studies (e.g., Stone 2003; Smultea et al. 2004; Moulton and Miller 2005; Bain and Williams 2006; Holst et al. 2006; Stone and Tasker 2006; Potter et al. 2007; Hauser et al. 2008; Holst and Smultea 2008; Weir 2008a; Barkaszi et al. 2009; Richardson et al. 2009).

Delphinids (Dolphins and similar) and Monodontids (Beluga).—Seismic operators and protected species observers on seismic vessels regularly see dolphins and other small toothed whales near operating airgun arrays, but in general there is a tendency for most delphinids to show some avoidance of operating seismic vessels (e.g., Goold 1996a,b,c; Calambokidis and Osmek 1998; Stone 2003; Moulton and Miller 2005; Holst et al. 2006; Stone and Tasker 2006; Weir 2008a; Richardson et al. 2009; see also Barkaszi et al. 2009). In most cases, the avoidance radii for delphinids appear to be small, on the order of 1 km or less, and some individuals show no apparent avoidance. Studies that have reported cases of small toothed whales close to the operating airguns include Duncan (1985), Arnold (1996), Stone (2003), and Holst et al. (2006). When a 3959 in³, 18-airgun array was firing off California, toothed whales behaved in a manner similar to that observed when the airguns were silent (Arnold 1996). Some dolphins seem to be attracted to the seismic vessel and floats, and some ride the bow wave of the seismic vessel even when a large array of airguns is firing (e.g., Moulton and Miller 2005). Nonetheless, small toothed whales more often tend to head away, or to maintain a somewhat greater distance from the vessel, when a large array of airguns is operating than when it is silent (e.g., Stone and Tasker 2006; Weir 2008a).

Weir (2008b) noted that a group of short-finned pilot whales initially showed an avoidance response to ramp up of a large airgun array, but that this response was limited in time and space. Although the ramp-up procedure is a widely-used mitigation measure, it remains uncertain how effective it is at alerting marine mammals (especially odontocetes) and causing them to move away from seismic operations (Weir 2008b).

Goold (1996a,b,c) studied the effects on common dolphins of 2D seismic surveys in the Irish Sea. Passive acoustic surveys were conducted from the “guard ship” that towed a hydrophone. The results indicated that there was a local displacement of dolphins around the seismic operation. However, observations indicated that the animals were tolerant of the sounds at distances outside a 1-km radius from the airguns (Goold 1996a). Initial reports of larger-scale displacement were later shown to represent a normal autumn migration of dolphins through the area, and were not attributable to seismic surveys (Goold 1996a,b,c).

The beluga is a species that (at least at times) shows long-distance avoidance of seismic vessels. Aerial surveys conducted in the southeastern Beaufort Sea in summer found that sighting rates of belugas were significantly lower at distances 10–20 km compared with 20–30 km from an operating airgun array (Miller et al. 2005). The low number of beluga sightings by protected species observers on the vessel seemed to confirm there was a strong avoidance response to the 2250 in³ airgun array. More recent seismic monitoring studies in the same area have confirmed that the apparent displacement effect on belugas extended farther than has been shown for other small odontocetes exposed to airgun pulses (e.g., Harris et al. 2007).

Observers stationed on seismic vessels operating off the U.K. from 1997 to 2000 have provided data on the occurrence and behavior of various toothed whales exposed to seismic pulses (Stone 2003;

Gordon et al. 2004; Stone and Tasker 2006). Dolphins of various species often showed more evidence of avoidance of operating airgun arrays than has been reported previously for small odontocetes. Sighting rates of white-sided dolphins, white-beaked dolphins, *Lagenorhynchus* spp., and all small odontocetes combined were significantly lower during periods when large-volume⁵ airgun arrays were shooting. Except for the pilot whale and bottlenose dolphin, CPA distances for all of the small odontocete species tested, including killer whales, were significantly farther from large airgun arrays during periods of shooting compared with periods of no shooting. Pilot whales were less responsive than other small odontocetes in the presence of seismic surveys (Stone and Tasker 2006). For small odontocetes as a group, and most individual species, orientations differed between times when large airgun arrays were operating vs. silent, with significantly fewer animals traveling towards and/or more traveling away from the vessel during shooting (Stone and Tasker 2006). Observers' records suggested that fewer cetaceans were feeding and fewer were interacting with the survey vessel (e.g., bow-riding) during periods with airguns operating, and small odontocetes tended to swim faster during periods of shooting (Stone and Tasker 2006). For most types of small odontocetes sighted by observers on seismic vessels, the median CPA distance was ≥ 0.5 km larger during airgun operations (Stone and Tasker 2006). Killer whales appeared to be more tolerant of seismic shooting in deeper waters.

Data collected during seismic operations in the Gulf of Mexico and off Central America show similar patterns. A summary of vessel-based monitoring data from the Gulf of Mexico during 2003–2008 showed that delphinids were generally seen farther from the vessel during seismic than during non-seismic periods (based on Barkaszi et al. 2009, excluding sperm whales). Similarly, during two NSF-funded L-DEO seismic surveys that used a large 20 airgun array (~ 7000 in³), sighting rates of delphinids were lower and initial sighting distances were farther away from the vessel during seismic than non-seismic periods (Smultea et al. 2004; Holst et al. 2005a, 2006; Richardson et al. 2009). Monitoring results during a seismic survey in the Southeast Caribbean showed that the mean CPA of delphinids was 991 m during seismic operations vs. 172 m when the airguns were not operational (Smultea et al. 2004). Surprisingly, nearly all acoustic detections via a towed passive acoustic monitoring (PAM) array, including both delphinids and sperm whales, were made when the airguns were operating (Smultea et al. 2004). Although the number of sightings during monitoring of a seismic survey off the Yucatán Peninsula, Mexico, was small ($n = 19$), the results showed that the mean CPA distance of delphinids there was 472 m during seismic operations vs. 178 m when the airguns were silent (Holst et al. 2005a). The acoustic detection rates were nearly 5 times higher during non-seismic compared with seismic operations (Holst et al. 2005a).

For two additional NSF-funded L-DEO seismic surveys in the Eastern Tropical Pacific, both using a large 36-airgun array (~ 6600 in³), the results are less easily interpreted (Richardson et al. 2009). During both surveys, the delphinid detection rate was lower during seismic than during non-seismic periods, as found in various other projects, but the mean CPA distance of delphinids was closer (not farther) during seismic periods (Hauser et al. 2008; Holst and Smultea 2008).

During two seismic surveys off Newfoundland and Labrador in 2004–05, dolphin sighting rates were lower during seismic periods than during non-seismic periods after taking temporal factors into account, although the difference was statistically significant only in 2004 (Moulton et al. 2005, 2006a). In 2005, the mean CPA distance of dolphins was significantly farther during seismic periods (807 vs. 652 m); in 2004, the corresponding difference was not significant.

⁵ Large volume means at least 1300 in³, with most (79%) at least 3000 in³.

Among Atlantic spotted dolphins off Angola ($n = 16$ useable groups), marked short-term and localized displacement was found in response to seismic operations conducted with a 24-airgun array (3147 in³ or 5085 in³) (Weir 2008a). Sample sizes were low, but CPA distances of dolphin groups were significantly larger when airguns were on (mean 1080 m) vs. off (mean 209 m). No Atlantic spotted dolphins were seen within 500 m of the airguns when they were operating, whereas all sightings when airguns were silent occurred within 500 m, including the only recorded “positive approach” behaviors.

Reactions of toothed whales to a single airgun or other small airgun source are not well documented, but tend to be less substantial than reactions to large airgun arrays (e.g., Stone 2003; Stone and Tasker 2006). During 91 site surveys off the U.K. in 1997–2000, sighting rates of all small odontocetes combined were significantly lower during periods the low-volume⁶ airgun sources were operating, and effects on orientation were evident for all species and groups tested (Stone and Tasker 2006). Results from four NSF-funded L-DEO seismic surveys using small arrays (up to 3 GI guns and 315 in³) were inconclusive. During surveys in the Eastern Tropical Pacific (Holst et al. 2005b) and in the Northwest Atlantic (Haley and Koski 2004), detection rates were slightly lower during seismic compared to non-seismic periods. However, mean CPAs were closer during seismic operations during one cruise (Holst et al. 2005b), and greater during the other cruise (Haley and Koski 2004). Interpretation of the data was confounded by the fact that survey effort and/or number of sightings during non-seismic periods during both surveys was small. Results from another two small-array surveys were even more variable (MacLean and Koski 2005; Smultea and Holst 2008).

Captive bottlenose dolphins and beluga whales exhibited changes in behavior when exposed to strong pulsed sounds similar in duration to those typically used in seismic surveys (Finneran et al. 2000, 2002, 2005). Finneran et al. (2002) exposed a captive bottlenose dolphin and beluga to single impulses from a water gun (80 in³). As compared with airgun pulses, water gun impulses were expected to contain proportionally more energy at higher frequencies because there is no significant gas-filled bubble, and thus little low-frequency bubble-pulse energy (Hutchinson and Detrick 1984). The captive animals sometimes vocalized after exposure and exhibited reluctance to station at the test site where subsequent exposure to impulses would be implemented (Finneran et al. 2002). Similar behaviors were exhibited by captive bottlenose dolphins and a beluga exposed to single underwater pulses designed to simulate those produced by distant underwater explosions (Finneran et al. 2000). It is uncertain what relevance these observed behaviors in captive, trained marine mammals exposed to single transient sounds may have to free-ranging animals exposed to multiple pulses. In any event, the animals tolerated rather high received levels of sound before exhibiting the aversive behaviors mentioned above.

Odontocete responses (or lack of responses) to noise pulses from underwater explosions (as opposed to airgun pulses) may be indicative of odontocete responses to very strong noise pulses. During the 1950s, small explosive charges were dropped into an Alaskan river in attempts to scare belugas away from salmon. Success was limited (Fish and Vania 1971; Frost et al. 1984). Small explosive charges were “not always effective” in moving bottlenose dolphins away from sites in the Gulf of Mexico where larger demolition blasts were about to occur (Klima et al. 1988). Odontocetes may be attracted to fish killed by explosions, and thus attracted rather than repelled by “scare” charges. Captive false killer whales showed no obvious reaction to single noise pulses from small (10 g) charges; the received level was ~185 dB re 1 μ Pa (Akamatsu et al. 1993). Jefferson and Curry (1994) reviewed several additional studies that found limited or no effects of noise pulses from small explosive charges on killer whales and

⁶ For low volume arrays, maximum volume was 820 in³, with most (87%) \leq 180 in³.

other odontocetes. Aside from the potential for causing auditory impairment (see below), the tolerance to these charges may indicate a lack of effect, or the failure to move away may simply indicate a stronger desire to feed, regardless of circumstances.

Phocoenids (Porpoises).—Porpoises, like delphinids, show variable reactions to seismic operations, and reactions apparently depend on species. The limited available data suggest that harbor porpoises show stronger avoidance of seismic operations than Dall's porpoises (Stone 2003; MacLean and Koski 2005; Bain and Williams 2006). In Washington State waters, the harbor porpoise—despite being considered a high-frequency specialist—appeared to be the species affected by the lowest received level of airgun sound (<145 dB re $1 \mu\text{Pa}_{\text{rms}}$ at a distance >70 km; Bain and Williams 2006). Similarly, during seismic surveys with large airgun arrays off the U.K. in 1997–2000, there were significant differences in directions of travel by harbor porpoises during periods when the airguns were shooting vs. silent (Stone 2003; Stone and Tasker 2006). A captive harbor porpoise exposed to single sound pulses from a small airgun showed aversive behavior upon receipt of a pulse with received level above 174 dB re $1 \mu\text{Pa}_{\text{pk-pk}}$ or SEL >145 dB re $1 \mu\text{Pa}^2 \cdot \text{s}$ (Lucke et al. 2009). In contrast, Dall's porpoises seem relatively tolerant of airgun operations (MacLean and Koski 2005; Bain and Williams 2006), although they too have been observed to avoid large arrays of operating airguns (Calambokidis and Osmek 1998; Bain and Williams 2006). The apparent tendency for greater responsiveness in the harbor porpoise is consistent with their relative responsiveness to boat traffic and some other acoustic sources (Richardson et al. 1995; Southall et al. 2007).

Beaked Whales.—There are almost no specific data on the behavioral reactions of beaked whales to seismic surveys. Most beaked whales tend to avoid approaching vessels of other types (e.g., Würsig et al. 1998). They may also dive for an extended period when approached by a vessel (e.g., Kasuya 1986), although it is uncertain how much longer such dives may be as compared to dives by undisturbed beaked whales, which also are often quite long (Baird et al. 2006; Tyack et al. 2006b). In any event, it is likely that most beaked whales would also show strong avoidance of an approaching seismic vessel, regardless of whether or not the airguns are operating. However, this has not been documented explicitly. Northern bottlenose whales sometimes are quite tolerant of slow-moving vessels not emitting airgun pulses (Reeves et al. 1993; Hooker et al. 2001). The few detections (acoustic or visual) of northern bottlenose whales from seismic vessels during recent seismic surveys off Nova Scotia have been during times when the airguns were shut down; no detections were reported when the airguns were operating (Moulton and Miller 2005; Potter et al. 2007). However, other visual and acoustic studies indicated that some northern bottlenose whales remained in the general area and continued to produce high-frequency clicks when exposed to sound pulses from distant seismic surveys (Gosselin and Lawson 2004; Laurinoli and Cochran 2005; Simard et al. 2005).

There are increasing indications that some beaked whales tend to strand when military exercises involving mid-frequency sonar operation are ongoing nearby (e.g., Simmonds and Lopez-Jurado 1991; Frantzis 1998; NOAA and USN 2001; Jepson et al. 2003; Barlow and Gisiner 2006; see also the “Strandings and Mortality” subsection, later). These strandings are apparently at least in part a disturbance response, although auditory or other injuries or other physiological effects may also be a factor. Whether beaked whales would ever react similarly to seismic surveys is unknown. Seismic survey sounds are quite different from those of the sonars in operation during the above-cited incidents. No conclusive link has been established between seismic surveys and beaked whale strandings. There was a stranding of two Cuvier's beaked whales in the Gulf of California (Mexico) in September 2002 when the R/V *Maurice Ewing* was conducting a seismic survey in the general area (e.g., Malakoff 2002; Hildebrand 2005). However, NMFS did not establish a cause and effect relationship between this stranding and the seismic survey activities (Hogarth 2002). Cox et al. (2006) noted the “lack of knowledge regard-

ing the temporal and spatial correlation between the [stranding] and the sound source”. Hildebrand (2005) illustrated the approximate temporal-spatial relationships between the stranding and the *Ewing*'s tracks, but the time of the stranding was not known with sufficient precision for accurate determination of the CPA distance of the whales to the *Ewing*. Another stranding of Cuvier's beaked whales in the Galápagos occurred during a seismic survey in April 2000; however “There is no obvious mechanism that bridges the distance between this source and the stranding site” (Gentry [ed.] 2002).

Sperm Whales.—All three species of sperm whales have been reported to show avoidance reactions to standard vessels not emitting airgun sounds (e.g., Richardson et al. 1995; Würsig et al. 1998; McAlpine 2002; Baird 2005). However, most studies of the sperm whale *Physeter macrocephalus* exposed to airgun sounds indicate that this species shows considerable tolerance of airgun pulses. The whales usually do not show strong avoidance (i.e., they do not leave the area) and they continue to call.

There were some early and limited observations suggesting that sperm whales in the Southern Ocean ceased calling during some (but not all) times when exposed to weak noise pulses from extremely distant (>300 km) seismic exploration. However, other operations in the area could also have been a factor (Bowles et al. 1994). This “quieting” was suspected to represent a disturbance effect, in part because sperm whales exposed to pulsed man-made sounds at higher frequencies often cease calling (Watkins and Schevill 1975; Watkins et al. 1985). Also, there was an early preliminary account of possible long-range avoidance of seismic vessels by sperm whales in the Gulf of Mexico (Mate et al. 1994). However, this has not been substantiated by subsequent more detailed work in that area (Gordon et al. 2006; Winsor and Mate 2006; Jochens et al. 2008; Miller et al. 2009).

Recent and more extensive data from vessel-based monitoring programs in U.K. waters and off Newfoundland and Angola suggest that sperm whales in those areas show little evidence of avoidance or behavioral disruption in the presence of operating seismic vessels (Stone 2003; Stone and Tasker 2006; Moulton et al. 2005, 2006a; Weir 2008a). Among sperm whales off Angola ($n = 96$ useable groups), there were no significant differences in encounter rates (sightings/hr) when a 24-airgun array (3147 in³ or 5085 in³) was operating vs. silent (Weir 2008a). There was also no significant difference in the CPA distances of the sperm whale sightings when airguns were on vs. off (means 3039 m vs. 2594 m, respectively). Encounter rate tended to increase over the 10-month duration of the seismic survey. These types of observations are difficult to interpret because the observers are stationed on or near the seismic vessel, and may underestimate reactions by some of the more responsive animals, which may be beyond visual range. However, these results do seem to show considerable tolerance of seismic surveys by at least some sperm whales. Also, a study off northern Norway indicated that sperm whales continued to call when exposed to pulses from a distant seismic vessel. Received levels of the seismic pulses were up to 146 dB re 1 $\mu\text{Pa}_{\text{p-p}}$ (Madsen et al. 2002).

Similarly, a study conducted off Nova Scotia that analyzed recordings of sperm whale vocalizations at various distances from an active seismic program did not detect any obvious changes in the distribution or behavior of sperm whales (McCall Howard 1999).

Sightings of sperm whales by observers on seismic vessels operating in the Gulf of Mexico during 2003–2008 were at very similar average distances regardless of the airgun operating conditions (Barkaszi et al. 2009). For example, the mean sighting distance was 1839 m when the airgun array was in full operation ($n=612$) vs. 1960 m when all airguns were off ($n=66$).

A controlled study of the reactions of tagged sperm whales to seismic surveys was done recently in the Gulf of Mexico — the Sperm Whale Seismic Study or SWSS (Gordon et al. 2006; Madsen et al. 2006; Winsor and Mate 2006; Jochens et al. 2008; Miller et al. 2009). During SWSS, D-tags (Johnson and Tyack 2003) were used to record the movement and acoustic exposure of eight foraging sperm whales

before, during, and after controlled exposures to sound from airgun arrays (Jochens et al. 2008; Miller et al. 2009). Whales were exposed to maximum received sound levels of 111–147 dB re 1 $\mu\text{Pa}_{\text{rms}}$ (131–162 dB re 1 $\mu\text{Pa}_{\text{pk-pk}}$) at ranges of ~1.4–12.8 km from the sound source (Miller et al. 2009). Although the tagged whales showed no discernible horizontal avoidance, some whales showed changes in diving and foraging behavior during full-array exposure, possibly indicative of subtle negative effects on foraging (Jochens et al. 2008; Miller et al. 2009; Tyack 2009). Two indications of foraging that they studied were oscillations in pitch and occurrence of echolocation buzzes, both of which tend to occur when a sperm whale closes-in on prey. "Oscillations in pitch generated by swimming movements during foraging dives were on average 6% lower during exposure than during the immediately following poSt. exposure period, with all 7 foraging whales exhibiting less pitching ($P = 0.014$). Buzz rates, a proxy for attempts to capture prey, were 19% lower during exposure..." (Miller et al. 2009). Although the latter difference was not statistically significant ($P = 0.141$), the percentage difference in buzz rate during exposure vs. poSt. exposure conditions appeared to be strongly correlated with airgun-whale distance (Miller et al. 2009: Fig. 5; Tyack 2009).

Discussion and Conclusions.—Dolphins and porpoises are often seen by observers on active seismic vessels, occasionally at close distances (e.g., bow riding). However, some studies near the U.K., Newfoundland and Angola, in the Gulf of Mexico, and off Central America have shown localized avoidance. Also, belugas summering in the Canadian Beaufort Sea showed larger-scale avoidance, tending to avoid waters out to 10–20 km from operating seismic vessels. In contrast, recent studies show little evidence of conspicuous reactions by sperm whales to airgun pulses, contrary to earlier indications.

There are almost no specific data on responses of beaked whales to seismic surveys, but it is likely that most if not all species show strong avoidance. There is increasing evidence that some beaked whales may strand after exposure to strong noise from sonars. Whether they ever do so in response to seismic survey noise is unknown. Northern bottlenose whales seem to continue to call when exposed to pulses from distant seismic vessels.

Overall, odontocete reactions to large arrays of airguns are variable and, at least for delphinids and some porpoises, seem to be confined to a smaller radius than has been observed for some mysticetes. However, other data suggest that some odontocetes species, including belugas and harbor porpoises, may be more responsive than might be expected given their poor low-frequency hearing. Reactions at longer distances may be particularly likely when sound propagation conditions are conducive to transmission of the higher-frequency components of airgun sound to the animals' location (DeRuiter et al. 2006; Goold and Coates 2006; Tyack et al. 2006a; Potter et al. 2007).

For delphinids, and possibly the Dall's porpoise, the available data suggest that ≥ 170 dB re 1 $\mu\text{Pa}_{\text{rms}}$ disturbance criterion (rather than ≥ 160 dB) would be appropriate. With a medium-to-large airgun array, received levels typically diminish to 170 dB within 1–4 km, whereas levels typically remain above 160 dB out to 4–15 km (e.g., Tolstoy et al. 2009). Reaction distances for delphinids are more consistent with the typical 170 dB re 1 $\mu\text{Pa}_{\text{rms}}$ distances. The 160 dB (rms) criterion currently applied by NMFS was developed based primarily on data from gray and bowhead whales. Avoidance distances for delphinids and Dall's porpoises tend to be shorter than for those two mysticete species. For delphinids and Dall's porpoises, there is no indication of strong avoidance or other disruption of behavior at distances beyond those where received levels would be ~ 170 dB re 1 $\mu\text{Pa}_{\text{rms}}$.

5.3 Pinnipeds

Few studies of the reactions of pinnipeds to noise from open-water seismic exploration have been published (for review of the early literature, see Richardson et al. 1995). However, pinnipeds have been

observed during a number of seismic monitoring studies. Monitoring in the Beaufort Sea during 1996–2002 provided a substantial amount of information on avoidance responses (or lack thereof) and associated behavior. Additional monitoring of that type has been done in the Beaufort and Chukchi Seas in 2006–2009. Pinnipeds exposed to seismic surveys have also been observed during seismic surveys along the U.S. west coast. Some limited data are available on physiological responses of pinnipeds exposed to seismic sound, as studied with the aid of radio telemetry. Also, there are data on the reactions of pinnipeds to various other related types of impulsive sounds.

Early observations provided considerable evidence that pinnipeds are often quite tolerant of strong pulsed sounds. During seismic exploration off Nova Scotia, gray seals exposed to noise from airguns and linear explosive charges reportedly did not react strongly (J. Parsons *in* Greene et al. 1985). An airgun caused an initial startle reaction among South African fur seals but was ineffective in scaring them away from fishing gear (Anonymous 1975). Pinnipeds in both water and air sometimes tolerate strong noise pulses from non-explosive and explosive scaring devices, especially if attracted to the area for feeding or reproduction (Mate and Harvey 1987; Reeves et al. 1996). Thus, pinnipeds are expected to be rather tolerant of, or to habituate to, repeated underwater sounds from distant seismic sources, at least when the animals are strongly attracted to the area.

In the U.K., a radio-telemetry study demonstrated short-term changes in the behavior of harbor (=common) and gray seals exposed to airgun pulses (Thompson et al. 1998). Harbor seals were exposed to seismic pulses from a 90-in³ array (3 × 30 in³ airguns), and behavioral responses differed among individuals. One harbor seal avoided the array at distances up to 2.5 km from the source and only resumed foraging dives after seismic stopped. Another harbor seal exposed to the same small airgun array showed no detectable behavioral response, even when the array was within 500 m. Gray seals exposed to a single 10-in³ airgun showed an avoidance reaction: they moved away from the source, increased swim speed and/or dive duration, and switched from foraging dives to predominantly transit dives. These effects appeared to be short-term as gray seals either remained in, or returned at least once to, the foraging area where they had been exposed to seismic pulses. These results suggest that there are interspecific as well as individual differences in seal responses to seismic sounds.

Off California, visual observations from a seismic vessel showed that California sea lions “typically ignored the vessel and array. When [they] displayed behavior modifications, they often appeared to be reacting visually to the sight of the towed array. At times, California sea lions were attracted to the array, even when it was on. At other times, these animals would appear to be actively avoiding the vessel and array” (Arnold 1996). In Puget Sound, sighting distances for harbor seals and California sea lions tended to be larger when airguns were operating; both species tended to orient away whether or not the airguns were firing (Calambokidis and Osmeck 1998). Bain and Williams (2006) also stated that their small sample of harbor seals and sea lions tended to orient and/or move away upon exposure to sounds from a large airgun array.

Monitoring work in the Alaskan Beaufort Sea during 1996–2001 provided considerable information regarding the behavior of seals exposed to seismic pulses (Harris et al. 2001; Moulton and Lawson 2002). Those seismic projects usually involved arrays of 6–16 airguns with total volumes 560–1500 in³. Subsequent monitoring work in the Canadian Beaufort Sea in 2001–2002, with a somewhat larger airgun system (24 airguns, 2250 in³), provided similar results (Miller et al. 2005). The combined results suggest that some seals avoid the immediate area around seismic vessels. In most survey years, ringed seal sightings averaged somewhat farther away from the seismic vessel when the airguns were operating than when they were not (Moulton and Lawson 2002). Also, seal sighting rates at the water surface were lower during airgun array operations than during no-airgun periods in each survey year except 1997.

However, the avoidance movements were relatively small, on the order of 100 m to (at most) a few hundreds of meters, and many seals remained within 100–200 m of the trackline as the operating airgun array passed by.

The operation of the airgun array had minor and variable effects on the behavior of seals visible at the surface within a few hundred meters of the airguns (Moulton and Lawson 2002). The behavioral data indicated that some seals were more likely to swim away from the source vessel during periods of airgun operations and more likely to swim towards or parallel to the vessel during non-seismic periods. No consistent relationship was observed between exposure to airgun noise and proportions of seals engaged in other recognizable behaviors, e.g., “looked” and “dove”. Such a relationship might have occurred if seals seek to reduce exposure to strong seismic pulses, given the reduced airgun noise levels close to the surface where “looking” occurs (Moulton and Lawson 2002).

Monitoring results from the Canadian Beaufort Sea during 2001–2002 were more variable (Miller et al. 2005). During 2001, sighting rates of seals (mostly ringed seals) were similar during all seismic states, including periods without airgun operations. However, seals tended to be seen closer to the vessel during non-seismic than seismic periods. In contrast, during 2002, sighting rates of seals were higher during non-seismic periods than seismic operations, and seals were seen farther from the vessel during non-seismic compared to seismic activity (a marginally significant result). The combined data for both years showed that sighting rates were higher during non-seismic periods compared to seismic periods, and that sighting distances were similar during both seismic states. Miller et al. (2005) concluded that seals showed very limited avoidance to the operating airgun array.

Vessel-based monitoring also took place in the Alaskan Chukchi and Beaufort seas during 2006–2008 (Reiser et al. 2009). Observers on the seismic vessels saw phocid seals less frequently while airguns were operating than when airguns were silent. Also, during airgun operations, those observers saw seals less frequently than did observers on nearby vessels without airguns. Finally, observers on the latter “no-airgun” vessels saw seals more often when the nearby source vessels’ airguns were operating than when they were silent. All of these observations are indicative of a tendency for phocid seals to exhibit localized avoidance of the seismic source vessel when airguns are firing (Reiser et al. 2009).

In summary, visual monitoring from seismic vessels has shown only slight (if any) avoidance of airguns by pinnipeds, and only slight (if any) changes in behavior. These studies show that many pinnipeds do not avoid the area within a few hundred meters of an operating airgun array. However, based on the studies with large sample size, or observations from a separate monitoring vessel, or radio telemetry, it is apparent that some phocid seals do show localized avoidance of operating airguns. The limited nature of this tendency for avoidance is a concern. It suggests that one cannot rely on pinnipeds to move away, or to move very far away, before received levels of sound from an approaching seismic survey vessel approach those that may cause hearing impairment (see below).

5.4 Sirenians, Sea Otter and Polar Bear

We are not aware of any information on the reactions of sirenians to airgun sounds

Behavior of sea otters along the California coast was monitored by Riedman (1983, 1984) while they were exposed to a single 100 in³ airgun and a 4089 in³ airgun array. No disturbance reactions were evident when the airgun array was as close as 0.9 km. Sea otters also did not respond noticeably to the single airgun. These results suggest that sea otters may be less responsive to marine seismic pulses than some other marine mammals, such as mysticetes and odontocetes (summarized above). Also, sea otters spend a great deal of time at the surface feeding and grooming (Riedman 1983, 1984). While at the

surface, the potential noise exposure of sea otters would be much reduced by pressure-release and interference (Lloyd's mirror) effects at the surface (Greene and Richardson 1988; Richardson et al. 1995).

Airgun effects on polar bears have not been studied. However, polar bears on the ice would be largely unaffected by underwater sound. Sound levels received by polar bears in the water would be attenuated because polar bears generally do not dive much below the surface and received levels of airgun sounds are reduced near the surface because of the aforementioned pressure release and interference effects at the water's surface.

6. Hearing Impairment and Other Physical Effects of Seismic Surveys

Temporary or permanent hearing impairment is a possibility when marine mammals are exposed to very strong sounds. Temporary threshold shift (TTS) has been demonstrated and studied in certain captive odontocetes and pinnipeds exposed to strong sounds (reviewed in Southall et al. 2007). However, there has been no specific documentation of TTS let alone permanent hearing damage, i.e. permanent threshold shift (PTS), in free-ranging marine mammals exposed to sequences of airgun pulses during realistic field conditions. Current NMFS policy regarding exposure of marine mammals to high-level sounds is that cetaceans and pinnipeds should not be exposed to impulsive sounds ≥ 80 and 190 dB re $1 \mu\text{Pa}_{\text{rms}}$, respectively (NMFS 2000). Those criteria have been used in establishing the safety (=shut-down) radii planned for numerous seismic surveys conducted under U.S. jurisdiction. However, those criteria were established before there was any information about the minimum received levels of sounds necessary to cause auditory impairment in marine mammals. As discussed below,

- the 180-dB criterion for cetaceans is probably quite precautionary, i.e., lower than necessary to avoid temporary auditory impairment let alone permanent auditory injury, at least for delphinids.
- TTS is not injury and does not constitute "Level A harassment" in U.S. MMPA terminology.
- the minimum sound level necessary to cause permanent hearing impairment ("Level A harassment") is higher, by a variable and generally unknown amount, than the level that induces barely-detectable TTS.
- the level associated with the onset of TTS is often considered to be a level below which there is no danger of permanent damage. The actual PTS threshold is likely to be well above the level causing onset of TTS (Southall et al. 2007).

Recommendations for new science-based noise exposure criteria for marine mammals, frequency-weighting procedures, and related matters were published recently (Southall et al. 2007). Those recommendations have not, as of late 2009, been formally adopted by NMFS for use in regulatory processes and during mitigation programs associated with seismic surveys. However, some aspects of the recommendations have been taken into account in certain EISs and small-take authorizations. NMFS has indicated that it may issue new noise exposure criteria for marine mammals that account for the now-available scientific data on TTS, the expected offset between the TTS and PTS thresholds, differences in the acoustic frequencies to which different marine mammal groups are sensitive, and other relevant factors. Preliminary information about possible changes in the regulatory and mitigation requirements, and about the possible structure of new criteria, was given by Wieting (2004) and NMFS (2005).

Several aspects of the monitoring and mitigation measures that are now often implemented during seismic survey projects are designed to detect marine mammals occurring near the airgun array, and to avoid exposing them to sound pulses that might, at least in theory, cause hearing impairment. In addition, many cetaceans and (to a limited degree) pinnipeds show some avoidance of the area where received levels of airgun sound are high enough such that hearing impairment could potentially occur. In those

cases, the avoidance responses of the animals themselves will reduce or (most likely) avoid the possibility of hearing impairment.

Non-auditory physical effects may also occur in marine mammals exposed to strong underwater pulsed sound. Possible types of non-auditory physiological effects or injuries that might (in theory) occur include stress, neurological effects, bubble formation, and other types of organ or tissue damage. It is possible that some marine mammal species (i.e., beaked whales) may be especially susceptible to injury and/or stranding when exposed to strong pulsed sounds. The following subsections summarize available data on noise-induced hearing impairment and non-auditory physical effects.

6.1 Temporary Threshold Shift (TTS)

TTS is the mildest form of hearing impairment that can occur during exposure to a strong sound (Kryter 1985). While experiencing TTS, the hearing threshold rises and a sound must be stronger in order to be heard. It is a temporary phenomenon, and (especially when mild) is not considered to represent physical damage or “injury” (Southall et al. 2007). Rather, the onset of TTS is an indicator that, if the animal is exposed to higher levels of that sound, physical damage is ultimately a possibility.

The magnitude of TTS depends on the level and duration of noise exposure, and to some degree on frequency, among other considerations (Kryter 1985; Richardson et al. 1995; Southall et al. 2007). For sound exposures at or somewhat above the TTS threshold, hearing sensitivity recovers rapidly after exposure to the noise ends. In terrestrial mammals, TTS can last from minutes or hours to (in cases of strong TTS) days. Only a few data have been obtained on sound levels and durations necessary to elicit mild TTS in marine mammals (none in mysticetes), and none of the published data concern TTS elicited by exposure to multiple pulses of sound during operational seismic surveys (Southall et al. 2007).

Toothed Whales.—There are empirical data on the sound exposures that elicit onset of TTS in captive bottlenose dolphins and belugas. The majority of these data concern non-impulse sound, but there are some limited published data concerning TTS onset upon exposure to a single pulse of sound from a watergun (Finneran et al. 2002). A detailed review of all TTS data from marine mammals can be found in Southall et al. (2007). The following summarizes some of the key results from odontocetes.

Recent information corroborates earlier expectations that the effect of exposure to strong transient sounds is closely related to the total amount of acoustic energy that is received. Finneran et al. (2005) examined the effects of tone duration on TTS in bottlenose dolphins. Bottlenose dolphins were exposed to 3 kHz tones (non-impulsive) for periods of 1, 2, 4 or 8 s, with hearing tested at 4.5 kHz. For 1-s exposures, TTS occurred with SELs of 197 dB, and for exposures >1 s, SEL >195 dB resulted in TTS (SEL is equivalent to energy flux, in dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$). At an SEL of 195 dB, the mean TTS (4 min after exposure) was 2.8 dB. Finneran et al. (2005) suggested that an SEL of 195 dB is the likely threshold for the onset of TTS in dolphins and belugas exposed to tones of durations 1–8 s (i.e., TTS onset occurs at a near-constant SEL, independent of exposure duration). That implies that, at least for non-impulsive tones, a doubling of exposure time results in a 3 dB lower TTS threshold.

The assumption that, in marine mammals, the occurrence and magnitude of TTS is a function of cumulative acoustic energy (SEL) is probably an oversimplification. Kastak et al. (2005) reported preliminary evidence from pinnipeds that, for prolonged non-impulse noise, higher SELs were required to elicit a given TTS if exposure duration was short than if it was longer, i.e., the results were not fully consistent with an equal-energy model to predict TTS onset. Mooney et al. (2009a) showed this in a bottlenose dolphin exposed to octave-band non-impulse noise ranging from 4 to 8 kHz at SPLs of 130 to 178 dB re 1 μPa for periods of 1.88 to 30 min. Higher SELs were required to induce a given TTS if exposure duration short than if it was longer. Exposure of the aforementioned bottlenose dolphin to a sequence of

brief sonar signals showed that, with those brief (but non-impulse) sounds, the received energy (SEL) necessary to elicit TTS was higher than was the case with exposure to the more prolonged octave-band noise (Mooney et al. 2009b). Those authors concluded that, when using (non-impulse) acoustic signals of duration ~ 0.5 s, SEL must be at least 210–214 dB re $1 \mu\text{Pa}^2 \cdot \text{s}$ to induce TTS in the bottlenose dolphin.

On the other hand, the TTS threshold for odontocetes exposed to a single impulse from a waterygun (Finneran et al. 2002) appeared to be somewhat lower than for exposure to non-impulse sound. This was expected, based on evidence from terrestrial mammals showing that broadband pulsed sounds with rapid rise times have greater auditory effect than do non-impulse sounds (Southall et al. 2007). The received energy level of a single seismic pulse that caused the onset of mild TTS in the beluga, as measured without frequency weighting, was ~ 186 dB re $1 \mu\text{Pa}^2 \cdot \text{s}$ or 186 dB SEL (Finneran et al. 2002).⁷ The rms level of an airgun pulse (in dB re $1 \mu\text{Pa}$ measured over the duration of the pulse) is typically 10–15 dB higher than the SEL for the same pulse when received within a few kilometers of the airguns. Thus, a single airgun pulse might need to have a received level of ~ 196 – 201 dB re $1 \mu\text{Pa}_{\text{rms}}$ in order to produce brief, mild TTS. Exposure to several strong seismic pulses that each has a flat-weighted received level near 190 dB_{rms} (175–180 dB SEL) could result in cumulative exposure of ~ 186 dB SEL (flat-weighted) or ~ 183 dB SEL (M_{mf} -weighted), and thus slight TTS in a small odontocete. That assumes that the TTS threshold upon exposure to multiple pulses is (to a first approximation) a function of the total received pulse energy, without allowance for any recovery between pulses.

The above TTS information for odontocetes is derived from studies on the bottlenose dolphin and beluga. For the one harbor porpoise tested, the received level of airgun sound that elicited onset of TTS was lower. The animal was exposed to single pulses from a small (20 in^3) airgun, and auditory evoked potential methods were used to test the animal's hearing sensitivity at frequencies of 4, 32, or 100 kHz after each exposure (Lucke et al. 2009). Based on the measurements at 4 kHz, TTS occurred upon exposure to one airgun pulse with received level ~ 200 dB re $1 \mu\text{Pa}_{\text{pk-pk}}$ or an SEL of 164.3 dB re $1 \mu\text{Pa}^2 \cdot \text{s}$. If these results from a single animal are representative, it is inappropriate to assume that onset of TTS occurs at similar received levels in all odontocetes (*cf.* Southall et al. 2007). Some cetaceans may incur TTS at lower sound exposures than are necessary to elicit TTS in the beluga or bottlenose dolphin.

Insofar as we are aware, there are no published data confirming that the auditory effect of a sequence of airgun pulses received by an odontocete is a function of their cumulative energy. Southall et al. (2007) consider that to be a reasonable, but probably somewhat precautionary, assumption. It is precautionary because, based on data from terrestrial mammals, one would expect that a given energy exposure would have somewhat less effect if separated into discrete pulses, with potential opportunity for partial auditory recovery between pulses. However, as yet there has been little study of the rate of recovery from TTS in marine mammals, and in humans and other terrestrial mammals the available data on recovery are quite variable. Southall et al. (2007) concluded that—until relevant data on recovery are available from marine mammals—it is appropriate not to allow for any assumed recovery during the intervals between pulses within a pulse sequence.

Additional data are needed to determine the received sound levels at which small odontocetes would start to incur TTS upon exposure to repeated, low-frequency pulses of airgun sound with variable received levels. To determine how close an airgun array would need to approach in order to elicit TTS, it

⁷ If the low-frequency components of the waterygun sound used in the experiments of Finneran et al. (2002) are downweighted as recommended by Southall et al. (2007) using their M_{mf} -weighting curve, the effective exposure level for onset of mild TTS was 183 dB re $1 \mu\text{Pa}^2 \cdot \text{s}$ (Southall et al. 2007).

is necessary to determine the total energy that a mammal would receive as an airgun array approaches, passes at various CPA distances, and moves away (e.g., Erbe and King 2009). At the present state of knowledge, it is also necessary to assume that the effect is directly related to total received energy even though that energy is received in multiple pulses separated by gaps. The lack of data on the exposure levels necessary to cause TTS in toothed whales when the signal is a series of pulsed sounds, separated by silent periods, remains a data gap, as is the lack of published data on TTS in odontocetes other than the beluga, bottlenose dolphin, and harbor porpoise.

Baleen Whales.—There are no data, direct or indirect, on levels or properties of sound that are required to induce TTS in any baleen whale. The frequencies to which mysticetes are most sensitive are assumed to be lower than those to which odontocetes are most sensitive, and natural background noise levels at those low frequencies tend to be higher. As a result, auditory thresholds of baleen whales within their frequency band of best hearing are believed to be higher (less sensitive) than are those of odontocetes at their best frequencies (Clark and Ellison 2004). From this, it is suspected that received levels causing TTS onset may also be higher in mysticetes (Southall et al. 2007). However, based on preliminary simulation modeling that attempted to allow for various uncertainties in assumptions and variability around population means, Gedamke et al. (2008) suggested that some baleen whales whose closest point of approach to a seismic vessel is 1 km or more could experience TTS or even PTS.

In practice during seismic surveys, few if any cases of TTS are expected given the strong likelihood that baleen whales would avoid the approaching airguns (or vessel) before being exposed to levels high enough for there to be any possibility of TTS (see above for evidence concerning avoidance responses by baleen whales). This assumes that the ramp-up (soft-start) procedure is used when commencing airgun operations, to give whales near the vessel the opportunity to move away before they are exposed to sound levels that might be strong enough to elicit TTS. As discussed earlier, single-airgun experiments with bowhead, gray, and humpback whales show that those species do tend to move away when a single airgun starts firing nearby, which simulates the onset of a ramp up.

Pinnipeds.—In pinnipeds, TTS thresholds associated with exposure to brief pulses (single or multiple) of underwater sound have not been measured. Two California sea lions did not incur TTS when exposed to single brief pulses with received levels of ~178 and 183 dB re 1 $\mu\text{Pa}_{\text{rms}}$ and total energy fluxes of 161 and 163 dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$ (Finneran et al. 2003). However, initial evidence from more prolonged (non-pulse) exposures suggested that some pinnipeds (harbor seals in particular) incur TTS at somewhat lower received levels than do small odontocetes exposed for similar durations (Kastak et al. 1999, 2005; Ketten et al. 2001). Kastak et al. (2005) reported that the amount of threshold shift increased with increasing SEL in a California sea lion and harbor seal. They noted that, for non-impulse sound, doubling the exposure duration from 25 to 50 min (i.e., a +3 dB change in SEL) had a greater effect on TTS than an increase of 15 dB (95 vs. 80 dB) in exposure level. Mean threshold shifts ranged from 2.9–12.2 dB, with full recovery within 24 hr (Kastak et al. 2005). Kastak et al. (2005) suggested that, for non-impulse sound, SELs resulting in TTS onset in three species of pinnipeds may range from 183 to 206 dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$, depending on the absolute hearing sensitivity.

As noted above for odontocetes, it is expected that—for impulse as opposed to non-impulse sound—the onset of TTS would occur at a lower cumulative SEL given the assumed greater auditory effect of broadband impulses with rapid rise times. The threshold for onset of mild TTS upon exposure of a harbor seal to impulse sounds has been estimated indirectly as being an SEL of ~171 dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$ (Southall et al. 2007). That would be approximately equivalent to a single pulse with received level ~181–186 dB re 1 $\mu\text{Pa}_{\text{rms}}$, or a series of pulses for which the highest rms values are a few dB lower.

At least for non-impulse sounds, TTS onset occurs at appreciably higher received levels in California sea lions and northern elephant seals than in harbor seals (Kastak et al. 2005). Thus, the former two species would presumably need to be closer to an airgun array than would a harbor seal before TTS is a possibility. Insofar as we are aware, there are no data to indicate whether the TTS thresholds of other pinniped species are more similar to those of the harbor seal or to those of the two less-sensitive species.

Sirenians, Sea Otter and Polar Bear.—There are no available data on TTS in sea otters and polar bears. However, TTS is unlikely to occur in sea otters or polar bears if they are on the water surface, given the pressure release and Lloyd's mirror effects at the water's surface. Furthermore, sea otters tend to inhabit shallow coastal habitats where large seismic survey vessels towing large spreads of streamers may be unable to operate. TTS is also considered unlikely to occur in sirenians as a result of exposure to sounds from a seismic survey. They, like sea otters, tend to inhabit shallow coastal habitats and rarely range far from shore, whereas seismic survey vessels towing large arrays of airguns and (usually) even larger arrays of streamers normally must remain farther offshore because of equipment clearance and maneuverability limitations. Exposures of sea otters and sirenians to seismic surveys are more likely to involve smaller seismic sources that can be used in shallow and confined waters. The impacts of these are inherently less than would occur from a larger source of the types often used farther offshore.

Likelihood of Incurring TTS.—Most cetaceans show some degree of avoidance of seismic vessels operating an airgun array (see above). It is unlikely that these cetaceans would be exposed to airgun pulses at a sufficiently high level for a sufficiently long period to cause more than mild TTS, given the relative movement of the vessel and the marine mammal. TTS would be more likely in any odontocetes that bow- or wake-ride or otherwise linger near the airguns. However, while bow- or wake-riding, odontocetes would be at the surface and thus not exposed to strong sound pulses given the pressure-release and Lloyd Mirror effects at the surface. But if bow- or wake-riding animals were to dive intermittently near airguns, they would be exposed to strong sound pulses, possibly repeatedly.

If some cetaceans did incur mild or moderate TTS through exposure to airgun sounds in this manner, this would very likely be a temporary and reversible phenomenon. However, even a temporary reduction in hearing sensitivity could be deleterious in the event that, during that period of reduced sensitivity, a marine mammal needed its full hearing sensitivity to detect approaching predators, or for some other reason.

Some pinnipeds show avoidance reactions to airguns, but their avoidance reactions are generally not as strong or consistent as those of cetaceans. Pinnipeds occasionally seem to be attracted to operating seismic vessels. There are no specific data on TTS thresholds of pinnipeds exposed to single or multiple low-frequency pulses. However, given the indirect indications of a lower TTS threshold for the harbor seal than for odontocetes exposed to impulse sound (see above), it is possible that some pinnipeds close to a large airgun array could incur TTS.

NMFS (1995, 2000) concluded that cetaceans should not be exposed to pulsed underwater noise at received levels >180 dB re $1 \mu\text{Pa}_{\text{rms}}$. The corresponding limit for pinnipeds has been set by NMFS at 190 dB, although the HESS Team (HESS 1999) recommended a 180-dB limit for pinnipeds in California. The 180 and 190 dB re $1 \mu\text{Pa}_{\text{rms}}$ levels have not been considered to be the levels above which TTS might occur. Rather, they were the received levels above which, in the view of a panel of bioacoustics specialists convened by NMFS before TTS measurements for marine mammals started to become available, one could not be certain that there would be no injurious effects, auditory or otherwise, to marine mammals. As summarized above, data that are now available imply that TTS is unlikely to occur in various odontocetes (and probably mysticetes as well) unless they are exposed to a sequence of several airgun pulses stronger than 190 dB re $1 \mu\text{Pa}_{\text{rms}}$. On the other hand, for the harbor seal, harbor porpoise, and perhaps

some other species, TTS may occur upon exposure to one or more airgun pulses whose received level equals the NMFS “do not exceed” value of 190 dB re 1 $\mu\text{Pa}_{\text{rms}}$. That criterion corresponds to a single-pulse SEL of 175–180 dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$ in typical conditions, whereas TTS is suspected to be possible in harbor seals and harbor porpoises with a cumulative SEL of ~ 171 and ~ 164 dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$, respectively.

It has been shown that most large whales and many smaller odontocetes (especially the harbor porpoise) show at least localized avoidance of ships and/or seismic operations (see above). Even when avoidance is limited to the area within a few hundred meters of an airgun array, that should usually be sufficient to avoid TTS based on what is currently known about thresholds for TTS onset in cetaceans. In addition, ramping up airgun arrays, which is standard operational protocol for many seismic operators, should allow cetaceans near the airguns at the time of startup (if the sounds are aversive) to move away from the seismic source and to avoid being exposed to the full acoustic output of the airgun array (see above). Thus, most baleen whales likely will not be exposed to high levels of airgun sounds provided the ramp-up procedure is applied. Likewise, many odontocetes close to the trackline are likely to move away before the sounds from an approaching seismic vessel become sufficiently strong for there to be any potential for TTS or other hearing impairment. Therefore, there is little potential for baleen whales or odontocetes that show avoidance of ships or airguns to be close enough to an airgun array to experience TTS. In the event that a few individual cetaceans did incur TTS through exposure to strong airgun sounds, this is a temporary and reversible phenomenon unless the exposure exceeds the TTS-onset threshold by a sufficient amount for PTS to be incurred (see below). If TTS but not PTS were incurred, it would most likely be mild, in which case recovery is expected to be quick (probably within minutes).

6.2 Permanent Threshold Shift (PTS)

When PTS occurs, there is physical damage to the sound receptors in the ear. In some cases, there can be total or partial deafness, whereas in other cases, the animal has an impaired ability to hear sounds in specific frequency ranges (Kryter 1985). Physical damage to a mammal’s hearing apparatus can occur if it is exposed to sound impulses that have very high peak pressures, especially if they have very short rise times. (Rise time is the interval required for sound pressure to increase from the baseline pressure to peak pressure.)

There is no specific evidence that exposure to pulses of airgun sound can cause PTS in any marine mammal, even with large arrays of airguns. However, given the likelihood that some mammals close to an airgun array might incur at least mild TTS (see above), there has been further speculation about the possibility that some individuals occurring very close to airguns might incur PTS (e.g., Richardson et al. 1995, p. 372ff; Gedamke et al. 2008). Single or occasional occurrences of mild TTS are not indicative of permanent auditory damage, but repeated or (in some cases) single exposures to a level well above that causing TTS onset might elicit PTS.

Relationships between TTS and PTS thresholds have not been studied in marine mammals, but are assumed to be similar to those in humans and other terrestrial mammals (Southall et al. 2007). Based on data from terrestrial mammals, a precautionary assumption is that the PTS threshold for impulse sounds (such as airgun pulses as received close to the source) is at least 6 dB higher than the TTS threshold on a peak-pressure basis, and probably >6 dB higher (Southall et al. 2007). The low-to-moderate levels of TTS that have been induced in captive odontocetes and pinnipeds during controlled studies of TTS have been confirmed to be temporary, with no measurable residual PTS (Kastak et al. 1999; Schlundt et al. 2000; Finneran et al. 2002, 2005; Nachtigall et al. 2003, 2004). However, very prolonged exposure to sound strong enough to elicit TTS, or shorter-term exposure to sound levels well above the TTS threshold, can cause PTS, at least in terrestrial mammals (Kryter 1985). In terrestrial mammals, the received sound level from a single non-impulsive sound exposure must be far above the TTS threshold for

any risk of permanent hearing damage (Kryter 1994; Richardson et al. 1995; Southall et al. 2007). However, there is special concern about strong sounds whose pulses have very rapid rise times. In terrestrial mammals, there are situations when pulses with rapid rise times (e.g., from explosions) can result in PTS even though their peak levels are only a few dB higher than the level causing slight TTS. The rise time of airgun pulses is fast, but not as fast as that of an explosion.

Some factors that contribute to onset of PTS, at least in terrestrial mammals, are as follows:

- exposure to single very intense sound,
- fast rise time from baseline to peak pressure,
- repetitive exposure to intense sounds that individually cause TTS but not PTS, and
- recurrent ear infections or (in captive animals) exposure to certain drugs.

Cavanagh (2000) reviewed the thresholds used to define TTS and PTS. Based on this review and SACLANT (1998), it is reasonable to assume that PTS might occur at a received sound level 20 dB or more above that inducing mild TTS. However, for PTS to occur at a received level only 20 dB above the TTS threshold, the animal probably would have to be exposed to a strong sound for an extended period, or to a strong sound with rather rapid rise time.

More recently, Southall et al. (2007) estimated that received levels would need to exceed the TTS threshold by at least 15 dB, on an SEL basis, for there to be risk of PTS. Thus, for cetaceans exposed to a sequence of sound pulses, they estimate that the PTS threshold might be an M-weighted SEL (for the sequence of received pulses) of ~ 198 dB re $1 \mu\text{Pa}^2 \cdot \text{s}$ (15 dB higher than the M_{mf} -weighted TTS threshold, in a beluga, for a watergun impulse). Additional assumptions had to be made to derive a corresponding estimate for pinnipeds, as the only available data on TTS-thresholds in pinnipeds pertained to non-impulse sound (see above). Southall et al. (2007) estimated that the PTS threshold could be a cumulative M_{pw} -weighted SEL of ~ 186 dB re $1 \mu\text{Pa}^2 \cdot \text{s}$ in the case of a harbor seal exposed to impulse sound. The PTS threshold for the California sea lion and northern elephant seal would probably be higher given the higher TTS thresholds in those species. Southall et al. (2007) also note that, regardless of the SEL, there is concern about the possibility of PTS if a cetacean or pinniped received one or more pulses with peak pressure exceeding 230 or 218 dB re $1 \mu\text{Pa}$, respectively. Thus, PTS might be expected upon exposure of cetaceans to either SEL ≥ 198 dB re $1 \mu\text{Pa}^2 \cdot \text{s}$ or peak pressure ≥ 230 dB re $1 \mu\text{Pa}$. Corresponding proposed dual criteria for pinnipeds (at least harbor seals) are ≥ 186 dB SEL and ≥ 218 dB peak pressure (Southall et al. 2007). These estimates are all first approximations, given the limited underlying data, assumptions, species differences, and evidence that the “equal energy” model is not be entirely correct.

Sound impulse duration, peak amplitude, rise time, number of pulses, and inter-pulse interval are the main factors thought to determine the onset and extent of PTS. Ketten (1994) has noted that the criteria for differentiating the sound pressure levels that result in PTS (or TTS) are location and species-specific. PTS effects may also be influenced strongly by the health of the receiver’s ear.

As described above for TTS, in estimating the amount of sound energy required to elicit the onset of TTS (and PTS), it is assumed that the auditory effect of a given cumulative SEL from a series of pulses is the same as if that amount of sound energy were received as a single strong sound. There are no data from marine mammals concerning the occurrence or magnitude of a potential partial recovery effect between pulses. In deriving the estimates of PTS (and TTS) thresholds quoted here, Southall et al. (2007) made the precautionary assumption that no recovery would occur between pulses.

The TTS section (above) concludes that exposure to several strong seismic pulses that each have flat-weighted received levels near 190 dB re $1 \mu\text{Pa}_{\text{rms}}$ (175–180 dB re $1 \mu\text{Pa}^2 \cdot \text{s}$ SEL) could result in cumulative exposure of ~ 186 dB SEL (flat-weighted) or ~ 183 dB SEL (M_{mf} -weighted), and thus slight

TTS in a small odontocete. Allowing for the assumed 15 dB offset between PTS and TTS thresholds, expressed on an SEL basis, exposure to several strong seismic pulses that each have flat-weighted received levels near 205 dB_{rms} (190–195 dB SEL) could result in cumulative exposure of ~198 dB SEL (M_{mf} -weighted), and thus slight PTS in a small odontocete. However, the levels of successive pulses that will be received by a marine mammal that is below the surface as a seismic vessel approaches, passes and moves away will tend to increase gradually and then decrease gradually, with periodic decreases superimposed on this pattern when the animal comes to the surface to breathe. To estimate how close an odontocete's CPA distance would have to be for the cumulative SEL to exceed 198 dB SEL (M_{mf} -weighted), one would (as a minimum) need to allow for the sequence of distances at which airgun shots would occur, and for the dependence of received SEL on distance in the region of the seismic operation (e.g., Erbe and King 2009).

It is unlikely that an odontocete would remain close enough to a large airgun array for sufficiently long to incur PTS. There is some concern about bowriding odontocetes, but for animals at or near the surface, auditory effects are reduced by Lloyd's mirror and surface release effects. The presence of the vessel between the airgun array and bow-riding odontocetes could also, in some but probably not all cases, reduce the levels received by bow-riding animals (e.g., Gabriele and Kipple 2009). The TTS (and thus PTS) thresholds of baleen whales are unknown but, as an interim measure, assumed to be no lower than those of odontocetes. Also, baleen whales generally avoid the immediate area around operating seismic vessels, so it is unlikely that a baleen whale could incur PTS from exposure to airgun pulses. The TTS (and thus PTS) thresholds of some pinnipeds (e.g., harbor seal) as well as the harbor porpoise may be lower (Kastak et al. 2005; Southall et al. 2007; Lucke et al. 2009). If so, TTS and potentially PTS may extend to a somewhat greater distance for those animals. Again, Lloyd's mirror and surface release effects will ameliorate the effects for animals at or near the surface.

Although it is unlikely that airgun operations during most seismic surveys would cause PTS in many marine mammals, caution is warranted given

- the limited knowledge about noise-induced hearing damage in marine mammals, particularly baleen whales, pinnipeds, and sea otters;
- the seemingly greater susceptibility of certain species (e.g., harbor porpoise and harbor seal) to TTS and presumably also PTS; and
- the lack of knowledge about TTS and PTS thresholds in many species, including various species closely related to the harbor porpoise and harbor seal.

The avoidance reactions of many marine mammals, along with commonly-applied monitoring and mitigation measures (visual and passive acoustic monitoring, ramp ups, and power downs or shut downs when mammals are detected within or approaching the "safety radii"), would reduce the already-low probability of exposure of marine mammals to sounds strong enough to induce PTS.

6.3 Strandings and Mortality

Marine mammals close to underwater detonations of high explosives can be killed or severely injured, and the auditory organs are especially susceptible to injury (Ketten et al. 1993; Ketten 1995). However, explosives are no longer used in marine waters for commercial seismic surveys or (with rare exceptions) for seismic research; they have been replaced by airguns and other non-explosive sources. Airgun pulses are less energetic and have slower rise times, and there is no specific evidence that they can cause serious injury, death, or stranding even in the case of large airgun arrays. However, the association of mass strandings of beaked whales with naval exercises and, in one case, a seismic survey (Malakoff 2002; Cox et al. 2006), has raised the possibility that beaked whales exposed to strong "pulsed" sounds

may be especially susceptible to injury and/or behavioral reactions that can lead to stranding (e.g., Hildebrand 2005; Southall et al. 2007). Hildebrand (2005) reviewed the association of cetacean strandings with high-intensity sound events and found that deep-diving odontocetes, primarily beaked whales, were by far the predominant (95%) cetaceans associated with these events, with 2% mysticete whales (minke). However, as summarized below, there is no definitive evidence that airguns can lead to injury, strandings, or mortality even for marine mammals in close proximity to large airgun arrays.

Specific sound-related processes that lead to strandings and mortality are not well documented, but may include (1) swimming in avoidance of a sound into shallow water; (2) a change in behavior (such as a change in diving behavior that might contribute to tissue damage, gas bubble formation, hypoxia, cardiac arrhythmia, hypertensive hemorrhage or other forms of trauma; (3) a physiological change such as a vestibular response leading to a behavioral change or stress-induced hemorrhagic diathesis, leading in turn to tissue damage; and (4) tissue damage directly from sound exposure, such as through acoustically mediated bubble formation and growth or acoustic resonance of tissues. Some of these mechanisms are unlikely to apply in the case of impulse sounds. However, there are increasing indications that gas-bubble disease (analogous to “the bends”), induced in supersaturated tissue by a behavioral response to acoustic exposure, could be a pathologic mechanism for the strandings and mortality of some deep-diving cetaceans exposed to sonar. The evidence for this remains circumstantial and associated with exposure to naval mid-frequency sonar, not seismic surveys (Cox et al. 2006; Southall et al. 2007).

Seismic pulses and mid-frequency sonar signals are quite different, and some mechanisms by which sonar sounds have been hypothesized to affect beaked whales are unlikely to apply to airgun pulses. Sounds produced by airgun arrays are broadband impulses with most of the energy below 1 kHz. Typical military mid-frequency sonars emit non-impulse sounds at frequencies of 2–10 kHz, generally with a relatively narrow bandwidth at any one time (though the frequency may change over time). Thus, it is not appropriate to assume that the effects of seismic surveys on beaked whales or other species would be the same as the apparent effects of military sonar. For example, resonance effects (Gentry 2002) and acoustically-mediated bubble-growth (Crum et al. 2005) are implausible in the case of exposure to broadband airgun pulses. Nonetheless, evidence that sonar signals can, in special circumstances, lead (at least indirectly) to physical damage and mortality (e.g., Balcomb and Claridge 2001; NOAA and USN 2001; Jepson et al. 2003; Fernández et al. 2004, 2005; Hildebrand 2005; Cox et al. 2006) suggests that caution is warranted when dealing with exposure of marine mammals to any high-intensity “pulsed” sound. One of the hypothesized mechanisms by which naval sonars lead to strandings might, in theory, also apply to seismic surveys: If the strong sounds sometimes cause deep-diving species to alter their surfacing–dive cycles in a way that causes bubble formation in tissue, that hypothesized mechanism might apply to seismic surveys as well as mid-frequency naval sonars. However, there is no specific evidence of this upon exposure to airgun pulses.

There is no conclusive evidence of cetacean strandings or deaths at sea as a result of exposure to seismic surveys, but a few cases of strandings in the general area where a seismic survey was ongoing have led to speculation concerning a possible link between seismic surveys and strandings. • Suggestions that there was a link between seismic surveys and strandings of humpback whales in Brazil (Engel et al. 2004) were not well founded (IAGC 2004; IWC 2007). • In Sept. 2002, there was a stranding of two Cuvier’s beaked whales in the Gulf of California, Mexico, when the L-DEO seismic vessel R/V *Maurice Ewing* was operating a 20-airgun, 8490-in³ airgun array in the general area. The evidence linking the stranding to the seismic survey was inconclusive and not based on any physical evidence (Hogarth 2002; Yoder 2002). The ship was also operating its multibeam echosounder at the same time, but this had much less potential than the aforementioned naval sonars to affect beaked whales, given its downward-directed beams, much shorter pulse durations, and lower duty cycle. Nonetheless, the Gulf of California incident

plus the beaked whale strandings near naval exercises involving use of mid-frequency sonar suggest a need for caution in conducting seismic surveys in areas occupied by beaked whales until more is known about effects of seismic surveys on those species (Hildebrand 2005).

6.4 Non-Auditory Physiological Effects

Based on evidence from terrestrial mammals and humans, sound is a potential source of stress (Wright and Kuczaj 2007; Wright et al. 2007a,b, 2009). However, almost no information is available on sound-induced stress in marine mammals, or on its potential (alone or in combination with other stressors) to affect the long-term well-being or reproductive success of marine mammals (Fair and Becker 2000; Hildebrand 2005; Wright et al. 2007a,b). Such long-term effects, if they occur, would be mainly associated with chronic noise exposure, which is characteristic of some seismic surveys and exposure situations (McCauley et al. 2000a:62ff; Nieuwkerk et al. 2009) but not of some others.

Available data on potential stress-related impacts of anthropogenic noise on marine mammals are extremely limited, and additional research on this topic is needed. We know of only two specific studies of noise-induced stress in marine mammals. (1) Romano et al. (2004) examined the effects of single underwater impulse sounds from a seismic water gun (source level up to 228 dB re 1 $\mu\text{Pa} \cdot \text{m}_{\text{p-p}}$) and single short-duration pure tones (sound pressure level up to 201 dB re 1 μPa) on the nervous and immune systems of a beluga and a bottlenose dolphin. They found that neural-immune changes to noise exposure were minimal. Although levels of some stress-released substances (e.g., catecholamines) changed significantly with exposure to sound, levels returned to baseline after 24 hr. (2) During playbacks of recorded drilling noise to four captive beluga whales, Thomas et al. (1990) found no changes in blood levels of stress-related hormones. Long-term effects were not measured, and no short-term effects were detected. For both studies, caution is necessary when extrapolating these results to wild animals and to real-world situations given the small sample sizes, use of captive animals, and other technical limitations of the two studies.

Aside from stress, other types of physiological effects that might, in theory, be involved in beaked whale strandings upon exposure to naval sonar (Cox et al. 2006), such as resonance and gas bubble formation, have not been demonstrated and are not expected upon exposure to airgun pulses (see preceding subsection). If seismic surveys disrupt diving patterns of deep-diving species, this might perhaps result in bubble formation and a form of “the bends”, as speculated to occur in beaked whales exposed to sonar. However, there is no specific evidence that exposure to airgun pulses has this effect.

In summary, very little is known about the potential for seismic survey sounds (or other types of strong underwater sounds) to cause non-auditory physiological effects in marine mammals. Such effects, if they occur at all, would presumably be limited to short distances and to activities that extend over a prolonged period. The available data do not allow identification of a specific exposure level above which non-auditory effects can be expected (Southall et al. 2007), or any meaningful quantitative predictions of the numbers (if any) of marine mammals that might be affected in these ways.

7. Literature Cited

- Akamatsu, T., Y. Hatakeyama, and N. Takatsu. 1993. Effects of pulsed sounds on escape behavior of false killer whales. **Nipp. Suis. Gakkaishi** 59(8):1297-1303.
- Angliss, R.P. and R.B. Outlaw. 2008. Alaska marine mammal stock assessments, 2007. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-180. 252 p.
- Anonymous. 1975. Phantom killer whales. **S. Afr. Ship. News & Fishing Indus. Rev.** 30(7):50-53.

- Arnold, B.W. 1996. Visual monitoring of marine mammal activity during the Exxon 3-D seismic survey: Santa Ynez unit, offshore California 9 November to 12 December 1995. Rep. from Impact Sciences Inc., San Diego, CA, for Exxon Co., U.S.A., Thousand Oaks, CA. 20 p.
- Au, W.W.L. 1993. *The Sonar of Dolphins*. Springer-Verlag, New York, NY. 277 p.
- Au, W.W.L., A.N. Popper, and R.R. Fay. 2000. Hearing by Whales and Dolphins. *Springer Handbook of Auditory Res.* Vol. 12. Springer-Verlag, New York, NY. 458 p.
- Au, W.W.L., A.A. Pack, M.O. Lammers, L.M. Herman, M.H. Deakos, and K. Andrews. 2006. Acoustic properties of humpback whale songs. **J. Acoust. Soc. Am.** 120(2):1103-1110.
- Backus, R.H. and W.E. Schevill. 1966. *Physeter* clicks. p. 510-528 in K.S. Norris (ed.), *Whales, dolphins, and porpoises*. Univ. Calif. Press, Berkeley, CA. 789 p
- Bain, D.E. and R. Williams. 2006. Long-range effects of airgun noise on marine mammals: responses as a function of received sound level and distance. Paper SC/58/E35 presented to the IWC Scient. Commit., IWC Annu. Meet., 1-13 June, St. Kitts.
- Baird, R.W. 2005. Sightings of dwarf (*Kogia sima*) and pygmy (*K. breviceps*) sperm whales from the main Hawaiian Islands. **Pacific Sci.** 59(3):461-466.
- Baird, R.W., D.L. Webster, D.J. McSweeney, A.D. Ligon, G.S. Schorr, and J. Barlow. 2006. Diving behavior and ecology of Cuvier's (*Ziphius cavirostris*) and Blainville's (*Mesoplodon densirostris*) beaked whales in Hawaii. **Can. J. Zool.** 84(8):1120-1128.
- Balcomb, K.C., III and D.E. Claridge. 2001. A mass stranding of cetaceans caused by naval sonar in the Bahamas. **Bahamas J. Sci.** 8(2):2-12.
- Barkaszi, M.J., D.M. Epperson, and B. Bennett. 2009. Six-year compilation of cetacean sighting data collected during commercial seismic survey mitigation observations throughout the Gulf of Mexico, USA. p. 24-25 *In: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Canada, Oct. 2009.* 306 p.
- Barlow, J. and R. Gisiner. 2006. Mitigating, monitoring and assessing the effects of anthropogenic sound on beaked whales. **J. Cetac. Res. Manage.** 7(3):239-249.
- Bauer, G.B., J.C. Gaspard, K. Dziuk, A. Cardwell, L. Read, R.L. Reep, and D.A. Mann. 2009. The manatee audiogram and auditory critical ratios. p. 27-28 *In: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Canada, Oct. 2009.* 306 p.
- Beale, C.M. and P. Monaghan. 2004. Behavioural responses to human disturbance: a matter of choice? **Anim. Behav.** 68(5):1065-1069.
- Beland, J.A., B. Haley, C.M. Reiser, D.M. Savarese, D.S. Ireland and D.W. Funk. 2009. Effects of the presence of other vessels on marine mammal sightings during multi-vessel operations in the Alaskan Chukchi Sea. p. 29 *In: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Oct. 2009:*29. 306 p.
- Berta, A., R. Racicot and T. Deméré. 2009. The comparative anatomy and evolution of the ear in *Balaenoptera* mysticetes. p. 33 *In: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Oct. 2009.* 306 p.
- Blackwell, S.B., R.G. Norman, C.R. Greene Jr., and W.J. Richardson. 2007. Acoustic measurements. p. 4-1 to 4-52 *In: Marine mammal monitoring and mitigation during open water seismic exploration by Shell Offshore Inc. in the Chukchi and Beaufort Seas, July-September 2006: 90-day report.* LGL Rep. P891-1. Rep. from LGL Alaska Res. Assoc. Inc., Anchorage, AK, and Greeneridge Sciences Inc., Santa Barbara, CA, for Shell Offshore Inc., Houston, TX, Nat. Mar. Fish. Serv., Silver Spring, MD, and U.S. Fish & Wildl. Serv., Anchorage, AK. 199 p.
- Blackwell, S.B., C.R. Greene, T.L. McDonald, C.S. Nations, R.G. Norman, and A. Thode. 2009a. Beaufort Sea bowhead whale migration route study. Chapter 8 *In: D.S. Ireland, D.W. Funk, R. Rodrigues, and W.R. Koski (eds.). 2009. Joint Monitoring Program in the Chukchi and Beaufort seas, open water seasons, 2006-2007.* LGL Alaska Rep. P971-2. Rep. from LGL Alaska Res. Assoc. Inc. (Anchorage, AK) et al. for Shell Offshore Inc. (Anchorage, AK) et al. 485 p. plus appendices.

- Blackwell, S.B., C.S. Nations, T.L. McDonald, A.M. Thode, K.H. Kim, C.R. Greene, and M.A. Macrander. 2009b. Effects of seismic exploration activities on the calling behavior of bowhead whales in the Alaskan Beaufort Sea. p. 35 *In: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Canada, 12-16 Oct. 2009.* 306 p.
- Bowles, A.E., M. Smultea, B. Würsig, D.P. DeMaster, and D. Palka. 1994. Relative abundance and behavior of marine mammals exposed to transmissions from the Heard Island Feasibility Test. **J. Acoust. Soc. Am.** 96(4):2469-2484.
- Bullock, T.H., T.J. Oshea, and M.C. McClune. 1982. Auditory evoked-potentials in the West Indian manatee (*Sirenia, Trichechus manatus*). **J. Comp. Physiol.** 148(4):547-554.
- Britto, M.K. and A. Silva Barreto. 2009. Marine mammal diversity registered on seismic surveys in Brazil, between 2000 and 2008. p. 41 *In: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Oct. 2009.* 306 p.
- Brodie, P.F. 1981. Energetic and behavioural considerations with respect to marine mammals and disturbance from underwater noise. p. 287-290 *In: N.M. Peterson (ed.), The question of sound from icebreaker operations: Proceedings of a workshop. Arctic Pilot Proj., Petro-Canada, Calgary, Alb.* 350 p.
- Burgess, W.C. and C.R. Greene, Jr. 1999. Physical acoustics measurements. p. 3-1 to 3-63 *In: W.J. Richardson (ed.), Marine mammal and acoustical monitoring of Western Geophysical's open-water seismic program in the Alaskan Beaufort Sea, 1998. LGL Rep. TA22303. Rep. from LGL Ltd., King City, Ont., and Greeneridge Sciences Inc., Santa Barbara, CA, for Western Geophysical, Houston, TX, and Nat. Mar. Fish. Serv., Anchorage, AK, and Silver Spring, MD.* 390 p.
- Calambokidis, J. and S.D. Osmek. 1998. Marine mammal research and mitigation in conjunction with air gun operation for the USGS 'SHIPS' seismic surveys in 1998. Rep. from Cascadia Res., Olympia, WA, for U.S. Geol. Surv., Nat. Mar. Fish. Serv., and Minerals Manage. Serv.
- Caldwell, J. and W. Dragoset. 2000. A brief overview of seismic air-gun arrays. **Leading Edge** 19(8):898-902.
- Cavanagh, R.C. 2000. Criteria and thresholds for adverse effects of underwater noise on marine animals. AFRL-HE-WP-TR-2000-0092. Rep. from Science Applications Intern. Corp., McLean, VA, for Air Force Res. Lab., Wright-Patterson AFB, OH.
- Christie, K., C. Lyons, W.R. Koski, D.S. Ireland, and D.W. Funk. 2009. Patterns of bowhead whale occurrence and distribution during marine seismic operations in the Alaskan Beaufort Sea. p. 55 *In: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Canada, 12-16 Oct. 2009.*
- Citta, J.J., L.T. Quakenbush, R.J. Small, and J.C. George. 2007. Movements of a tagged bowhead whale in the vicinity of a seismic survey in the Beaufort Sea. Poster Paper, Soc. Mar. Mammal. 17th Bienn. Meet., Cape Town, South Africa.
- Clark, C.W. and W.T. Ellison. 2004. Potential use of low-frequency sounds by baleen whales for probing the environment: Evidence from models and empirical measurements. p. 564-589 *In: J.A. Thomas, C.F. Moss and M. Vater (eds.), Echolocation in Bats and Dolphins. Univ. Chicago Press, Chicago, IL.* 604 p.
- Clark, C.W. and G.C. Gagnon. 2006. Considering the temporal and spatial scales of noise exposures from seismic surveys on baleen whales. Intern. Whal. Commis. Working Pap. SC/58/E9. 9 p.
- Cook, M.L.H., R.A. Varela, J.D. Goldstein, S.D. McCulloch, G.D. Bossart, J.J. Finneran, D. Houser, and A. Mann. 2006. Beaked whale auditory evoked potential hearing measurements. **J. Comp. Physiol.** A 192:489-495.
- Cox, T.M., T.J. Ragen, A.J. Read, E. Vos, R.W. Baird, K. Balcomb, J. Barlow, J. Caldwell, T. Cranford, L. Crum, A. D'Amico, G. D'Spain, A. Fernández, J. Finneran, R. Gentry, W. Gerth, F. Gulland, J. Hildebrand, D. Houser, R. Hullar, P.D. Jepson, D. Ketten, C.D. Macleod, P. Miller, S. Moore, D.C. Mountain, D. Palka, P. Ponganis, S. Rommel, T. Rowles, B. Taylor, P. Tyack, D. Wartzok, R. Gisiner, J. Meads, and L. Benner. 2006. Understanding the impacts of anthropogenic sound on beaked whales. **J. Cetac. Res. Manage.** 7(3):177-187.
- Crum, L.A., M.R. Bailey, J. Guan, P.R. Hilmo, S.G. Kargl, and T.J. Matula. 2005. Monitoring bubble growth in supersaturated blood and tissue ex vivo and the relevance to marine mammal bioeffects. **Acoustic Res. Lett. Online** 6(3):214-220.

- Dahlheim, M.E. 1987. Bio-acoustics of the gray whale (*Eschrichtius robustus*). Ph.D. Dissertation, Univ. British Columbia, Vancouver, BC. 315 p.
- DeRuiter, S.L., P.L. Tyack, Y.-T. Lin, A.E. Newhall, J.F. Lynch, and P.J.O. Miller. 2006. Modeling acoustic propagation of airgun array pulses recorded on tagged sperm whales (*Physeter macrocephalus*). **J. Acoust. Soc. Am.** 120(6):4100-4114.
- Di Iorio, L. and C.W. Clark. 2009. Exposure to seismic survey alters blue whale acoustic communication. **Biol. Lett.** doi: 10.1098/rsbl.2009.0651.
- Dolman, S.J. and M.P. Simmonds. 2006. An updated note on the vulnerability of cetaceans to acoustic disturbance. Paper SC/58/E22 presented to the IWC Scient. Commit., IWC Annu. Meet., 1-13 June, St. Kitts.
- Duncan, P.M. 1985. Seismic sources in a marine environment. p. 56-88 *In: Proceedings of the Workshop on Effects of Explosives Use in the Marine Environment*, Jan. 1985, Halifax, N.S. Tech. Rep. 5. Can. Oil & Gas Lands Admin., Environ. Prot. Branch, Ottawa, Ont.
- Dunn, R.A. and O. Hernandez. 2009. Tracking blue whales in the eastern tropical Pacific with an ocean-bottom seismometer and hydrophone array. **J. Acoust. Soc. Am.** 126(3):1084-1094.
- Engel, M.H., M.C.C. Marcondes, C.C.A. Martins, F.O. Luna, R.P. Lima, and A. Campos. 2004. Are seismic surveys responsible for cetacean strandings? An unusual mortality of adult humpback whales in Abrolhos Bank, northeastern coast of Brazil. Paper SC/56/E28 presented to the IWC Scient. Commit., IWC Annu. Meet., 19-22 July, Sorrento, Italy.
- Erbe, C. and A.R. King. 2009. Modeling cumulative sound exposure around marine seismic surveys. **J. Acoust. Soc. Am.** 125(4):2443-2451.
- Fair, P.A. and P.R. Becker. 2000. Review of stress in marine mammals. **J. Aquat. Ecosyst. Stress Recov.** 7:335-354.
- Fernández, A., M. Arbelo, R. Deaville, I.A.P. Patterson, P. Castro, J.R. Baker, E. Degollada, H.M. Ross, P. Herráez, A.M. Pocknell, E. Rodríguez, F.E. Howie, A. Espinosa, R.J. Reid, J.R. Jaber, V. Martin, A.A. Cunningham, and P.D. Jepson. 2004. Pathology: whales, sonar and decompression sickness (reply). **Nature** 428(6984, 15 Apr.). doi: 10.1038/nature02528a.
- Fernández, A., J.F. Edwards, F. Rodriguez, A.E. de los Monteros, P. Herráez, P. Castro, J.R. Jaber, V. Martin, and M. Arbelo. 2005. "Gas and fat embolic syndrome" involving a mass stranding of beaked whales (Family Ziphiidae) exposed to anthropogenic sonar signals. **Veterin. Pathol.** 42(4):446-457.
- Finneran, J.J. and C.E. Schlundt. 2004. Effects of intense pure tones on the behavior of trained odontocetes. Tech. Rep. 1913. Space and Naval Warfare (SPAWAR) Systems Center, San Diego, CA. 15 p.
- Finneran, J.J., C.E. Schlundt, D.A. Carder, J.A. Clark, J.A. Young, J.B. Gaspin, and S.H. Ridgway. 2000. Auditory and behavioral responses of bottlenose dolphins (*Tursiops truncatus*) and beluga whale (*Delphinapterus leucas*) to impulsive sounds resembling distant signatures of underwater explosions. **J. Acoust. Soc. Am.** 108(1):417-431.
- Finneran, J.J., C.E. Schlundt, R. Dear, D.A. Carder, and S.H. Ridgway. 2002. Temporary shift in masked hearing thresholds in odontocetes after exposure to single underwater impulses from a seismic watergun. **J. Acoust. Soc. Am.** 111(6):2929-2940.
- Finneran, J.J., R. Dear, D.A. Carder, and S.H. Ridgway. 2003. Auditory and behavioral responses of California sea lions (*Zalophus californianus*) to single underwater impulses from an arc-gap transducer. **J. Acoust. Soc. Am.** 114(3):1667-1677.
- Finneran, J.J., D.A. Carder, C.E. Schlundt, and S.H. Ridgway. 2005. Temporary threshold shift in bottlenose dolphins (*Tursiops truncatus*) exposed to mid-frequency tones. **J. Acoust. Soc. Am.** 118(4):2696-2705.
- Finneran, J.J., D.S. Houser, B. Mase-Guthrie, R.Y. Ewing and R.G. Lingenfelter. 2009. Auditory evoked potentials in a stranded Gervais' beaked whale (*Mesoplodon europaeus*). **J. Acoust. Soc. Am.** 126(1):484-490.
- Fish, J.F. and J.S. Vania. 1971. Killer whale, *Orcinus orca*, sounds repel white whales, *Delphinapterus leucas*. **Fish. Bull.** 69(3):531-535.

- Fox, C.G., R.P. Dziak, and H. Matsumoto. 2002. NOAA efforts in monitoring of low-frequency sound in the global ocean. **J. Acoust. Soc. Am.** 112(5, Pt. 2):2260 (Abstract).
- Frankel, A. 2005. Gray whales hear and respond to a 21-25 kHz high-frequency whale-finding sonar. p. 97 *In*: Abstr. 16th Bienn. Conf. Biol. Mar. Mamm., San Diego, CA, Dec. 2005. 306 p.
- Frantzis, A. 1998. Does acoustic testing strand whales? **Nature** 392(6671):29.
- Frost, K.J., L.F. Lowry, and R.R. Nelson. 1984. Belukha whale studies in Bristol Bay, Alaska. p. 187-200 *In*: B.R. Melteff and D.H. Rosenberg (eds.), Proceedings of the Workshop on Biological Interactions among Marine Mammals and Commercial Fisheries in the Southeastern Bering Sea, Oct. 1983, Anchorage, AK. Univ. Alaska Sea Grant Rep. 84-1. Univ. Alaska, Fairbanks, AK.
- Gabriele, C.M. and B. Kipple. 2009. Measurements of near-surface, near-bow underwater sound from cruise ships. p. 86 *In*: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Oct. 2009. 306 p.
- Gailey, G., B. Würsig, and T.L. McDonald. 2007. Abundance, behavior, and movement patterns of western gray whales in relation to a 3-D seismic survey, northeast Sakhalin Island, Russia. **Environ. Monit. Assessm.** 134(1-3):75-91.
- Gedamke, J., S. Frydman, and N. Gales. 2008. Risk of baleen whale hearing loss from seismic surveys: preliminary results from simulations accounting for uncertainty and individual variation. Intern. Whal. Comm. Working Pap. SC/60/E9. 10 p.
- Gentry, R. (ed.). 2002. Report of the workshop on acoustic resonance as a source of tissue trauma in cetaceans. 24-25 April, Nat. Mar. Fish. Serv., Silver Spring, MD. 19 p. Available at <http://www.nmfs.noaa.gov/pr/acoustics/reports.htm>
- Gerstein, E.R., L.A. Gerstein, S.E. Forsythe, and J.E. Blue. 1999. The underwater audiogram of a West Indian manatee (*Trichechus manatus*). **J. Acoust. Soc. Am.** 105(6):3575-3583.
- Gerstein, E., L. Gerstein, S. Forsythe and J. Blue. 2004. Do manatees utilize infrasonic communication or detection? **J. Acoust. Soc. Am.** 115(5, Pt. 2):2554-2555 (Abstract).
- Ghoul, A., C. Reichmuth, and J. Mulsow. 2009. Source levels and spectral analysis of southern sea otter (*Enhydra lutris nereis*) scream vocalizations. p. 90 *In*: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Canada, Oct. 2009. 306 p.
- Gisiner, R.C. (ed.). 1999. Proceedings – Workshop on the Effects of Anthropogenic Noise in the Marine Environment, Bethesda, MD, 10-12 Feb. 1998. Office of Naval Res., Arlington, VA. Available (as of Nov. 2009) at http://www.onr.navy.mil/sci_tech/34/341/docs/proceed.pdf
- Goold, J.C. 1996a. Acoustic assessment of common dolphins off the West Wales coast, in conjunction with 16th round seismic surveying. Rep. from School of Ocean Sciences, Univ. Wales, Bangor, Wales, for Chevron UK Ltd., Repsol Exploration (UK) Ltd., and Aran Energy Exploration Ltd. 22 p.
- Goold, J.C. 1996b. Acoustic assessment of populations of common dolphin *Delphinus delphis* in conjunction with seismic surveying. **J. Mar. Biol. Assoc. U.K.** 76:811-820.
- Goold, J.C. 1996c. Acoustic cetacean monitoring off the west Wales coast. Rep. from School of Ocean Sciences, Univ. Wales, Bangor, Wales, for Chevron UK Ltd, Repsol Explor. (UK) Ltd, and Aran Energy Explor. Ltd. 20 p.
- Goold, J.C. and R.F.W. Coates. 2006. Near source, high frequency air-gun signatures. Paper SC/58/E30 presented to the IWC Scient. Commit., IWC Annu. Meet., 1-13 June, St. Kitts.
- Goold, J.C. and P.J. Fish. 1998. Broadband spectra of seismic survey air-gun emissions, with reference to dolphin auditory thresholds. **J. Acoust. Soc. Am.** 103(4):2177-2184.
- Gordon, J., D. Gillespie, J. Potter, A. Frantzis, M.P. Simmonds, R. Swift, and D. Thompson. 2004. A review of the effects of seismic surveys on marine mammals. **Mar. Technol. Soc. J.** 37(4):16-34.
- Gordon, J., R. Antunes, N. Jaquet and B. Würsig. 2006. An investigation of sperm whale headings and surface behaviour before, during and after seismic line changes in the Gulf of Mexico. Intern. Whal. Comm. Working Pap. SC/58/E45. 10 p.

- Gosselin, J.-F. and J. Lawson. 2004. Distribution and abundance indices of marine mammals in the Gully and two adjacent canyons of the Scotian Shelf before and during nearby hydrocarbon seismic exploration programmes in April and July 2003. Res. Doc. 2004/133. Can. Sci. Advis. Secretariat, Fisheries & Oceans Canada. 24 p. Available at http://www.dfo-mpo.gc.ca/csas/Csas/DocREC/2004/RES2004_133_e.pdf
- Greene, C.R., Jr. 1997. Physical acoustics measurements. p. 3-1 to 3-63 *In*: W.J. Richardson (ed.), Northstar marine mammal monitoring program, 1996: marine mammal and acoustical monitoring of a seismic program in the Alaskan Beaufort Sea. LGL Rep. 2121-2. Rep. from LGL Ltd., King City, Ont., and Greeneridge Sciences Inc., Santa Barbara, CA, for BP Explor. (Alaska) Inc., Anchorage, AK, and Nat. Mar. Fish. Serv., Anchorage, AK, and Silver Spring, MD. 245 p.
- Greene, C.R., Jr. and W.J. Richardson. 1988. Characteristics of marine seismic survey sounds in the Beaufort Sea. **J. Acoust. Soc. Am.** 83(6):2246-2254.
- Greene, G.D., F.R. Engelhardt, and R.J. Paterson (eds.). 1985. Proceedings of the Workshop on Effects of Explosives Use in the Marine Environment, Jan. 1985, Halifax, NS. Tech. Rep. 5. Can. Oil & Gas Lands Admin., Environ. Prot. Branch, Ottawa, Ont.
- Greene, C.R., Jr., N.S. Altman, and W.J. Richardson. 1999a. Bowhead whale calls. p. 6-1 to 6-23 *In*: W.J. Richardson (ed.), Marine mammal and acoustical monitoring of Western Geophysical's open-water seismic program in the Alaskan Beaufort Sea, 1998. LGL Rep. TA2230-3. Rep. from LGL Ltd., King City, Ont., and Greeneridge Sciences Inc., Santa Barbara, CA, for Western Geophysical, Houston, TX, and Nat. Mar. Fish. Serv., Anchorage, AK, and Silver Spring, MD. 390 p.
- Greene, C.R., Jr., N.S. Altman and W.J. Richardson. 1999b. The influence of seismic survey sounds on bowhead whale calling rates. **J. Acoust. Soc. Am.** 106(4, Pt. 2):2280 (Abstract).
- Guerra, M., A.M. Thode, S.B. Blackwell, C.R. Greene Jr. and M. Macrander. 2009. Quantifying masking effects of seismic survey reverberation off the Alaskan North Slope. **J. Acoust. Soc. Am.** 126(4, Pt. 2):2230 (Abstract).
- Gunn, L.M. 1988. A behavioral audiogram of the North American river otter (*Lutra canadensis*). M.S. thesis, San Diego State Univ., San Diego, CA. 40 p.
- Haley, B., and W.R. Koski. 2004. Marine mammal monitoring during Lamont-Doherty Earth Observatory's seismic program in the Northwest Atlantic Ocean, July–August 2004. LGL Rep. TA2822-27. Rep. from LGL Ltd., King City, Ont., for Lamont-Doherty Earth Observatory, Columbia Univ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD. November. 80 p.
- Hanser, S.F., L.R. Doyle, A.R. Szabo, F.A. Sharpe and B. McCowan. 2009. Bubble-net feeding humpback whales in Southeast Alaska change their vocalization patterns in the presence of moderate vessel noise. p. 105 *In*: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Canada, Oct. 2009. 306 p.
- Harris, R.E., G.W. Miller, and W.J. Richardson. 2001. Seal responses to airgun sounds during summer seismic surveys in the Alaskan Beaufort Sea. **Mar. Mamm. Sci.** 17:795-812.
- Harris, R.E., [R.E.] T. Elliott, and R.A. Davis. 2007. Results of mitigation and monitoring program, Beaufort Span 2-D marine seismic program, open-water season 2006. LGL Rep. TA4319-1. Rep. from LGL Ltd., King City, Ont., for GX Technol. Corp., Houston, TX. 48 p.
- Hauser, D.D.W., M Holst, and V.D. Moulton. 2008. Marine mammal and sea turtle monitoring during Lamont-Doherty Earth Observatory's marine seismic program in the Eastern Tropical Pacific, April–August 2008. LGL Rep. TA4656/7-1. Rep. from LGL Ltd., King City., Ont., and St. John's, Nfld, for Lamont-Doherty Earth Observatory of Columbia Univ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD. 98 p.
- HESS Team. 1999. High Energy Seismic Survey review process and interim operational guidelines for marine surveys offshore Southern California. Rep. from High Energy Seismic Survey Team for Calif. State Lands Commis. and Minerals Manage. Serv., Camarillo, CA. 39 p. + Appendices.
Available at www.mms.gov/omm/pacific/lease/fullhessrept.pdf

- Hildebrand, J.A. 2005. Impacts of anthropogenic sound. p. 101-124 *In*: J.E. Reynolds, W.F. Perrin, R.R. Reeves, S. Montgomery, and T. Ragen (eds.), *Marine Mammal Research: Conservation Beyond Crisis*. Johns Hopkins Univ. Press, Baltimore, MD. 223 p.
- Hogarth, W.T. 2002. Declaration of William T. Hogarth in opposition to plaintiff's motion for temporary restraining order, 23 Oct. Civ. No. 02-05065-JL. U.S. District Court, Northern District of Calif., San Francisco Div.
- Holst, M. and M.A. Smultea. 2008. Marine mammal and sea turtle monitoring during Lamont-Doherty Earth Observatory's marine seismic program off Central America, February – April 2008. LGL Rep. TA4342-3. Rep. from LGL Ltd., King City, Ont., for Lamont-Doherty Earth Observatory of Columbia Univ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD. 133 p.
- Holst, M., M.A. Smultea, W.R. Koski, and B. Haley. 2005a. Marine mammal and sea turtle monitoring during Lamont-Doherty Earth Observatory's marine seismic program off the Northern Yucatán Peninsula in the Southern Gulf of Mexico, January–February 2005. LGL Rep. TA2822-31. Rep. from LGL Ltd., King City, Ont., for Lamont-Doherty Earth Observatory of Columbia Univ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD.
- Holst, M., M.A. Smultea, W.R. Koski, and B. Haley. 2005b. Marine mammal and sea turtle monitoring during Lamont-Doherty Earth Observatory's marine seismic program in the Eastern Tropical Pacific Ocean off Central America, November–December 2004. LGL Rep. TA2822-30. Rep. from LGL Ltd., King City, Ont., for Lamont-Doherty Earth Observatory of Columbia Univ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD.
- Holst, M., W.J. Richardson, W.R. Koski, M.A. Smultea, B. Haley, M.W. Fitzgerald, and M. Rawson. 2006. Effects of large- and small-source seismic surveys on marine mammals and sea turtles. **Eos**, Trans. Am. Geophys. Union 87(36), Joint Assembly Suppl., Abstract OS42A-01. 23-26 May, Baltimore, MD.
- Hooker, S.K., R.W. Baird, S. Al-Omari, S. Gowans, and H. Whitehead. 2001. Behavioral reactions of northern bottlenose whales (*Hyperoodon ampullatus*) to biopsy darting and tag attachment procedures. **Fish. Bull.** 99(2):303-308.
- Hutchinson, D.R. and R.S. Detrick. 1984. Water gun vs. air gun: a comparison. **Mar. Geophys. Res.** 6(3):295-310.
- IAGC. 2004. Further analysis of 2002 Abrolhos Bank, Brazil humpback whale strandings coincident with seismic surveys. Intern. Assoc. Geophys. Contractors, Houston, TX. 12 p.
- Ireland, D., M. Holst, and W.R. Koski. 2005. Marine mammal monitoring during Lamont-Doherty Earth Observatory's seismic program off the Aleutian Islands, Alaska, July-August 2005. LGL Rep. TA4089-3. Rep. from LGL Ltd., King City, Ont., for Lamont-Doherty Earth Observatory of Columbia Univ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD. 67 p.
- IWC. 2007. Report of the standing working group on environmental concerns. Annex K to Report of the Scientific Committee. **J. Cetac. Res. Manage.** 9(Suppl.):227-260.
- Jefferson, T.A. and B.E. Curry. 1994. Review and evaluation of potential acoustic methods of reducing or eliminating marine mammal-fishery interactions. Rep. from the Mar. Mamm. Res. Progr., Texas A & M Univ., College Station, TX, for U.S. Mar. Mamm. Commis., Washington, DC. 59 p. NTIS PB95-100384.
- Jepson, P.D., M. Arbelo, R. Deaville, I.A.P. Patterson, P. Castro, J.R. Baker, E. Degollada, H.M. Ross, P. Herráez, A.M. Pocknell, F. Rodríguez, F.E. Howie, A. Espinosa, R.J. Reid, J.R. Jaber, V. Martin, A.A. Cunningham, and A. Fernández. 2003. Gas-bubble lesions in stranded cetaceans. **Nature** 425(6958):575-576.
- Jochens, A., D. Biggs, K. Benoit-Bird, D. Engelhaupt, J. Gordon, C. Hu, N. Jaquet, M. Johnson, R. Leben, B. Mate, P. Miller, J. Ortega-Ortiz, A. Thode, P. Tyack, and B. Würsig. 2008. Sperm whale seismic study in the Gulf of Mexico/Synthesis report. OCS Study MMS 2008-006. Rep. from Dep. Oceanogr., Texas A & M Univ., College Station, TX, for U.S. Minerals Manage. Serv., Gulf of Mexico OCS Reg., New Orleans, LA. 323 p.

- Johnson, M.P. and P.L. Tyack. 2003. A digital acoustic recording tag for measuring the response of wild marine mammals to sound. **IEEE J. Oceanic Eng.** 28(1):3-12.
- Johnson, S.R., W.J. Richardson, S.B. Yazvenko, S.A. Blokhin, G. Gailey, M.R. Jenkerson, S.K. Meier, H.R. Melton, M.W. Newcomer, A.S. Perlov, S.A. Rutenko, B. Würsig, C.R. Martin, and D.E. Egging. 2007. A western gray whale mitigation and monitoring program for a 3-D seismic survey, Sakhalin Island, Russia. **Environ. Monit. Assessm.** 134(1-3):1-19.
- Kastak, D. and R.J. Schusterman. 1999. In-air and underwater hearing sensitivity of a northern elephant seal (*Mirounga angustirostris*). **Can. J. Zool.** 77(11):1751-1758.
- Kastak, D., R.L. Schusterman, B.L. Southall, and C.J. Reichmuth. 1999. Underwater temporary threshold shift induced by octave-band noise in three species of pinnipeds. **J. Acoust. Soc. Am.** 106(2):1142-1148.
- Kastak, D., B.L. Southall, R.J. Schusterman, and C. Reichmuth. 2005. Underwater temporary threshold shift in pinnipeds: effects of noise level and duration. **J. Acoust. Soc. Am.** 118(5):3154-3163.
- Kastelein, R.A., P. Mosterd, B. van Santen, M. Hagedoorn, and D. de Haan. 2002. Underwater audiogram of a Pacific walrus (*Odobenus rosmarus divergens*) measured with narrow-band frequency-modulated signals. **J. Acoust. Soc. Am.** 112(5):2173-2182.
- Kastelein, R.A., W.C. Verboom, N. Jennings, and D. de Haan. 2008. Behavioral avoidance threshold level of a harbor porpoise (*Phocoena phocoena*) for a continuous 50 kHz pure tone (L). **J. Acoust. Soc. Am.** 123(4): 1858-1861.
- Kastelein, R.A., P.J. Wensveen, L. Hoek, W.C. Verboom and J.M. Terhune. 2009. Underwater detection of tonal signals between 0.125 and 100 kHz by harbor seals (*Phoca vitulina*). **J. Acoust. Soc. Am.** 125(2):1222-1229.
- Kasuya, T. 1986. Distribution and behavior of Baird's beaked whales off the Pacific coast of Japan. **Sci. Rep. Whales Res. Inst.** 37:61-83.
- Ketten, D.R. 1991. The marine mammal ear: specializations for aquatic audition and echolocation. p. 717-750 *In*: D. Webster, R. Fay and A. Popper (eds.), *The Biology of Hearing*. Springer-Verlag, Berlin.
- Ketten, D.R. 1992. The cetacean ear: form, frequency, and evolution. p. 53-75 *In*: J.A. Thomas, R.A. Kastelein, and A. Ya Supin (eds.), *Marine Mammal Sensory Systems*. Plenum, New York, NY.
- Ketten, D.R. 1994. Functional analysis of whale ears: adaptations for underwater hearing. **IEEE Proc. Underwater Acoust.** 1:264-270.
- Ketten, D.R. 1995. Estimates of blast injury and acoustic trauma zones for marine mammals from underwater explosions. p. 391-407 *In*: R.A. Kastelein, J.A. Thomas, and P.E. Nachtigall (eds.), *Sensory Systems of Aquatic Mammals*. De Spil Publishers, Woerden, Netherlands. 588 p.
- Ketten, D.R. 1998. Marine mammal auditory systems: a summary of audiometric and anatomical data and its implications for underwater acoustic impacts. NOAA Tech. Memo. NOAA-TM-NMFS-SWFSC-256. Southwest Fisheries Sci. Cent., La Jolla, CA. 74 p.
- Ketten, D.R. 2000. Cetacean ears. p. 43-108 *In*: W.W.L. Au, A.N. Popper, and R.R. Fay (eds.), *Hearing by Whales and Dolphins*. Springer-Verlag, New York, NY. 485 p.
- Ketten, D.R., J. Lien and S. Todd. 1993. Blast injury in humpback whale ears: evidence and implications. **J. Acoust. Soc. Am.** 94(3, Pt. 2):1849-1850 (Abstract).
- Ketten, D.R., J. O'Malley, P.W.B. Moore, S. Ridgway, and C. Merigo. 2001. Aging, injury, disease, and noise in marine mammal ears. **J. Acoust. Soc. Am.** 110(5, Pt. 2):2721 (Abstract).
- Klima, E.F., G.R. Gitschlag, and M.L. Renaud. 1988. Impacts of the explosive removal of offshore petroleum platforms on sea turtles and dolphins. **Mar. Fish. Rev.** 50(3):33-42.
- Koski, W.R., D.W. Funk, D.S. Ireland, C. Lyons, K. Christie, A.M. Macrander and S.B. Blackwell. 2009. An update on feeding by bowhead whales near an offshore seismic survey in the central Beaufort Sea. Intern. Whal. Comm. Working Pap. SC/61/BRG3. 15 p

- Kraus, S., A. Read, A. Solov, K. Baldwin, T. Spradlin, E. Anderson, and J. Williamson. 1997. Acoustic alarms reduce porpoise mortality. **Nature** 388(6642):525.
- Kremser, U., P. Klemm, and W.D. Kötz. 2005. Estimating the risk of temporary acoustic threshold shift, caused by hydroacoustic devices, in whales in the Southern Ocean. **Antarctic Sci.** 17(1):3-10.
- Kryter, K.D. 1985. *The Effects of Noise on Man*. 2nd ed. Academic Press, Orlando, FL. 688 p.
- Kryter, K.D. 1994. *The Handbook of Hearing and the Effects of Noise*. Academic Press, Orlando, FL. 673 p.
- Laurinolli, M.H. and N.A. Cochrane. 2005. Hydroacoustic analysis of marine mammal vocalization data from ocean bottom seismometer mounted hydrophones in the Gully. p. 89-95 *In*: K. Lee, H. Bain and G.V. Hurley (eds.), *Acoustic monitoring and marine mammal surveys in The Gully and Outer Scotian Shelf before and during active seismic surveys*. Environ. Stud. Res. Funds Rep. 151. 154 p. Published 2007.
- Lesage, V., C. Barrette, M.C.S. Kingsley, and B. Sjare. 1999. The effect of vessel noise on the vocal behavior of belugas in the St. Lawrence River estuary, Canada. **Mar. Mamm. Sci.** 15(1):65-84.
- Ljungblad, D.K., B. Würsig, S.L. Swartz, and J.M. Keene. 1988. Observations on the behavioral responses of bowhead whales (*Balaena mysticetus*) to active geophysical vessels in the Alaskan Beaufort Sea. **Arctic** 41(3):183-194.
- Lucke, K., U. Siebert, P.A. Lepper and M.-A. Blanchet. 2009. Temporary shift in masked hearing thresholds in a harbor porpoise (*Phocoena phocoena*) after exposure to seismic airgun stimuli. **J. Acoust. Soc. Am.** 125(6):4060-4070.
- Lusseau, D. and L. Bejder. 2007. The long-term consequences of short-term responses to disturbance experience from whalewatching impact assessment. **Intern. J. Compar. Psychol.** 20(2-3):228-236.
- MacGillivray, A.O. and D. Hannay. 2007a. Summary of noise assessment. p. 3-1 to 3-21 *In*: Marine mammal monitoring and mitigation during open water seismic exploration by ConocoPhillips Alaska, Inc., in the Chukchi Sea, July-October 2006. LGL Rep. P903-2 (Jan. 2007). Rep. from LGL Alaska Res. Assoc. Inc., Anchorage, AK, and JASCO Res. Ltd., Victoria, B.C., for ConocoPhillips Alaska Inc., Anchorage, AK, and Nat. Mar. Fish. Serv., Silver Spring, MD. 116 p.
- MacGillivray, A. and D. Hannay. 2007b. Field measurements of airgun array sound levels. p. 4-1 to 4-19 *In*: Marine mammal monitoring and mitigation during open water seismic exploration by GX Technology in the Chukchi Sea, October-November 2006: 90-day report. LGL Rep. P891-1 (Feb. 2007). Rep. from LGL Alaska Res. Assoc. Inc., Anchorage, AK, and JASCO Res. Ltd., Victoria, B.C., for GX Technology, Houston, TX, and Nat. Mar. Fish. Serv., Silver Spring, MD. 118 p.
- MacLean, S.A. and B. Haley. 2004. Marine mammal monitoring during Lamont-Doherty Earth Observatory's seismic study in the Støregga Slide area of the Norwegian Sea, August - September 2003. LGL Rep. TA2822-20. Rep. from LGL Ltd., King City, Ont., for Lamont-Doherty Earth Observatory, Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD. 59 p.
- MacLean, S.A. and W.R. Koski. 2005. Marine mammal monitoring during Lamont-Doherty Earth Observatory's seismic program in the Gulf of Alaska, August-September 2004. LGL Rep. TA2822-28. Rep. from LGL Ltd., King City, Ont., for Lamont-Doherty Earth Observatory, Columbia Univ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD. 102 p.
- Madsen, P.T. 2005. Marine mammals and noise: problems with root mean square sound pressure levels for transients. **J. Acoust. Soc. Am.** 117(6):3952-3957.
- Madsen, P.T., B. Mohl, B.K. Nielsen, and M. Wahlberg. 2002. Male sperm whale behavior during exposures to distant seismic survey pulses. **Aquat. Mamm.** 28(3):231-240.
- Madsen, P.T., M. Johnson, P.J.O. Miller, N. Aguilar de Soto, J. Lynch, and P.L. Tyack. 2006. Quantitative measures of air gun pulses recorded on sperm whales (*Physeter macrocephalus*) using acoustic tags during controlled exposure experiments. **J. Acoust. Soc. Am.** 120(4):2366-2379.
- Malakoff, D. 2002. Suit ties whale deaths to research cruise. **Science** 298(5594):722-723.

- Malme, C.I. and P.R. Miles. 1985. Behavioral responses of marine mammals (gray whales) to seismic discharges. p. 253-280 *In*: G.D. Greene, F.R. Engelhard, and R.J. Paterson (eds.), Proc. Workshop on Effects of Explosives Use in the Marine Environment, Jan. 1985, Halifax, NS. Tech. Rep. 5. Can. Oil & Gas Lands Admin., Environ. Prot. Br., Ottawa, Ont. 398 p.
- Malme, C.I., P.R. Miles, C.W. Clark, P. Tyack, and J.E. Bird. 1984. Investigations of the potential effects of underwater noise from petroleum industry activities on migrating gray whale behavior/Phase II: January 1984 migration. BBN Rep. 5586. Rep. from Bolt Beranek & Newman Inc., Cambridge, MA, for MMS, Alaska OCS Region, Anchorage, AK. NTIS PB86-218377.
- Malme, C.I., P.R. Miles, P. Tyack, C.W. Clark, and J.E. Bird. 1985. Investigation of the potential effects of underwater noise from petroleum industry activities on feeding humpback whale behavior. BBN Rep. 5851; OCS Study MMS 85-0019. Rep. from BBN Labs Inc., Cambridge, MA, for MMS, Anchorage, AK. NTIS PB86-218385.
- Malme, C.I., B. Würsig, J.E. Bird, and P. Tyack. 1986. Behavioral responses of gray whales to industrial noise: feeding observations and predictive modeling. BBN Rep. 6265. OCS Study MMS 88-0048. Outer Contin. Shelf Environ. Assess. Progr., Final Rep. Princ. Invest., NOAA, Anchorage 56(1988): 393-600. NTIS PB88-249008.
- Malme, C.I., B. Würsig, B., J.E. Bird, and P. Tyack. 1988. Observations of feeding gray whale responses to controlled industrial noise exposure. p. 55-73 *In*: W.M. Sackinger, M.O. Jeffries, J.L. Imm, and S.D. Treacy (eds.), Port and Ocean Engineering Under Arctic Conditions. Vol. II. Symposium on Noise and Marine Mammals. Univ. Alaska Fairbanks, Fairbanks, AK. 111 p.
- Manly, B.F.J., V.D. Moulton, R.E. Elliott, G.W. Miller and W.J. Richardson. 2007. Analysis of covariance of fall migrations of bowhead whales in relation to human activities and environmental factors, Alaskan Beaufort Sea: Phase I, 1996-1998. LGL Rep. TA2799-2; OCS Study MMS 2005-033. Rep. from LGL Ltd., King City, Ont., and WEST Inc., Cheyenne, WY, for U.S. Minerals Manage. Serv., Herndon, VA, and Anchorage, AK. 128 p.
- Mate, B.R. and J.T. Harvey. 1987. Acoustical deterrents in marine mammal conflicts with fisheries. ORESU-W-86-001. Oregon State Univ., Sea Grant Coll. Prog., Corvallis, OR. 116 p.
- Mate, B.R., K.M. Stafford, and D.K. Ljungblad. 1994. A change in sperm whale (*Physeter macrocephalus*) distribution correlated to seismic surveys in the Gulf of Mexico. **J. Acoust. Soc. Am.** 96(5, Pt. 2):3268-3269 (Abstract).
- McAlpine, D.F. 2002. Pygmy and dwarf sperm whales. p. 1007-1009 *In*: W.F. Perrin, B. Würsig, and J.G.M. Thewissen (eds.), Encyclopedia of Marine Mammals. Academic Press, San Diego, CA. 1414 p.
- McCall Howard, M.P. 1999. Sperm whales *Physeter macrocephalus* in the Gully, Nova Scotia: Population, distribution, and response to seismic surveying. B.Sc. (Honours) Thesis. Dalhousie Univ., Halifax, NS.
- McCauley, R.D., M.-N. Jenner, C. Jenner, K.A. McCabe, and J. Murdoch. 1998. The response of humpback whales (*Megaptera novaeangliae*) to offshore seismic survey noise: preliminary results of observations about a working seismic vessel and experimental exposures. **APPEA J.** 38:692-707.
- McCauley, R.D., J. Fewtrell, A.J. Duncan, C. Jenner, M.-N. Jenner, J.D. Penrose, R.I.T. Prince, A. Adhitya, J. Murdoch, and K. McCabe. 2000a. Marine seismic surveys: Analysis of airgun signals; and effects of air gun exposure on humpback whales, sea turtles, fishes and squid. Rep. from Centre for Marine Science and Technology, Curtin Univ., Perth, Western Australia, for Australian Petrol. Produc. & Explor. Association, Sydney, NSW. 188 p.
- McCauley, R.D., J. Fewtrell, A.J. Duncan, M.-N. Jenner, M.-N., C. Jenner, R.I.T. Prince, A. Adhitya, K. McCabe and J. Murdoch. 2000b. Marine seismic surveys – a study of environmental implications. **APPEA J.** 40: 692-708.
- McDonald, M.A., J.A. Hildebrand, and S.C. Webb. 1995. Blue and fin whales observed on a seafloor array in the Northeast Pacific. **J. Acoust. Soc. Am.** 98(2, Pt. 1):712-721.

- McShane, L.J., J.A. Estes, M.L. Riedman, and M.M. Staedler. 1995. Repertoire, structure, and individual variation of vocalizations in the sea otter. **J. Mammal.** 76(2):414-427.
- Meier, S.K., S.B. Yazvenko, S.A. Blokhin, P. Wainwright, M.K. Maminov, Y.M. Yakovlev, and M.W. Newcomer. 2007. Distribution and abundance of western gray whales off northeastern Sakhalin Island, Russia, 2001-2003. **Environ. Monit. Assessm.** 134(1-3):107-136.
- Miller, G.W., R.E. Elliott, W.R. Koski, V.D. Moulton, and W.J. Richardson. 1999. Whales. p. 5-1 to 5-109 *In*: W.J. Richardson (ed.), Marine mammal and acoustical monitoring of Western Geophysical's open-water seismic program in the Alaskan Beaufort Sea, 1998. LGL Rep. TA2230-3. Rep. from LGL Ltd., King City, Ont., and Greeneridge Sciences Inc., Santa Barbara, CA, for Western Geophysical, Houston, TX, and Nat. Mar. Fish. Serv., Anchorage, AK, and Silver Spring, MD. 390 p.
- Miller, G.W., V.D. Moulton, R.A. Davis, M. Holst, P. Millman, A. MacGillivray, and D. Hannay. 2005. Monitoring seismic effects on marine mammals—southeastern Beaufort Sea, 2001-2002. p. 511-542 *In*: S.L. Armsworthy, P.J. Cranford, and K. Lee (eds.), Offshore Oil and Gas Environmental Effects Monitoring/Approaches and Technologies. Battelle Press, Columbus, OH.
- Miller, P.J.O., M.P. Johnson, P.T. Madsen, N. Biassoni, M. Quero, and P.L. Tyack. 2009. Using at-sea experiments to study the effects of airguns on the foraging behavior of sperm whales in the Gulf of Mexico. **Deep-Sea Res. I** 56(7):1168-1181.
- Mooney, T.A., P.E. Nachtigall, M. Breese, S. Vlachos, and W.W.L. Au. 2009a. Predicting temporary threshold shifts in a bottlenose dolphin (*Tursiops truncatus*): the effects of noise level and duration. **J. Acoust. Soc. Am.** 125(3):1816-1826.
- Mooney, T.A., P.E. Nachtigall and S. Vlachos. 2009b. Sonar-induced temporary hearing loss in dolphins. **Biol. Lett.** 4(4):565-567.
- Moore, S.E. and Angliss, R.P. 2006. Overview of planned seismic surveys offshore northern Alaska, July-October 2006. Paper SC/58/E6 presented to IWC Scient. Commit., IWC Annu. Meet., 1-13 June, St Kitts.
- Morton A.B. and H.K. Symonds. 2002. Displacement of *Orcinus orca* (L.) by high amplitude sound in British Columbia, Canada. **ICES J. Mar. Sci.** 59(1):71-80
- Moulton, V.D. and J.W. Lawson. 2002. Seals, 2001. p. 3-1 to 3-48 *In*: W.J. Richardson (ed.), Marine mammal and acoustical monitoring of WesternGeco's open water seismic program in the Alaskan Beaufort Sea, 2001. LGL Rep. TA2564-4. Rep. from LGL Ltd., King City, Ont., and Greeneridge Sciences Inc., Santa Barbara, CA, for WesternGeco, Houston, TX, and Nat. Mar. Fish. Serv., Anchorage, AK, and Silver Spring, MD. 95 p.
- Moulton, V.D. and G.W. Miller. 2005. Marine mammal monitoring of a seismic survey on the Scotian Slope, 2003. p. 29-40 *In*: K. Lee, H. Bain, and G.V. Hurley (eds.), Acoustic monitoring and marine mammal surveys in the Gully and outer Scotian Shelf before and during active seismic programs. Environ. Stud. Res. Funds Rep. 151. 154 p (Published 2007).
- Moulton, V.D., B.D. Mactavish, and R.A. Buchanan. 2005. Marine mammal and seabird monitoring of Chevron Canada Resources' 3-D seismic program on the Orphan Basin, 2004. LGL Rep. SA817. Rep. by LGL Ltd., St. John's, NL, for Chevron Canada Resources, Calgary, Alb., ExxonMobil Canada Ltd., St. John's, Nfld., and Imperial Oil Resources Ventures Ltd., Calgary, Alb. 90 p. + appendices.
- Moulton, V.D., B.D. Mactavish, R.E. Harris, and R.A. Buchanan. 2006a. Marine mammal and seabird monitoring of Chevron Canada Limited's 3-D seismic program on the Orphan Basin, 2005. LGL Rep. SA843. Rep. by LGL Ltd., St. John's, Nfld., for Chevron Canada Resources, Calgary, Alb., ExxonMobil Canada Ltd., St. John's, Nfld., and Imperial Oil Resources Ventures Ltd., Calgary, Alb. 111 p. + appendices.
- Moulton, V.D., B.D. Mactavish, and R.A. Buchanan. 2006b. Marine mammal and seabird monitoring of ConocoPhillips' 3-D seismic program in the Laurentian Sub-basin, 2005. LGL Rep. SA849. Rep. by LGL Ltd., St. John's, Nfld., for ConocoPhillips Canada Resources Corp., Calgary, Alb. 97 p. + appendices.
- Nachtigall, P.E., J.L. Pawloski, and W.W.L. Au. 2003. Temporary threshold shifts and recovery following noise exposure in the Atlantic bottlenose dolphin (*Tursiops truncatus*). **J. Acoust. Soc. Am.** 113(6):3425-3429.

- Nachtigall, P.E., A.Y. Supin, J. Pawloski, and W.W.L. Au. 2004. Temporary threshold shifts after noise exposure in the bottlenose dolphin (*Tursiops truncatus*) measured using evoked auditory potentials. **Mar. Mamm. Sci.** 20(4):673-687
- Nachtigall, P.E., A.Y. Supin, M. Amundin, B. Röken., T. Møller, A. Mooney, K.A. Taylor, and M. Yuen. 2007. Polar bear *Ursus maritimus* hearing measured with auditory evoked potentials. **J. Exp. Biol.** 210(7):1116-1122.
- Nations, C.S., S.B. Blackwell, K.H. Kim, A.M. Thode, C.R. Greene Jr., A.M. Macrander, and T.L. McDonald. 2009. Effects of seismic exploration in the Beaufort Sea on bowhead whale call distributions. **J. Acoust. Soc. Am.** 126(4, Pt. 2):2230 (Abstract).
- Nieukirk, S.L., K.M. Stafford, D.K. Mellinger, R.P. Dziak, and C.G. Fox. 2004. Low-frequency whale and seismic airgun sounds recorded in the mid-Atlantic Ocean. **J. Acoust. Soc. Am.** 115(4):1832-1843.
- Nieukirk, S.L., D.K. Mellinger, J.A. Hildebrand, M.A. McDonald, and R.P. Dziak. 2005. Downward shift in the frequency of blue whale vocalizations. p. 205 *In*: Abstr. 16th Bienn. Conf. Biol. Mar. Mamm., San Diego, CA, 12-16 Dec. 2005.
- Nieukirk, S.L., S.L. Heimlich, S.E. Moore, K.M. Stafford, R.P. Dziak, M. Fowler, J. Haxel, J. Goslin and D.K. Mellinger. 2009. Whales and airguns: an eight-year acoustic study in the central North Atlantic. p. 181-182 *In*: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Oct. 2009. 306 p.
- NMFS. 1995. Small takes of marine mammals incidental to specified activities; offshore seismic activities in southern California. **Fed. Regist.** 60(200):53753-53760.
- NMFS. 2000. Small takes of marine mammals incidental to specified activities; marine seismic-reflection data collection in southern California. **Fed. Regist.** 65(20):16374-16379.
- NMFS. 2001. Small takes of marine mammals incidental to specified activities; oil and gas exploration drilling activities in the Beaufort Sea/Notice of issuance of an incidental harassment authorization. **Fed. Regist.** 66(26):9291-9298.
- NMFS. 2005. Endangered Fish and Wildlife; Notice of Intent to Prepare an Environmental Impact Statement. **Fed. Regist.** 70(7):1871-1875.
- NOAA and U.S. Navy. 2001. Joint interim report: Bahamas marine mammal stranding event of 15-16 March 2000. Nat. Mar. Fish. Serv., Silver Spring, MD, and Assistant Secretary of the Navy, Installations & Environ., Washington, DC. 61 p. Available at <http://www.nmfs.noaa.gov/pr/acoustics/reports.htm>
- Nowacek, D.P., L.H. Thorne, D.W. Johnston, and P.L. Tyack. 2007. Responses of cetaceans to anthropogenic noise. **Mammal Rev.** 37(2):81-115.
- NRC. 2005. Marine Mammal Populations and Ocean Noise: Determining When Noise Causes Biologically Significant Effects. U. S. Nat. Res. Council., Ocean Studies Board. (Authors D.W. Wartzok, J. Altmann, W. Au, K. Ralls, A. Starfield, and P.L. Tyack). Nat. Acad. Press, Washington, DC. 126 p.
- Parente, C.L., M.C.C. Marcondes, and M.H. Engel. 2006. Humpback whale strandings and seismic surveys in Brazil from 1999 to 2004. Intern. Whal. Commis. Working Pap. SC/58/E41. 16 p.
- Parente, C.L., J.P. de Araújo and M.E. de Araújo. 2007. Diversity of cetaceans as tool in monitoring environmental impacts of seismic surveys. **Biota Neotrop.** 7(1):1-7.
- Parks, S.E., C.W. Clark, and P.L. Tyack. 2007a. Short- and long-term changes in right whale calling behavior: the potential effects of noise on acoustic communication. **J. Acoust. Soc. Am.** 122(6):3725-3731.
- Parks, S.E., D.R. Ketten, J.T. O'Malley and J. Arruda. 2007b. Anatomical predictions of hearing in the North Atlantic right whale. **Anat. Rec.** 290(6):734-744.
- Parks, S.E., I. Urazghildiiev and C.W. Clark. 2009. Variability in ambient noise levels and call parameters of North Atlantic right whales in three habitat areas. **J. Acoust. Soc. Am.** 125(2):1230-1239.
- Potter, J.R., M. Thillet, C. Douglas, M.A. Chitre, Z. Doborzynski, and P.J. Seekings. 2007. Visual and passive acoustic marine mammal observations and high-frequency seismic source characteristics recorded during a seismic survey. **IEEE J. Oceanic Eng.** 32(2):469-483.

- Reeves, R.R. 1992. Whale responses to anthropogenic sounds: A literature review. Sci. & Res. Ser. 47. New Zealand Dep. Conserv., Wellington. 47 p.
- Reeves, R.R., E. Mitchell, and H. Whitehead. 1993. Status of the northern bottlenose whale, *Hyperoodon ampullatus*. **Can. Field-Nat.** 107(4):490-508.
- Reeves, R.R., R.J. Hofman, G.K. Silber, and D. Wilkinson. 1996. Acoustic deterrence of harmful marine mammal-fishery interactions: proceedings of a workshop held in Seattle, Washington, 20-22 March 1996. NOAA Tech. Memo. NMFS-OPR-10. Nat. Mar. Fish. Serv., Northwest Fisheries Sci. Cent., Seattle, WA. 70 p.
- Reiser, C.M., B. Haley, J. Beland, D.M. Savarese, D.S. Ireland, and D.W. Funk. 2009. Evidence of short-range movements by phocid species in reaction to marine seismic surveys in the Alaskan Chukchi and Beaufort seas. p. 211 *In*: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Canada, Oct. 2009. 306 p.
- Richardson, W.J. and C.I. Malme. 1993. Man-made noise and behavioral responses. p. 631-700 *In*: J.J. Burns, J.J. Montague, and C.J. Cowles (eds.), *The Bowhead Whale*. Spec. Publ. 2, Soc. Mar. Mammal., Lawrence, KS. 787 p.
- Richardson, W.J. and B. Würsig. 1997. Influences of man-made noise and other human actions on cetacean behaviour. **Mar. Freshw. Behav. Physiol.** 29(1-4):183-209.
- Richardson, W.J., B. Würsig, and C.R. Greene. 1986. Reactions of bowhead whales, *Balaena mysticetus*, to seismic exploration in the Canadian Beaufort Sea. **J. Acoust. Soc. Am.** 79(4):1117-1128.
- Richardson, W.J., R.A. Davis, C.R. Evans, D.K. Ljungblad, and P. Norton. 1987. Summer distribution of bowhead whales, *Balaena mysticetus*, relative to oil industry activities in the Canadian Beaufort Sea, 1980-84. **Arctic** 40(2):93-104.
- Richardson, W.J., C.R. Greene, Jr., C.I. Malme, and D.H. Thomson. 1995. *Marine Mammals and Noise*. Academic Press, San Diego, CA. 576 p.
- Richardson, W.J., G.W. Miller, and C.R. Greene, Jr. 1999. Displacement of migrating bowhead whales by sounds from seismic surveys in shallow waters of the Beaufort Sea. **J. Acoust. Soc. Am.** 106(4, Pt. 2):2281 (Abstract).
- Richardson, W.J., M. Holst, W.R. Koski and M. Cummings. 2009. Responses of cetaceans to large-source seismic surveys by Lamont-Doherty Earth Observatory. p. 213 *In*: Abstr. 18th Bienn. Conf. Biol. Mar. Mamm., Québec, Oct. 2009. 306 p.
- Riedman, M.L. 1983. Studies of the effects of experimentally produced noise associated with oil and gas exploration and development on sea otters in California. Rep. from Center for Coastal Marine Studies, Univ. Calif., Santa Cruz, CA, for MMS, Anchorage, AK. 92 p. NTIS PB86-218575.
- Riedman, M.L. 1984. Effects of sounds associated with petroleum industry activities on the behavior of sea otters in California. p. D-1 to D-12 *In*: C.I. Malme, P.R. Miles, C.W. Clark, P. Tyack, and J.E. Bird. Investigations of the potential effects of underwater noise from petroleum industry activities on migrating gray whale behavior/Phase II: January 1984 migration. BBN Rep. 5586. Rep. from BBN Inc., Cambridge, MA, for Minerals Manage. Serv. Anchorage, AK. NTIS PB86-218377.
- Romano, T.A., M.J. Keogh, C.Kelly, P. Feng, L. Berk, C.E. Schlundt, D.A. Carder, and J.J. Finneran. 2004. Anthropogenic sound and marine mammal health: measures of the nervous and immune systems before and after intense sound exposure. **Can. J. Fish. Aquat. Sci.** 61(7):1124-1134.
- SACLANT. 1998. Estimation of cetacean hearing criteria levels. Section II, Chapter 7 *In*: SACLANTCEN Bioacoustics Panel Summary Record and Report. Rep. from NATO Undersea Res. Center. Available at <http://enterprise.spawar.navy.mil/nepa/whales/pdf/doc2-7.pdf>
- Scheifele, P.M., S. Andrew, R.A. Cooper, M. Darre, F.E. Musiek, and L. Max. 2005. Indication of a Lombard vocal response in the St. Lawrence River beluga. **J. Acoust. Soc. Am.** 117(3, Pt. 1):1486-1492.
- Schlundt, C.E., J.J. Finneran, D.A. Carder, and S.H. Ridgway. 2000. Temporary shift in masking hearing thresholds of bottlenose dolphins, *Tursiops truncatus*, and white whales, *Delphinapterus leucas*, after exposure to intense tones. **J. Acoust. Soc. Am.** 107(6):3496-3508.

- Simard, Y., F. Samaran and N. Roy. 2005. Measurement of whale and seismic sounds in the Scotian Gully and adjacent canyons in July 2003. p. 97-115 *In*: K. Lee, H. Bain and C.V. Hurley (eds.), Acoustic monitoring and marine mammal surveys in The Gully and Outer Scotian Shelf before and during active seismic surveys. Environ. Stud. Res. Funds Rep. 151. 154 p (Published 2007).
- Simmonds, M. P. and L.F. Lopez-Jurado. 1991. Whales and the military. **Nature** 351(6326):448.
- Smultea, M.A. and M. Holst. 2008. Marine mammal monitoring during a University of Texas Institute for Geophysics seismic survey in the Northeast Pacific Ocean, July 2008. LGL Rep. TA4584-2. Rep. from LGL Ltd., King City, Ont., for Lamont-Doherty Earth Observatory of Columbia Univ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD. 80 p.
- Smultea, M.A., M. Holst, W.R. Koski, and S. Stoltz. 2004. Marine mammal monitoring during Lamont-Doherty Earth Observatory's seismic program in the Southeast Caribbean Sea and adjacent Atlantic Ocean, April-June 2004. LGL Rep. TA2822-26. Rep. from LGL Ltd., King City, Ont., for Lamont-Doherty Earth Observatory of Columbia Univ., Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD. 106 p.
- Sodal, A. 1999. Measured underwater acoustic wave propagation from a seismic source. Proc. Airgun Environmental Workshop, 6 July, London, UK.
- Southall, B.L., A.E. Bowles, W.T. Ellison, J.J. Finneran, R.L. Gentry, C.R. Greene Jr., D. Kastak, D.R. Ketten, J.H. Miller, P.E. Nachtigall, W.J. Richardson, J.A. Thomas, and P.L. Tyack. 2007. Marine mammal noise exposure criteria: initial scientific recommendations. **Aquat. Mamm.** 33(4):411-522.
- Stone, C.J. 2003. The effects of seismic activity on marine mammals in UK waters 1998-2000. JNCC Rep. 323. Joint Nature Conserv. Commit., Aberdeen, Scotland. 43 p.
- Stone, C.J. and M.L. Tasker. 2006. The effects of seismic airguns on cetaceans in UK waters. **J. Cetac. Res. Manage.** 8(3):255-263.
- Terhune, J.M. 1999. Pitch separation as a possible jamming-avoidance mechanism in underwater calls of bearded seals (*Erignathus barbatus*). **Can. J. Zool.** 77(7):1025-1034.
- Thomas, J.A., R.A. Kastelein and F.T. Awbrey. 1990. Behavior and blood catecholamines of captive belugas during playbacks of noise from an oil drilling platform. **Zoo Biol.** 9(5):393-402.
- Thompson, D., M. Sjöberg, E.B. Bryant, P. Lovell, and A. Bjørge. 1998. Behavioural and physiological responses of harbour (*Phoca vitulina*) and grey (*Halichoerus grypus*) seals to seismic surveys. p. 134 *In*: Abstr. 12th Bienn. Conf. and World Mar. Mamm. Sci. Conf., 20-25 Jan., Monte Carlo, Monaco. 160 p.
- Thomson, D.H. and W.J. Richardson. 1995. Marine mammal sounds. p. 159-204 *In*: W.J. Richardson, C.R. Greene, Jr., C.I. Malme, and D.H. Thomson. Marine Mammals and Noise. Academic Press, San Diego, CA. 576 p.
- Tolstoy, M., J. Diebold, S. Webb, D. Bohnenstiehl, and E. Chapp. 2004a. Acoustic calibration measurements. Chapter 3 *In*: W.J. Richardson (ed.), Marine mammal and acoustic monitoring during Lamont-Doherty Earth Observatory's acoustic calibration study in the northern Gulf of Mexico, 2003. Revised Rep. from LGL Ltd., King City, Ont., for Lamont-Doherty Earth Observatory, Palisades, NY, and Nat. Mar. Fish. Serv., Silver Spring, MD.
- Tolstoy, M., J.B. Diebold, S.C. Webb, D.R. Bohnenstiehl, E. Chapp, R.C. Holmes, and M. Rawson. 2004b. Broadband calibration of R/V *Ewing* seismic sources. **Geophys. Res. Lett.** 31:L14310. doi: 10.1029/2004GL020234
- Tolstoy, M., J. Diebold, L. Doermann, S. Nooner, S.C. Webb, D.R. Bohnenstiehl, T.J. Crone and R.C. Holmes. 2009. Broadband calibration of the R/V *Marcus G. Langseth* four-string seismic sources. **Geochem. Geophys. Geosyst.** 10(8):1-15. Q08011.
- Tyack, P.L. 2008. Implications for marine mammals of large-scale changes in the marine acoustic environment. **J. Mammal.** 89(3):549-558.
- Tyack, P.L. 2009. Human-generated sound and marine mammals. **Phys. Today** 62(11, Nov.):39-44.

- Tyack, P., M. Johnson, and P. Miller. 2003. Tracking responses of sperm whales to experimental exposures of airguns. p. 115-120 *In*: A.E. Jochens and D.C. Biggs (eds.), Sperm whale seismic study in the Gulf of Mexico/Annual Report: Year 1. OCS Study MMS 2003-069. Rep. from Texas A&M Univ., College Station, TX, for U.S. Minerals Manage. Serv., Gulf of Mexico OCS Region, New Orleans, LA.
- Tyack, P.L., M.P. Johnson, P.T. Madsen, P.J. Miller, and J. Lynch. 2006a. Biological significance of acoustic impacts on marine mammals: examples using an acoustic recording tag to define acoustic exposure of sperm whales, *Physeter catodon*, exposed to airgun sounds in controlled exposure experiments. **Eos**, Trans. Am. Geophys. Union 87(36), Joint Assembly Suppl., Abstract OS42A-02. 23-26 May, Baltimore, MD.
- Tyack, P.L., M. Johnson, N. Aguilar Soto, A. Sturlese, and P.T. Madsen. 2006b. Extreme diving of beaked whales. **J. Exp. Biol.** 209(21):4238-4253.
- Urick, R.J. 1983. Principles of Underwater Sound. 3rd ed. Peninsula Publ., Los Altos, CA. 423 p.
- van der Woude, S. 2007. Assessing effects of an acoustic marine geophysical survey on the behaviour of bottlenose dolphins *Tursiops truncatus*. *In*: Abstr. 17th Bienn. Conf. Biol. Mar. Mamm., 29 Nov.–3 Dec., Cape Town, South Africa.
- Wartzok, D., A.N. Popper, J. Gordon, and J. Merrill. 2004. Factors affecting the responses of marine mammals to acoustic disturbance. **Mar. Technol. Soc. J.** 37(4):6-15.
- Watkins, W.A. 1977. Acoustic behavior of sperm whales. **Oceanus** 20(2):50-58.
- Watkins, W.A. 1986. Whale reactions to human activities in Cape Cod waters. **Mar. Mamm. Sci.** 2(4):251-262.
- Watkins, W.A. and W.E. Schevill. 1975. Sperm whales (*Physeter catodon*) react to pingers. **Deep-Sea Res.** 22(3):123-129.
- Watkins, W.A., K.E. Moore, and P. Tyack. 1985. Sperm whale acoustic behaviors in the southeast Caribbean. **Cetology** 49:1-15.
- Weilgart, L.S. 2007. A brief review of known effects of noise on marine mammals. **Intern. J. Comp. Psychol.** 20:159-168.
- Weir, C.R. 2008a. Overt responses of humpback whales (*Megaptera novaeangliae*), sperm whales (*Physeter macrocephalus*), and Atlantic spotted dolphins (*Stenella frontalis*) to seismic exploration off Angola. **Aquat. Mamm.** 34(1):71-83.
- Weir, C.R. 2008b. Short-finned pilot whales (*Globicephala macrorhynchus*) respond to an airgun ramp-up procedure off Gabon. **Aquat. Mamm.** 34(3):349-354.
- Weller, D.W., Y.V. Ivashchenko, G.A. Tsidulko, A.M. Burdin, and R.L. Brownell, Jr. 2002. Influence of seismic surveys on western gray whales off Sakhalin Island, Russia in 2001. Paper SC/54/BRG14, IWC, Western Gray Whale Working Group Meet., 22-25 Oct., Ulsan, South Korea. 12 p.
- Weller, D.W., S.H. Rickards, A.L. Bradford, A.M. Burdin, and R.L. Brownell, Jr. 2006a. The influence of 1997 seismic surveys on the behavior of western gray whales off Sakhalin Island, Russia. Paper SC/58/E4 presented to the IWC Scient. Commit., IWC Annu. Meet., 1-13 June, St. Kitts.
- Weller, D.W., G.A. Tsidulko, Y.V. Ivashchenko, A.M. Burdin and R.L. Brownell Jr. 2006b. A re-evaluation of the influence of 2001 seismic surveys on western gray whales off Sakhalin Island, Russia. Paper SC/58/E5 presented to the IWC Scient. Commit., IWC Annu. Meet., 1-13 June, St. Kitts.
- Wieting, D. 2004. Background on development and intended use of criteria. p. 20 *In*: S. Orenstein, L. Langstaff, L. Manning, and R. Maund (eds.), Advisory Committee on Acoustic Impacts on Marine Mammals, Final Meet. Summary. Second Meet., April 28-30, 2004, Arlington, VA. Sponsored by the Mar. Mamm. Commis., 10 Aug.
- Winsor, M.H. and B.R. Mate. 2006. Seismic survey activity and the proximity of satellite tagged sperm whales. Intern. Whal. Comm. Working Pap. SC/58/E16. 8 p.
- Wright, A.J. and S. Kuczaj. 2007. Noise-related stress and marine mammals: An Introduction. **Intern. J. Comp. Psychol.** 20(2-3):iii-viii.

- Wright, A.J., N. Aguilar Soto, A.L. Baldwin, M. Bateson, C.M. Beale, C. Clark, T. Deak, E.F. Edwards, A. Fernández, A. Godinho, L.T. Hatch, A. Kakuschke, D. Lusseau, D. Martineau, L.M. Romero, L.S. Weilgart, B.A. Wintle, G. Notarbartolo-di-Sciara, and V. Martin. 2007a. Do marine mammals experience stress related to anthropogenic noise? **Intern. J. Comp. Psychol.** 20(2-3):274-316.
- Wright, A.J., N. Aguilar Soto, A.L. Baldwin, M. Bateson, C.M. Beale, C. Clark, T. Deak, E.F. Edwards, A. Fernández, A. Godinho, L.T. Hatch, A. Kakuschke, D. Lusseau, D. Martineau, L.M. Romero, L.S. Weilgart, B.A. Wintle, G. Notarbartolo-di-Sciara and V. Martin. 2007b. Anthropogenic noise as a stressor in animals: A multidisciplinary perspective. **Intern. J. Comp. Psychol.** 20(2-3): 250-273.
- Wright, A.J., T. Deak and E.C.M. Parsons. 2009. Concerns related to chronic stress in marine mammals. Intern. Whal. Comm. Working Pap. SC/61/E16. 7 p.
- Würsig, B., S.K. Lynn, T.A. Jefferson, and K.D. Mullin. 1998. Behaviour of cetaceans in the northern Gulf of Mexico relative to survey ships and aircraft. **Aquat. Mamm.** 24(1):41-50.
- Würsig, B.G., D.W. Weller, A.M. Burdin, S.H. Reeve, A.L. Bradford, S.A. Blokhin, and R.L. Brownell, Jr. 1999. Gray whales summering off Sakhalin Island, Far East Russia: July-October 1997. A joint U.S.-Russian scientific investigation. Final Report. Rep. from Texas A&M Univ., College Station, TX, and Kamchatka Inst. Ecol. & Nature Manage., Russian Acad. Sci., Kamchatka, Russia, for Sakhalin Energy Investment Co. Ltd and Exxon Neftegaz Ltd, Yuzhno-Sakhalinsk, Russia. 101 p.
- Yazvenko, S.B., T.L. McDonald, S.A. Blokhin, S.R. Johnson, S.K. Meier, H.R. Melton, M.W. Newcomer, R.M. Nielson, V.L. Vladimirov, and P.W. Wainwright. 2007a. Distribution and abundance of western gray whales during a seismic survey near Sakhalin Island, Russia. **Environ. Monit. Assessm.** 134(1-3):45-73.
- Yazvenko, S. B., T.L. McDonald, S.A. Blokhin, S.R. Johnson, H.R. Melton, and M.W. Newcomer. 2007b. Feeding activity of western gray whales during a seismic survey near Sakhalin Island, Russia. **Environ. Monit. Assessm.** 134(1-3):93-106.
- Yoder, J.A. 2002. Declaration James A. Yoder in opposition to plaintiff's motion for temporary restraining order, 28 October 2002. Civ. No. 02-05065-JL. U.S. District Court, Northern District of Calif., San Francisco Div

APPENDIX E

REVIEW OF THE EFFECTS OF AIRGUN SOUNDS ON FISH⁸

Here we review literature about the effects of airgun sounds on fishes during seismic surveys. The potential effect of seismic sounds on fish has been studied with a variety of taxa, including marine, freshwater, and anadromous species (reviewed by Fay and Popper 2000; Ladich and Popper 2004; Hastings and Popper 2005; Popper and Hastings 2009a,b).

It is sometimes difficult to interpret studies on the effects of underwater sound on marine animals because authors often do not provide enough information, including received sound levels, source sound levels, and specific characteristics of the sound. Specific characteristics of the sound include units and references, whether the sound is continuous or impulsive, and its frequency range. Underwater sound pressure levels are typically reported as a number of decibels referenced to a reference level, usually 1 micro-Pascal (μPa). However, the sound pressure dB number can represent multiple types of measurements, including “zero to peak”, “peak to peak”, or averaged (“rms”). Sound exposure levels (SEL) may also be reported as dB. The SEL is the integration of all the acoustic energy contained within a single sound event. Unless precise measurement types are reported, it can be impossible to directly compare results from two or more independent studies.

1. Acoustic Capabilities

Sensory systems – like those that allow for hearing – provide information about an animal’s physical, biological, and social environments, in both air and water. Extensive work has been done to understand the structures, mechanisms, and functions of animal sensory systems in aquatic environments (Atema et al. 1988; Kapoor and Hara 2001; Collin and Marshall 2003). All fish species have hearing and skin-based mechanosensory systems (inner ear and lateral line systems, respectively) that provide information about their surroundings (Fay and Popper 2000). Fay (2009) and some others refer to the ambient sounds to which fishes are exposed as ‘underwater soundscapes’. Anthropogenic sounds can have important negative consequences for fish survival and reproduction if they disrupt an individual’s ability to sense its soundscape, which often tells of predation risk, prey items, or mating opportunities. Potential negative effects include masking of key environmental sounds or social signals, displacement of fish from their habitat, or interference with sensory orientation and navigation.

Fish hearing via the inner ear is typically restricted to low frequencies. As with other vertebrates, fish hearing involves a mechanism whereby the beds of hair cells (Howard et al. 1988; Hudspeth and Markin 1994) located in the inner ear are mechanically affected and cause a neural discharge (Popper and Fay 1999). At least two major pathways for sound transmittance between sound source and the inner ear have been identified for fishes. The most primitive pathway involves direct transmission to the inner ear’s otolith, a calcium carbonate mass enveloped by sensory hairs. The inertial difference between the dense otolith and the less-dense inner ear causes the otolith to stimulate the surrounding sensory hair cells. This motion differential is interpreted by the central nervous system as sound.

⁸ By **W. John Richardson** and **Valerie D. Moulton**, with subsequent updates (to November 2009) by WJR and VDM plus **Patrick Abgrall**, **William E. Cross**, **Meike Holst**, and **Mari A. Smultea**, all of LGL Ltd., environmental research associates

The second transmission pathway between sound source and the inner ear of fishes is via the swim bladder, a gas-filled structure that is much less dense than the rest of the fish's body. The swim bladder, being more compressible and expandable than either water or fish tissue, will differentially contract and expand relative to the rest of the fish in a sound field. The pulsating swim bladder transmits this mechanical disturbance directly to the inner ear (discussed below). Such a secondary source of sound detection may be more or less effective at stimulating the inner ear depending on the amplitude and frequency of the pulsation, and the distance and mechanical coupling between the swim bladder and the inner ear (Popper and Fay 1993).

A recent paper by Popper and Fay (2010) discusses the designation of fishes based on sound detection capabilities. They suggest that the designations 'hearing specialist' and 'hearing generalist' no longer be used for fishes because of their vague and sometimes contradictory definitions, and that there is instead a range of hearing capabilities across species that is more like a continuum, presumably based on the relative contributions of pressure to the overall hearing capabilities of a species.

According to Popper and Fay (2010), one end of this continuum is represented by fishes that only detect particle motion because they lack pressure-sensitive gas bubbles (e.g., swim bladder). These species include elasmobranchs (e.g., sharks) and jawless fishes, and some teleosts including flatfishes. Fishes at this end of the continuum are typically capable of detecting sound frequencies below 1500 Hz.

The other end of the fish hearing continuum is represented by fishes with highly specialized otophysical connections between pressure receptive organs, such as the swim bladder, and the inner ear. These fishes include some squirrelfish, mormyrids, herrings, and otophysan fishes (freshwater fishes with Weberian apparatus, an articulated series of small bones that extend from the swim bladder to the inner ear). Rather than being limited to 1.5 kHz or less in hearing, these fishes can typically hear up to several kHz. One group of fish in the anadromous herring sub-family Alosinae (shads and menhaden) can detect sounds to well over 180 kHz (Mann et al. 1997, 1998, 2001). This may be the widest hearing range of any vertebrate that has been studied to date. While the specific reason for this very high frequency hearing is not totally clear, there is strong evidence that this capability evolved for the detection of the ultrasonic sounds produced by echolocating dolphins to enable the fish to detect, and avoid, predation (Mann et al. 1997; Plachta and Popper 2003).

All other fishes have hearing capabilities that fall somewhere between these two extremes of the continuum. Some have unconnected swim bladders located relatively far from the inner ear (e.g., salmonids, tuna) while others have unconnected swim bladders located relatively close to the inner ear (e.g., Atlantic cod, *Gadus morhua*). There has also been the suggestion that Atlantic cod can detect 38 kHz (Astrup and Møhl 1993). However, the general consensus was that this was not hearing with the ear; probably the fish were responding to exceedingly high pressure signals from the 38-kHz source through some other receptor in the skin, such as touch receptors (Astrup and Møhl 1998).

It is important to recognize that the swim bladder itself is not a sensory end organ, but rather an intermediate part of the sound pathway between sound source and the inner ear of some fishes. The inner ear of fishes is ultimately the organ that translates the particle displacement component into neural signals for the brain to interpret as sound.

A third mechanosensory pathway found in most bony fishes and elasmobranchs (i.e., cartilaginous fishes) involves the lateral line system. It too relies on sensitivity to water particle motion. The basic sensory unit of the lateral line system is the neuromast, a bundle of sensory and supporting cells whose projecting cilia, similar to those in the ears, are encased in a gelatinous cap. Neuromasts detect distorted sound waves in the immediate vicinity of fishes. Generally, fishes use the lateral line system to detect the particle displacement component of low frequency acoustic signals (up to 160 to 200 Hz) over a distance

of one to two body lengths. The lateral line is used in conjunction with other sensory systems, including hearing (Sand 1981; Coombs and Montgomery 1999).

2. Potential Effects on Fishes

Review papers on the effects of anthropogenic sources of underwater sound on fishes have been published recently (Popper 2009; Popper and Hastings 2009a,b). These papers consider various sources of anthropogenic sound, including seismic airguns. For the purposes of this review, only the effects of seismic airgun sound are considered.

2.1 Marine Fishes

Evidence for airgun-induced damage to fish ears has come from studies using pink snapper *Pagrus auratus* (McCauley et al. 2000a,b, 2003). In these experiments, fish were caged and exposed to the sound of a single moving seismic airgun every 10 s over a period of 1 h and 41 min. The source SPL at 1 m was about 223 dB re 1 $\mu\text{Pa} \cdot \text{m}_{\text{p-p}}$, and the received SPLs ranged from 165 to 209 dB re 1 $\mu\text{Pa}_{\text{p-p}}$. The sound energy was highest over the 20–70 Hz frequency range. The pink snapper were exposed to more than 600 airgun discharges during the study. In some individual fish, the sensory epithelium of the inner ear sustained extensive damage as indicated by ablated hair cells. Damage was more extensive in fish examined 58 days post-exposure compared to those examined 18 h post-exposure. There was no evidence of repair or replacement of damaged sensory cells up to 58 days post-exposure. McCauley et al. (2000a,b, 2003) included the following caveats in the study reports: (1) fish were caged and unable to swim away from the seismic source, (2) only one species of fish was examined, (3) the impact on the ultimate survival of the fish is unclear, and (4) airgun exposure specifics required to cause the observed damage were not obtained (i.e., a few high SPL signals or the cumulative effect of many low to moderate SPL signals).

The fish exposed to sound from a single airgun in this study also exhibited startle responses to short range start up and high-level airgun signals (i.e., with received SPLs of 182 to 195 dB re 1 $\mu\text{Pa}_{\text{rms}}$ (McCauley et al. 2000a,b). Smaller fish were more likely to display a startle response. Responses were observed above received SPLs of 156 to 161 dB re 1 $\mu\text{Pa}_{\text{rms}}$. The occurrence of both startle response (classic C-turn response) and alarm responses (e.g., darting movements, flash school expansion, fast swimming) decreased over time. Other observations included downward distributional shift that was restricted by the 10 m x 6 m x 3 m cages, increase in swimming speed, and the formation of denser aggregations. Fish behavior appeared to return to pre-exposure state 15–30 min after cessation of seismic firing.

Pearson et al. (1992) investigated the effects of seismic airgun sound on the behavior of captive rockfishes (*Sebastes* sp.) exposed to the sound of a single stationary airgun at a variety of distances. The airgun used in the study had a source SPL at 1 m of 223 dB re 1 $\mu\text{Pa} \cdot \text{m}_{0\text{-p}}$, and measured received SPLs ranged from 137 to 206 dB re 1 $\mu\text{Pa}_{0\text{-p}}$. The authors reported that rockfishes reacted to the airgun sounds by exhibiting varying degrees of startle and alarm responses, depending on the species of rockfish and the received SPL. Startle responses were observed at a minimum received SPL of 200 dB re 1 $\mu\text{Pa}_{0\text{-p}}$, and alarm responses occurred at a minimum received SPL of 177 dB re 1 $\mu\text{Pa}_{0\text{-p}}$. Other observed behavioral changes included the tightening of schools, downward distributional shift, and random movement and orientation. Some fishes ascended in the water column and commenced to mill (i.e., “eddy”) at increased speed, while others descended to the bottom of the enclosure and remained motionless. Pre-exposure behavior was reestablished from 20 to 60 min after cessation of seismic airgun discharge. Pearson et al. (1992) concluded that received SPL thresholds for overt rockfish behavioral response and more subtle rockfish behavioral response are 180 dB re 1 $\mu\text{Pa}_{0\text{-p}}$ and 161 dB re 1 $\mu\text{Pa}_{0\text{-p}}$, respectively.

Using an experimental hook and line fishery approach, Skalski et al. (1992) studied the potential effects of seismic airgun sound on the distribution and catchability of rockfishes. The source SPL of the single airgun used in the study was 223 dB re $1 \mu\text{Pa} \cdot \text{m}_{0-p}$, and the received SPLs at the bases of the rockfish aggregations ranged from 186 to 191 dB re $1 \mu\text{Pa}_{0-p}$. Characteristics of the fish aggregations were assessed using echosounders. During long-term stationary seismic airgun discharge, there was an overall downward shift in fish distribution. The authors also observed a significant decline in total catch of rockfishes during seismic discharge. It should be noted that this experimental approach was quite different from an actual seismic survey, in that duration of exposure was much longer.

In another study, caged European sea bass (*Dicentrarchus labrax*) were exposed to multiple discharges from a moving seismic airgun array with a source SPL of about 256 dB re $1 \mu\text{Pa} \cdot \text{m}_{0-p}$ (unspecified measure type) (Santulli et al. 1999). The airguns were discharged every 25 s during a 2-h period. The minimum distance between fish and seismic source was 180 m. The authors did not indicate any observed pathological injury to the sea bass. Blood was collected from both exposed fish (6 h post-exposure) and control fish (6 h pre-exposure) and subsequently analyzed for cortisol, glucose, and lactate levels. Levels of cortisol, glucose, and lactate were significantly higher in the sera of exposed fish compared to sera of control fish. The elevated levels of all three chemicals returned to pre-exposure levels within 72 h of exposure (Santulli et al. 1999).

Santulli et al. (1999) also used underwater video cameras to monitor fish response to seismic airgun discharge. Resultant video indicated slight startle responses by some of the sea bass when the seismic airgun array discharged as far as 2.5 km from the cage. The proportion of sea bass that exhibited startle response increased as the airgun sound source approached the cage. Once the seismic array was within 180 m of the cage, the sea bass were densely packed at the middle of the enclosure, exhibiting random orientation, and appearing more active than they had been under pre-exposure conditions. Normal behavior resumed about 2 h after airgun discharge nearest the fish (Santulli et al. 1999).

Boeger et al. (2006) reported observations of coral reef fishes in field enclosures before, during and after exposure to seismic airgun sound. This Brazilian study used an array of eight airguns that was presented to the fishes as both a mobile sound source and a static sound source. Minimum distances between the sound source and the fish cage ranged from 0 to 7 m. Received sound levels were not reported by Boeger et al. (2006). Neither mortality nor external damage to the fishes was observed in any of the experimental scenarios. Most of the airgun array discharges resulted in startle responses although these behavioral changes lessened with repeated exposures, suggesting habituation.

Chapman and Hawkins (1969) investigated the reactions of free ranging whiting (silver hake), *Merluccius bilinearis*, to an intermittently discharging stationary airgun with a source SPL of 220 dB re $1 \mu\text{Pa} \cdot \text{m}_{0-p}$. Received SPLs were estimated to be 178 dB re $1 \mu\text{Pa}_{0-p}$. The whiting were monitored with an echosounder. Prior to any airgun discharge, the fish were located at a depth range of 25 to 55 m. In apparent response to the airgun sound, the fish descended, forming a compact layer at depths greater than 55 m. After an hour of exposure to the airgun sound, the fish appeared to have habituated as indicated by their return to the pre-exposure depth range, despite the continuing airgun discharge. Airgun discharge ceased for a time and upon its resumption, the fish again descended to greater depths, indicating only temporary habituation.

Hassel et al. (2003, 2004) studied the potential effects of exposure to airgun sound on the behavior of captive lesser sandeel, *Ammodytes marinus*. Depth of the study enclosure used to hold the sandeel was about 55 m. The moving airgun array had an estimated source SPL of 256 dB re $1 \mu\text{Pa} \cdot \text{m}$ (unspecified measure type). Received SPLs were not measured. Exposures were conducted over a 3-day period in a $10 \text{ km} \times 10 \text{ km}$ area with the cage at its center. The distance between airgun array and fish cage ranged

from 55 m when the array was overhead to 7.5 km. No mortality attributable to exposure to the airgun sound was noted. Behavior of the fish was monitored using underwater video cameras, echosounders, and commercial fishery data collected close to the study area. The approach of the seismic vessel appeared to cause an increase in tail-beat frequency although the sandeels still appeared to swim calmly. During seismic airgun discharge, many fish exhibited startle responses, followed by flight from the immediate area. The frequency of occurrence of startle response seemed to increase as the operating seismic array moved closer to the fish. The sandeels stopped exhibiting the startle response once the airgun discharge ceased. The sandeel tended to remain higher in the water column during the airgun discharge, and none of them were observed burying themselves in the soft substrate. The commercial fishery catch data were inconclusive with respect to behavioral effects.

Various species of demersal fishes, blue whiting, and some small pelagic fishes were exposed to a moving seismic airgun array with a source SPL of about 250 dB re 1 $\mu\text{Pa} \cdot \text{m}$ (unspecified measure type) (Dalen and Knutsen 1986). Received SPLs estimated using the assumption of spherical spreading ranged from 200 to 210 dB re 1 μPa (unspecified measure type). Seismic sound exposures were conducted every 10 s during a one week period. The authors used echosounders and sonars to assess the pre- and post-exposure fish distributions. The acoustic mapping results indicated a significant decrease in abundance of demersal fish (36%) after airgun discharge but comparative trawl catches did not support this. Non-significant reductions in the abundances of blue whiting and small pelagic fish were also indicated by post-exposure acoustic mapping.

La Bella et al. (1996) studied the effects of exposure to seismic airgun sound on fish distribution using echosounder monitoring and changes in catch rate of hake by trawl, and clupeoids by gill netting. The seismic array used was composed of 16 airguns and had a source SPL of 256 dB re 1 $\mu\text{Pa} \cdot \text{m}_{0-p}$. The shot interval was 25 s, and exposure durations ranged from 4.6 to 12 h. Horizontal distributions did not appear to change as a result of exposure to seismic discharge, but there was some indication of a downward shift in the vertical distribution. The catch rates during experimental fishing did not differ significantly between pre- and post-seismic fishing periods.

Wardle et al. (2001) used video and telemetry to make behavioral observations of marine fishes (primarily juvenile saithe, adult pollock, juvenile cod, and adult mackerel) inhabiting an inshore reef off Scotland before, during, and after exposure to discharges of a stationary airgun. The received SPLs ranged from about 195 to 218 dB re 1 μPa_{0-p} . Pollock did not move away from the reef in response to the seismic airgun sound, and their diurnal rhythm did not appear to be affected. However, there was an indication of a slight effect on the long-term day-to-night movements of the pollock. Video camera observations indicated that fish exhibited startle responses (“C-starts”) to all received levels. There were also indications of behavioral responses to visual stimuli. If the seismic source was visible to the fish, they fled from it. However, if the source was not visible to the fish, they often continued to move toward it.

The potential effects of exposure to seismic sound on fish abundance and distribution were also investigated by Slotte et al. (2004). Twelve days of seismic survey operations spread over a period of 1 month used a seismic airgun array with a source SPL of 222.6 dB re 1 $\mu\text{Pa} \cdot \text{m}_{p-p}$. The SPLs received by the fish were not measured. Acoustic surveys of the local distributions of various kinds of pelagic fish, including herring, blue whiting, and mesopelagic species, were conducted during the seismic surveys. There was no strong evidence of short-term horizontal distributional effects. With respect to vertical distribution, blue whiting and mesopelagics were distributed deeper (20 to 50 m) during the seismic survey compared to pre-exposure. The average densities of fish aggregations were lower within the

seismic survey area, and fish abundances appeared to increase in accordance with increasing distance from the seismic survey area.

Fertilized capelin (*Mallotus villosus*) eggs and monkfish (*Lophius americanus*) larvae were exposed to seismic airgun sound and subsequently examined and monitored for possible effects of the exposure (Payne et al. 2009). The laboratory exposure studies involved a single airgun. Approximate received SPLs measured in the capelin egg and monkfish larvae exposures were 199 to 205 dB re 1 μPa_{p-p} and 205 dB re 1 μPa_{p-p} , respectively. The capelin eggs were exposed to either 10 or 20 airgun discharges, and the monkfish larvae were exposed to either 10 or 30 discharges. No statistical differences in mortality/morbidity between control and exposed subjects were found at 1 to 4 days post-exposure in any of the exposure trials for either the capelin eggs or the monkfish larvae.

In uncontrolled experiments, Kostyvchenko (1973) exposed the eggs of numerous fish species (anchovy, red mullet, crucian carp, blue runner) to various sound sources, including seismic airguns. With the seismic airgun discharge as close as 0.5 m from the eggs, over 75% of them survived the exposure. Egg survival rate increased to over 90% when placed 10 m from the airgun sound source. The range of received SPLs was about 215 to 233 dB re 1 μPa_{0-p} .

Eggs, yolk sac larvae, post-yolk sac larvae, post-larvae, and fry of various commercially important fish species (cod, saithe, herring, turbot, and plaice) were exposed to received SPLs ranging from 220 to 242 dB re 1 μPa (unspecified measure type) (Booman et al. 1996). These received levels corresponded to exposure distances ranging from 0.75 to 6 m. The authors reported some cases of injury and mortality but most of these occurred as a result of exposures at very close range (i.e., <15 m). The rigor of anatomical and pathological assessments was questionable.

Saetre and Ona (1996) applied a “worst-case scenario” mathematical model to investigate the effects of seismic sound on fish eggs and larvae. They concluded that mortality rates caused by exposure to seismic airgun sound are so low compared to the natural mortality that the impact of seismic surveying on recruitment to a fish stock must be regarded as insignificant.

2.2 Freshwater Fishes

Popper et al. (2005) tested the hearing sensitivity of three Mackenzie River fish species after exposure to five discharges from a seismic airgun. The mean received peak SPL was 205 to 209 dB re 1 μPa per discharge, and the approximate mean received SEL was 176 to 180 dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$ per discharge. While the broad whitefish showed no Temporary Threshold Shift (TTS) as a result of the exposure, adult northern pike and lake chub exhibited TTSs of 10 to 15 dB, followed by complete recovery within 24 h of exposure. The same animals were also examined to determine whether there were observable effects on the sensory cells of the inner ear as a result of exposure to seismic sound (Song et al. 2008). No damage to the ears of the fishes was found, including those that exhibited TTS.

In another part of the same Mackenzie River project, Jorgenson and Gyselman (2009) investigated the behavioral responses of arctic riverine fishes to seismic airgun sound. They used hydroacoustic survey techniques to determine whether fish behavior upon exposure to airgun sound can either mitigate or enhance the potential impact of the sound. The study indicated that fish behavioral characteristics were generally unchanged by the exposure to airgun sound. The tracked fish did not exhibit herding behavior in front of the mobile airgun array and, therefore, were not exposed to sustained high sound levels.

2.3 Anadromous Fishes

In uncontrolled experiments using a very small sample of different groups of young salmonids, including Arctic cisco, fish were caged and exposed to various types of sound. One sound type was either a

single firing or a series of four firings 10 to 15 s apart of a 300-in³ seismic airgun at 2000 to 2200 psi (Falk and Lawrence 1973). Swim bladder damage was reported but no mortality was observed when fish were exposed within 1 to 2 m of an airgun source with source level, as estimated by Turnpenny and Nedwell (1994), of ~230 dB re 1 $\mu\text{Pa}\cdot\text{m}$ (unspecified measure).

Thomsen (2002) exposed rainbow trout and Atlantic salmon held in aquaculture enclosures to the sounds from a small airgun array. Received SPLs were 142 to 186 dB re 1 $\mu\text{Pa}_{\text{p-p}}$. The fish were exposed to 124 pulses over a 3-day period. In addition to monitoring fish behavior with underwater video cameras, the authors also analyzed cod and haddock catch data from a longline fishing vessel operating in the immediate area. Only eight of the 124 shots appeared to evoke behavioral reactions by the salmonids, but overall impacts were minimal. No fish mortality was observed during or immediately after exposure. The author reported no significant effects on cod and haddock catch rates, and the behavioral effects were hard to differentiate from normal behavior.

Weinhold and Weaver (1972, cited in Turnpenny et al. 1994) exposed caged coho salmon smolts to impulses from 330 and 660-in³ airguns at distances ranging from 1 to 10 m, resulting in received levels estimated at ~214 to 216 dB (units not given). No lethal effects were observed.

It should be noted that, in a recent and comprehensive review, Hastings and Popper (2005) take issue with many of the authors cited above for problems with experimental design and execution, measurements, and interpretation. Hastings and Popper (2005) deal primarily with possible effects of pile-driving sounds (which, like airgun sounds, are impulsive and repetitive). However, that review provides an excellent and critical review of the impacts to fish from other underwater anthropogenic sounds.

3. Indirect Effects on Fisheries

The most comprehensive experimentation on the effects of seismic airgun sound on catchability of fishes was conducted in the Barents Sea by Engås et al. (1993, 1996). They investigated the effects of seismic airgun sound on distributions, abundances, and catch rates of cod and haddock using acoustic mapping and experimental fishing with trawls and longlines. The maximum source SPL was about 248 dB re 1 $\mu\text{Pa}\cdot\text{m}_{0-p}$ based on back-calculations from measurements collected via a hydrophone at depth 80 m. No measurements of the received SPLs were made. Davis et al. (1998) estimated the received SPL at the sea bottom immediately below the array and at 18 km from the array to be 205 dB re 1 μPa_{0-p} and 178 dB re 1 μPa_{0-p} , respectively. Engås et al. (1993, 1996) concluded that there were indications of distributional change during and immediately following the seismic airgun discharge (45 to 64% decrease in acoustic density according to sonar data). The lowest densities were observed within 9.3 km of the seismic discharge area. The authors indicated that trawl catches of both cod and haddock declined after the seismic operations. While longline catches of haddock also showed decline after seismic airgun discharge, those for cod increased.

Løkkeborg (1991), Løkkeborg and Soldal (1993), and Dalen and Knutsen (1986) also examined the effects of seismic airgun sound on demersal fish catches. Løkkeborg (1991) examined the effects on cod catches. The source SPL of the airgun array used in his study was 239 dB re 1 $\mu\text{Pa}\cdot\text{m}$ (unspecified measure type), but received SPLs were not measured. Approximately 43 h of seismic airgun discharge occurred during an 11-day period, with a five-second interval between pulses. Catch rate decreases ranging from 55 to 80% within the seismic survey area were observed. This apparent effect persisted for at least 24 h within about 10 km of the survey area.

Turnpenny et al. (1994) examined results of these studies as well as the results of other studies on rockfish. They used rough estimations of received SPLs at catch locations and concluded that catchability is reduced when received SPLs exceed 160 to 180 dB re 1 μPa_{0-p} . They also concluded that reaction

thresholds of fishes lacking a swim bladder (e.g., flatfish) would likely be about 20 dB higher. Given the considerable variability in sound transmission loss between different geographic locations, the SPLs that were assumed in these studies were likely quite inaccurate.

Turnpenny and Nedwell (1994) also reported on the effects of seismic airgun discharge on inshore bass fisheries in shallow U.K. waters (5 to 30 m deep). The airgun array used had a source level of 250 dB re 1 $\mu\text{Pa} \cdot \text{m}_{0-p}$. Received levels in the fishing areas were estimated to be 163–191 dB re 1 μPa_{0-p} . Using fish tagging and catch record methodologies, they concluded that there was not any distinguishable migration from the ensonified area, nor was there any reduction in bass catches on days when seismic airguns were discharged. The authors concluded that effects on fisheries would be smaller in shallow nearshore waters than in deep water because attenuation of sound is more rapid in shallow water.

Skalski et al. (1992) used a 100-in³ airgun with a source level of 223 dB re 1 $\mu\text{Pa} \cdot \text{m}_{0-p}$ to examine the potential effects of airgun sound on the catchability of rockfishes. The moving airgun was discharged along transects in the study fishing area, after which a fishing vessel deployed a set line, ran three echosounder transects, and then deployed two more set lines. Each fishing experiment lasted 1 h 25 min. Received SPLs at the base of the rockfish aggregations ranged from 186 to 191 dB re 1 μPa_{0-p} . The catch-per-unit-effort (CPUE) for rockfish declined on average by 52.4% when the airguns were operating. Skalski et al. (1992) believed that the reduction in catch resulted from a change in behavior of the fishes. The fish schools descended towards the bottom and their swimming behavior changed during airgun discharge. Although fish dispersal was not observed, the authors hypothesized that it could have occurred at a different location with a different bottom type. Skalski et al. (1992) did not continue fishing after cessation of airgun discharge. They speculated that CPUE would quickly return to normal in the experimental area, because fish behavior appeared to normalize within minutes of cessation of airgun discharge. However, in an area where exposure to airgun sound might have caused the fish to disperse, the authors suggested that a lower CPUE might persist for a longer period.

European sea bass were exposed to sound from seismic airgun arrays with a source SPL of 262 dB re 1 $\mu\text{Pa} \cdot \text{m}_{0-p}$ (Pickett et al. 1994). The seismic survey was conducted over a period of 4 to 5 months. The study was intended to investigate the effects of seismic airgun discharge on inshore bass fisheries. Information was collected through a tag and release program, and from the logbooks of commercial fishermen. Most of the 152 recovered fish from the tagging program were caught within 10 km of the release site, and it was suggested that most of these bass did not leave the area for a prolonged period. With respect to the commercial fishery, no significant changes in catch rate were observed (Pickett et al. 1994).

4. Literature Cited

- Atema, J., R.R. Fay, A.N. Popper, and W.N. Tavolga. 1988. The sensory biology of aquatic animals. Springer-Boeger, W.A., M.R. Pie, A. Ostrensky, and M.F. Cardoso. 2006. The effect of exposure to seismic prospecting on coral reef fishes. **Braz. J. Oceanog.** 54(4):235-239.
- Booman, C., J. Dalen, H. Leivestad, A. Levsen, T. van der Meeren, and K. Toklum. 1996. Effeter av luftkanon-skyting på egg, larver og yngel. **Fisken Og Havet** 1996(3):1-83 (Norwegian with English summary).
- Chapman, C.J. and A.D. Hawkins. 1969. The importance of sound in fish behaviour in relation to capture by trawls. **FAO Fish. Rep.** 62:717-729.
- Collin, S.P. and N.J. Marshall (eds.). 2003. Sensory processing in aquatic environments. Springer-Verlag, New York, NY. 446 p.

- Coombs, S. and J.C. Montgomery. 1999. The enigmatic lateral line system. p. 319-362 *In*: R.R. Fay and A.N. Popper (eds.), *Comparative hearing: fish and amphibians*. Springer Handbook of Auditory Research 11. Springer-Verlag, New York, NY. 438 p.
- Dalen, J. and G.M. Knutsen. 1986. Scaring effects in fish and harmful effects on eggs, larvae and fry by offshore seismic explorations. Symposium on Underwater Acoustics, Halifax.
- Davis, R.A., D. Thomson, and C.I. Malme. 1998. Environmental assessment of seismic exploration of the Scotian Shelf. Rep. by LGL Ltd., King City, Ont., and Charles I. Malme, Engineering and Scientific Services, Hingham, MA, for Mobil Oil Canada Properties Ltd., Shell Canada Ltd., and Imperial Oil Ltd.
- Engås, A., S. Løkkeborg, A.V. Soldal, and E. Ona. 1993. Comparative trials for cod and haddock using commercial trawl and longline at two different stock levels. **J. Northw. Atl. Fish. Sci.** 19:83-90.
- Engås, A., S. Løkkeborg, E. Ona, and A.V. Soldal. 1996. Effects of seismic shooting on local abundance and catch rates of cod (*G. morhua*) and haddock (*M. aeglefinus*). **Can. J. Fish. Aquat. Sci.** 53(10):2238-2249.
- Falk, M.R. and M.J. Lawrence. 1973. Seismic exploration: its nature and effects on fish. Tech. Rep. Ser. CEN/T-73-9. Can. Dep. Environ., Fisheries & Marine Serv., Resource Manage. Br., Fisheries Operations Directorate, Central Region (Environment), Winnipeg, Man.
- Fay, R. 2009. Soundscapes and the sense of hearing of fishes. **Integr. Zool.** 4(1):26-32.
- Fay, R.R. and A.N. Popper. 2000. Evolution of hearing in vertebrates: The inner ears and processing. **Hearing Res.** 149(1):1-10.
- Hassel, A., T. Knutsen, J. Dalen, S. Løkkeborg, K. Skaar, Ø. Østensen, E.K. Haugland, M. Fonn, Å. Høines, and O.A. Misund. 2003. Reaction of sandeel to seismic shooting: a field experiment and fishery statistics study. Institute of Marine Research, Bergen, Norway.
- Hassel, A., T. Knutsen, J. Dalen, K. Skaar, S. Løkkeborg, O.A. Misund, O. Ostensen, M. Fonn, and E.K. Haugland. 2004. Influence of seismic shooting on the lesser sandeel (*Ammodytes marinus*). **ICES J. Mar. Sci.** 61(7):1165-1173.
- Hastings, M.C. and A.N. Popper. 2005. Effects of sound on fish. Rep. from Jones & Stokes, Sacramento, CA, for California Department of Transportation, Sacramento, CA. 28 January.
- Howard J, W.M. Roberts, and A.J. Hudspeth. 1988. Mechanoelectrical transduction by hair cells. **Annu. Rev. Biophys. Chem.** 17:99-124.
- Hudspeth, A.J. and V.S. Markin. 1994. The ear's gears: mechanical transduction by hair cells. **Physics Today** 47(2):22-28.
- Jorgenson, J.K. and E.C. Gyselman. 2009. Hydroacoustic measurements of the behavioral response of arctic riverine fishes to seismic airguns. **J. Acoust. Soc. Am.** 126(3):1598-1606.
- Kapoor, B.G. and T.J. Hara (eds.). 2001. *Sensory biology of jawed fishes: new insights*. Science Publishers, Inc., Enfield, NH. 404 p.
- Kostyvchenko, L.P. 1973. Effects of elastic waves generated in marine seismic prospecting on fish eggs in the Black Sea. **Hydrobiol. J.** 9:45-48.
- La Bella, G., S. Cannata, C. Frogliola, A. Modica, S. Ratti, and G. Rivas. 1996. First assessment of effects of air-gun seismic shooting on marine resources in the Central Adriatic Sea. p. 227-238 *In*: Society of Petroleum Engineers, Intern. Conf. on Health, Safety and Environ., New Orleans, LA, 9-12 June.
- Ladich, F. and A.N. Popper. 2004. Parallel evolution in fish hearing organs. p. 95-127 *In*: G.A. Manley, A.N. Popper, and R.R. Fay (eds.), *Evolution of the vertebrate auditory system*. Springer-Verlag, New York, NY. 415 p.
- Løkkeborg, S. 1991. Effects of geophysical survey on catching success in longline fishing. Paper presented at Intern. Council for the Exploration of the Sea (ICES) Annual Science Conf. **ICES CM B** 40:1-9.
- Løkkeborg, S. and A.V. Soldal. 1993. The influence of seismic explorations on cod (*Gadus morhua*) behaviour and catch rates. **ICES Mar. Sci. Symp.** 196:62-67.

- Mann, D.A., Z. Lu, and A.N. Popper. 1997. A clupeid fish can detect ultrasound. **Nature** 389(6649):341.
- Mann, D.A., Z. Lu, M.C. Hastings, and A.N. Popper. 1998. Detection of ultrasonic tones and simulated dolphin echolocation clicks by a teleost fish, the American shad (*Alosa sapidissima*). **J. Acoust. Soc. Am.** 104(1):562-568.
- Mann, D.A., D.M. Higgs, W.N. Tavolga, M.J. Souza, and A.N. Popper. 2001. Ultrasound detection by clupeiform fishes. **J. Acoust. Soc. Am.** 109(6):3048-3054.
- McCauley, R.D., J. Fewtrell, A.J. Duncan, C. Jenner, M.-N. Jenner, J.D. Penrose, R.I.T. Prince, A. Adhitya, J. Murdoch, and K. McCabe. 2000a. Marine seismic surveys: analysis of airgun signals; and effects of air gun exposure on humpback whales, sea turtles, fishes and squid. Rep. from Centre for Marine Science and Technology, Curtin University, Perth, WA, for Australian Petroleum Production Association, Sydney, NSW.
- McCauley, R.D., J. Fewtrell, A.J. Duncan, C. Jenner, M.-N. Jenner, J.D. Penrose, R.I.T. Prince, A. Adhitya, J. Murdoch, and K. McCabe. 2000b. Marine seismic surveys – a study of environmental implications. **APPEA J.** 40:692-706.
- McCauley, R.D., J. Fewtrell, and A.N. Popper. 2003. High intensity anthropogenic sound damages fish ears. **J. Acoust. Soc. Am.** 113(1):638-642.
- Payne, J.F., J. Coady, and D. White. 2009. Potential effects of seismic airgun discharges on monkfish eggs (*Lophius americanus*) and larvae. Environ. Stud. Res. Funds Rep. 170. St. John's, NL. 35 p.
- Pearson, W.H., J.R. Skalski, and C.I. Malme. 1992. Effects of sounds from a geophysical survey device on behavior of captive rockfish (*Sebastes* spp.). **Can. J. Fish. Aquat. Sci.** 49(7):1343-1356.
- Pickett, G.D., D.R. Eaton, R.M.H. Seaby, and G.P. Arnold. 1994. Results of bass tagging in Poole Bay during 1992. Laboratory Leaflet Number 74. Ministry of Agriculture, Fisheries and Food, Directorate of Fisheries Research, Lowestoft, UK.
- Popper, A.N. 2009. Are we drowning out fish in a sea of noise? **Marine Scientist** 27:18-20.
- Popper, A.N. and R.R. Fay. 1993. Sound detection and processing by fish: critical review and major research questions. **Brain Behav. Evol.** 41(1):14-38.
- Popper, A.N. and R.R. Fay. 1999. The auditory periphery in fishes. p. 43-100 *In*: R.R. Fay and A.N. Popper (eds.), Comparative hearing: fish and amphibians. Springer-Verlag, New York, NY. 438 p.
- Popper, A.N. and R.R. Fay. 2010. Rethinking sound detection by fishes. **Hear. Res.** (in press) doi: 10.1016/j.heares.2009.12.023.
- Popper, A.N. and M.C. Hastings. 2009a. The effects of human-generated sound on fish. **Integr. Zool.** 4(1):43-52.
- Popper, A.N. and M.C. Hastings. 2009b. The effects of anthropogenic sources of sound on fishes. **J. Fish Biol.** 75(3):455-489.
- Popper, A.N., M.E. Smith, P.A. Cott, B.W. Hanna, A.O. MacGillivray, M.E. Austin, and D.A. Mann. 2005. Effects of exposure to seismic airgun use on hearing of three fish species. **J. Acoust. Soc. Am.** 117(6):3958-3971.
- Saetre, R. and E. Ona. 1996. Seismiske undersøkelser og skader på fiskeegg og -larver en vurdering av mulige effekter pa bestandsniv. [Seismic investigations and damages on fish eggs and larvae; an evaluation of possible effects on stock level] **Fisken og Havet** 1996:1-17, 1-8. (in Norwegian with English summary).
- Sand, O. 1981. The lateral line and sound reception. p. 459-478 *In*: W.N. Tavolga, A.N. Popper, and R.R. Fay (eds.), Hearing and sound communication in fishes. Springer-Verlag, New York, NY.
- Santulli, A., C. Messina, L. Ceffa, A. Curatolo, G. Rivas, G. Fabi, and V. Damelio. 1999. Biochemical responses of European sea bass (*Dicentrarchus labrax*) to the stress induced by offshore experimental seismic prospecting. **Mar. Poll. Bull.** 38(12):1105-1114.
- Skalski, J.R., W.H. Pearson, and C.I. Malme. 1992. Effects of sounds from a geophysical survey device on catch-per-unit-effort in a hook-and-line fishery for rockfish (*Sebastes* spp.). **Can. J. Fish. Aquat. Sci.** 49(7):1357-1365.

- Slotte, A., K. Hansen, J. Dalen, and E. Ona. 2004. Acoustic mapping of pelagic fish distribution and abundance in relation to a seismic shooting area off the Norwegian west coast. **Fish. Res.** 67(2):143-150.
- Song, J., D.A. Mann, P.A. Cott, B.W. Hanna, and A.N. Popper. 2008. The inner ears of Northern Canadian freshwater fishes following exposure to seismic air gun sounds. **J. Acoust. Soc. Am.** 124(2):1360-1366.
- Thomsen, B. 2002. An experiment on how seismic shooting affects caged fish. Thesis, Faroese Fisheries Laboratory, University of Aberdeen, Aberdeen, Scotland. 16 August.
- Turnpenny, A.W.H. and J.R. Nedwell. 1994. Consultancy Report: The effects on marine fish, diving mammals and birds of underwater sound generated by seismic surveys. FCR 089/94. Rep. from Fawley Aquatic Research Laboratories Ltd. for U.K. Offshore Operators Association (UKOOA).
- Turnpenny, A.W.H., K.P. Thatcher, and J.R. Nedwell. 1994. Research report: the effects on fish and other marine animals of high-level underwater sound. FRR 127/94. Rep. from Fawley Aquatic Research Laboratories, Ltd. for the Defence Research Agency.
- Wardle, C.S., T.J. Carter, G.G. Urquhart, A.D.F. Johnstone, A.M. Ziolkowski, G. Hampson, and D. Mackie. 2001. Effects of seismic airguns on marine fish. **Cont. Shelf Res.** 21(8-10):1005-1027.

APPENDIX F

REVIEW OF THE EFFECTS OF AIRGUN SOUNDS ON MARINE INVERTEBRATES⁹

This review provides a detailed summary of the limited data and available literature on the observed effects (or lack of effects) of exposure to airgun sound on marine invertebrates. Specific conditions and results of the studies, including sound exposure levels and sound thresholds of responses, are discussed when available.

Sound caused by underwater seismic survey equipment results in energy pulses with very high peak pressures (Richardson et al. 1995). This was especially true when chemical explosives were used for underwater surveys. Virtually all underwater seismic surveying conducted today uses airguns which typically have lower peak pressures and longer rise times than chemical explosives. However, sound levels from underwater airgun discharges might still be high enough to potentially injure or kill animals located close to the source. Also, there is a potential for disturbance to normal behavior upon exposure to airgun sound. The following sections provide an overview of sound production and detection in marine invertebrates, and information on the effects of exposure to sound on marine invertebrates, with an emphasis on seismic survey sound. In addition, Fisheries and Oceans Canada has published two internal documents that provide a literature review of the effects of seismic and other underwater sound on invertebrates (Moriyasu et al. 2004; Payne et al. 2008). The available information as reviewed in those documents and here includes results of studies of varying degrees of scientific rigor as well as anecdotal information.

1. Sound Production

Much of the available information on acoustic abilities of marine invertebrates pertains to crustaceans, specifically lobsters, crabs and shrimps. Other acoustic-related studies have been conducted on cephalopods. Many invertebrates are capable of producing sound, including barnacles, amphipods, shrimp, crabs, and lobsters (Au and Banks 1998; Tolstoganova 2002). Invertebrates typically produce sound by scraping or rubbing various parts of their bodies, although they also produce sound in other ways. Sounds made by marine invertebrates may be associated with territorial behavior, mating, courtship, and aggression. On the other hand, some of these sounds may be incidental and not have any biological relevance. Sounds known to be produced by marine invertebrates have frequencies ranging from 87 Hz to 200 kHz, depending on the species.

Both male and female American lobsters *Homarus americanus* produce a buzzing vibration with the carapace when grasped (Pye and Watson III 2004; Henninger and Watson III 2005). Larger lobsters vibrate more consistently than smaller lobsters, suggesting that sound production may be involved with mating behavior. Sound production by other species of lobsters has also been studied. Among deep-sea lobsters, sound level was more variable at night than during the day, with the highest levels occurring at the lowest frequencies.

While feeding, king crab *Paralithodes camtschaticus* produce impulsive sounds that appear to stimulate movement by other crabs, including approach behavior (Tolstoganova 2002). King crab also appeared to produce 'discomfort' sounds when environmental conditions were manipulated. These discomfort sounds differ from the feeding sounds in terms of frequency range and pulse duration.

⁹ By **John R. Christian**, LGL Ltd., Environmental Research Associates (revised Nov. 2009).

Snapping shrimp *Synalpheus parneomeris* are among the major sources of biological sound in temperate and tropical shallow-water areas (Au and Banks 1998). By rapidly closing one of its frontal chelae (claws), a snapping shrimp generates a forward jet of water and the cavitation of fast moving water produces a sound. Both the sound and the jet of water may function in feeding and territorial behaviors of alpheididae shrimp. Measured source sound pressure levels (SPLs) for snapping shrimp were 183–189 dB re $1 \mu\text{Pa} \cdot \text{m}_{\text{p-p}}$ and extended over a frequency range of 2–200 kHz.

2. Sound Detection

There is considerable debate about the hearing capabilities of aquatic invertebrates. Whether they are able to hear or not depends on how underwater sound and underwater hearing are defined. In contrast to the situation in fish and marine mammals, no physical structures have been discovered in aquatic invertebrates that are stimulated by the pressure component of sound. However, vibrations (i.e., mechanical disturbances of the water) are also characteristic of sound waves. Rather than being pressure-sensitive, aquatic invertebrates appear to be most sensitive to the vibrational component of sound (Breithaupt 2002). Statocyst organs may provide one means of vibration detection for aquatic invertebrates.

More is known about the acoustic detection capabilities in decapod crustaceans than in any other marine invertebrate group, although cephalopod acoustic capabilities are now becoming a focus of study. Crustaceans appear to be most sensitive to sounds of low frequencies, i.e., <1000 Hz (Budelmann 1992; Popper et al. 2001). A study by Lovell et al. (2005) suggests greater sensitivity of the prawn *Palaemon serratus* to low-frequency sound than previously thought. Lovell et al. (2006) showed that *P. serratus* is capable of detecting a 500 Hz tone regardless of the prawn's body size and the related number and size of statocyst hair cells. Studies of American lobsters suggest that these crustaceans are more sensitive to higher frequency sounds than previously realized (Pye and Watson III 2004).

It is possible that statocyst hair cells of cephalopods are directionally sensitive in a way that is similar to the responses of hair cells of the vertebrate vestibular and lateral line systems (Budelmann and Williamson 1994; Budelmann 1996). Kaifu et al. (2008) provided evidence that the cephalopod *Octopus ocellatus* detects particle motion with its statocyst. Studies by Packard et al. (1990), Rawizza (1995) and Komak et al. (2005) have tested the sensitivities of various cephalopods to water-borne vibrations, some of which were generated by low-frequency sound. Using the auditory brainstem response (ABR) approach, Hu et al. (2009) showed that auditory evoked potentials can be obtained in the frequency ranges 400 to 1500 Hz for the squid *Sepiotheutis lessoniana* and 400 to 1000 Hz for the octopus *Octopus vulgaris*, higher than frequencies previously observed to be detectable by cephalopods.

In summary, only a few studies have been conducted on the sensitivity of certain invertebrate species to underwater sound. Available data suggest that they are capable of detecting vibrations but they do not appear to be capable of detecting pressure fluctuations.

3. Potential Seismic Effects

In marine invertebrates, potential effects of exposure to sound can be categorized as pathological, physiological, and behavioral. Pathological effects include lethal and sub-lethal injury to the animals, physiological effects include temporary primary and secondary stress responses, and behavioral effects refer to changes in exhibited behaviors (i.e., disturbance). The three categories should not be considered as independent of one another and are likely interrelated in complex ways.

Pathological Effects.—In water, acute injury or death of organisms as a result of exposure to sound appears to depend on two features of the sound source: (1) the received peak pressure, and (2) the time required for the pressure to rise and decay. Generally, the higher the received pressure and the less

time it takes for the pressure to rise and decay, the greater the chance of acute pathological effects. Considering the peak pressure and rise/decay time characteristics of seismic airgun arrays used today, the associated pathological zone for invertebrates would be expected to be small (i.e., within a few meters of the seismic source, at most). Few studies have assessed the potential for pathological effects on invertebrates from exposure to seismic sound.

The pathological impacts of seismic survey sound on marine invertebrates were investigated in a pilot study on snow crabs *Chionoecetes opilio* (Christian et al. 2003, 2004). Under controlled field experimental conditions, captive adult male snow crabs, egg-carrying female snow crabs, and fertilized snow crab eggs were exposed to variable SPLs (191–221 dB re 1 μPa_{0-p}) and sound energy levels (SELs) (<130–187 dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$). Neither acute nor chronic (12 weeks post-exposure) mortality was observed for the adult crabs. However, a significant difference in development rate was noted between the exposed and unexposed fertilized eggs/embryos. The egg mass exposed to seismic energy had a higher proportion of less-developed eggs than did the unexposed mass. It should be noted that both egg masses came from a single female and any measure of natural variability was unattainable (Christian et al. 2003, 2004).

In 2003, a collaborative study was conducted in the southern Gulf of St. Lawrence, Canada, to investigate the effects of exposure to sound from a commercial seismic survey on egg-bearing female snow crabs (DFO 2004). This study had design problems that impacted interpretation of some of the results (Chadwick 2004). Caged animals were placed on the ocean bottom at a location within the survey area and at a location outside of the survey area. The maximum received SPL was ~195 dB re 1 μPa_{0-p} . The crabs were exposed for 132 hr of the survey, equivalent to thousands of seismic shots of varying received SPLs. The animals were retrieved and transferred to laboratories for analyses. Neither acute nor chronic lethal or sub-lethal injury to the female crabs or crab embryos was indicated. DFO (2004) reported that some exposed individuals had short-term soiling of gills, antennules and statocysts, bruising of the hepatopancreas and ovary, and detached outer membranes of oocytes. However, these differences could not be linked conclusively to exposure to seismic survey sound. Boudreau et al. (2009) presented the proceedings of a workshop held to evaluate the results of additional studies conducted to answer some questions arising from the original study discussed in DFO (2004). Proceedings of the workshop did not include any more definitive conclusions regarding the original results.

Payne et al. (2007) recently conducted a pilot study of the effects of exposure to airgun sound on various health endpoints of the American lobster. Adult lobsters were exposed either 20 to 200 times to 202 dB re 1 μPa_{p-p} or 50 times to 227 dB re 1 μPa_{p-p} , and then monitored for changes in survival, food consumption, turnover rate, serum protein level, serum enzyme levels, and serum calcium level. Observations extended over a period of a few days to several months. Results showed no delayed mortality or damage to the mechanosensory systems associated with animal equilibrium and posture (as assessed by turnover rate).

In a field study, Pearson et al. (1994) exposed Stage II larvae of the Dungeness crab *Cancer magister* to single discharges from a seven-airgun array and compared their mortality and development rates with those of unexposed larvae. No statistically significant differences were found in immediate survival, long-term survival, or time to molt between the exposed and unexposed larvae, even those exposed within 1 m of the seismic source.

In 2001 and 2003, there were two incidents of multiple strandings of the giant squid *Architeuthis dux* on the north coast of Spain, and there was speculation that the strandings were caused by exposure to geophysical seismic survey sounds occurring at about the same time in the Bay of Biscay (Guerra et al. 2004). A total of nine giant squid, either stranded or moribund and floating at the surface, were collected at these times. However, Guerra et al. (2004) did not present any evidence that conclusively links the

giant squid strandings and floaters to seismic activity in the area. Based on necropsies of seven (six females and one male) specimens, there was evidence of acute tissue damage. The authors speculated that one female with extensive tissue damage was affected by the impact of acoustic waves. However, little is known about the impact of strong airgun signals on cephalopods and the authors did not describe the seismic sources, locations, and durations of the Bay of Biscay surveys. In addition, there were no controls, the observations were circumstantial, and the examined animals had been dead long enough for commencement of tissue degradation.

McCauley et al. (2000a,b) exposed caged cephalopods to noise from a single 20-in³ airgun with maximum SPLs of >200 dB re 1 μPa_{0-p} . Statocysts were removed and preserved, but at the time of publication, results of the statocyst analyses were not available. No squid or cuttlefish mortalities were reported as a result of these exposures.

Physiological Effects.—Biochemical responses by marine invertebrates to acoustic exposure have also been studied to a limited degree. Such studies of stress responses could possibly provide some indication of the physiological consequences of acoustic exposure and perhaps any subsequent chronic detrimental effects. Stress responses could potentially affect animal populations by reducing reproductive capacity and adult abundance.

Stress indicators in the haemolymph of adult male snow crabs were monitored immediately after exposure of the animals to seismic survey sound (Christian et al. 2003, 2004) and at various intervals after exposure. No significant acute or chronic differences were found between exposed and unexposed animals in which various stress indicators (e.g., proteins, enzymes, cell type count) were measured.

Payne et al. (2007), in their study of the effects of exposure of adult American lobsters to airgun sound, noted decreases in the levels of serum protein, particular serum enzymes and serum calcium, in the haemolymph of animals exposed to the sound pulses. Statistically significant differences ($P=0.05$) were noted in serum protein at 12 days post-exposure, serum enzymes at 5 days post-exposure, and serum calcium at 12 days post-exposure. During the histological analysis conducted 4 months post-exposure, Payne et al. (2007) noted more deposits of PAS-stained material, likely glycogen, in the hepatopancreas of some of the exposed lobsters. Accumulation of glycogen could be due to stress or disturbance of cellular processes.

Price (2007) found that blue mussels *Mytilus edulis* responded to a 10 kHz pure tone continuous signal by decreasing respiration. Smaller mussels did not appear to react until exposed for 30 min whereas larger mussels responded after 10 min of exposure. The oxygen uptake rate tended to be reduced to a greater degree in the larger mussels than in the smaller animals.

In general, the limited studies done to date on the effects of acoustic exposure on marine invertebrates have not demonstrated any serious pathological and physiological effects.

Behavioral Effects.—Some recent studies have focused on potential behavioral effects on marine invertebrates.

Christian et al. (2003) investigated the behavioral effects of exposure to airgun sound on snow crabs. Eight animals were equipped with ultrasonic tags, released, and monitored for multiple days prior to exposure and after exposure. Received SPL and SEL were ~191 dB re 1 μPa_{0-p} and <130 dB re 1 $\mu\text{Pa}^2 \cdot \text{s}$, respectively. The crabs were exposed to 200 discharges over a 33-min period. None of the tagged animals left the immediate area after exposure to the seismic survey sound. Five animals were captured in the snow crab commercial fishery the following year, one at the release location, one 35 km from the release location, and three at intermediate distances from the release location.

Another study approach used by Christian et al. (2003) involved monitoring snow crabs with a remote video camera during their exposure to airgun sound. The caged animals were placed on the ocean bottom at a depth of 50 m. Received SPL and SEL were ~ 202 dB re $1 \mu\text{Pa}_{0-p}$ and 150 dB re $1 \mu\text{Pa}^2 \cdot \text{s}$, respectively. The crabs were exposed to 200 discharges over a 33-min period. They did not exhibit any overt startle response during the exposure period.

Christian et al. (2003) also investigated the pre- and post-exposure catchability of snow crabs during a commercial fishery. Received SPLs and SELs were not measured directly and likely ranged widely considering the area fished. Maximum SPL and SEL were likely similar to those measured during the telemetry study. There were seven pre-exposure and six post-exposure trap sets. Unfortunately, there was considerable variability in set duration because of poor weather. Results indicated that the catch-per-unit-effort did not decrease after the crabs were exposed to seismic survey sound.

Parry and Gason (2006) statistically analyzed data related to rock lobster *Jasus edwardsii* commercial catches and seismic surveying in Australian waters from 1978 to 2004. They did not find any evidence that lobster catch rates were affected by seismic surveys.

Caged female snow crabs exposed to airgun sound associated with a recent commercial seismic survey conducted in the southern Gulf of St. Lawrence, Canada, exhibited a higher rate of ‘righting’ than those crabs not exposed to seismic survey sound (J. Payne, Research Scientist, DFO, St. John’s, Nfld., pers. comm.). ‘Righting’ refers to a crab’s ability to return itself to an upright position after being placed on its back. Christian et al. (2003) made the same observation in their study.

Payne et al. (2007), in their study of the effects of exposure to airgun sound on adult American lobsters, noted a trend for increased food consumption by the animals exposed to seismic sound.

Andriguetto-Filho et al. (2005) attempted to evaluate the impact of seismic survey sound on artisanal shrimp fisheries off Brazil. Bottom trawl yields were measured before and after multiple-day shooting of an airgun array. Water depth in the experimental area ranged between 2 and 15 m. Results of the study did not indicate any significant deleterious impact on shrimp catches. Anecdotal information from Newfoundland, Canada, indicated that catch rates of snow crabs showed a significant reduction immediately following a pass by a seismic survey vessel (G. Chidley, Newfoundland fisherman, pers. comm.). Additional anecdotal information from Newfoundland indicated that a school of shrimp observed via a fishing vessel sonar shifted downwards and away from a nearby seismic airgun sound source (H. Thorne, Newfoundland fisherman, pers. comm.). This observed effect was temporary.

Caged brown shrimp *Crangon crangon* reared under different acoustical conditions exhibited differences in aggressive behavior and feeding rate (Lagardère 1982). Those exposed to a continuous sound source showed more aggression and less feeding behavior. It should be noted that behavioral responses by caged animals may differ from behavioral responses of animals in the wild.

McCauley et al. (2000a,b) provided the first evidence of the behavioral response of southern calamari squid *Sepioteuthis australis* exposed to seismic survey sound. McCauley et al. reported on the exposure of caged cephalopods (50 squid and two cuttlefish) to noise from a single 20-in³ airgun. The cephalopods were exposed to both stationary and mobile sound sources. The two-run total exposure times during the three trials ranged from 69 to 119 min. at a firing rate of once every 10–15 s. The maximum SPL was >200 dB re $1 \mu\text{Pa}_{0-p}$. Some of the squid fired their ink sacs apparently in response to the first shot of one of the trials and then moved quickly away from the airgun. In addition to the above-described startle responses, some squid also moved towards the water surface as the airgun approached. McCauley et al. (2000a,b) reported that the startle and avoidance responses occurred at a received SPL of 174 dB re $1 \mu\text{Pa}_{\text{rms}}$. They also exposed squid to a ramped approach-depart airgun signal whereby the received SPL was gradually increased over time. No strong startle response (i.e., ink discharge) was

observed, but alarm responses, including increased swimming speed and movement to the surface, were observed once the received SPL reached a level in the 156–161 dB re 1 $\mu\text{Pa}_{\text{rms}}$ range.

Komak et al. (2005) also reported the results of a study of cephalopod behavioral responses to local water movements. In this case, juvenile cuttlefish *Sepia officinalis* exhibited various behavioral responses to local sinusoidal water movements of different frequencies between 0.01 and 1000 Hz. These responses included body pattern changing, movement, burrowing, reorientation, and swimming. Similarly, the behavioral responses of the octopus *Octopus ocellatus* to non-impulse sound have been investigated by Kaifu et al. (2007). The sound stimuli, reported as having levels 120 dB re 1 $\mu\text{Pa}_{\text{rms}}$, were at various frequencies: 50, 100, 150, 200 and 1000 Hz. The respiratory activity of the octopus changed when exposed to sound in the 50–150 Hz range but not for sound at 200–1,000 Hz. Respiratory suppression by the octopus might have represented a means of escaping detection by a predator.

Low-frequency sound (<200 Hz) has also been used as a means of preventing settling/fouling by aquatic invertebrates such as zebra mussels *Dreissena polymorpha* (Donskoy and Ludyanskiy 1995) and balanoid barnacles *Balanus* sp. (Branscomb and Rittschof 1984). Price (2007) observed that blue mussels *Mytilus edulis* closed their valves upon exposure to 10 kHz pure tone continuous sound.

Although not demonstrated in the invertebrate literature, masking can be considered a potential effect of anthropogenic underwater sound on marine invertebrates. Some invertebrates are known to produce sounds (Au and Banks 1998; Tolstoganova 2002; Latha et al. 2005). The functionality and biological relevance of these sounds are not understood (Jeffs et al. 2003, 2005; Lovell et al. 2005; Radford et al. 2007). If some of the sounds are of biological significance to some invertebrates, then masking of those sounds or of sounds produced by predators, at least the particle displacement component, could potentially have adverse effects on marine invertebrates. However, even if masking does occur in some invertebrates, the intermittent nature of airgun sound is expected to result in less masking effect than would occur with continuous sound.

4. Literature Cited

- Andriguetto-Filho, J.M., A. Ostrensky, M.R. Pie, U.A. Silva, and W.A. Boeger. 2005. Evaluating the impact of seismic prospecting on artisanal shrimp fisheries. **Cont. Shelf Res.** 25:1720-1727.
- Au, W.W.L. and K. Banks. 1998. The acoustics of snapping shrimp *Synalpheus parneomeris* in Kaneohe Bay. **J. Acoust. Soc. Am.** 103:41-47.
- Boudreau, M., S.C. Courtenay, and K. Lee (eds.). 2009. Proceedings of a workshop held 23 January 2007 at the Gulf Fisheries Center, Potential impacts of seismic energy on snow crab: An update to the September 2004 review. **Can. Tech. Rep. Fish. Aquat. Sci.** 2836.
- Branscomb, E.S. and D. Rittschof. 1984. An investigation of low frequency sound waves as a means of inhibiting barnacle settlement. **J. Exp. Mar. Biol. Ecol.** 79:149-154.
- Breithaupt, T. 2002. Sound perception in aquatic crustaceans. p. 548-558 In: K. Wiese (ed.), *The crustacean nervous system*. Springer-Verlag, Berlin-Heidelberg, Germany. 623 p.
- Budelmann, B.U. 1992. Hearing in crustacea. p. 131-139 In: D.B. Webster, R.R. Fay, and A.N. Popper (eds.), *Evolutionary biology of hearing*. Springer-Verlag, New York, NY.
- Budelmann, B.U. 1996. Active marine predators: the sensory world of cephalopods. **Mar. Freshw. Behav. Physiol.** 27:59-75.
- Budelmann, B.U. and R. Williamson. 1994. Directional sensitivity of hair cell afferents in the octopus statocyst. **J. Exp. Biol.** 187:245-259.
- Chadwick, M. 2004. Proceedings of the peer review on potential impacts of seismic energy on snow crab. Gulf Region, Department of Fisheries and Oceans Canada, Science Advisory Secretariat Proceedings Series 2004/045.

- Christian, J.R., A. Mathieu, D.H. Thomson, D. White, and R.A. Buchanan. 2003. Effect of seismic energy on snow crab (*Chionoecetes opilio*). Environ. Stud. Res. Funds Rep. 144. Calgary, Alberta.
- Christian, J.R., A. Mathieu, and R.A. Buchanan. 2004. Chronic effects of seismic energy on snow crab (*Chionoecetes opilio*). Environ. Stud. Res. Funds Rep. 158, Calgary, Alberta.
- DFO. 2004. Potential impacts of seismic energy on snow crab. Canadian Science Advisory Secretariat Habitat Status Report 2004/003.
- Donskoy, D.M. and M.L. Ludyanskiy. 1995. Low frequency sound as a control measure for zebra mussel fouling. Proc. 5th Int. Zebra Mussel and Other Aquatic Nuisance Organisms Conference, February 1995, Toronto, Ont.
- Guerra, A., A.F. González, and F. Rocha. 2004. A review of the records of giant squid in the north-eastern Atlantic and severe injuries in *Architeuthis dux* stranded after acoustic explorations. Paper presented at the International Council for the Exploration of the Sea (ICES) Annual Science Conference, 22–25 Sept. 2004, Vigo, Spain. ICES CM 2004/CC:29.
- Henninger, H.P. and W.H. Watson, III. 2005. Mechanisms underlying the production of carapace vibrations and associated waterborne sounds in the American lobster, *Homarus americanus*. **J. Exp. Biol.** 208:3421-3429.
- Hu, M.Y., H.Y. Yan, W-S Chung, J-C Shiao, and P-P Hwang. 2009. Acoustically evoked potentials in two cephalopods inferred using the auditory brainstem response (ABR) approach. **Comp. Biochem. Physiol. A** 153:278-283.
- Jeffs, A., N. Tolimieri, and J.C. Montgomery. 2003. Crabs on cue for the coast: the use of underwater sound for orientation by pelagic crab stages. **Mar. Freshw. Res.** 54:841-845.
- Jeffs, A.G., J.C. Montgomery, and C.T. Tindle. 2005. How do spiny lobster post-larvae find the coast? **N.Z. J. Mar. Fresh. Res.** 39:605-617.
- Kaifu, K., S. Segawa, and K. Tsuchiya. 2007. Behavioral responses to underwater sound in the small benthic octopus *Octopus ocellatus*. **J. Mar. Acoust. Soc. Japan** 34:46-53.
- Kaifu, K., T. Akamatsu, and S. Segawa. 2008. Underwater sound detection by cephalopod statocyst. **Fish. Sci.** 74:781-786.
- Komak, S., J.G. Boal, L. Dickel, and B.U. Budelmann. 2005. Behavioural responses of juvenile cuttlefish (*Sepia officinalis*) to local water movements. **Mar. Freshw. Behav. Physiol.** 38:117-125.
- Lagardère, J.P. 1982. Effects of noise on growth and reproduction of *Crangon crangon* in rearing tanks. **Mar. Biol.** 71:177-186.
- Latha, G., S. Senthilvadivu, R. Venkatesan, and V. Rajendran. 2005. Sound of shallow and deep water lobsters: measurements, analysis, and characterization (L). **J. Acoust. Soc. Am.** 117: 2720-2723.
- Lovell, J.M., M.M. Findley, R.M. Moate, and H.Y. Yan. 2005. The hearing abilities of the prawn *Palaemon serratus*. **Comp. Biochem. Physiol. A** 140:89-100.
- Lovell, J.M., R.M. Moate, L. Christiansen, and M.M. Findlay. 2006. The relationship between body size and evoked potentials from the statocysts of the prawn *Palaemon serratus*. **J. Exp. Biol.** 209:2480-2485.
- McCauley, R.D., J. Fewtrell, A.J. Duncan, C. Jenner, M.-N. Jenner, J.D. Penrose, R.I.T. Prince, A. Adhitya, J. Murdoch, and K. McCabe. 2000a. Marine seismic surveys: analysis of airgun signals; and effects of air gun exposure on humpback whales, sea turtles, fishes and squid. Rep. from Centre for Marine Science and Technology, Curtin University, Perth, Western Australia, for Australian Petroleum Production Association, Sydney, NSW.
- McCauley, R.D., J. Fewtrell, A.J. Duncan, C. Jenner, M.-N. Jenner, J.D. Penrose, R.I.T. Prince, A. Adhitya, J. Murdoch, and K. McCabe. 2000b. Marine seismic surveys – a study of environmental implications. **APPEA J.** 40:692-706.
- Moriyasu, M., R. Allain, K. Benhalima, and R. Claytor. 2004. Effects of seismic and marine noise on invertebrates: A literature review. Fisheries and Oceans Canada, Science. Canadian Science Advisory Secretariat Research Document 2004/126.

- Packard, A., H.E. Karlsen, and O. Sand. 1990. Low frequency hearing in cephalopods. **J. Comp. Physiol. A** 166: 501-505.
- Parry, G.D. and A. Gason. 2006. The effect of seismic surveys on catch rates of rock lobsters in western Victoria, Australia. **Fish. Res.** 79:272-284.
- Payne, J.F., C.A. Andrews, L.L. Fancey, A.L. Cook, and J.R. Christian. 2007. Pilot study on the effects of seismic air gun noise on lobster (*Homarus americanus*). Can. Tech. Rep. Fish. Aquatic Sci. 2712.
- Payne, J.F., C. Andrews, L. Fancey, D. White, and J. Christian. 2008. Potential effects of seismic energy on fish and shellfish: An update since 2003. Fisheries and Oceans Canada Science, Canadian Science Advisory Secretariat Research Document 2008/060.
- Pearson, W., J. Skalski, S. Sulkin, and C. Malme. 1994. Effects of seismic energy releases on the survival and development of zoeal larvae of Dungeness crab (*Cancer magister*). **Mar. Environ. Res.** 38:93-113.
- Popper, A.N., M. Salmon, and K.W. Horch. 2001. Acoustic detection and communication by decapod crustaceans. **J. Comp. Physiol. A** 187:83-89.
- Price, A. 2007. The effects of high frequency, high intensity underwater sound on the oxygen uptakes of *Mytilus edulis* (L.). B.Sc.(Hons.) Thesis, Heriot-Watt Univ., Scotland.
- Pye, H.J., and W.H. Watson, III. 2004. Sound detection and production in the American lobster, *Homarus americanus*: sensitivity range and behavioral implications. **J. Acoust. Soc. Am.** 115 (Part 2):2486.
- Radford, C.A., A.G. Jeffs, and J.C. Montgomery. 2007. Orientated swimming behavior of crab postlarvae in response to reef sound. Poster at First Intern. Conf. on Effects of Noise on Aquatic Life, Nyborg, Denmark, Aug. 2007.
- Rawizza, H.E. 1995. Hearing and associative learning in cuttlefish, *Sepia officinalis*. Hopkins Marine Station Student Paper. Stanford Univ., Palo Alto, CA.
- Richardson, W.J., C.R. Greene, Jr., C.I. Malme, and D.H. Thomson. 1995. Marine mammals and noise. Academic Press, San Diego, CA. 576 p.
- Tolstoganova, L.K. 2002. Acoustical behavior in king crab (*Paralithodes camtschaticus*). p. 247-254 In: A.J. Paul, E.G. Dawe, R. Elner, G.S. Jamieson, G.H. Kruse, R.S. Otto, B. Sainte-Marie, T.C. Shirley, and D. Woodby (eds.), Crabs in cold water regions: biology, management, and economics. University of Alaska Sea Grant, AK-SG-02-01, Fairbanks, AK.

Addendum to the Draft Environmental Assessment of a Marine Seismic Survey of the Arctic Ocean U.S. Geological Survey in 2010

This addendum supplements the environmental assessment (EA) for the proposed marine seismic survey of portions of the Arctic Ocean to be conducted by the U.S. Geological Survey (USGS) in the late summer-early fall of 2010. USGS conducted early coordination with the National Marine Fisheries Service (NMFS) and solicited their comments on the preliminary draft of the subject EA. Supplemental information to the draft EA was requested by NMFS to address potential marine mammal “takes” from icebreaking activity intrinsic to the project.

Icebreaking is considered by NMFS to be a continuous sound and NMFS (2005) indicates the existing threshold for Level B harassment by continuous sounds is a received sound level of 120 dB SPL. Potential takes of marine mammals may ensue from the icebreaking activity in which the USCGC *Healy* is expected to engage outside of U.S. waters, i.e. north of ~74.1°N. While breaking ice, the noise from the ship, including impact with ice, engine noise, and propeller cavitation, will exceed 120 dB continuously. The draft EA presents take estimates based exclusively on the seismic survey component of the project within U.S. waters. If icebreaking does occur in U.S. waters, we expect it will occur during seismic operations. The safety radius for the marine mammal Level B harassment threshold during the proposed seismic activities is greater than the calculated radius during icebreaking. Therefore, if the *Healy* breaks ice during seismic operations within the U.S. waters, the greater radius, i.e. that for seismic operations, supersedes that for icebreaking, so no additional takes have been estimated within U.S. waters. This addendum presents calculations of exposures to marine mammals due to icebreaking only outside U.S. waters when the USCG *Healy* will be breaking ice for the *Louis S. St. Laurent*.

It is important to note that non-icebreaking vessels, as well as natural sounds such as those arising from sea ice motion and whale flukes hitting the ocean surface, also present similar sound impacts. Underwater noise from various vessels, including tug boats, oceanographic research vessels, and fisheries research vessels in open water, as well as icebreakers traversing sea ice, often exceed 120 dB, the existing threshold for Level B harassment set by NMFS (2005).

The sound level and other estimates provided in this addendum are for information purposes only and do not represent any conclusions with regard to harassment. Further studies are needed before a precedent can be established.

The objectives and plans of the proposed project remain unchanged. The following includes specifics of the estimation of trackline while the USCGC *Healy* breaks ice outside U.S. waters and the calculation of the resulting potential takes. The supplemental information has been organized in a manner consistent with the draft EA. The estimated takes provided in this addendum are in addition to the number of estimated takes due to seismic activities within U.S. waters that are presented in the Incidental Harassment Authorization (IHA) Application submitted to the NMFS on 28 May 2010.

II. Alternatives Including Proposed Action

Proposed Action

(2) Proposed Activities

The proposed geophysical survey will be conducted for ~28 days from approximately 7 August to 3 September 2010. Icebreaking outside U.S. waters will occur between the latitudes of ~74 to 84 °N. Vessel operations and ice conditions from similar survey activities and timing in 2008 and 2009 were used to estimate the amount of icebreaking (in trackline km) that is likely to occur in 2010.

We expect that the *Louis S. St. Laurent* and the *Healy* will be working in tandem through the ice for a maximum of 23–25 days while outside U.S. waters. The average distance travelled in 2008 and 2009 when the *Healy* broke ice for the *Louis S. St. Laurent* was 135 km/d (Table Add-1). Based on the 23–25 day period of icebreaking, we calculate that, at most ~3102–3372 km of vessel trackline may involve icebreaking. This calculation is likely an overestimation because icebreakers often follow leads when they are available and thus do not break ice at all times.

TABLE Add-1. Projected 2010 icebreaking effort for USGS/ GSC 2010 Extended Continental Shelf Survey in the northern Beaufort Sea and Arctic Ocean.

| | 2-ship operations (days) | 2-ship operations (km) | km/day |
|-----------------------|-----------------------------|---------------------------|--------|
| 2008 | 19 | 2469 | 130 |
| 2009 | 27 | 3774 | 140 |
| Avg. 2008-2009 | 23 | 3122 | 135 |
| Projected 2010 | 23-25 | 3102-3372 | -- |

III. Affected Environment

Within the latitudes of the proposed survey when the *Healy* will be breaking ice outside of U.S. waters, no cetaceans were observed by marine mammal observers (MMOs) along approximately 21,322 km of effort during projects in 2005, 2006, 2008 and 2009 (Haley and Ireland 2006, Haley 2006, Jackson and DesRoches 2008, Mosher et al. 2009). The estimated maximum amount of icebreaking outside of U.S. waters for this project, i.e. 3372 line km, is considerably less than the combined trackline for the aforementioned projects. At least one MMO will stand watch at all times while the *Healy* is breaking ice for the *Louis S. St. Laurent*. We do not expect that MMOs will observe any cetaceans during the proposed survey.

Seals and polar bears were reported by MMOs during the 2005, 2006, 2008 and 2009 effort within the latitudes of the proposed survey (Table Add-2).

TABLE Add-2. Number of marine mammals reported during 2005, 2006, 2008 and 2009 projects within the latitudes where the *Healy* will be breaking ice outside of U.S. waters for the proposed Arctic Ocean survey (Haley and Ireland 2006, Haley 2006, Geological Survey of Canada [GSC] unpubl. data 2008, Mosher et al. 2009).

| Species | No. of Sightings | No. of Individuals |
|-------------------------|-------------------------|---------------------------|
| <i>Pinnipeds</i> | | |
| Ringed seal | 116 | 125 |
| Bearded seal | 24 | 26 |
| Unidentified seal | 128 | 140 |
| <i>Ursidea</i> | | |
| Polar bear | 39 | 53 |

IV. ENVIRONMENTAL CONSEQUENCES OF PROPOSED ACTION

(5) Possible Effects of Icebreaking Activities

The *Healy* is designed for continuous passage at 3 kt through ice 1.4 m thick. During this project the *Healy* will typically encounter first- or second-year ice while avoiding thicker ice floes, particularly large intact multi-year ice, whenever possible. In addition, the icebreaker will follow leads when possible while following the survey route. As the icebreaker passes through the ice, the ship causes the ice to part and travel alongside the hull. This ice typically returns to fill the wake as the ship passes. The effects are transitory, i.e. hours at most, and localized, i.e. constrained to a relatively narrow swath perhaps 10 m to each side of the vessel (Fig. Add-1).

Healy's maximum beam is 25 m (Appendix D of the original application). Applying the maximum estimated amount of icebreaking, i.e. 3372 km, to the corridor opened by the ship, we anticipate that a maximum of ~152 km² of ice may be disturbed. This encompasses an insignificant amount (<0.005%) of the total Arctic ice extent in Aug and Sep of 2008 and 2009 which ranged from 3.24 million km² to 4.1 million km².



FIGURE ADD-1. Icebreakers *Healy* and *Louis S. St-Laurent* transiting 10/10 ice pack on 2 September 2009, showing minimal disturbance to the ice pack abeam of the ship's path, small jog to avoid larger ice body, and closing of the pack ice in the ship wake.

Icebreaking will create temporary leads in the ice and could possibly destroy unoccupied seal lairs. Seal pups are born in the spring, therefore, pupping and nursing will have concluded and the lairs will be vacated at the time of the proposed survey. Breaking ice may damage seal breathing holes and will also reduce the haulout area in the immediate vicinity of the ship's track.

Icebreaking along a maximum of 3372 km of trackline will alter local ice conditions in the immediate vicinity of the vessel. This has the potential to temporarily lead to a reduction of suitable seal haul-out habitat. However the dynamic sea-ice environment requires that seals be able to adapt to changes in sea, ice, and snow conditions, and they therefore create new breathing holes and lairs throughout winter and spring (Hammill and Smith 1989). In addition, seals often use open leads and cracks in the ice to surface and breathe (Smith and Stirling 1975). Disturbance to the ice will occur in a very small area (<0.005%) relative to the Arctic icepack and no significant impact on marine mammals is anticipated by icebreaking during the proposed project.

(7) Numbers of Marine Mammals that May be “Taken by Harassment”

All anticipated takes would be “takes by harassment”, as described in § V of the original application, involving temporary changes in behavior. The mitigation measures to be applied will minimize the possibility of injurious takes. (However, as noted in Appendix D of the original application, there is no specific information demonstrating that injurious “takes” would occur even in the absence of the planned mitigation measures.) The sections below describe methods used to estimate “take by harassment” and present estimates of the numbers of marine mammals that might be affected during the proposed seismic study in the Arctic Ocean.

Few data (systematic or otherwise) are available on the distribution and numbers of marine mammals in the northern Beaufort Sea or offshore water of the Arctic Ocean. Both “maximum estimates” as well as “best estimates” of marine mammal densities (Table Add-3) and the numbers of marine mammals potentially exposed to underwater sound (Table Add-4) were calculated as described below. The best (or average) estimate is based on available distribution and abundance data and represents the most likely number of animals that may be encountered during the survey, assuming no avoidance of the airguns or vessel. The maximum estimate is either the highest estimate from applicable distribution and abundance data or the average estimate increase by a multiplier intended to produce a very conservative (over) estimate of the number of animals that may be present in the survey area. There is some uncertainty about how representative the available data are and the assumptions used below to estimate the potential “take by harassment”. However, the approach used here is accepted by NMFS as the best available at this time.

The following estimates are based on a consideration of the number of marine mammals that might be disturbed appreciably over the ~3102–3372 line kilometers of icebreaking that may occur during the proposed project as described above .

(a) Marine Mammal Density Estimates

Numbers of marine mammals that might be present and potentially disturbed are estimated below based on available data about mammal distribution and densities in the area. “Take by harassment” is calculated by multiplying the expected densities of marine mammals likely to occur in the survey area by the area of water potentially ensonified to sound levels ≥ 120 dB re 1 μ Pa (rms). This section provides descriptions of the estimated densities of marine mammals that may occur in the survey area.

No published densities of marine mammals are available for the region of the proposed survey between 74°N and 84°N where the *Healy* will be breaking ice outside U.S. waters. However, vessel-based surveys through the general area in 2005, 2006, 2008 and 2009 encountered few marine mammals as described in § IV in the original application. MMOs recorded 268 sightings of 291 individual seals along ~21,322 km of monitored trackline between 74°N and 84°N (Haley and Ireland 2006, Haley 2006, GSC unpubl. data 2008, Mosher et al. 2009). Thirty-nine sightings of 53 individual polar bears were also reported. No cetaceans were observed during the surveys between 74°N and 84°N.

Given the few sightings of marine mammals along the ~21,322 km vessel trackline in previous years, we estimate that the densities of marine mammals encountered while breaking ice will be 1/10 of the estimated densities of mammals that may be encountered within the ice margin habitat described in the original application (Table Add-3).

TABLE Add-3. Expected summer densities of marine mammals in ice margin (from the original application) and polar pack ice habitats in the Arctic Ocean. Densities are corrected for $f(0)$ and $g(0)$ biases. Species listed as endangered are in italics.

| Species | Ice Margin | | Polar Pack | |
|-----------------------|--|--|--|--|
| | Average Density (# / km ²) | Maximum Density (# / km ²) | Average Density (# / km ²) | Maximum Density (# / km ²) |
| Odontocetes | | | | |
| Beluga | 0.0354 | 0.0709 | 0.0035 | 0.0071 |
| Narwhal | 0.0000 | 0.0002 | 0.0000 | 0.0001 |
| Delphinidae | | | | |
| Killer whale | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| Phocoenidae | | | | |
| Harbor porpoise | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| Mysticetes | | | | |
| <i>Bowhead whale</i> | 0.0061 | 0.0122 | 0.0006 | 0.0012 |
| Gray whale | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| Minke whale | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| <i>Fin whale</i> | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| <i>Humpback whale</i> | 0.0000 | 0.0001 | 0.0000 | 0.0001 |
| Pinnipeds | | | | |
| Bearded seal | 0.0128 | 0.0512 | 0.0013 | 0.0051 |
| Spotted seal | 0.0001 | 0.0004 | 0.0000 | 0.0000 |
| Ringed seal | 0.2510 | 1.0040 | 0.0251 | 0.1004 |
| Pacific Walrus | 0.0001 | 0.0004 | 0.0000 | 0.0000 |
| Ursidae | | | | |
| Polar Bear | 0.0001 | 0.0016 | 0.0000 | 0.0002 |

(b) Estimation of Area Ensonified to Sound Levels ≥ 120 dB rms

The area potentially exposed to received levels ≥ 120 dB due to icebreaking operations was estimated by multiplying the anticipated trackline distance breaking ice by the estimated cross-track distance to received levels of 120 dB caused by icebreaking.

In 2008, Scripps Institute of Oceanography Marine Physical Laboratory conducted measurements of sound pressure levels (SPL) of *Healy* icebreaking under various conditions (Roth and Schmidt 2010). The results indicated that the highest mean sound pressure level (SPL; 185 dB) was measured at survey speeds of 4 to 4.5 kt in conditions of 5/10 ice and greater. Mean SPL under conditions where the ship was breaking heavy ice by backing and ramming was actually lower (180 dB). In addition, when backing and ramming, the vessel is essentially stationary, so the ensonified area is limited for a short period (on the order of minutes to tens of minutes) to the immediate vicinity of the boat until the ship breaks free and once again makes headway.

Although the report by Roth and Schmidt has not yet been reviewed externally nor peer-reviewed for publication, the SPL results reported are consistent with previous studies (Thiele, 1981, 1988; LGL and Greeneridge, 1986, Richardson and others, 1995).

NMFS (2005) indicates the existing threshold for Level B harassment for continuous sounds is a received sound level of 120 dB SPL. Therefore, we estimated the 120 dB received sound level radius around the *Healy* while icebreaking. Using a spherical spreading model, a source level of 185 dB decays to 120 dB in about 1750 m. This model is corroborated by Roth and Schmidt (2010). Therefore, as the ship travels through the ice, a swath 3500 m wide would be subject to sound levels ≥ 120 dB. This results in the potential exposure of 11,802 km² to sounds ≥ 120 dB from icebreaking.

(c) Potential Number of Marine Mammal “Exposures” to Sound Levels ≥ 120

Numbers of marine mammals that might be present and potentially disturbed are estimated below based on available data about mammal distribution and densities in the Arctic Ocean during the summer as described above.

The number of individuals of each species potentially exposed to received levels ≥ 120 dB re 1 μ Pa (rms) by icebreaking was estimated by multiplying

- the anticipated area to be ensonified to ≥ 120 dB, by
- the expected species density

Some of the animals estimated to be exposed to sound levels ≥ 120 dB re 1 μ Pa, particularly migrating bowhead whales, might show avoidance reactions before actual exposure to this sound level (Appendix D). Thus, these calculations actually estimate the number of individuals potentially exposed to ≥ 120 dB rms that would occur if there were no avoidance of the area ensonified to that level.

Based on the operational plans and marine mammal densities described above, the estimates of marine mammals potentially exposed to sounds ≥ 120 dB during the maximum estimation of icebreaking outside U.S. waters, i.e. 3372 km, are presented in Table Add-4. For the common species, the requested numbers are calculated as described above and based on the average densities from the data reported in the different studies mentioned above. For less common species, estimates were set to minimal values to allow for chance encounters.

TABLE ADD-4. Estimates of the numbers of marine mammals potentially exposed to received sound levels ≥ 120 dB during USGS's proposed seismic program while breaking ice outside of U.S. waters. Species in italics are listed under the U.S. ESA as endangered.

| Number of Exposures to Sound Levels ≥ 120 dB | | |
|---|-------------------|----------------|
| Species | Polar Pack | |
| | Average | Maximum |
| Odontocetes | | |
| Monodontidae | | |
| Beluga | 42 | 84 |
| Narwhal | 0 | 1 |
| Delphinidae | | |
| Killer whale | 0 | 1 |
| Phocoenidae | | |
| Harbor porpoise | 0 | 1 |
| Mysticetes | | |
| <i>Bowhead whale</i> | 7 | 1 |
| Gray whale | 0 | 1 |
| Minke whale | 0 | 1 |
| <i>Fin whale</i> | 0 | 1 |
| <i>Humpback whale</i> | 0 | 0 |
| Total Cetaceans | 49 | 92 |
| Pinnipeds | | |
| Bearded seal | 15 | 60 |
| Spotted seal | 0 | 0 |
| Ringed seal | 296 | 1185 |
| Pacific walrus | 0 | 0 |
| Total Pinnipeds | 311 | 1245 |
| Ursidea | | |
| Polar Bear | 0 | 2 |

Literature Cited

- Haley, B. 2006. Marine mammal monitoring during University of Texas at Austin's marine geophysical survey of the western Canada Basin, Chukchi Borderland and Mendeleev Ridge, Arctic Ocean, July–August 2006. Report from LGL Alaska Research Associates, Inc., Anchorage AK, and LGL Ltd., King City, Ont., for the University of Texas at Austin, the Nat. Mar. Fish. Serv., Silver Springs, MD, and the U.S. Fish and Wildl. Serv., Anchorage, AK.
- Haley, B. and D. Ireland. 2006. Marine mammal monitoring during University of Alaska Fairbanks' marine geophysical survey across the Arctic Ocean, August-September 2005. LGL Rep. TA4122-3. Rep. from LGL Ltd., King City, Ont., for Univ. Alaska Fairbanks, Fairbanks, AK, and Nat. Mar. Fish. Serv., Silver Spring, MD. 80 p.
- Hammill, M.O. and T.G. Smith. 1989. Factors affecting the distribution and abundance of ringed seal structures in Barrow Strait, Northwest Territories. *Canadian Journal of Zoology* 67(9): 2212-2219.
- Jackson, H.R. and K.J. DesRoches, eds. 200., 2008 Louis S. St-Laurent field report, August 22 - October 3, 2008: Geological Survey of Canada Open File 6275, 180 pp.
- Mosher, D.C., J.W. Shimeld, and D.R. Hutchinson. 2009. 2009 Canada Basin seismic reflection and refraction survey, western Arctic Ocean: CCGS Louis S. St. Laurent expedition report. Geological Survey of Canada open file 6343.
- NMFS. 2005. Endangered Fish and Wildlife; Notice of Intent to Prepare an Environmental Impact Statement. **Fed. Regist.** 70(7, 11 Jan.):1871-1875.
- Roth, E. H, and V. Schmidt. 2010. Noise levels generated by research icebreakers and marine seismic sources in the deep-water, Arctic Ocean. MPL Tech. Mem. 527.
- Smith, T.G. and I. Stirling. 1975. The breeding habitat of the ringed seal (*Phoca hispida*). The birth lair and associated structures. *Can. J. Zool* 53:1297-1305.