


Prepared for

MARINE CORPS AIR STATION CHERRY POINT, NORTH CAROLINA
ENVIRONMENTAL AFFAIRS DEPARTMENT

February 2009



**FINAL MARINE MAMMAL PROTECTION ACT
COMPLIANCE REPORT
FOR BOMBING TARGETS BT-9 AND BT-11**

**AT MARINE CORPS AIR STATION CHERRY POINT
NORTH CAROLINA**

This page is intentionally blank.

**MARINE MAMMAL PROTECTION ACT
COMPLIANCE REPORT FOR ACTIVITIES AT
MARINE CORPS AIR STATION CHERRY POINT,
NORTH CAROLINA**

FINAL REPORT

PREPARED FOR:

**MARINE CORPS AIR STATION CHERRY POINT
ENVIRONMENTAL AFFAIRS DEPARTMENT
MR. CARMEN LOMBARDO
CHERRY POINT, NORTH CAROLINA
CARMEN.LOMBARDO@USMC.MIL
TELEPHONE (252) 466-5870**

**NAVAL FACILITIES ENGINEERING COMMAND ATLANTIC
ENVIRONMENTAL/BIOLOGICAL SCIENCES BRANCH
MS. DEANNA REES
DEANNA.REES@NAVY.MIL
TELEPHONE (757) 322-4940**

February 2009

This page is intentionally blank.

TABLE OF CONTENTS

	<u>PAGE</u>
LIST OF TABLES	ii
LIST OF FIGURES.....	ii
LIST OF ACRONYMS, ABBREVIATIONS, AND SYMBOLS.....	iii
1. DESCRIPTION OF ACTIVITIES	1-1
1.1 Introduction.....	1-1
1.2 Munitions Firing.....	1-3
1.2.1 Surface-to-Surface.....	1-7
1.2.2 Air-to-Surface.....	1-7
1.2.3 Quantity and Accuracy of Ordnance Used	1-8
1.3 Small Boat Maneuvers	1-9
2. LOCATION OF THE ACTIVITIES.....	2-1
2.1 Bombing Target-9	2-1
2.2 Bombing Target-11	2-2
3. AFFECTED SPECIES, STATUS, AND DISTRIBUTION.....	3-1
3.1 Bombing Target-11 Marine Mammal Species of the MCAS Bombing Targets	3-1
3.1.1 Common Bottlenose Dolphin (<i>Tursiops truncatus</i>).....	3-1
3.1.2 West Indian Manatee (<i>Trichechus manatus</i>).....	3-5
4. MARINE MAMMAL DENSITY ESTIMATES	4-1
4.1 Exposure Modeling Considerations	4-1
4.2 Impacts from Explosive Munitions Firing: Underwater Noise	4-2
4.2.1 Regulatory Framework.....	4-2
4.2.2 Explosive Sources	4-3
4.2.3 Criteria and Impact Thresholds for Underwater Noise on Marine Mammals	4-5
4.3 Underwater Noise Impacts from Explosive Ordnance.....	4-8
4.4 Estimating Impacts of Inert Ordnance.....	4-9
4.5 Impacts from Small Boat Maneuvers.....	4-10
4.6 Indirect Impacts.....	4-11
5. IMPACTS ON SUBSISTENCE USE.....	5-1
6. IMPACTS TO MARINE MAMMAL HABITAT AND RESTORATION	6-1
7. MEANS OF EFFECTING THE LEAST PRACTICABLE ADVERSE IMPACTS.....	7-1
7.1 Additional measures considered.....	7-2
8. MONITORING AND REPORTING MEASURES.....	8-1
9. RESEARCH.....	9-1
10. IMPACT TO CETACEAN SPECIES OR STOCKS	10-1
11. LIST OF PREPARERS.....	11-1
12. LITERATURE CITED.....	3

LIST OF TABLES

Table 1-1. MCAS Cherry Point Water Training Range Descriptions 1-3
Table 1-2. Munitions Authorized for Use at the MCAS Cherry Point Range Complex..... 1-5
Table 1-3. Proposed Level of Live and Inert Munitions Expended at BT-9..... 1-8
Table 1-4. Proposed Level of Live and Inert Munitions Expended at BT-11..... 1-8
Table 1-5. Annual Estimates for Munitions that May Land in Water Ranges**Error! Bookmark not defined.**
Table 1-6. Current Boat Operation at BT-9 and BT-11 1-10
Table 4-1. Source Weights and Peak Source Levels 4-4
Table 4-2. Effects, Criteria, and Thresholds for Impacts to Marine Mammals from Small Explosives 4-8
Table 4-3. Potential Bottlenose Dolphin Exposures to Underwater Noise 4-9
Table 4-4. Estimated number of potential direct strikes on cetaceans from inert ordnance at the BTs
expected annually..... 4-10
Table 10-1. Density of Marine Mammals in the Cherry Point Sites Area..... 10-1

LIST OF FIGURES

Figure 1-1. MCAS Cherry Point bombing targets BT-9, Brant Island Bombing Target, and BT-11, Piney
Island Bombing Range/Complex (33 CFR 334.420; USMC 2001b). 1-2
Figure 2-1. BTs at Rattan Bay, Piney Island Bombing Range/Complex (BT-11) (USMC 2001b; USMC
2004). 2-3
Figure 10-1. The Read et al. (2003) Bottlenose Dolphin Study Area Used to Calculate Density. 10-1

LIST OF ACRONYMS, ABBREVIATIONS, AND SYMBOLS

BT	Bombing Target
°C	Degree(s) Celsius
cal	Caliber
CETAP	Cetacean and Turtle Assessment Program
CFR	Code of Federal Regulations
Cm	Centimeters
CURRS	Consolidated Utilization Range Report System
dB	Decibel
dB re: 1 µPa	Decibels with reference pressure of one microPascal
dB re 1 µPa²-m	Decibels with reference to microPascals squared per meter
dB re 1 µPa²-s	Decibels with reference to microPascals squared per second
DoN	Department of the Navy
EFD	Energy flux density
ESA	Endangered Species Act
°F	Degree(s) Fahrenheit
FEIS	Final Environmental Impact Statement
FLMI	Florida Marine Research Institute
ft	Foot
FY	Fiscal year
HE	High explosive
hz	Hertz
in	Inches
in-lb/in²	Inch-pound per square inch
kg	Kilogram (s)
kHz	kiloHertz
km	Kilometer (s)
km²	Square kilometer(s)
lb(s)	Pound (s)
LGTR	Laser Guided Training Round
m	Meter(s)
m²	Square meter(s)
MCAS	Marine Corps Air Station
MMPA	Marine Mammal Protection Act
mi	Mile (s)
mi²	Square miles
mm	Millimeter (s)
msec	Millisecond
MU	Management Unit
N/A	Not applicable
NC	North Carolina
NCWM	North Carolina Winter Mixed
NEW	Net explosive weight
nm	Nautical mile(s)
NM	Northern Migratory
NMFS	National Marine Fisheries Service
NNC	Northern North Carolina
NOAA	National Oceanic and Atmospheric Administration

List of Acronyms, Abbreviations and Symbols

OTO	One-third-octave
psi	Pounds per square inch
psu	Practical salinity unit
PTS	Permanent threshold shift
SAMs	Surface-to-air missiles
SM	Statute-mile
SNC	Southern North Carolina
sq ft	Square feet
sq km	Square kilometers
sq mi	Square miles
TL	Transmission loss
TM	Tympanic membrane
TNT	Trinitrotoluene
TTS	Temporary threshold shift
U.S.	United States
USC	U.S. Code
USFWS	U.S. Fish and Wildlife Service
USMC	U.S. Marine Corps
ZOI	Zone of Influence

1. DESCRIPTION OF ACTIVITIES

1.1 INTRODUCTION


The United States Marine Corps (USMC) has prepared this Marine Mammal Compliance Report in accordance with the applicable regulations and the Marine Mammal Protection Act (MMPA), as amended by the National Defense Authorization Act for Fiscal Year 2004 (Public Law 108-136). This report analyzes the potential effects to marine mammals associated with bombing and target training within two in-water bombing targets (BTs) located in Pamlico Sound, North Carolina (NC), also known as the Brant Island Target (BT-9) and the Rattan Bay BTs of the Piney Island Bombing Range/Complex (BT-11) (Figure 1-1). These bombing targets are under the control and management of Marine Corps Air Station (MCAS), Cherry Point.

The MMPA of 1972, as amended (16 United States Code [USC] Section [§] 1371[a][5][A] and [D]), authorizes the issuance of regulations, Letters of Authorization (LOAs), and Incidental Harassment Authorizations (IHAs) for the incidental taking of marine mammals by a specified activity. The issuance of an LOA or IHA occurs when the Secretary of Commerce, after notice has been published in the Federal Register and opportunity for comment has been provided, finds that such takes will have a negligible impact on the species and stocks of marine mammals and will not have an unmitigable adverse impact on their availability for subsistence uses. The National Marine Fisheries Service (NMFS) has promulgated implementing regulations under 50 Code of Federal Regulations (CFR) § 216.101–106 that provide a mechanism for allowing the incidental, but not intentional, taking of marine mammals while engaged in a specified activity.

The United States Marine Corps (USMC) proposes to support and conduct operations at the MCAS Cherry Point Range Complex (Figure 1-1). Emerging training requirements reflect an increase in the amount of training operations conducted. Some of the mission increases and current training activities are not relevant to this compliance document as they have no potential to affect marine mammals. Thus, the focus of this document includes only those training missions occurring on the water ranges or with impact areas over the water because of their potential to affect marine mammals. These missions include:

- **Munitions Firing.** Units conduct air-to-ground, surface-to-surface, and air-to-surface munitions delivery at targets that are located on land or in water. Air-to-ground firing does not impact the water and is not analyzed or discussed further in this document.
- **Small Boat Maneuvers.** Units operate Small Unit River Craft, Combat Rubber Raiding Craft, Rigid Hull Inflatable Boats, Patrol Craft, and many other versions of these types of boats. These boats use inboard or outboard engines with either propeller or water jet propulsion.

Water range areas addressed within this report include the BTs located in Pamlico Sound, NC, known as the Brant Island Target (BT-9) and the Rattan Bay BTs of the Piney Island Bombing Range/Complex (BT-11) [Figure 1-1]. These BTs are used to train military personnel to deliver ordnance on a target. Ordnance is primarily delivered from aircraft but is also occasionally delivered from small military watercraft. Table 1-1 provides a general description of the training activities and munitions used on BT-9 and BT-11.



1

Figure 1-1. MCAS Cherry Point bombing targets BT-9, Brant Island Bombing Target, and BT-11, Piney Island Bombing Range/Complex (33 CFR 334.420; USMC 2001b).

Table 1-1. MCAS Cherry Point Water Training Range Descriptions

Range Asset	Training Operation	Type of Munitions Used
BT-9 (Brant Island Target)	Water-based target range for air-to-surface and surface-to-surface warfare training, including bombing, strafing, special (laser systems) weapons, and surface fires, using non-explosive and explosive ordnance; also provides a mining exercise area	Small Arms, Large Arms (live and inert), Bombs (live and inert), and Pyrotechnics
BT-11 (Piney Island Bombing Range)	Complex of land- and water-based targets designed to provide training in the delivery of conventional (non-explosive) and special (laser systems) weapons; secondary use for surface-to-surface training by small military watercraft	Small Arms, Large Arms (inert), Bombs (inert), and Pyrotechnics

1
2 The following subsections provide detailed information about the three MCAS Cherry Point
3 mission categories analyzed within this compliance report. Chapter 2 provides specific
4 information about the geographic location of this training.

5 **1.2 MUNITIONS FIRING**

6 Munitions firing training conducted on the water ranges include surface-to-surface firing (from
7 ship or boat to surface targets) and air-to-surface firing (from aircraft to surface water targets).
8 This activity occurs year round with no seasonal restrictions. There are many types of ordnance
9 used at BT-9 and BT-11 including practice bombs, rockets, flares, chaff, gun ammunition, and
10 grenades (Table 1-2 and Appendix A). Explosive ordnance is authorized only at BT-9; the
11 standard operating procedure sets the maximum limit of 100 pounds (lbs) trinitrotoluene (TNT)
12 equivalent for explosive ordnance at BT-9. Net explosive weights (NEW) used at BT-9 range
13 from 0.1019 to 15 lbs.

14 Practice bombs are non-explosive and are used at both targets. Lighter practice bombs (less than
15 500 lbs) contain a small amount of an explosive marking charge in a signal cartridge that allows
16 the target hit to be detected by range scorekeepers and the pilot. Explosives in the signal
17 cartridge are less than 1 lb TNT equivalent (USMC 2001a). Table 1-2 lists the types of
18 munitions expended at the MCAS Cherry Point Range Complex. It should be noted that all
19 munitions fired on the BT-11 range are non-explosive with the exception of the small explosives
20 in the single charges.

This page is intentionally blank.

Table 1-2. Munitions Authorized for Use at the MCAS Cherry Point Range Complex.

Small Arms	Large Arms	Missiles ¹	Rockets	Bombs	Pyrotechnics	
.22 cal-Live	20 mm-Inert	Hellfire	2.75-inch Rocket-Inert	G911 Grenade-Live	Chaff	
5.56 mm-Live	25 mm-Inert	Tube-launched, optically tracked, wire-guided (TOW)	2.75-inch Rocket Illumination-Inert	Hand Grenade-Inert	LUU-2	
7.62 mm-Live	30 mm-Inert		2.75-inch Rocket White Phosphorous-Inert	Non-Lethal Stun Grenade-Inert	MI27A1-Parachute Flare	
9 mm-Live	30 mm-Live		2.75-inch Rocket-Live	BDU-48 10 lb-Inert	Self Protection Flare	
.40 cal-Live	40 mm-Inert		5-inch Rocket-Inert	BDU-33 25 lb-Inert	Signal Illuminations-Inert	
.45 cal-Live	40 mm-Live		5-inch Rocket White Phosphorous-Inert	MK-48-Inert	Simulated Booby Traps-Inert	
.50 cal-Live	40 mm Illumination-Inert		5-inch Rocket-Live	MK-76 25 lb-Inert	Smokey Sams	
12 Gauge-Live	105 mm Target Practice-Inert 105mm 40 lb-Live			LGTR 90 lb-Inert		
				BDU-45 500 lb-Inert		
				BDU-50 500 lb-Inert		
				GBU-12 500 lb-Inert		
			MK-82 500 lb-Inert			
			BDU-38 750 lb-Inert			
			GBU-16 1,000 lb-Inert			
				MK-83 1,000 lb-Inert		

Note: 1. Two types of missiles, Hellfire and TOW, were previously approved for use at BT-9 per Air Station Order P3570.2R; however, use of these missiles at MCAS Cherry Point has been cancelled since Fiscal Year (FY) 2005 due to operational limitations imposed by an insufficient weapon safety footprint at the water range.

This page is intentionally blank.

1.2.1 Surface-to-Surface

Gunnery exercises are the only category of surface-to-surface activities currently conducted within the MCAS Cherry Point BTs.

- **Gunnery Exercise:** A small boat, typically operated by Special Boat Team personnel, uses a machine gun to attack and disable or destroy a surface target that simulates another ship, boat, swimmer, floating mine or near shore land targets. Boats conducting surface-to-surface firing activities will typically use 7.62 millimeter (mm) or .50 caliber (cal) machine guns; 40 mm Grenade machine guns or G911 Concussion hand grenades. This exercise is usually a live-fire exercise, but at times blanks may be used so that the boat crews can practice their ship handling skills. The most common exercise target used is BT-9. A target is not used for the G911 Concussion grenade, as the goal is to learn how to throw it into the water.

1.2.2 Air-to-Surface

There are four types of air-to-surface activities conducted within the MCAS Cherry Point BTs. These training activities are grouped into either mine laying; bombing; gunnery or rocket exercises.

- **Mine Laying:** These activities involve the use of a fixed wing aircraft deploying inert mine shapes in an offensive or defensive pattern. Mine laying operations are regularly conducted in the waters in the vicinity of BT-9.
- **Bombing Exercise:** During this exercise fixed wing aircraft deliver bombs against surface maritime targets, day or night, with the goal of destroying or disabling enemy ships or boats. Air-to-surface bombing exercises employ either unguided or precision-guided munitions. Unguided munitions include MK-76 and BDU-45 inert training bombs, and MK-80 series of inert bombs (no cluster munitions authorized). Precision-guided munitions consist of laser-guided bombs (inert) and laser-guided training rounds (inert).
- **Gunnery Exercise:** Rotary wing gunnery and fixed wing exercises are also conducted. Rotary wing exercises involve either CH-53, UH-1, CH-46, MV-22, or H-60 rotary-wing aircraft with mounted 7.62 mm or .50 cal machine guns. Each gunner expends approximately 800 rounds of 7.62 mm and 200 rounds of .50 cal ammunition in each exercise.

Fixed wing gunnery exercises involve the flight of two aircraft that begin to descend to the target from an altitude of approximately 914 meters (m) (3,000 feet [ft]) while still several miles away. Within a distance of 1,219 m (4,000 ft) from the target, each aircraft fires a burst of approximately 30 rounds before reaching an altitude of 305 m (1,000 ft), then breaks off and repositions for another strafing run until each aircraft expends its exercise ordnance allowance of approximately 250 rounds. Typically these fixed wing exercise events involve an F/A-18 and AH-1 with Vulcan M61A1/A2, 20 mm cannon; AV-8 with GAU-12, 25 mm cannon.

- **Rocket Exercise:** Fixed- and rotary-wing aircraft crews launch rockets at surface maritime targets, day and night, with the goal of destroying or disabling enemy ships or boats. These operations employ 2.75-inch and 5-inch rockets.

1.2.3 Quantity and Accuracy of Ordnance Used

The proposed amounts of ordnance to be expended at BT-9 and BT-11 under the proposed action are 897,932 and 1,109,955, respectively (Table 1-3 and Table 1-4). The amounts of ordnance expended at the BTs account for all use of the targets, including Navy use. There are five types of explosive sources used at BT-9: 2.75" Rocket High Explosives, 5" Rocket High Explosives, 30 mm High Explosives, 40 mm High Explosives, and G911 grenades. No high explosive munitions are used at BT-11.

Table 1-3. Proposed Level of Live and Inert Munitions Expended at BT-9

Proposed Munitions	Proposed Total No. of Rounds	Proposed Number of Explosive Rounds Having an Impact on the Water	Net Explosive Weight (lb)
Small Arms Rounds Excluding .50 cal	525,610	N/A	N/A
.50 Cal	257,067	N/A	N/A
Large Arms Rounds – Live	12,592	30mm HE: 3,120 40mm HE: 9,472	0.1019 0.1199
Large Arms Rounds – Inert	93,024	N/A	N/A
Rockets – Live	241	2.75" Rocket: 184 5" Rocket: 57	4.8 15.0
Rockets – Inert	703	N/A	N/A
Bombs and Grenades – Live	144	G911 Grenade: 144	0.5
Bombs and Grenades – Inert	4,055	N/A	N/A
Pyrotechnics	4,496	N/A	N/A
Total	897,932	12,977	N/A

Note: Increased munitions estimated using FY 2007 Consolidated Utilization Range Report System (CURRS) data on a per sortie-operation basis. HE = High Explosive. All live rounds are not necessarily high explosive rounds. N/A = Not applicable.

Table 1-4. Proposed Level of Live and Inert Munitions Expended at BT-11

Proposed Munitions	Proposed Total No. of Rounds ¹
Small Arms Rounds Excluding .50 Cal	507,812
.50 Cal	326,234
Large Arms Rounds	240,334
Rockets	4,549
Bombs and Grenades	22,114
Pyrotechnics	8,912
Total	1,109,955

Note: 1. Increased munitions estimated using FY 2007 CURRS data on a per sortie-operation basis.

The main target areas used by MCAS Cherry Point are located in or relatively near to bodies of water resulting in the potential for rounds of munitions impacting or detonating in the water. BT-

1 9 comprises two hulls anchored in the water on the Brant Island Shoals. Because it is located and
2 surrounded completely by water, DoN assumes that 100 percent of the munitions rounds
3 expended at BT-9 would impact the water. The DoN also assumes that 100 percent of the
4 munitions rounds expended at BT-9 would detonate in the water at a depth of approximately 1.2
5 m (3.9 ft).

6 BT-11 has three targets that are located either within or immediately adjacent to Rattan Bay. In
7 addition, several other targets at BT-11 are located and used such that the associated Safety
8 Danger Zones (SDZs) extend into the water. Thirty-six percent of the composite weapons SDZs
9 fall over water. Based on the location of the SDZs, the DoN conservatively estimates that 36
10 percent of the total number of rounds expended at BT-11 would impact in the water.

11 **1.3 SMALL BOAT MANEUVERS**

12 A number of different types of boats are used depending on the unit using the boat and their
13 mission. Boats are most often used by Naval Special Warfare teams, Navy Expeditionary
14 Combat Command units (Naval Coastal Warfare, Inshore Boat Units, Mobile Security
15 Detachments, Explosive Ordnance Disposal, and Riverine Forces), and US Coast Guard units.
16 These units have missions to protect ships in harbors and high value units, such as aircraft
17 carriers, nuclear submarines, liquid natural gas tankers, etc., while entering and leaving ports, as
18 well as to conduct riverine operations, insertions and extractions, and various Naval Special
19 Warfare operations. The boats used by these units include: Small Unit River Craft, Combat
20 Rubber Raiding Craft, Rigid Hull Inflatable Boats, Patrol Craft, and many other versions of these
21 types of boats. These boats use inboard or outboard, diesel or gasoline engines with either
22 propeller or water jet propulsion.

23 Table 1-6 shows current surface-to-surface operations at BT-9 and BT-11. In FY 2007, a total of
24 216 boat sorties were conducted at BT-9 and BT-11. These sorties occurred in all seasons, with
25 approximately the same number of sorties occurring in each season. The majority of boat sorties
26 at BT-9 originated from MCB Camp Lejeune. Boats are transported on trailers from MCB Camp
27 Lejeune to the Pamlico Sound for weapons training at BT-9, where live fire of 7.62 mm, .50 cal,
28 and 40 mm grenades is allowed, as well as use of G911 concussion grenades.

Table 1-5. Current Boat Operations at BT-9 and BT-11

Range	Sorties ¹	Munitions Type	Munitions Expended ²
BT-9	165	5.56 mm	1,468
		7.62 mm	218,500
		.50 cal	166,900
		40 mm Grenade - Inert	15,734
		40 mm Grenade – Live (HE)	9,472
		G911 Grenade	144
BT-11	51	7.62 mm	44,100
		.40 cal	4,600
		40 mm Grenade - Inert	1,517
		40 mm Illumination-Inert	9

1

Notes: 1. Sorties are from FY 2007 CURRS data.

2. Munitions expenditures for all munitions types except for the G911 grenade are from FY 2007 CURRS data. G911 grenades are not recorded as being used in FY 2007, so FY 2006 CURRS data were determined to be representative for this munitions type.

2. LOCATION OF THE ACTIVITIES

The BTs are located at the convergence of the Neuse River with Pamlico Sound (Figure 1-1). Pamlico Sound and its tributaries represent one of the largest intracoastal, estuarine ecosystems found in the United States (U.S.). Pamlico Sound is a shallow estuary with a mean water depth of 4.5 m (14.7 ft) and a low tidal range (approximately 10 centimeters [cm] [3.9 inches]) (Eisma et al. 1997; Paerl et al. 2001). The circulation in the sound is wind-driven and freshwater input, predominantly from the Neuse River, is low, resulting in long residence times for sound waters. Fine sand and silt cover the bottom of Pamlico Sound (Wells 1989; Eisma et al. 1997). Salinities in Pamlico Sound are highly variable, ranging from 10 to 31 practical salinity units (psu), and fluctuate seasonally with highest salinities found in the fall (15 to 31 psu) and lowest salinities found in spring (10 to 19 psu) (Wells 1989; National Ocean Service 2001). Surface water temperatures for Pamlico Sound vary seasonally with the average lowest temperatures (15 to 17 degrees Celsius [°C]) [59 to 62.6 degrees Fahrenheit [°F]) found in winter and the highest (26° to 27°C [78.8 to 80.6 °F]) occurring in summer (DoN 2003).

2.1 BOMBING TARGET-9

BT-9, also known as Brant Island Target (Figure 1-1), is a water-based target area located approximately 52 kilometers (km) (28 nautical miles [nm]) northeast of MCAS Cherry Point in Pamlico Sound, Pamlico County. It consists of a ship hull grounded on Brant Island Shoals. Brant Island Shoals is located approximately 4.8 km (3 miles [mi]) southeast of Goose Creek Island. Inert (non-explosive) ordnance (practice bombs) up to 454 kilograms (kg) (1,000 lbs) and live (explosive) ordnance up to 45.4 kg (100 lbs) TNT equivalent, including ordnance released during strafing, are authorized for use at this target range.

The target is defined by a 6 statute-mile (SM) diameter prohibited area designated by the U.S. Army Corps of Engineers, Wilmington District (33 CFR 334.420). Surface vessels are not permitted within the prohibited area, which is delineated by large signs located on pilings surrounding the perimeter of the BT.

Brant Island Shoal is oriented roughly northwest to southeast through the center of the circular target area. Hydrographic survey data collected for the vicinity of BT-9 indicate that water depths within the BT range from approximately 1.2 m along the crest of the shoal to 6 m at the perimeter of the prohibited area; bottom substrate exhibits a gradient ranging from harder-packed sediments (sand; possibly hard bottom) along the shoal crest in the shallowest water (1.2 to 3 m [4-10 ft]) to soft sediments (mud or silts) in the deeper regions off the flank of the shoal (depths greater than 4.9 m [16 ft]) (USMC 2001a). Surveys of the BT also indicate that areas of submerged debris appear to be clustered along the eastern half of the prohibited or target area (USMC 2001a).

Water temperatures at BT-9 average 8°C (46.4 °F) for winter (December through March), 18 °C (64.4 °F) for spring and fall (May and October, respectively), and 27°C (80.6 °F) for summer (June through September), while salinities remained a fairly constant 16.8 psu throughout the year (USMC 2001a).

2.2 BOMBING TARGET-11

BT-11 is a 50.6 square kilometers (sq km) (19.5 square miles [sq mi]) complex of land- and water-based targets on Piney Island. It includes both land (all of Piney Island) and surrounding water areas in the Pamlico Sound. The in-water stationary targets of BT-11 consist of a barge and PT boat located in roughly the center of Rattan Bay (Figure 2-1). All practice and live-fire exercises at MCAS Cherry Point ranges (BT-9 and BT-11) are conducted so that all ammunition and other ordnance strike and/or fall within the existing danger zones (water) (water prohibited areas) or water restricted areas for each of the bombing target ranges. A danger zone (water) is a defined water area that is closed to the public on an intermittent or full-time basis for use by military forces for hazardous operations such as target practice and ordnance firing. A water restricted area is a defined water area where public access is prohibited or limited in order to provide security for Government property and/or to protect the public from the risks of injury or damage that could occur from the Government's use of that area (33 CFR 334.2).

Surface danger zones are designated areas of rocket firing, target practice, or other hazardous operations (33 CFR 334.420). A surface danger zone is designated with a 2.9 km (1.8 sm) radius centered on a target in Rattan Bay. BT-11 also has three water restricted areas within 0.8 km (0.5 sm) radius areas located west of Point of Marsh and at Newstump Point and Jacks Bay. The prohibited area centered on the Rattan Bay target is closed to navigation at all times unless vessels are authorized by MCAS Cherry Point to enter the target area. The in-water target area (prohibited area) in Rattan Bay includes approximately 9.3 sq km (3.6 sq mi) of water surface. A currently proposed additional intermittent water restricted area would be designated around the already established danger zone (water) at BT-11 to better accommodate training in .50 cal weapons delivery fired from helicopters and boats.

Water depths in the center of Rattan Bay are estimated as 2.4 to 3 m (8 to 10 ft) with bottom depths ranging from 0.3 to 1.5 m (1 to 5 ft) adjacent to the shoreline of Piney Island (USMC 2001a). A shallow ledge, with substrate expected to be hard-packed to hard bottom, surrounds Piney Island.




Figure 2-1. BTs at Rattan Bay, Piney Island Bombing Range/Complex (BT-11) (USMC 2001b; USMC 2004).

This page is intentionally blank.

3. AFFECTED SPECIES, STATUS, AND DISTRIBUTION

While there are records showing the occurrence of 40 marine mammal species in the nearshore and offshore waters of North Carolina, the vast majority of these species are oceanic in distribution. Only one marine mammal species, the common bottlenose dolphin (*Tursiops truncatus*), has been repeatedly sighted in Pamlico Sound, while an additional species, the endangered West Indian manatee (*Trichechus manatus*), has been sighted rarely (Lefebvre et al. 2001; DoN 2003). A review of the records contained in the North Atlantic Right Whale Database, maintained by the National Marine Fisheries Service (NMFS), shows that no sightings of the endangered North Atlantic right whale (*Eubalaena glacialis*) have ever been recorded within Pamlico Sound or in vicinity of the BTs (Kenney 2006). No suitable habitat exists for this species in the shallow Pamlico Sound or BT vicinity. Therefore, no further information regarding this species is included in this document.

Marine mammal distribution is affected by demographic, evolutionary, ecological, habitat-related, and anthropogenic factors (Bjørge 2002; Stevick et al. 2002). Marine mammal movements are often related to feeding or breeding activity (Stevick et al. 2002). Cetacean movements are often a reflection of the distribution and abundance of prey and changes in cetacean distributions have been correlated with shifts in the distribution and abundance of prey (Kenney et al. 1996). Marine mammal movements have also been linked to environmental parameters indirectly indicative of prey distribution such as temperature variations (oceanic fronts), sea-surface chlorophyll a concentrations, and bottom depth (Fiedler 2002).

In fulfillment of the MMPA, the NMFS has identified certain cetacean stocks as strategic. All marine mammal species listed under the Endangered Species Act (ESA) are strategic stocks. Marine mammal stocks are listed as strategic if non-natural mortalities or serious injuries (e.g., fishery takes) have either exceeded the predicted maximum the stock can withstand or insufficient information exists to make such a determination. Marine mammal stocks not listed under the ESA are listed as strategic under the same guidelines. When applicable to a stock, such information has been included with the species' description that follows.

3.1 BOMBING TARGET-11 MARINE MAMMAL SPECIES OF THE MCAS BOMBING TARGETS

Two marine mammal species, the common bottlenose dolphin and the West Indian manatee, occur with an expected or rare occurrence, respectively, in the vicinity of the BTs; all other species potentially occurring in the southwestern area of Pamlico Sound are considered extralimital.

3.1.1 Common Bottlenose Dolphin (*Tursiops truncatus*)

Bottlenose dolphins are large and relatively robust, varying in color from light gray to charcoal. The genus *Tursiops* is named for its short, stocky snout that is distinct from the melon (Jefferson et al., 1993). The dorsal fin is tall, curved, and pointed. Striking regional variations exist in body size, with adult body lengths ranging from 1.9 to 3.8 m (6.2 to 12.5 ft) (Jefferson et al. 1993). *Tursiops* use a wide variety of feeding strategies and feed opportunistically on an assortment of fish, cephalopod, and shrimp species (Shane 1990; Wells and Scott 1999).

1 Scientists recognize two separate stocks of bottlenose dolphins in the western North Atlantic
2 Ocean: nearshore (coastal) and offshore forms. The stocks can be distinguished by genetics, diet,
3 blood characteristics, and outward appearance (Duffield et al., 1983; Hersh and Duffield, 1990;
4 Mead and Potter, 1995; Curry and Smith, 1997). Within the nearshore stock, NMFS further
5 segregates the stock into seven substocks or management units (MU) with discrete spatial and
6 temporal components (Waring et al., 2008). Discrete MUs exhibit seasonal migrations regulated
7 by temperature and prey availability (Torres et al., 2005; Waring et al., 2007), traveling as far
8 north as New Jersey in summer and as far south as central Florida in winter (Waring et al.,
9 2007). Two of the nearshore management units occur in the waters associated with Pamlico
10 Sound, the Northern Migratory (NM) management unit and the Northern North Carolina (NNC)
11 management unit. During the summer, the NM MU occurs from the New York/New Jersey
12 border to the Virginia/North Carolina border. The NNC MU ranges from the Virginia/North
13 Carolina border to Cape Lookout, North Carolina, during the summer months. In the winter
14 months, these two MUs and a third, the Southern North Carolina (SNC) MU overlap along the
15 coast of North Carolina and southern Virginia. A recent analysis suggests that a separate stock of
16 bottlenose dolphins may exist in Pamlico Sound, NC. One suggestion is that there is a resident
17 group of bottlenose dolphins in Pamlico Sound, which move into nearby nearshore waters in the
18 winter; while an alternative suggestion is that this group is a subset of the migrating dolphins that
19 overwinter in North Carolina waters (NMFS 2005a). Additional research is necessary to further
20 delineate this North Carolina stock.

21 NMFS provides abundance estimates for each MU by season. During the summer, the best
22 estimate of abundance for the NM and NNC MUs are 17,466 and 7,079 individuals respectively
23 (Waring et al., 2008). During the winter, an estimated 16,913 individuals comprise the North
24 Carolina Winter Mixed (NCWM) management unit, which is comprised of individuals from the
25 NNC MU, the SNC MU, and the migratory MUs that overwinter in North Carolina coastal
26 waters. The MUs making up the coastal stock are considered depleted under the MMPA (Waring
27 et al., 2007).

28 Dive durations for Atlantic bottlenose dolphins lasting as long as 15 minutes are recorded for
29 trained individuals (Ridgway et al., 1969). Typical dives, however, are more shallow and shorter
30 in duration. Mean dive durations of Atlantic bottlenose dolphins typically range from 20 to 40
31 seconds at shallow depths (Mate et al., 1995) and can last longer than 5 minutes during deep
32 offshore dives (Klatsky et al., 2005). Offshore bottlenose dolphins regularly dive to 450 m (1,476
33 ft) and possibly dive as deep as 700 m (2,297 ft) (Klatsky et al., 2005). Dive behavior of
34 bottlenose dolphins may correlate with the 24-hour cycle of day and night, or diel cycle (Mate et
35 al., 1995; Klatsky et al., 2005); this may be especially true for offshore stocks, which dive deeper
36 and more frequently at night to feed on the deep scattering layer (Klatsky et al., 2005).

37 Sounds emitted by bottlenose dolphins have been classified into two broad categories: pulsed
38 sounds (including clicks and burst-pulses) and narrow-band continuous sounds (whistles), which
39 usually are frequency modulated. Clicks have a dominant frequency range of 110 to 130
40 kiloHertz (kHz) and a source level of 218 to 228 decibels referenced to 1 microPascal at 1 meter
41 (dB re 1 μ Pa-m) peak-to-peak (Au, 1993) and 3.4 to 14.5 kHz and 125 to 173 dB re 1 μ Pam
42 peak-to-peak, respectively (Ketten, 1998). Whistles are primarily associated with communication
43 and can serve to identify specific individuals (i.e., signature whistles) (Caldwell and Caldwell,
44 1965; Janik et al., 2006). Up to 52 percent of whistles produced by bottlenose dolphin groups
45 with mother-calf pairs can be classified as signature whistles (Cook et al., 2004). Sound

1 production is also influenced by group type (single or multiple individuals), habitat, and behavior
2 (Nowacek, 2005). Bray calls (low-frequency vocalizations; majority of energy below 4 kHz), for
3 example, are used when capturing fish, specifically sea trout (*Salmo trutta*) and Atlantic salmon
4 (*Salmo salar*), in some regions (i.e., Moray Firth, Scotland) (Janik, 2000). Additionally, whistle
5 production has been observed to increase while feeding (Acevedo-Gutiérrez and Stienessen,
6 2004; Cook et al., 2004). Furthermore, both whistles and clicks have been demonstrated to vary
7 geographically in terms of overall vocal activity, group size, and specific context (e.g., feeding,
8 milling, traveling, and socializing) (Jones and Sayigh, 2002; Zaretsky et al., 2005; Baron, 2006).
9 For example, preliminary research indicates that characteristics of whistles from populations in
10 the northern Gulf of Mexico significantly differ (i.e., in frequency and duration) from those in
11 the western North Atlantic (Zaretsky et al., 2005; Baron, 2006).

12 Bottlenose dolphins can typically hear within a broad frequency range of 0.04 to 160 kHz (Au,
13 1993; Turl, 1993). Electrophysiological experiments suggest that the bottlenose dolphin brain
14 has a dual analysis system: one specialized for ultrasonic clicks and another for lower-frequency
15 sounds, such as whistles (Ridgway, 2000). Scientists have reported a range of highest sensitivity
16 between 25 and 70 kHz, with peaks in sensitivity at 25 and 50 kHz (Nachtigall et al., 2000).
17 Recent research on the same individuals indicates that auditory thresholds obtained by
18 electrophysiological methods correlate well with those obtained in behavior studies, except at
19 some lower (10 kHz) and higher (80 and 100 kHz) frequencies (Finneran and Houser, 2006).

20 Temporary threshold shifts (TTS) in hearing have been experimentally induced in captive
21 bottlenose dolphins using a variety of noises (i.e., broad-band, pulses) (Ridgway et al., 1997;
22 Schlundt et al., 2000; Nachtigall et al., 2003; Finneran et al., 2005; Mooney et al., 2005;
23 Mooney, 2006). For example, TTS has been induced with exposure to a 3 kHz, one-second pulse
24 with sound exposure level of 195 decibels referred to 1 micro-Pascal squared per second (dB re 1
25 $\mu\text{Pa}^2\text{-s}$) (Finneran et al., 2005), one-second pulses from 3 to 20 kHz at 192 to 201 decibels
26 referenced to 1 microPascal per meter (dB re 1 $\mu\text{Pa-m}$) (Schlundt et al., 2000), and octave band
27 noise (4 to 11 kHz) for 50 minutes at 179 dB re 1 $\mu\text{Pa-m}$ (Nachtigall et al., 2003). Preliminary
28 research indicates that TTS and recovery after noise exposure are frequency dependent and that
29 an inverse relationship exists between exposure time and sound pressure level associated with
30 exposure (Mooney et al., 2005; Mooney, 2006). Observed changes in behavior were induced
31 with an exposure to a 75 kHz one-second pulse at 178 dB re 1 $\mu\text{Pa-m}$ (Ridgway et al., 1997;
32 Schlundt et al., 2000). Finneran et al. (2005) concluded that a sound exposure level of 195 dB re
33 1 $\mu\text{Pa}^2\text{-s}$ is a reasonable threshold for the onset of TTS in bottlenose dolphins exposed to mid-
34 frequency tones.

35 The overall range of the bottlenose dolphin is worldwide in tropical and temperate waters. This
36 species occurs in all three major oceans and many seas. Dolphins of the genus *Tursiops* generally
37 do not range poleward of 45° latitude, except around the United Kingdom and northern Europe
38 (Jefferson et al., 1993). Climate changes can contribute to range extensions as witnessed in
39 association with the 1982/83 El Niño event when the range of some bottlenose dolphins known
40 to the San Diego, California area was extended 600 km (324 nm) northward to Monterey Bay
41 (Wells et al., 1990). Bottlenose dolphins continue to occur in Monterey Bay to this day. In the
42 western North Atlantic, bottlenose dolphins occur as far north as Nova Scotia but are most
43 common in coastal waters from New England to Florida, the Gulf of Mexico, the Caribbean, and
44 southward to Venezuela and Brazil (Würsig et al., 2000). Bottlenose dolphins occur seasonally in
45 estuaries and coastal embayments as far north as Delaware Bay (Kenney, 1990) and in waters

1 over the outer continental shelf and inner slope, as far north as Georges Bank [Cetacean and
2 Turtle Assessment Program (CETAP), 1982; Kenney, 1990].

3 Genetic analyses and spatial patterns observed from aerial surveys indicate regional and seasonal
4 distribution differences between the coastal and offshore stocks. North of Cape Hatteras, the
5 coastal stock is thought to be restricted to waters less than 25 m (82 ft) in depth, while offshore
6 dolphins generally range beyond the 50-m (164-ft) isobath (CETAP, 1982; Kenney, 1990;
7 Waring et al., 2007). Mitochondrial DNA and spatial analyses from dolphins south of Cape
8 Hatteras suggest individuals sighted within 7.5 km (4 nm) of shore are of the coastal form and
9 those beyond 34 km (18 nm) from shore and in waters with a bottom depth greater than 34 m
10 (112 ft) are of the offshore form (Torres et al., 2003). However, Torres et al. (2003) also found
11 an extensive region of overlap between the coastal and offshore stocks between 7.5 (4.0 nm) and
12 34 km (18 nm) from shore. Garrison et al. (2003b) also noted, that in this region, coastal
13 dolphins may be found in waters as deep as 31 m (102 ft) and 75 km (40 nm) from shore while
14 offshore dolphins may occur in waters as shallow as 13 m (43 ft). Additional aerial surveys and
15 genetic sampling are required to better understand the distribution of the two stocks throughout
16 the year.

17 Coastal bottlenose dolphins along the western Atlantic coast may exhibit either resident or
18 migratory patterns (Waring et al., 2008). Photo-identification studies support evidence of year-
19 round resident bottlenose dolphin populations in Beaufort and Wilmington, North Carolina
20 (Koster et al., 2000; Waring et al., 2008); these are the northernmost documented sites of year-
21 round residency for bottlenose dolphins in the western North Atlantic (Koster et al., 2000). A
22 high rate of exchange occurs between the Beaufort and Wilmington sites as well (Waring et al.,
23 2008). Individuals from the NM MU may enter these areas seasonally as well, as evidenced by a
24 bottlenose dolphin tagged in 2001 in Virginia Beach who overwintered in waters between Cape
25 Hatteras and Cape Lookout (NMFS- Southeast Fisheries Science Center [SEFSC], 2001).

26 Common bottlenose dolphins in North Carolina concentrate in shallow water habitats along
27 shorelines, and few, if any, individuals are present in the central portions of the sounds (Gannon
28 2003; Read et al. 2003a, 2003b). Bottlenose dolphins in the BT area select shallow habitats, such
29 as tributary creeks and the edges of the Neuse River, where the bottom depth is less than 3.5 m
30 (Gannon 2003). Dolphins use the downstream portion of the Neuse River estuary and lateral
31 creeks more than the upstream area (Gannon 2003). Dolphin density is highest during the spring
32 (May and June) and lowest during the summer (July and August) in the Neuse River estuary
33 (Gannon 2003). Fine-scale distribution of dolphins seems to relate to the presence of topography
34 or vertical structure, such as the steeply-sloping bottom near the shore and oyster reefs, which
35 may be used to facilitate prey capture (Gannon 2003).

36 Common bottlenose dolphins have been sighted during 11 boat-based surveys from 2002 through
37 2003 conducted for the USMC by Duke University in the waters surrounding BT-9 and BT-11;
38 during this time, one sighting in the restricted area surrounding BT-9 and two sightings in
39 proximity to BT-11 were observed, as well as seven sightings in adjacent waters for a total of
40 276 bottlenose dolphins sighted (Read et al. 2002, 2003b, 2003c). Some large groups of dolphins
41 have been reported in the restricted BT areas. In October 2002, a sighting of 50 dolphins was
42 made within BT-9 while two months later in December 2002, a group of 70 dolphins moved into
43 the prohibited area of the in-water targets of BT-11 (Read et al. 2003b). The NMFS-Southeast
44 Fisheries Science Center (SEFSC) in Beaufort, NC has aerially surveyed the area of

1 southwestern Pamlico Sound encompassed by the restricted airspace R-5306A, including both
2 in-water BTs. From July 2004 through June 2005, the NMFS-SEFSC conducted 23 aerial
3 surveys of the waters beneath the restricted airspace, during which time no dolphins were
4 observed at or near BT-9 or BT-11, although a total of 296 common bottlenose dolphins were
5 observed in the waters within the restricted airspace (Goodman et al. 2004, 2005a, 2005b,
6 2005c).

7 **3.1.2 West Indian Manatee (*Trichechus manatus*)**

8 The West Indian manatee is a rotund, slow-moving animal reaching a maximum length of 3.9 m
9 (12.8 ft) (Jefferson et al. 1993). The manatee has a small head, a squarish snout containing two
10 semi-circular nostrils at the front, and fleshy mobile lips. The tail is horizontal, rounded, and
11 paddle-shaped. The body is gray or gray-brown and covered with fine hairs that are sparsely
12 distributed. The back is often covered with distinctive scars resulting from collisions with boat
13 propellers (Moore 1956).

14 West Indian manatees are classified as endangered under the ESA and afforded additional
15 protection under the MMPA. West Indian manatee numbers are assessed by aerial surveys during
16 the winter months when manatees are concentrated in warm-water refuges. Aerial surveys
17 conducted in February 2006 by the Florida Marine Research Institute (FMRI) produced a
18 preliminary abundance estimate of 3,113 individuals (FMRI, 2006). Along Florida's Gulf Coast,
19 observers counted 1,474 West Indian manatees, while observers on the Atlantic coast counted
20 1,639. In the most recent revision of the West Indian manatee recovery plan, scientists concluded
21 that, based on movement patterns, manatees around Florida should be divided into four relatively
22 discrete management units or subpopulations. These subpopulations each represent a significant
23 portion of the species' range (U.S. Fish and Wildlife Service [USFWS], 2001). West Indian
24 manatees found along the Atlantic U.S. coast are of the Atlantic subpopulation (USFWS, 2001).
25 The other three subpopulations in Florida are the Upper St. Johns River, Northwest and
26 Southwest subpopulations (USFWS, 2001).

27 Manatees are shallow divers. The distribution of preferred seagrasses is mostly limited to areas
28 of high light; therefore, manatees are fairly restricted to shallower nearshore waters (Lefebvre et
29 al., 2001). It is unlikely that manatees descend much deeper than 20 m (66 ft), and manatees do
30 not usually remain submerged for longer than 2 to 3 minutes. However, when resting on the
31 bottom, manatees have been known to stay submerged for up to 24 minutes (Reynolds III, 1981).

32 West Indian manatees produce a variety of squeak-like sounds that have a typical frequency
33 range of 0.6 to 12 kHz (dominant frequency range from 2 to 5 kHz), and last 0.25 to 0.5 s (Steel
34 and Morris, 1982; Thomson and Richardson, 1995; Niezrecki et al., 2003). Recently,
35 vocalizations below 0.1 kHz have also been recorded (Frisch and Frisch, 2003; Frisch, 2006).
36 Overall, West Indian manatee vocalizations are considered relatively stereotypic, with little
37 variation between isolated populations examined (i.e., Florida and Belize) (Nowacek et al.,
38 2003). However, vocalizations have been shown to possess nonlinear dynamic characteristics
39 (e.g., subharmonics or abrupt, unpredictable transitions between frequencies). These
40 characteristics could aid in individual recognition and mother-calf communication (Mann et al.,
41 2006). Average source levels for vocalizations have been calculated to range from 90 to 138 dB
42 re: 1 μ Pa (average: 100 to 112 dB re: 1 μ Pa) (Nowacek et al., 2003; Phillips et al., 2004).
43 Behavioral data on two animals indicate an underwater hearing range of approximately 0.4 to 46

1 kHz, with best sensitivity between 16 and 18 kHz (Gerstein et al., 1999), while earlier
2 electrophysiological studies indicated best sensitivity from 1 to 1.5 kHz (Bullock et al., 1982).

3 West Indian manatees occur in warm, subtropical, and tropical waters of the western North
4 Atlantic Ocean, from the southeastern U.S. to Central America, northern South America, and the
5 West Indies (Lefebvre et al., 2001). West Indian manatees occur along both the Atlantic and Gulf
6 coasts of Florida. West Indian manatees are sometimes reported in the Florida Keys; these
7 sightings are typically in the upper Florida Keys, with some reports as far south as Key West
8 (Moore, 1951a, 1951b; Beck, 2006a). During winter months, the West Indian manatee
9 population confines itself to inshore and inner shelf waters of the southern half of peninsular
10 Florida and to springs and warm water outfalls (e.g., power plant cooling water outfalls). As
11 water temperatures rise in spring, West Indian manatees disperse from winter aggregation areas.
12 West Indian manatees are frequently reported in coastal rivers of Georgia and South Carolina
13 during warmer months (Lefebvre et al., 2001).

14 Historically, West Indian manatees were likely restricted to southernmost Florida during winter
15 and expanded their distribution northward during summer. However, industrial development has
16 made warm-water refuges available (e.g., power plant effluent plumes), and the introduction of
17 several exotic aquatic plant species has expanded the available food supply. These factors have
18 enabled an expansion of the manatee's winter range (USFWS, 2001; Laist and Reynolds, 2005).
19 Several patterns of seasonal movement are known along the Atlantic coast ranging from year-
20 round residence to long-distance migration (Deutsch et al., 2003). Deutsch et al., (2003) found
21 that manatees are highly consistent in seasonal movement patterns and show strong fidelity to
22 warm and winter ranges, both within and across years.

23 Although West Indian manatees are expected to inhabit nearshore areas, a few individuals have
24 been sighted offshore. West Indian manatees off the east coast of Florida have been documented
25 to occasionally make their way further offshore. For example, "Xoshi" was radio-tagged and
26 released near Biscayne Beach in March 1999. A few weeks later, she was "rescued" 60 km (32
27 nm) offshore of Port Canaveral, Florida in the Gulf Stream (Reid et al., 1991). Perhaps the most
28 famous long-distance movements of any West Indian manatee were exhibited by the animal
29 known as "Chessie," who gained fame in the summer of 1995 by swimming to Rhode Island,
30 returning to Florida for the winter, and traveling north again to Virginia where he was last seen
31 in 1996 (U.S. Geological Survey, 2001). In early September 2001, "Chessie" was once again
32 sighted in Virginia. More recently, in August 2006, a West Indian manatee was sighted in waters
33 off Rhode Island and Massachusetts, as well as in the Hudson River in New York City
34 (Anonymous, 2006; Beck, 2006b).

35 In general, manatee sightings along the Atlantic coast drop off markedly north of South Carolina
36 (Lefebvre et al. 2001). Manatees were first sighted in North Carolina waters in 1919 and have
37 been intermittently observed there since (Schwartz 1995). Most reported sightings in the inshore
38 waters of North Carolina occur during the summer months but sightings have been reported from
39 June through October (DoN 2003; North Carolina Natural Heritage Program 2004). In the
40 vicinity of the BTs, manatees have not been observed during the last 20 years in the waters
41 adjacent to Pamlico County but are known to occur in the waters of Carteret County (North
42 Carolina Natural Heritage Program 2004). For example, on 13 August 2002, the BT-11 range
43 operators sighted a manatee in the canal directly adjacent to the range complex.

This page is intentionally blank.

4. MARINE MAMMAL DENSITY ESTIMATES

All MCAS Cherry Point sites are within the shallow waters of Pamlico Sound (Figure 10-1). Within this area, the only marine mammal species that will occur regularly and for which reliable densities are available is the bottlenose dolphin (*Tursiops truncatus*). The West Indian manatee occurs so rarely in the vicinity of the BTs that no density estimate can be derived for this species in the waters of Pamlico Sound.

Aerial surveys specifically for dolphins and turtles were conducted in Pamlico and Core sounds from July 2004 to April 2006 (Goodman et al., 2007). Densities for bottlenose dolphins in the western part of Pamlico Sound (overlying the MCAS Cherry Point sites) ranged from 0.0272/km² (0.0105 mi²) in winter to 0.2158/km² (0.0833 mi²) in autumn (Goodman et al., 2007). The authors noted that animals were likely to be missed during the aerial surveys compared to boat-based surveys due to the short amount of time that observers have to make a sighting, resulting in underestimated densities. Goodman et al. (2007) were unable to incorporate correction factors for animals residing at the surface but not sighted in surveys, and animals not sighted because they were below the surface during surveys. A mark-recapture survey was conducted in the estuaries, bays and sounds of North Carolina by Read et al. (2003a) in summer 2000, which yielded a density of 0.183/ km² (0.071 mi²) (based on an estimate of 919 dolphins for the northern inshore waters divided by an estimated 5,015 km² (1,936 mi²) survey area, as measured in ArcGIS; Figure 5) for dolphins in North Carolina inland waters. Although this density is slightly lower than the highest density from Goodman et al. (2007), it is likely more precise and incorporates less bias than estimates derived from aerial surveys and is therefore a more conservative and applicable estimate. Surveys were done only in summer, but this population is considered resident and likely does not leave the estuary and sounds, therefore this density would apply to all MCAS Cherry Point sites year round.

Bottlenose dolphin densities for Pamlico Sound are listed in Table 10-1.

4.1 EXPOSURE MODELING CONSIDERATIONS

To increase the accuracy of modeling when estimating potential exposures of marine mammals from specific sources, a three-dimensional density estimate is utilized. Typically, density is reported for an area in the following form: animals/square kilometers (km²). Analyses of survey results generally include correction factors for animals at the surface as well as animals that may be below the surface and not seen. Therefore, although the area (e.g., km²) appears to represent only the surface of the water (two-dimensions), density actually implicitly includes animals anywhere within the water column under that surface area. By combining marine mammal densities with depth distribution information, a three-dimensional density estimate is achieved.

Generally, these density estimates assume that animals are uniformly distributed within the prescribed area, even though this may not be entirely accurate in nature. For instance, marine mammals are often clumped in areas of greater importance such as areas of high productivity, lower predation, safe calving, basking, etc. Density can occasionally be calculated for smaller areas that are used regularly by marine mammals, but more often than not, there is insufficient data to calculate density for small areas. Tagging and other technologies have also expanded our

1 understanding of marine mammal behavior and the physiological parameters which drive the
2 ways marine mammals utilize the water column. For instance, some species are capable of
3 regular deep dives (>800 meters [m] [2,625 ft]), while others dive to <200 m (656 ft) or
4 shallower, regardless of the bottom depth. There is limited depth distribution data for most
5 marine mammals, especially cetaceans. This is a result of the difficulties associated with tracking
6 cetaceans, which must be tagged at-sea using a tag that is either implanted into the skin/blubber
7 or adheres to the skin. While some suitable depth distribution data exist for some marine
8 mammal species, sample sizes are usually extremely small (nearly always fewer than 10 animals,
9 and most often only one or two). Often, depth distribution information is interpolated from
10 similar species for which more data exist or from other indirect methodologies, such as stomach
11 content analysis, habitat preference analysis, or preferred prey characteristics. However, due to
12 the complexities of animal behavior and the limited availability of fine-scale information, when
13 assuming an even distribution, modeling remains the standard for describing animal densities.
14 While assuming this may not be the most accurate depiction, and can present a distorted view of
15 marine mammal distribution in a region, it remains the best available science.

16 For the marine mammal species present in the MCAS Cherry Point sites (i.e., the bottlenose
17 dolphin) sufficient data exist to describe depth information more accurately than assuming an
18 even distribution. In the modeling, all bottlenose dolphins within the Cherry Point site areas
19 were distributed in waters with less than 10 m (33 ft) of depth.

20 **4.2 IMPACTS FROM EXPLOSIVE MUNITIONS FIRING: UNDERWATER NOISE**

21 **4.2.1 Regulatory Framework**

22 All marine mammals are protected under the MMPA. The MMPA prohibits, with certain
23 exceptions, the take of marine mammals in U.S. waters and by U.S. citizens on the high seas, and
24 the importation of marine mammals and marine mammal products into the United States. The
25 Endangered Species Act of 1973 (ESA) provides for the conservation of species that are
26 endangered or threatened throughout all or a significant portion of their range, and the
27 conservation of their ecosystems. A “species” is considered endangered if it is in danger of
28 extinction throughout all or a significant portion of its range. A species is considered threatened
29 if it is likely to become an endangered species within the foreseeable future. There are marine
30 mammals, already protected under MMPA, listed as either endangered or threatened under ESA,
31 and afforded special protections.

32 Actions involving sound in the water include the potential to harass marine animals in the
33 surrounding waters. Demonstration of compliance with MMPA and the ESA, using best
34 available science, has been assessed using criteria and thresholds accepted or negotiated, and
35 described here.

36 Sections of the MMPA (16 U.S.C. 1361 et seq.) direct the Secretary of Commerce to allow, upon
37 request, the incidental, but not intentional, taking of small numbers of marine mammals by U.S.
38 citizens who engage in a specified activity, other than commercial fishing, within a specified
39 geographical region. Through a specific process, if certain findings are made and regulations are
40 issued or, if the taking is limited to harassment, notice of a proposed authorization is provided to
41 the public for review.

1 Authorization for incidental takings may be granted if NMFS finds that the taking will have no
2 more than a negligible impact on the species or stock(s), will not have an immitigable adverse
3 impact on the availability of the species or stock(s) for subsistence uses, and that the permissible
4 methods of taking, and requirements pertaining to the mitigation, monitoring and reporting of
5 such taking are set forth. NMFS has defined negligible impact in 50 CFR 216.103 as an impact
6 resulting from the specified activity that cannot be reasonably expected to, and is not reasonably
7 likely to adversely affect the species or stock through effects on annual rates of recruitment or
8 survival.

9 Subsection 101(a)(5)(D) of the MMPA established an expedited process by which citizens of the
10 United States can apply for an authorization to incidentally take small numbers of marine
11 mammals by harassment. The National Defense Authorization Act of 2004 (Public Law 108-
12 136) removed the small numbers limitation and amended the definition of “harassment” as it
13 applies to a military readiness activity to read as follows:

- 14 *(i) any act that injures or has the significant potential to injure a marine mammal or*
15 *marine mammal stock in the wild [Level A Harassment]; or*
16 *(ii) any act that disturbs or is likely to disturb a marine mammal or marine mammal stock*
17 *in the wild by causing disruption of natural behavioral patterns, including, but not*
18 *limited to, migration, surfacing, nursing, breeding, feeding, or sheltering, to a point*
19 *where such behavioral patterns are abandoned or significantly altered [Level B*
20 *Harassment].*

21 The primary potential impact to marine mammals from underwater acoustics is Level B
22 harassment from noise.

23 **4.2.2 Explosive Sources**

24 The acoustic sources employed in the MCAS Cherry Point Exercise Areas are categorized as
25 broadband (producing sound over a wide frequency band) explosives. Broadband explosives
26 produce significant acoustic energy across several frequency decades of bandwidth. Propagation
27 loss is sufficiently sensitive to frequency as to require model estimates at several frequencies
28 over such a wide band.

29 Explosives are impulsive sources that produce a shock wave that dictates additional pressure-
30 related metrics (peak pressure and positive impulse). Detailed descriptions of the sources in the
31 MCAS Cherry Point Exercise Areas are provided in this subsection.

32 Explosives detonated underwater introduce loud, impulsive, broadband sounds into the marine
33 environment. Three source parameters influence the effect of an explosive: the weight of the
34 explosive material, the type of explosive material, and the detonation depth. The net explosive
35 weight (or NEW) accounts for the first two parameters. The NEW of an explosive is the weight
36 of TNT required to produce an equivalent explosive power.

37 The detonation depth of an explosive is particularly important due to a propagation effect known
38 as surface-image interference. For sources located near the sea surface, a distinct interference
39 pattern arises from the coherent sum of the two paths that differ only by a single reflection from
40 the pressure-release surface. As the source depth and/or the source frequency decreases, these
41 two paths increasingly, destructively interfere with each other, reaching total cancellation at the

1 surface (barring surface-reflection scattering loss). For MCAS Cherry Point there are five types
 2 of explosive sources: 2.75” Rocket High Explosives, 5” Rocket High Explosives, 30 mm High
 3 Explosives, 40 mm High Explosives, and G911 grenades.

4 The harassments expected to result from these sources are computed on a per in-water explosive
 5 basis; to estimate the number of harassments for multiple explosives, consider the following. Let
 6 A represent the impact area (that is, the area in which the chosen metric exceeds the threshold)
 7 for a single explosive. The cumulative effect of a series of explosives is then dictated by the
 8 spacing of the explosives relative to the movement of the marine wildlife. If the detonations are
 9 spaced widely in time or space, allowing for sufficient animal movements as to ensure a different
 10 population of animals is considered for each detonation, then the cumulative impact area of N
 11 explosives is merely NA regardless of the metric. This leads to a worst case estimate of
 12 harassments and is the method used in this analysis.

13 At the other extreme is the case where the detonations occur at essentially the same time and
 14 location (but not close enough to require the source emissions to be coherently summed). In this
 15 case, the pressure metrics (peak pressure and positive impulse) are constant regardless of the
 16 number of pings, while the energy metrics increase at a rate of $N^{1/2}$ (under spherical spreading
 17 loss only) or less.

18 The firing sequence for some of the munitions consists of a number of rapid bursts, often lasting
 19 a second or less. Due to the tight spacing in time, each burst can be treated as a single
 20 detonation. For the energy metrics the impact area of a burst is computed using a source energy
 21 spectrum that is the source spectrum for a single detonation scaled by the number of rounds in a
 22 burst. For the pressure metrics, the impact area for a burst is the same as the impact area of a
 23 single round. For all metrics, the cumulative impact area of an event consisting of N bursts is
 24 merely the product of the impact area of a single burst and the number of bursts, as would be the
 25 case if the bursts are sufficiently spaced in time or location as to insure that each burst is
 26 affecting a different set of marine wildlife.

27 All explosives are modeled as detonating at a 1.2 m (3.9 ft) depth. The NEW for these sources
 28 are provided in Table 4-1. Included in this table are the peak one-third-octave (OTO) source
 29 level and the approximate frequency at which the peak occurs.

Table 4-1. Source Weights and Peak Source Levels

Source Type	NEW	Peak OTO SL	Frequency of Peak OTO SL	Rounds per Burst
2.75” Rocket	4.8 lbs	223.9 dB re: 1µPa	~ 1500 Hertz (Hz)	1
5” Rocket	15.0 lbs	228.9 dB re: 1µPa	~ 1000 Hz	1
30 mm	0.1019 lbs	212.1 dB re: 1µPa	~ 2500 Hz	30
40 mm	0.1199 lbs	227.8 dB re: 1µPa	~ 1100 Hz	5
G911 Grenade	0.5	213.9 dB re: 1 µPa	~ 2500 Hz	1

30

31

1 For sources that are detonated at shallow depths, it is frequently the case that the explosion may
2 breach the surface with some of the acoustic energy escaping the water column. The source
3 levels presented in the table above have not been adjusted for possible venting nor does the
4 subsequent analysis attempt to take this into account. Because of the low source net explosive
5 weights and sufficient depths (1.2 m [3.9 ft]) venting of noise energy through the surface is
6 expected to be minor.

7 **4.2.3 Criteria and Impact Thresholds for Underwater Noise on Marine Mammals**

8 **Criteria and Thresholds for Small Explosives**

9 Criteria and thresholds for estimating the exposures from a single explosive activity on marine
10 mammals were established for the Seawolf Submarine Shock Test Final Environmental Impact
11 Statement (FEIS) (“Seawolf”) and subsequently used in the USS *Winston S. Churchill* (DDG-81)
12 Ship Shock FEIS (“Churchill”) (DoN, 1998 and 2001). Due to the spatial and temporal
13 characteristics, these detonations are treated as individual explosions with non-overlapping
14 sound fields for the purpose of this analysis. NMFS adopted these criteria and thresholds in its
15 final rule on unintentional taking of marine animals occurring incidental to the shock testing
16 (National Oceanic and Atmospheric Administration [NOAA], 1998). In addition, this section
17 reflects a revised acoustic criterion for small underwater explosions (i.e., 23 pounds per square
18 inch [psi] instead of previous acoustic criteria of 12 psi for peak pressure over all exposures),
19 which is based on an incidental harassment authorization (IHA) issued to the U.S. Air Force
20 (NOAA, 2006a). Criteria and thresholds are summarized in Table 4-2.

21 ***Criteria and Thresholds for Injurious Physiological Effects***

22 *Single Explosions*

23 The approach to risk assessment for impulsive sound in the water was derived from the
24 Seawolf/Churchill approach. For injury, MCAS Cherry Point uses dual criteria: eardrum rupture
25 (i.e. tympanic membrane [TM] rupture) and onset of slight lung injury. These criteria are
26 considered indicative of the onset of injury. The threshold for TM rupture corresponds to a 50
27 percent rate of rupture (i.e., 50 percent of animals exposed to the level are expected to suffer TM
28 rupture); this is stated in terms of an Energy Flux Density (EFD) Level value of 1.17 inch pounds
29 per square inch (in-lb/in²) (about 205 dB re 1 $\mu\text{Pa}^2\text{-s}$). This recognizes that TM rupture is not
30 necessarily a serious or life-threatening injury, but it is a useful index of possible injury that is
31 well correlated with measures of permanent hearing impairment [e.g., Ketten (1998) indicates a
32 30 percent incidence of permanent threshold shift (PTS) at the same threshold].

33 The threshold for onset of slight lung injury is calculated for a small animal (a dolphin calf
34 weighing 29.6 lbs) and is given in terms of the “Goertner modified positive impulse,” indexed to
35 13 psi-millisecond (msec) (DoN, 2001). This threshold is conservative since the positive impulse
36 needed to cause injury is proportional to animal mass, and therefore, larger animals require a
37 higher impulse to cause the onset of injury. This analysis assumed the marine species
38 populations were 100 percent small animals. The criterion with the largest potential impact range
39 (most conservative), either TM rupture (energy threshold) or onset of slight lung injury (peak
40 pressure threshold), will be used in the analysis to determine injurious physiological exposures.

1 For mortality, MCAS Cherry Point uses the criterion corresponding to the onset of extensive
2 lung injury. For small mammals, the threshold is given in terms of the Goertner modified
3 positive impulse, indexed to 30.5 pounds per square inch-millisecond (psi-ms). Since the
4 Goertner approach depends on propagation, source/animal depths, and animal mass in a complex
5 way, the actual impulse value corresponding to the 30.5 psi-msec index is a complicated
6 calculation. To be conservative, the analysis used the mass of a dolphin calf (at 12.2 kg [26.9
7 lbs]) for 100 percent of the populations. This analysis was also conservative in that it
8 corresponds to a 1 percent chance of mortal injury, and yet any animal experiencing onset severe
9 lung injury is counted as a lethal exposure.

10 *Multiple Explosions*

11 For multiple successive explosions, the acoustic criteria were developed from an extension of the
12 Churchill approach. For multiple explosions, accumulated energy over the entire training time is
13 the natural extension for energy thresholds since energy accumulates with each subsequent shot
14 (explosion); this is consistent with the treatment of multiple arrivals in Churchill. For positive
15 impulse, it is consistent with Churchill to use the maximum value over all impulses received.

16 **Criteria and Thresholds for Non-injurious Physiological Effects**

17 The MCAS Cherry Point criterion for non-injurious physiological effects is temporary threshold
18 shift (TTS) – a slight, recoverable loss of hearing sensitivity (DoN, 2001). For this assessment,
19 there are dual thresholds for TTS, an energy threshold and a peak pressure threshold. The
20 criterion with the largest potential impact range (most conservative) either the energy threshold,
21 or peak pressure threshold, will be used in the analysis to determine non-injurious physiological
22 (TTS) exposures.

23 *Single Explosion – TTS Energy Threshold*

24 The TTS energy threshold is a 182 dB re 1 $\mu\text{Pa}^2\text{-s}$ maximum energy flux density level in any 1/3-
25 octave band at frequencies above 100 Hz for toothed whales and in any 1/3-octave band above
26 10 Hz for baleen whales. For large explosives, the latter limits at 100 and 10 Hz make a
27 difference in the impact range estimates. NMFS has defined large explosives in prior rulemaking
28 as greater than 907 kg (2,000 lbs) Net Explosive Weight (NEW) (NMFS, 2006b). For small
29 explosives (<1500 lbs NEW), as what was modeled for this analysis, the spectrum of the shot
30 arrival is broad, and there is essentially no difference in exposure ranges resulting from the 10
31 and 100 Hz frequency range cutoffs for the two classes of whales.

32 The TTS energy threshold for explosives is derived from the Space and Naval Warfare Systems
33 Center pure-tone tests for TTS (Schlundt et al., 2000; Finneran and Schlundt, 2004). The pure-
34 tone threshold (192 decibels (dB) as the lowest value) is modified for explosive by (a)
35 interpreting it as an energy metric, (b) reducing it by 10 dB to account for the time constant of
36 the mammal ear, and (c) measuring the energy in 1/3 octave bands, the natural filter band of the
37 ear. The resulting threshold is 182 dB re 1 $\mu\text{Pa}^2\text{-s}$ in any 1/3 octave band. The energy threshold
38 usually dominates over peak pressure threshold and is used in the analysis to determine potential
39 non-injurious physiological (TTS) for single explosion ordnance.

40

1 *Single Explosion - TTS Peak Pressure Threshold*

2 The TTS peak pressure threshold applies to all cetacean species and is stated in terms of peak
3 pressure at 23 psi (NOAA, 2006b). This threshold was adopted for Precision Strike Weapons
4 (PSW) Testing and Training by Eglin Air Force Base in the Gulf of Mexico (NMFS, 2005). It is
5 important to note that for small shots near the surface (such as in this analysis), the 23-psi peak
6 pressure threshold will generally produce longer exposure ranges than the 182-dB energy metric.
7 Furthermore, it is not unusual for the TTS exposure range for the 23-psi pressure metric to
8 actually exceed the disturbance exposure range for the 177-dB energy metric.

9 *Multiple Explosions – TTS*

10 For multiple explosions, accumulated energy over the entire training time is the natural extension
11 for energy thresholds since energy accumulates with each subsequent shot/detonation. This is
12 consistent with the energy argument in Churchill. For peak pressure, it is consistent with
13 Churchill to use the maximum value over all impulses received.

14 **Criteria and Thresholds for Behavioral Effects**

15 *Single Explosion*

16 For a single explosion, to be consistent with Churchill, TTS is the criterion for non-injurious
17 physiological exposure. In other words, because behavioral disturbance for a single explosion is
18 likely to be limited to a short-lived startle reaction, use of the TTS criterion is considered
19 sufficient protection and, therefore, behavioral effects are not considered for single explosions.

20 *Multiple Explosions – without TTS*

21 For this analysis, because multiple explosions would occur within a discrete time period, a new
22 acoustic criterion – behavioral disturbance – is used to account for behavioral effects significant
23 enough to be judged as harassment, but occurring at lower noise levels than those that may cause
24 TTS.

25 The threshold is based on test results published in Schlundt et al. (2000), with derivation
26 following the approach of the Churchill EIS for the energy-based TTS threshold. The original
27 Schlundt et al. (2000) data and the report of Finneran and Schlundt (2004) are the basis for
28 thresholds for behavioral disturbance. As reported by Schlundt et al. (2000), instances of altered
29 behavior generally began at lower exposures than those causing TTS; however, there were many
30 instances when subjects exhibited no altered behavior at levels above the onset-TTS levels.
31 Regardless of reactions at higher or lower levels, all instances of altered behavior were included
32 in the statistical summary.

33 The behavioral disturbance threshold for tones is derived from the SSC tests, and is found to be
34 five dB below the threshold for TTS, or 177 dB re $1 \mu\text{Pa}^2\text{-s}$ maximum energy flux density level
35 in any 1/3 octave band at frequencies above 100 Hz for toothed whales, and in any 1/3 octave
36 band above 10 Hz for baleen whales. As stated previously for TTS, for small explosives (<
37 1m,500 lbs NEW), as what was modeled for this analysis, the spectrum of the shot arrival is
38 broad and there is essentially no difference in exposure ranges for toothed or baleen whales. In
39 shallower water, the behavioral disturbance exposure range can be about twice the exposure

1 range for TTS. However, in deeper water, the TTS exposure criteria (23 psi) exposure range can
 2 result in a longer exposure range than the behavioral disturbance criteria. This is due to the fact
 3 that in a deeper water environment, it is more likely that there is a direct path for the shockwave
 4 to propagate, which results in a larger peak pressure range. In shallow water, there is reflection,
 5 absorption, and cancellation of the shockwave propagation due to interactions with the seafloor,
 6 which can limit the peak pressure range.

7 **Summary of Thresholds and Criteria for Impulsive Sounds**

8 Table 4-2 summarizes the effects, criteria, and thresholds used in the assessment for impulsive
 9 sounds. Non-injurious effects are determined by either the dual physiological criteria for single
 10 detonations, or by the behavioral criteria for multiple detonations. The criterion for behavioral
 11 disturbance used in this analysis is based on the use of multiple explosives.

Table 4-2. Effects, Criteria, and Thresholds for Impacts to Marine Mammals from Small Explosives

Effect	Criteria	Metric	Threshold	MMPA Harassment Level
Onset Mortality	Onset extensive lung injury	Goertner modified positive impulse	30.5 psi-ms	Mortality
Injurious Physiological	50 percent TM rupture	Energy flux density	205 dB re $1\mu\text{ Pa}^2\text{-s}$ (1.17 in-lb/in ²)	Level A
Injurious Physiological	Onset slight lung injury	Goertner modified positive impulse	Indexed to 13 psi-ms	Level A
Non-injurious Physiological	TTS (for single explosions)	Greatest energy flux density level in any 1/3-octave band for total energy over all exposures	182dB re $1\mu\text{ Pa}^2\text{-s}$	Level B
Non-injurious Physiological	TTS (for multiple successive explosions)	Peak pressure over all exposures	23 psi	Level B
Non-injurious Behavioral	Behavioral Disturbance	Greatest energy flux density level in any 1/3 octave band for total energy over all exposures (multiple explosions only)	177 dB re $1\mu\text{ Pa}^2\text{-s}$	Level B

12 dB re $1\mu\text{ Pa}^2\text{-s}$ = decibel referenced to 1 micropascal squared second; Hz = hertz; psi-ms = pounds per square inch-
 13 millisecond; TM = tympanic membrane; TTS = temporary threshold shift

14 **4.3 UNDERWATER NOISE IMPACTS FROM EXPLOSIVE ORDNANCE**

15 Table 4-3 lists the range in meters of the explosive effects associated with a particular
 16 threshold. Table 4- presents the number of bottlenose dolphins potentially affected by
 17 underwater noise using the methodology described previously in this section. An estimate of the
 18 number of animals exposed to noise above established thresholds from a given type of detonation
 19 (exposure) was calculated by first multiplying the area of noise associated with a particular effect
 20 or harassment threshold by the annual density of bottlenose dolphins for Pamlico Sound. As
 21 previously discussed the area exposed to explosive noise is affected by such factors as amount of
 22 net explosive weight, depth of explosion and environmental province. The volume of the area
 23 exposed to the sound relative to each threshold level was multiplied by the dolphin density and

1
2**Table 4-3. Explosive Noise Maximum Effect Ranges for Impact Thresholds**

	Behavioral Disturbance (177 dB Energy)	TTS (23 psi)	Level A (13 psi-msec)	Mortality (31 psi-ms)
2.75" Rocket HE	N/A	172 m (564 ft)	47 m (154 ft)	27 m (89 ft)
5" Rocket HE	N/A	255 m (837 ft)	61 m (200 ft)	39 m (128 ft)
30mm HE	209 m (686 ft)	N/A	10 m (33 ft)	5 m (16 ft)
40mm HE	144 m (472 ft)	N/A	10 m (33 ft)	5 m (16 ft)
G911 Grenade	N/A	83 m (272 ft)	21 m (33 ft)	10 m (33 ft)

3
4**Table 4-4. Potential Bottlenose Dolphin Exposures to Underwater Noise**

	Behavioral Disturbance	TTS	Level A	Mortality
2.75" Rocket HE	N/A	4.97	0.17	0.06
5" Rocket HE	N/A	3.39	0.09	0.03
30mm HE	2.55	N/A	0.05	0.00
40mm HE	12.60	N/A	0.16	0.01
G911 Grenade	N/A	0.87	0.03	0.01
Total	15.15	9.23	0.5	0.11

5
6
N/A = The criteria are not applicable.7 **4.4 ESTIMATING IMPACTS OF INERT ORDNANCE**

8 The potential risk to marine mammals from non-explosive ordnance without any conservation
9 measures entails two possible impacts, from noise or from the ordnance physically hitting an
10 animal. Estimates of the noise fields generated in water by the impact of non-explosive ordnance
11 indicate that the energy radiated is about 1 to 2 percent of the total kinetic energy of the impact.
12 This energy level (and likely peak pressure levels) is well below the impact threshold, even at 1-
13 m from the impact. Therefore, the noise generated by the in-water impact of non-explosive
14 ordnance will not pose a risk to marine life; the only potential risk to marine mammals from non-
15 explosive ordnance is from the very small possibility that the ordnance will hit (strike) an animal.

The potential risk of a direct hit to an animal in the target area is estimated to be so low it is discountable. A Range Air Installation Compatible Use Zone (RAICUZ) study generated the surface area or footprints of weapon impact areas associated with air-to-ground ordnance delivery (USMC 2001a). Statistically, a weapon safety footprint describes the area needed to contain 99.99 percent of initial and ricochet impacts at the 95 percent confidence interval for each type of aircraft and ordnance utilized on the BTs. At both BT-9 and BT-11 the probability of deployed ordnance landing in the impact footprint is essentially 1.0, since the footprints were designed to contain 99.99 percent of impacts, including ricochets. However, only 36 percent of the weapon footprint for BT-11 is over water in Rattan Bay, so the likelihood of a weapon striking an animal at the BT in Rattan Bay is 64 percent less. Water depths in Rattan Bay range from 3 m (10 ft) in the deepest part of the bay to 0.5 m (1.6 m) close to shore, so that nearly the entire habitat in Rattan Bay suitable for marine mammal use (or 36 percent of the weapon footprint).

The probability of hitting a bottlenose dolphin at the BTs can be derived by:

1 Probability = dolphin's dorsal surface area * density of dolphins

2 Example: Probability for BT-9 = $1.425 \text{ m}^2 * 0.183 \text{ km}^2 = 2.61 \times 10^{-7}$

3 The estimate of the dorsal surface area of a bottlenose dolphin is assumed to be 1.425 m^2 (or the
4 average length of 2.85 m times the average body width of 0.5 m). Thus, using Read et al.'s
5 (2003d) density estimates (0.183 km^2), the probability of a dolphin being hit in the waters of BT-
6 9 is 2.61×10^{-7} and of BT-11 is 9.4×10^{-8} (or 36 percent of the BT-11 probability of 2.61×10^{-7}).
7 Using the proposed levels of ordnance expenditures at each in-water BT (Tables 1-3 and 1-4)
8 and taking into account that only 36 percent of the ordnance deployed at BT-11 is over water, the
9 estimated potential number of ordnance strikes on a marine mammal per year is estimated as
10 0.263 at BT-9 and 0.034 at BT-11 (Table 4-3). It would take approximately three years of
11 ordnance deployment at the BTs before it would be likely or probable that one bottlenose
12 dolphin would be struck by deployed inert ordnance.

Table 4-3. Estimated number of potential direct strikes on cetaceans from inert ordnance at the BTs expected annually.

	Estimated Ordnance Levels using FY 2007 data at BT-9 and in-water BT-11	Probability of Striking Dolphin/Ordnance Deployment	Number Ordnance Strikes/Year
BT-9	1,007,931	2.61×10^{-7}	0.263
BT-11	359,985	9.4×10^{-8}	0.034

13

14 4.5 IMPACTS FROM SMALL BOAT MANEUVERS

15 The bottlenose dolphin is the most common marine mammal species within the CHPT study
16 area. Small boats operating at high speeds have the potential to strike a bottlenose dolphin,
17 though Wells and Scott (1997) note that in the busy boating waters of Sarasota Bay, Florida,
18 bottlenose dolphins suffer few injuries from boat traffic. Still, boat and dolphin collisions can
19 occur. Wells and Scott (1997) have documented a few injuries and note that mothers with
20 calves, and unhealthy dolphins may be particularly susceptible as they are limited in their
21 movements. Wells and Scott (1997) noted obvious propeller-caused injuries to dolphins in

1 Sarasota Bay during periods of high recreational boat usage and coinciding with high speed
2 powerboat races, though they did not correlate boat races nor a particular boat speed with
3 dolphin strikes.

4 Small military boats operating within the CHPT water ranges may travel at speeds up to 74
5 km/hr (40 knots). However, the bottlenose dolphin is least susceptible to serious impacts from
6 vessels because of its swimming speed and ability to maneuver around moving vessels.

7 In addition to direct strike, the presence of boats can result in disturbance, causing dolphins to
8 alter their swimming patterns and behavior. Impacts such as disturbance are difficult to assess
9 given the affinity this species has to bowride in front of moving vessels. Some researchers have
10 noted changes in behavior of bottlenose dolphins to vessels as far away as 100 meters (328 ft).
11 These alterations include changes in direction, changes in breathing rate and decreased spacing
12 between dolphins traveling in the same group (Lemon et al., 2006).

13 The West Indian manatee is susceptible to injury because it is slow moving, and when surfacing
14 to breathe, may be struck by boats and propellers. Because this species rarely occurs within the
15 study area, the likelihood of an encounter with a small boat is very low.

16 **4.6 INDIRECT IMPACTS**

17 Explosive ordnance may have an impact not only on marine mammals but also on other species
18 and the ecosystem of BT-9. Explosive and non-explosive ordnance delivery can also impact the
19 bottom sediments at both BTs as can the routine target replacement or equipment/system
20 maintenance. Sediment disturbance effects from ordnance delivery are difficult to predict but are
21 expected to be initially disturbed after ordnance delivery with quick settlement afterwards. Thus
22 the turbidity impacts on marine biota in the vicinity of the BTs are expected to be of short
23 duration and negligible.

24 The distributions of various fishes and invertebrates in the vicinity of BT-9 may be impacted by
25 explosive ordnance but the ecosystem impact is expected to be of a brief duration and result
26 primarily in fish evacuating the vicinity of the blast. Since little time is expected to occur before
27 fish or invertebrates move back into the detonation area, no adverse effect is expected on the
28 trophic regime or ecosystem at BT-9. The likelihood of any permanent ecosystem impact at BT-
29 11 due to inert ordnance use is also extremely unlikely.

30

This page is intentionally blank.

1 **5. IMPACTS ON SUBSISTENCE USE**

2 Subsistence use pertains to the use of those species sought in traditional Arctic hunting areas.
3 None of the marine mammal species discussed in this document is sought for subsistence use in
4 the U.S. There would be no impacts on the availability of the species or stocks for subsistence
5 use as identified in MMPA Section 101(a)(5)(A)(i).

This page is intentionally blank.

1

1 **6. IMPACTS TO MARINE MAMMAL HABITAT AND RESTORATION**

2 Potential impacts to marine mammal habitat at the BTs have previously been evaluated during a
3 Section 7 consultation with the NMFS (USMC 2001a) and were determined to be minimal
4 (NMFS 2002). Delivery of explosive or non-explosive ordnance may involve a temporary
5 disturbance to the substrate (sediments) and a subsequent increase in localized turbidity.
6 Turbidity may fluctuate based upon rainfall, wind, tides, and season. It is difficult to predict the
7 amount of disturbance that occurs immediately after ordnance hits the substrate. The amount of
8 sediment disturbance will be influenced by whether or not the ordnance hits the target first
9 before hitting the substrate as well as the orientation of the ordnance (i.e., whether it lands
10 upright or on its side) upon impact with the sediments (USMC 2001a). Sediments in the action
11 area are primarily soft sediments such as sand and mud, which should experience only minimal
12 disturbance and quick resettlement. Impacts to the substrate are considered temporary and
13 negligible with no adverse or long-term effects anticipated. Studies at the BTs of water quality
14 following ordnance deployment have shown that turbidity levels are well below the state's water
15 quality levels (USMC 2001a). Turbidity impacts are not likely to disturb the feeding or transiting
16 behavior of dolphins in the vicinity of the BTs since this species does not rely on vision for
17 maneuvering.

18 Physical effects associated with pressure waves generated by underwater detonations of
19 explosives might affect fishes within proximity of BT-9. In particular, the rapid oscillation
20 between high and low-pressure peaks has the potential to burst the swim bladders and other gas-
21 containing organs of fishes (Keevin and Hempen 1997). Sublethal effects, such as changes in
22 behavior of fishes, have been observed on several occasions as a result of noise produced by
23 explosives (Wright 1982; National Resource Council 2003). The abundances of various fishes
24 and invertebrates near the detonation point could be altered for a few hours before repopulation
25 occurs. As a result, a short-term and localized effect on prey availability may be expected in the
26 vicinity of BT-9.

27 Chemical effects of using explosive ordnance during training are considered negligible here as
28 they were during the shock trial of the Winston S. Churchill (DoN 2001). Initial concentrations
29 of the chemical by-products of ordnance detonations are not hazardous to marine life (O'Keeffe
30 and Young 1984; DoN 2001). Water quality parameters at BT-9 and BT-11 are all within the
31 limits set by North Carolina Water Quality Standards for Saltwater Classifications (for those
32 parameters with Standards). Elevated aluminum concentrations or visual detections of chaff in
33 either the water or sediment samples from any of the tested areas have not been detected with
34 previous exercises (USMC 2001a).

35 While debris associated with ordnance delivery occurs at the BTs (e.g., parachutes, strands of
36 chaff and ordnance remains), no reports of ingestion of debris has been reported nor have any
37 stranded animals been reported with such contents in their stomachs (USMC 2001a). It is most
38 likely that ordnance debris will sink to the bottom and become incorporated into the sediments
39 and is thus not likely to be ingested by marine mammals.

40 Target establishment and maintenance occurs infrequently (i.e., every five years), and MCAS
41 personnel are required to ensure that new targets are free of environmental contaminants prior to
42 placing them in the water for use. Small boat operations or target replacement could also
43 occasionally result in temporary and infrequent sediment disturbance, as explained above with
44 explosive impacts to sediments.

This page is intentionally blank.

1

Means of Effecting the Least Practicable Adverse Impacts

- 1 • Acoustic deterrent devices that produce sound (e.g., Dukane NetMark™ 1000
2 [fundamental frequency 10 to 12 kHz, sound pressure level 132 dB re: 1 µPa, pulse
3 duration of 300 msec]) have been demonstrated to be an effective means in repelling
4 delphinids (Barlow and Cameron 2003).
5
- 6 • Active acoustic detection involves using sound to detect an animal. Target strengths
7 (reflection energy) have been identified for some small cetaceans (i.e., spinner
8 dolphins [*Stenella longirostris*] and dusky dolphin [*Lagenorhynchus obscurus*]) using
9 common fish finders (i.e., Computrol Tournament Master Fishfinder NCC 5300)
10 (Benoit-Bird and Au 2003; Benoit-Bird et al. 2004). The advantage of active acoustic
11 detection is that an animal does not need to be producing sound in order to be
12 identified.

13 **7.1 ADDITIONAL MEASURES CONSIDERED**

14 Protective measures will include thorough monitoring of the impact area before, during, and after
15 the event, in addition to establishing a sequence of procedures to achieve the least practicable
16 adverse impacts on protected resources. Other mitigation alternatives considered during
17 development of the protective measures described in Section 8.0 but dismissed because of their
18 impact to military readiness, include:

- 19 ➤ Since the ranges operate continuously and the bottlenose dolphin population is presumed
20 to be stable throughout the year, time or seasonal restrictions on the use of explosive
21 ordnance are impracticable and would severely compromise training flexibility. The use
22 of explosive ordnance use is permitted at BT-9 at any time during the year, although the
23 majority of the explosive ordnance use occurs during large training exercises. Two to
24 three of these large exercises occur throughout the year (they do not occur more heavily
25 in any one season).
26
- 27 ➤ Visual monitoring of the impact area using a boat or vessel would not be possible due to
28 personnel safety.
29

1 **8. MONITORING AND REPORTING MEASURES**

2 As described in the previous section, the Marine Corps will monitor the bombing ranges to
3 reduce the potential impacts to marine mammals. Information associated with the presence of
4 any marine mammals observed during training, including the number, general location of
5 animals, and time of sighting will be recorded. The compiled sighting information and
6 description of any action taken will be maintained by MCAS Cherry Point.

7 Any observations of stranded or injured marine mammals within the BT-9 and BT-11 vicinity
8 will be immediately reported to the NMFS stranding network. MCAS Cherry Point personnel
9 will be responsible for ensuring that the MCAS local Chain of Command and the MCI-East are
10 notified of any occurrence.

This page is intentionally blank.

1

1 **9. RESEARCH**

2 In an effort to improve the availability of data on the abundance and seasonal distribution of
3 marine mammals in the BTs, the MCAS Cherry Point has provided funding support to augment
4 the surveys conducted by the NMFS Southeast Fisheries Science Centers. For example, to
5 provide additional information on the distribution and abundance of marine mammals and sea
6 turtles in the mid-Atlantic region, the NMFS Southeast Fisheries Science Center has been
7 contracted by the MCAS to conduct aerial surveys covering the R-5306A restricted airspace that
8 encompasses portions of Pamlico and Core Sounds (Goodman et al. 2004, 2005a, 2005b, 2005c).
9 Previous access to these areas has been restricted, but these studies are providing valuable
10 information on the use of this region by protected species and may provide seasonality
11 information that could be used to plan exercises to least impact protected species.

12 The three objectives of these studies are to:

- 13 1. Survey the waters of Core and Pamlico Sounds within the R-5306A airspace and
14 document seasonal distribution and number of sea turtles and marine mammals over a 16-
15 month period (Final report was expected in February 2006);
- 16 2. Provide an index of abundance for the protected species that can be compared among
17 seasons; and
- 18 3. Provide observer training to MCAS Cherry Point personnel sufficient to allow these
19 personnel to perform pre-and post-exercise surveys for protected species.

20 The MCAS Cherry Point has funded sighting surveys performed by Duke University from July
21 2002 to June 2003 targeting common bottlenose dolphins. These boat-based surveys additionally
22 photo-identify sighted dolphins and dolphins identities are compiled and cataloged by Duke
23 (Read et al. 2002, 2003b, 2003c, and 2003d). The MCAS has also contracted Duke University to
24 conduct acoustic monitoring studies of the use of BT-9 and BT-11 by common bottlenose
25 dolphins. Since August 2005, two pop-up hydrophone buoys have been deployed in BT-9 and
26 one at Rattan Bay of BT-11 (Urian 2005a, 2005b).


27 The MCAS Cherry Point has funded additional work with Duke University to develop and test a
28 real-time passive acoustic monitoring system that will allow automated detection of bottlenose
29 dolphin whistles. The work will be in two distinct phases. Phase I will develop an automated
30 signal detector (a software program) to recognize the whistles of dolphins in the BT-9 and BT-11
31 ranges. Phase II will assemble and deploy a prototype real-time monitoring unit on one of the
32 towers in the BT-9 range. The success of this effort will help direct future monitoring initiatives
33 and activities within the BTs.

This page is intentionally blank.

1

1 **10. IMPACT TO CETACEAN SPECIES OR STOCKS**

2 This section addresses potential impacts to marine mammal species or stocks from underwater
 3 noise and inert munitions (Figure 10-1 and Table 10-1). Based on the best available science, the
 4 USMC concludes that exposures to marine mammal species and stocks would result in only
 5 short-term effects and would not affect annual rates of recruitment or survival based on the
 6 analysis presented in the previous chapters.
 7



8 **Figure 10-1. The Read et al. (2003) Bottlenose Dolphin Study Area Used to Calculate Density.**

9 **Table 10-1. Density of Marine Mammals in the Cherry Point Sites Area**

Common Name	Species Name	Status	Density (animals/km ²)	Season	Depth (<10 m)
Bottlenose dolphin	<i>Tursiops truncatus</i>	Not listed	0.183 ¹	year round	100 percent of bottlenose dolphin time would be at <10 m

10 ¹ Read et al. (2003a)

This page is intentionally blank.

11. LIST OF PREPARERS

Name/Title	Qualifications	Project Role
Deanna Rees Naval Facilities Engineering Command, Atlantic Division Norfolk, Virginia	B.S. Wildlife Resources, University of Idaho	Navy Technical Representative and Navy Technical Review
Carmen Lombardo Natural Resources Manager Environmental Affairs Department MCAS Cherry Point Cherry Point, North Carolina	M.S. Wildlife and Fisheries Science University of Tennessee, Knoxville B.S. Environmental Studies Unity College, Maine	Point of Contact for MCAS Cherry Point/Technical Review
Richard Weaver Natural Resource Specialist Environmental Affairs Department MCAS Cherry Point Cherry Point, North Carolina	M.S. Marine Sciences, University of Delaware B.S. Chemistry, Villanova University	Secondary Point of Contact for MCAS Cherry Point/Technical Review
Russ Piovesan SAIC, Environmental Scientist	M.S. Environmental Management/ B.S. Biology - 13 years environmental science	Project Manager Client Liaison Author
W. James McKee SAIC, Environmental Scientist	B.S. Marine Biology - 23 years environmental science	Author
Mark Lockwood SAIC, Noise Analyst	M.S. Mathematics/B.S. Mathematics – 4 years acoustics	Underwater Noise Analysis
Janet Clarke SAIC, Environmental Scientist	M.A. Anthropology/B.S. Wildlife Management – 25 years environmental science	Author
Angie Toole SAIC, Technical Editor	B.S. Journalism - 25 years experience in editing and production	Editor
Bud Albee TEC Inc., Project Manager	M.S. Limnology- 16 years experience in management of environmental compliance studies	TEC Inc. – Project Manager
Lori Thursby TEC Inc., Deputy Project Manager	M.A.H. Master of Architectural History – 12 years experience in regulatory compliance	TEC Inc. - Deputy Project Manager/Technical Review
Jaclyn Johnson TEC Inc., Environmental Planner	B.S. Ecology – 9 years experience in NEPA analysis and regulatory compliance	TEC Inc. - Environmental Planner/Technical Review

This page is intentionally blank.

12. LITERATURE CITED

- Acevedo-Gutiérrez, A., and S. C. Stienessen, 2004. Bottlenose dolphins (*Tursiops truncatus*) increase number of whistles when feeding. *Aquatic Mammals*, Vol 3, No 3, pp 357–362.
- Anonymous, 2006. Accidental tourist: Manatee cruises Hudson River. Retrieved from <http://www.cnn.com/2006/US/08/07/manatee.hudson.river.ap/index.html> on 15 August 2006.
- Au, W. W. L., 1993. The sonar of dolphins. Springer-Verlag: New York.
- Barlow, J., and G. A. Cameron, 2003. Field experiments show that acoustic pingers reduce marine mammal bycatch in the California drift gill net fishery. *Marine Mammal Science* 19(2):265–283.
- Baron, S., 2006. Personal communication via email between Dr. Susan Baron, National Marine Fisheries Service, Southeast Fisheries Science Center, Miami, Florida, and Dr. Amy R. Scholik, Geo-Marine, Inc., Hampton, Virginia, on 31 August 2006.
- Beck, C., 2006a. Personal communication via email between Ms. Cathy Beck, U.S. Geological Survey, Sirenia Project, Gainesville, Florida, and Ms. Dagmar Fertl, Geo-Marine, Inc., Plano, Texas, on 6 September, and 27 October 2006.
- Beck, C., 2006b. Florida manatee travels to Cape Cod, Massachusetts. *SireNews*, Vol 46, pp 15–16.
- Benoit-Bird, K. J., and W. W. L. Au, 2003. Prey dynamics affect foraging by a pelagic predator (*Stenella longirostris*) over a range of spatial and temporal scales. *Behavioral Ecology and Sociobiology* 53:364–373.
- Benoit-Bird, K. J., and B. Würsig, and C. J. McFadden, 2004. Dusky dolphin (*Lagenorhynchus obscurus*) foraging in two different habitats: active acoustic detection of dolphins and their prey. *Marine Mammal Science* 20(2):215–231.
- Bjørge, A., 2002. How persistent are marine mammal habitats in an ocean of variability? Pages 63–91 in *Marine mammals: Biology and conservation*. P. G. H. Evans and J. A. Raga. New York, New York, Kluwer Academic/Plenum Publishers.
- Bullock, T. H., T. J. O’Shea, and M. C. McClune, 1982. Auditory evoked potentials in the West Indian manatee (*Sirenia Trichechus manatus*). *Journal of Comparative Physiology*, Vol 148, pp 547–554. Caldwell and Caldwell, 1965.
- Caldwell, D. K., and M.C. Caldwell, 1965. Individualized whistle contours in bottlenose dolphins (*Tursiops truncatus*), *Nature*, Vol 207, pp 434–435.
- Cetacean and Turtle Assessment Program (CETAP), 1982. Characterization of marine mammals and turtles in the mid- and North Atlantic areas of the U.S. outer continental shelf. Final report to the U.S. Bureau of Land Management, Washington, D.C., from the Graduate School of Oceanography, University of Rhode Island, Kingston. NTIS PB83-215855.
- Cook, M. L. H., L. S. Sayigh, J. E. Blum, and R. S. Wells, 2004. Signature-whistle production in undisturbed free-ranging bottlenose dolphins (*Tursiops truncatus*). *Proceedings of the Royal Society B: Biological Sciences*, Vol 271, pp 1043–1049. Curry and Smith, 1997.
- Curry, B. E., and J. Smith, 1997. Phylogeographic structure of the bottlenose dolphin (*Tursiops truncatus*): Stock identification and implications for management. *Molecular genetics of marine mammals*.

List of Preparers

- Society for Marine Mammalogy Special Publication 3, D.E. Dizon, S.J. Chivers, and W.F. Perrin, eds. Lawrence, Kansas. pp 227–247.
- Deutsch, C. J., J. P. Reid, R. K. Bonde, D. E. Easton, H. I. Kochman, and T. J. O’Shea, 2003. Seasonal movements, migratory behavior, and site fidelity of West Indian manatees along the Atlantic coast of the United States. *Wildlife Monographs*, Vol 151, pp 1–77.
- Department of the Navy (DoN), 1998. Final environmental impact statement, shock testing the SEAWOLF submarine. Washington, D.C.: Naval Sea Systems Command.
- , 2001. Final environmental impact statement, shock trial of the Winston S. Churchill (DDG 81). Washington, D.C.: Naval Sea Systems Command.
- , 2003. Marine Resource Assessment for the Cherry Point and Southern Virginia Capes (VACAPES) Inshore and Estuarine Areas. Final Report. Naval Facilities Engineering Command, Norfolk, VA. Cotntract #N62470-95-D-1160, CTO 0030. Prepared by Geo-Marine, Inc. Plano, Texas.
- Duffield, D. A., S. H. Ridgway, and L. H. Cornell, 1983. Hematology distinguishes coastal and offshore forms of dolphins (*Tursiops*). *Canadian Journal of Zoology*, Vol 61, pp 930–933.
- Eisma, D., P. L. de Boer, G. C. Cadée, K. Dijkema, H. Ridderinkhof, and C. Philippart, 1997. Intertidal deposits: River mouths, tidal flats, and coastal lagoons. Washington, D.C.: CRC Press.
- Fiedler, P. C., 2002. Ocean environment. Pages 824-830 in *Encyclopedia of marine mammals*, eds. W. F. Perrin, B. Würsig, and J. G. M. Thewissen. San Diego, California, Academic Press.
- Finneran, J. J., D. A. Carder, C. E. Schlundt and S. H. Ridgway, 2005. Temporary threshold shift in bottlenose dolphins (*Tursiops truncatus*) exposed to mid-frequency tones. *Journal of Acoustical Society of America*. 118:2696–2705.
- Finneran, J. J. and D. S. Houser, 2006. Comparison of in-air evoked potential and underwater behavioral hearing thresholds in four bottlenose dolphins (*Tursiops truncatus*). *Journal of the Acoustical Society of America*, Vol 119, No 5, pp 3181–3192.
- Finneran, J. J., and C. E. Schlundt, 2004. Effects of intense pure tones on the behavior of trained odontocetes. Space and Naval Warfare Systems Center, San Diego, Technical Document. September.
- Florida Marine Research Institute (FMRI), 2006. Manatee synoptic surveys. Retrieved from http://research.myfwc.com/features/print_article.asp?id=15246, on 20 October 2006
- Frisch, S., 2006. Personal communication via e-mail between Dr. Stefan Frisch, University of South Florida, Tampa, Florida and Dr. Amy Scholk, Geo-Marine, Inc., Hampton, Virginia, on 11 January 2006.
- Frisch, S., and K. Frisch, 2003. Low frequency vocalizations in the Florida manatee (*Trichechus manatus latirostris*), in Abstracts, Fifteenth Biennial Conference on the Biology of Marine Mammals. 14–19 December 2003. Greensboro, North Carolina. pp 55–56.
- Gannon, D. P., 2003. Behavioral ecology of an acoustically mediated predator-prey system: Bottlenose dolphins and sciaenid fishes. Ph.D. thesis, Duke University.
- Garrison, L. P., P. E. Rosel, A. Hohn, R. Baird, and W. Hoggard, 2003. Abundance of the coastal morphotype of bottlenose dolphin, *Tursiops truncatus*, in U.S. continental shelf waters between New Jersey and Florida during winter and summer 2002. Unpublished document prepared for the Take Reduction Team on Coastal Bottlenose Dolphins in the Western Atlantic.

List of Preparers

- Gerstein, E. R., L. Gerstein, S. E. Forsythe, and J. E. Blue, 1999. The underwater audiogram of the West Indian manatee (*Trichechus manatus*). *Journal of the Acoustical Society of America*, Vol 105, No 6, pp 3575–3583.
- Goertner, J. F., 1982. Prediction of underwater explosion safe ranges for sea mammals. Report NSWC/WOL TR 82-188. Silver Spring, Maryland: Naval Ordnance Laboratory.
- Goodman, M. A., J. B. McNeill, and A. A. Hohn, 2004. Protected species aerial survey data collection and analysis in waters underlying the R-5306A airspace, 1st quarterly report submitted to the U.S. Marine Corps, MCAS Cherry Point, 27 October 2004. Beaufort, North Carolina: NOAA Beaufort Lab, NMFS, Southeast Fisheries Science Center.
- , 2005a. Protected species aerial survey data collection and analysis in waters underlying the R-5306A airspace, 2nd quarterly report submitted to the U.S. Marine Corps, MCAS Cherry Point, 31 January 2005. Beaufort, North Carolina: NOAA Beaufort Lab, NMFS, Southeast Fisheries Science Center.
- , 2005b. Protected species aerial survey data collection and analysis in waters underlying the R-5306A airspace, 3rd quarterly report submitted to the U.S. Marine Corps, MCAS Cherry Point, 15 April 2005. Beaufort, North Carolina: NOAA Beaufort Lab, NMFS, Southeast Fisheries Science Center.
- , 2005c. Protected species aerial survey data collection and analysis in waters underlying the R-5306A airspace, 4th quarterly report submitted to the U.S. Marine Corps, MCAS Cherry Point, 28 July 2005. Beaufort, North Carolina: NOAA Beaufort Lab, NMFS, Southeast Fisheries Science Center.
- , 2007. NOAA Technical Memorandum NMFS-SEFSC-551 Protected Species Aerial Survey Data Collection and Analysis in Waters Underlying the R-5306A Airspace. Final report submitted to U.S. Marine Corps, MCAS Cherry Point. January 2007. Beaufort, North Carolina: NOAA Beaufort Lab, NMFS, Southeast Fisheries Science Center.
- Hersh, S. L., and D. A. Duffield, 1990. Distinction between northwest Atlantic offshore and coastal bottlenose dolphins based on hemoglobin profile and morphometry, in *The Bottlenose Dolphin*, S. Leatherwood and R.R. Reeves, eds. Academic Press: San Diego. pp 129–139.
- Janik, V. M., 2000. Food-related bray calls in wild bottlenose dolphins (*Tursiops truncatus*). *Proceedings of the Royal Society B: Biological Sciences*, Vol 267, pp 923–927.
- Janik, V. M., L. S. Sayigh, and R. S. Wells, 2006. Signature whistle shape conveys identity information to bottlenose dolphins. *Proceedings of the National Academy of Sciences of the United States of America*, Vol 103, No 2, pp 8293–8297.
- Jefferson, T.A., S. Leatherwood, and M.A. Webber, 1993. *FAO species identification guide. Marine mammals of the world*. Rome: Food and Agriculture Organization of the United Nations.
- Jones, G. J. and L. S. Sayigh, 2002. Geographic variation in rates of vocal production of free-ranging bottlenose dolphins. *Marine Mammal Science*, Vol 18, No 2, pp 374–393.
- Keevin, T. M., and G. L. Hempen, 1997. *The environmental effects of underwater explosions with methods to mitigate impacts*. St. Louis, Missouri: U.S. Army Corps of Engineers.
- Kenney, R. D., 1990. Bottlenose dolphins off the northeastern United States, in *Leatherwood, in The Bottlenose Dolphin*, S. and R.R. Reeves, eds. Academic Press: San Diego, California. pp 369–386.

List of Preparers

- Kenney, R., 2006. Personal communication via email between Dr. Robert Kenney, North Atlantic Right Whale Consortium/University of Rhode Island, Narragansett, Rhode Island, and Ms Cheryl Schroeder, Geo-Marine, Inc., Hampton, Virginia, 6 January 2006.
- Kenney, R. D., P. M. Payne, D. W. Heinemann, and H. E. Winn, 1996. Pages 169-196 in Shifts in northeast shelf cetacean distributions relative to trends in Gulf of Maine/Georges Bank finfish abundance. The northeast shelf ecosystem: Assessment, sustainability, and management. K. Sherman, N. A. Jaworski and T. J. Smayda. Boston, Massachusetts, Blackwell Science.
- Ketten, D. R., 1998. Marine mammal auditory systems: A summary of audiometric and anatomical data and its implications for underwater acoustic impacts. NOAA Technical Memorandum NOAA-NMFS-SWFSC-256:1-74.
- Klatsky, L., R. Wells, and J. Sweeney, 2005. Bermuda's deep diving dolphins - Movements and dive behavior of offshore bottlenose dolphins in the Northwest Atlantic Ocean near Bermuda. Page 152 in Abstracts, Sixteenth Biennial Conference on the Biology of Marine Mammals. 12–16 December 2005. San Diego, California.
- Koster, D., L. Sayigh, K. Urian, and A. Read, 2000. Evidence for year-round residency and extended home ranges by bottlenose dolphins in North Carolina. Page 3 in Eighth Annual Atlantic Coastal Dolphin Conference. 24–26 March 2000. Wilmington, North Carolina.
- Laist, D. W. and J.E. Reynolds III, 2005. Influence of power plants and other warm-water refuges on Florida manatees. Marine Mammal Science, Vol 21, No 4, pp 739–764.
- Lefebvre, L. W., M. Marmontel, J. P. Reid, G. B. Rathbun, and D. P. Domning, 2001. Distribution, status, and biogeography of the West Indian manatee. Pages 425-474 in C.A. Woods and F.E. Sergile, eds. Biogeography of the West Indies: Patterns and perspectives. 2d ed. Boca Raton, Florida: CRC Press.
- Lemon, M., T. Lynch, D.H. Cato, R.G. Harcourt. 2006. Response of travelling bottlenose dolphins (*Tursiops aduncus*) to experimental approaches by a powerboat in Jervis Bay, New South Wales, Australia. Biological Conservation 127: 363-372.
- Mann, D.A., T.J. O'Shea, and D.P. Nowacek, 2006. Nonlinear dynamics in manatee vocalization. Marine Mammal Science, Vol 22, No 3, pp 548–555.
- Mate, B. R., K. A. Rossbach, S. L. Nieukirk, R. S. Wells, A. B. Irvine, M. D. Scott, and A. J. Read, 1995. Satellite-monitored movements and dive behavior of a bottlenose dolphin (*Tursiops truncatus*) in Tampa Bay, Florida. Marine Mammal Science, Vol 11, No 4, pp 452–463.
- Mead, J. G. and C. W. Potter, 1995. Recognizing two populations of the bottlenose dolphin (*Tursiops truncatus*) off the Atlantic Coast of North America: Morphologic and ecologic considerations. IBI Reports, Vol 5, pp 1–44.
- Mooney, T. A., 2006. Personal communication via email between Dr. Aran Mooney, University of Hawaii, Marine Mammal Research Program, Kane'ohe, Hawaii, and Dr. Amy R. Scholik, Geo-Marine, Inc., Hampton, Virginia, on 29 August 2006.
- Mooney, T. A., P. E. Nachtigall, W. W. L. Au, M. Breese, and S. Vlachos, 2005. Bottlenose dolphin: Effects of noise duration, intensity, and frequency, in Abstracts, Sixteenth Biennial Conference on the Biology of Marine Mammals. 12–16 December 2005. San Diego, California. p 197.
- Moore, J. C., 1951a. The range of the Florida manatee. Quarterly Journal of the Florida Academy of Sciences, Vol 14, No 1, pp 1–19.

List of Preparers

- , 1951b. The status of the manatee in the Everglades National Park, with notes on its natural history. *Journal of Mammalogy*, Vol 32, No 1, pp 22–36.
- , 1956. Observations of manatees in aggregations. *American Museum Novitates* 1811:1–24.
- Nachtigall, P. E., D. W. Lemonds, and H. L. Roitblat, 2000. Psychoacoustic studies of dolphin and whale hearing, in *Hearing by Whales and Dolphins*, Au, W.W.L., A.N. Popper, and R.R. Fay, eds. Springer-Verlag: New York. pp 330–363.
- Nachtigall, P. E., J. L. Pawloski, and W. W. L. Au, 2003. Temporary threshold shift and recovery following noise exposure in the Atlantic bottlenose dolphin (*Tursiops truncatus*). *Journal of the Acoustical Society of America*. 113:3425–3429.
- National Marine Fisheries Service (NMFS), 2002. Endangered Species Act—Section Consultation Biological Opinion for Ongoing Ordnance Delivery at Bombing Target 9 (BT-9) and Bombing Target 11 (BT-11) at Marine Corps Air Station, Cherry Point, North Carolina (Consultation No. F/SER/2002/00130). St. Petersburg, Florida: National Marine Fisheries Service, Southeast Regional Office, Protected Resources Division.
- , 2005a. Bottlenose dolphin (*Tursiops truncatus*): Western north Atlantic coastal morphotype stocks; Draft 2005 stock assessment report. Silver Spring, MD: National Marine Fisheries Service, Office of Protected Species.
- , 2005b. Taking and importing marine mammals; taking marine mammals incidental to conducting the Precision Strike Weapon (PSW) testing and training by Eglin Air Force Base in the Gulf of Mexico--Notice of issuance of an incidental harassment authorization. *Federal Register* 70(160):48575–48691.
- , 2006a. *Environmental Impact Statement to Implement the Operational Measures of the North Atlantic Right Whale Ship Strike Reduction Strategy, Draft Environmental Impact Statement*. Retrieved from <http://www.nmfs.noaa.gov/pr/pdfs/shipstrike/deis.pdf>, on 27 September 2007.
- , 2006b. *Draft U.S. Atlantic marine mammal stock assessments - 2006*. Silver Spring, Maryland: National Marine Fisheries Service.
- National Marine Fisheries Service-Southeast Fisheries Science Center (NMFS-SEFSC), 2005. Geographic coordinates for sea turtles strandings in North Carolina from 1998-2005. Data received May 2006 from Ms. Wendy Teas. Miami, Florida: National Marine Fisheries Service, Southeast Fisheries Science Center.
- National Ocean Service, 2001. Environmental Sensitivity Index Atlases: North Carolina. Data on CD-ROM. Seattle: National Ocean Service, Office of Response and Restoration, Hazardous Materials Response Division.
- National Oceanic and Atmospheric Administration (NOAA), 1998. December 1, 1998. Final Rule for the incidental taking of marine mammals, Naval activities; USS Seawolf submarine shock testing. *Federal Register*, Vol 63, No 230, 66069–66077.
- National Oceanic and Atmospheric Administration (NOAA), 2006a. NOAA proposes regulations to reduce risk of whale collisions. *Marine Pollution Bulletin*, Vol 52, p 837.
- National Oceanic and Atmospheric Administration (NOAA), 2006b. Proposed Rule Taking and Importing Marine Mammals; Taking Marine Mammals Incidental to Conducting Precision Strike Weapons Testing and Training by Eglin Air Force Base in the Gulf of Mexico, Department of Commerce,

List of Preparers

- NOAA National Marine Fisheries Service, *Federal Register*, Vol 71, No 149, pp 44001–44012. 3 August 2006.
- National Resource Council, 2003. Ocean noise and marine mammals. Washington, D.C.: National Academies Press.
- Niezrecki, C., R. Phillips, M. Meyer, and D.O. Beusse, 2003. Acoustic detection of manatee vocalizations. *Journal of the Acoustical Society of America*, Vol 114, No 3, pp 1640–1647.
- North Carolina Natural Heritage Program, 2004. Natural heritage program list of the rare animal species of North Carolina. Raleigh, North Carolina: North Carolina Natural Heritage Program, North Carolina Department of Environment and Natural Resources.
- Nowacek, D. P., 2005. Acoustic ecology of foraging bottlenose dolphins (*Tursiops truncatus*), habitat-specific use of three sound types. *Marine Mammal Science*, Vol 21, No 4, pp 587–602.
- Nowacek, D. P., B. M. Casper, R. S. Wells, S. M. Nowacek, and D. A. Mann, 2003. Intraspecific and geographic variation of West Indian manatee (*Trichechus manatus* spp.) vocalizations (L). *Journal of the Acoustical Society of America*, Vol 114, No 1, pp 66–69.
- O’Keeffe, D. J., and G. A. Young, 1984. Handbook on the environmental effects of underwater explosions. Report NSWC TR 83-240. Dahlgren, Virginia: Naval Surface Warfare Center.
- Paerl, H. W., J. D. Bales, L. W. Ausley, C. P. Buzzelli, L. B. Crowder, L. A. Eby, J. M. Fear, M. Go, B. L. Peierls, T. L. Richardson, and J. S. Ramus, 2001. Ecosystem impacts of three sequential hurricanes (Dennis, Floyd, and Irene) on the United States largest lagoonal estuary, Pamlico Sound, NC. *Proceedings of the National Academy of Sciences* 98(10): 5655–5660.
- Phillips, R., C. Niezrecki, and D.O. Beusse, 2004. Determination of West Indian manatee vocalization levels and rate. *Journal of the Acoustical Society of America*, Vol 115, No 1, pp 422–428.
- Read, A., K. W. Urian, and D. Waples, 2002. Monitoring bottlenose dolphin use of the Brant Island Shoal Bombing Target (BT-9) and the Piney Island Bombing Range (BT-11) and adjacent waters; July- September, 2002. Progress report prepared for the MCAS Cherry Point by Duke University Marine Laboratory.
- Read, A. J., K. W. Urian, B. Wilson, and D. M. Waples, 2003a. Abundance of bottlenose dolphins in the bays, sounds and estuaries of North Carolina. *Marine Mammal Science* 19(1):59–73.
- Read, A., K. W. Urian, and D. M. Waples, 2003b. Monitoring bottlenose dolphin use of the Brant Island Shoal Bombing Target (BT-9) and the Piney Island Bombing Range (BT-11) and adjacent waters; October-December 2002. Progress report prepared for the MCAS Cherry Point by Duke University Marine Laboratory.
- Read, A., K. W. Urian, and D. M. Waples, 2003c. Monitoring bottlenose dolphin use of the Brant Island Shoal Bombing Target (BT-9) and the Piney Island Bombing Range (BT-11) and adjacent waters; January-March 2003. Progress report prepared for the MCAS Cherry Point by Duke University Marine Laboratory.
- Read, A., K. W. Urian, and D. M. Waples, 2003d. Monitoring bottlenose dolphin use of the Brant Island Shoal Bombing Target (BT-9) and the Piney Island Bombing Range (BT-11) and adjacent waters. Final report prepared for the MCAS Cherry Point by Duke University Marine Laboratory.
- Reid, J. P., G. B. Rathbun, and J. R. Wilcox, 1991. Distribution patterns of individually identifiable West Indian manatees (*Trichechus manatus*) in Florida. *Marine Mammal Science*, Vol 7, No 2, pp 180–190.

List of Preparers

- Reynolds III., J. E., 1981. Behavior patterns in the West Indian manatee with emphasis on feeding and diving. *Florida Scientist*, Vol 44, pp 233–242.
- Ridgway, S. H., B. L. Scronce, and J. Kanwisher, 1969a. Respiration and deep diving in the bottlenose porpoise. *Science*, Vol 166, pp 1651–1654.
- Ridgway, S. H., 2000. The auditory central nervous system, in *Hearing by Whales and Dolphins*, W.W.L. Au, A.N. Popper, and R.R. Fay, eds. Springer-Verlag: New York. pp 273–293.
- Ridgway, S. H., D. A. Carder, R. R. Smith, T. Kamolnick, C. E. Schlundt, and W. R. Elsberry, 1997. Behavioral responses and temporary shift in masked hearing threshold of bottlenose dolphins, *Tursiops truncatus*, to 1-second tones of 141 to 201 dB re 1 μ Pa. Technical Report 1751, Revision 1. San Diego, California: Naval Sea Systems Command.
- Schlundt, C. E., J. J. Finneran, D. A. Carder, and S. H. Ridgway, 2000. Temporary shift in masked hearing thresholds of bottlenose dolphins, *Tursiops truncatus*, and white whales, *Delphinapterous leucas*, after exposure to intense tones. *Journal of the Acoustical Society of America*. 107:3496–3508.
- Schwartz, F. J., 1995. Florida manatees, *Trichechus manatus* (Sirenia: Trichechidae), in North Carolina 1919-1994. *Brimlyeyana* 22:53–60.
- Shane, S. H., 1990. Comparison of bottlenose dolphin behavior in Texas and Florida, with a critique of methods for studying dolphin behavior. Pages 541–558 in S. Leatherwood and R.R. Reeves, eds. The bottlenose dolphin. San Diego: Academic Press.
- Steel, C., and J. G. Morris, 1982. The West Indian manatee: An acoustic analysis. *American Zoologist*, Vol 22, No 4, pp 925–926.
- Stevick, P. T., B. J. McConnell, and P.S. Hammond, 2002. Pages 185-216 in *Patterns of movement. Marine mammal biology: An evolutionary approach*. A. R. Hoelzel. Oxford, United Kingdom, Blackwell Science.
- Thomson, D. H. and W. J. Richardson, 1995. Marine mammal sounds, in *Marine Mammals and Noise*, W. J. Richardson, C. R. Greene, Jr., C. I. Malme, and D. H. Thomson, eds. Academic Press: San Diego. pp 159–204.
- Torres, L. G., P. E. Rosel, C. D'Agrosa, and A. J. Read, 2003. Improving management of overlapping bottlenose dolphin ecotypes through spatial analysis and genetics. *Marine Mammal Science*, Vol 19, No 3, pp 502–514.
- Torres, L. G., W. A. McLellan, E. Meagher, and D. A. Pabst, 2005. Seasonal distribution and relative abundance of bottlenose dolphins, *Tursiops truncatus*, along the US mid-Atlantic Coast. *Journal of Cetacean Research and Management*, Vol 7, No 2, pp 153–161.
- Turl, C. W., 1993. Low-frequency sound detection by a bottlenose dolphin. *Journal of the Acoustical Society of America*, Vol 94, No 5, pp 3006–3008.
- Urian, K., 2005a. Personal communication via email between Ms Kim Urian, Duke University, Beaufort, North Carolina, and Mr. William Rogers, Marine Corps Air Station Cherry Point, Cherry Point, North Carolina, 24 October.
- Urian, K., 2005b. Personal communication via email between Ms Kim Urian, Duke University, Beaufort, North Carolina, and Mr. William Rogers, Marine Corps Air Station Cherry Point, Cherry Point, North Carolina, 7 November.

List of Preparers

- U.S. Fish and Wildlife Service (USFWS), 2001. Florida manatee recovery plan (*Trichechus manatus latirostris*), third revision. Atlanta, Georgia: U.S. Fish and Wildlife Service.
- U.S. Geological Survey (USGS), 2001. Chessie the manatee on a comeback tour after 5-year hiatus. Accessed 7 August 2004. <http://fl.water.usgs.gov/News/ChessieManatee.html>.
- U.S. Marine Corps (USMC), 2001a. Biological Assessment for Ongoing Ordnance Delivery at Bombing Target 9 and Bombing Target 11. Cherry Point, North Carolina: Marine Corps Air Station Cherry Point, Environmental Affairs Department.
- , 2001b. Marine Corps Air Station Cherry Point GIS boundary shapefile. Data received 12 December 2001 from GIS manager. Cherry Point, North Carolina: Marine Corps Air Station Cherry Point.
- , 2004. Orthophotograph of Piney Island, North Carolina. GIS image data received from Denise Smith. Cherry Point, North Carolina: US Marine Corps Air Station Cherry Point.
- Waring, G. T., E. Josephson, C. P. Fairfield, K. Maze-Foley, eds., 2007. U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments—2006. NOAA Technical Memorandum, NMFS-NE-201. April 2007.
- Waring G. T., E. Josephson, C. P. Fairfield-Walsh, and K. Maze-Foley. 2008. US Atlantic and Gulf of Mexico Marine Mammal Stock Assessments – 2007. U.S. Department of Commerce. National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Northeast Fisheries Science Center. NOAA Technical Memorandum NMFS-NE-205. Woods Hole. January.
- Wells, J. T., 1989. A scoping study of the distribution, composition, and dynamics of water-column and bottom sediments: Albemarle-Pamlico estuarine system. Albemarle-Pamlico Estuarine Study Project No. 89-05. Raleigh, North Carolina: North Carolina Department of Environment, Health, and Natural Resources.
- Wells, R. S. & SCOTT, M.D. 1997. Seasonal incidence of boat strikes of bottlenose dolphin near Sarasota, Florida. *Marine Mammal Science* 13: 475–480.
- Wells, R. S., and M. D. Scott, 1999. Bottlenose dolphin—*Tursiops truncatus* (Montagu, 1821). Pages 137–182 in S. H. Ridgway and R. Harrison, eds. *Handbook of marine mammals*. Volume 6: The second book of dolphins and the porpoises. San Diego: Academic Press.
- Wells, R. S., L. J. Hansen, A. Baldrige, T. P. Dohl, D. L. Kelly, and R. H. Defran, 1990. Northward extension of the range of bottlenose dolphins along the California coast, in *The bottlenose Dolphin*, S. Leatherwood and R. R. Reeves, eds. Academic Press: San Diego. pp 421–431.
- Wright, D. G., 1982. A discussion paper on the effects of explosives on fish and marine mammals in the waters of the Northwest Territories. Canadian Technical Report of Fisheries and Aquatic Sciences 1052:1–16.
- Würsig, B., T. A. Jefferson, and D. J. Schmidly, 2000. *The Marine Mammals of the Gulf of Mexico*. Texas A&M University Press: College Station.
- Zaretsky, S. C., A. Martinez, L. P. Garrison, and E. O. Keith, 2005. Differences in acoustic signals from marine mammals in the western North Atlantic and northern Gulf of Mexico, in Abstracts, Sixteenth Biennial Conference on the Biology of Marine Mammals. 12-16 December 2005. S