

List of Core Measures

Core Measures, Data Sources, and Availability for Select Groups

Measure	Data source	Black	Hispanic	Asian or API	AI/AN	Poor
Quality						
Women age 40 and over who reported they had a mammogram within the past 2 years	NHIS	✓	✓	✓	✓	✓
Breast cancer incidence per 100,000 women age 40 and over diagnosed at advanced stage	SEER	✓	✓	✓	✓	
Breast cancer deaths per 100,000 female population per year	NVSS-M	✓	✓	✓	✓	
Adults age 50 and over who received a colonoscopy, sigmoidoscopy, proctoscopy, or fecal occult blood test	NHIS	✓	✓	✓	✓	✓
Colorectal cancer diagnosed at advanced stage per 100,000 population age 50 and over	SEER	✓	✓	✓	✓	
Colorectal cancer deaths per 100,000 population per year	NVSS-M	✓	✓	✓	✓	
Adults age 40 and over with diabetes who received all three exams in the calendar year: hemoglobin A1c measurement, dilated eye examination, and foot examination	MEPS	✓	✓			✓
Adult dialysis patients registered on a waiting list for transplantation	USRDS	✓	✓	✓	✓	
Hemodialysis patients with adequate dialysis	ESRD CPMP	✓	✓	✓	✓	
Hospital care for heart attack patients	QIO	✓	✓	✓	✓	
Hospital care for heart failure patients	QIO	✓	✓	✓	✓	
Deaths per 1,000 adult hospital admissions with acute myocardial infarction	HCUP	✓	✓	✓		✓
New AIDS cases per 100,000 population ages 13 and over	CDC AIDS Surveillance	✓	✓	✓	✓	
Children ages 19-35 months who received all recommended vaccines	NIS	✓	✓	✓	✓	✓

✓ Indicates that reliable data on measure are available for this group and included in summary across measures of quality and access for this group.

Key: API=Asian or Pacific Islander; AI/AN=American Indian or Alaska Native; Poor=individuals with household incomes <100% of Federal poverty thresholds.

Core Measures, Data Sources, and Availability for Select Groups (continued)

Measure	Data source	Black	Hispanic	Asian or API	AI/AN	Poor
Children ages 2-17 for whom a health provider gave advice about physical activity	MEPS	✓	✓	✓		✓
Children ages 3-6 who had their vision checked by a health provider	MEPS	✓	✓			✓
Children ages 2-17 who received advice about healthy eating	MEPS	✓	✓	✓		✓
Children ages 2-17 who had a dental visit in the past year	MEPS	✓	✓	✓		✓
Suicide deaths per 100,000 population	NVSS-M	✓	✓		✓	
Adults with a major depressive episode who received treatment for depression	NSDUH	✓	✓			✓
People age 12 and over who needed treatment for illicit drug use or alcohol problem and who received such treatment at a specialty facility	NSDUH	✓	✓		✓	✓
Adult smokers who received advice to quit smoking	MEPS	✓	✓			✓
Adults with obesity who received advice to exercise more	MEPS	✓	✓			✓
Adults with obesity who received advice about healthy eating	MEPS	✓	✓			✓
Adults age 65 and over who ever received pneumococcal vaccination	NHIS	✓	✓	✓		✓
Hospital care for pneumonia patients	QIO	✓	✓	✓	✓	
Patients with tuberculosis who completed a curative course of treatment within 1 year of initiation of treatment	CDC TB Surveillance	✓	✓	✓	✓	
Older women screened for osteoporosis	MCBS	✓	✓	✓	✓	✓
Long-stay nursing home residents with physical restraints	CMS MDS	✓	✓	✓	✓	
High-risk long-stay nursing home residents with pressure sores	CMS MDS	✓	✓	✓	✓	
Short-stay nursing home residents with pressure sores	CMS MDS	✓	✓	✓	✓	
Adult home health care patients whose ability to walk or move around improved	OASIS	✓	✓	✓	✓	
Adult home health care patients who were admitted to the hospital	OASIS	✓	✓	✓	✓	
Adult surgery patients with postoperative pneumonia or venous thromboembolic event	MPSMS	✓				
Adult surgery patients who received appropriate timing of antibiotics	QIO	✓	✓	✓	✓	

✓ Indicates that reliable data on measure are available for this group and included in summary across measures of quality and access for this group.

Key: API=Asian or Pacific Islander; AI/AN=American Indian or Alaska Native; Poor=individuals with household incomes <100% of Federal poverty thresholds.

Core Measures, Data Sources, and Availability for Select Groups (continued)

Measure	Data source	Black	Hispanic	Asian or API	AI/AN	Poor
Bloodstream infections or mechanical adverse events associated with central venous catheter placements	MPSMS	✓				
Deaths per 1,000 discharges following complications of care	HCUP	✓	✓	✓		✓
Adults age 65 and over who received potentially inappropriate prescription medications	MEPS	✓	✓			✓
Adults who sometimes or never got care for illness or injury as soon as wanted	MEPS	✓	✓	✓		✓
Emergency department visits in which patients left without being seen	NHAMCS	✓		✓		
Adults whose health providers sometimes or never listened carefully, explained things clearly, respected what they had to say, and spent enough time with them	MEPS	✓	✓	✓		✓
Children whose health providers sometimes or never listened carefully, explained things clearly, respected what they or their parents had to say, and spent enough time with them	MEPS	✓	✓	✓		✓
Access						
People under age 65 with health insurance	NHIS	✓	✓	✓	✓	✓
People under age 65 who were uninsured all year	MEPS	✓	✓	✓	✓	✓
People with a specific source of ongoing care	NHIS	✓	✓	✓	✓	✓
People with a usual primary care provider	MEPS	✓	✓	✓	✓	✓
People without a usual source of care who indicated a financial or insurance reason for not having a source of care	MEPS	✓	✓	✓		✓
People who were unable to get or delayed in getting needed care	MEPS	✓	✓	✓	✓	✓

✓ Indicates that reliable data on measure are available for this group and included in summary across measures of quality and access for this group.

Key: API=Asian or Pacific Islander; AI/AN=American Indian or Alaska Native; Poor=individuals with household incomes <100% of Federal poverty thresholds.