

The American Urological Association's Symptom Score for Benign Prostatic Hyperplasia (BPH)						
	Not at all	Less than 1 time in 5 times	Less than half of the time	About half of the time	More than half of the time	Almost always
During the past month or so, how often did you have the feeling of not having completely emptied your bladder after urinating?	0	1	2	3	4	5
During the past month or so, how often did you have to urinate again less than 2 hours after you finished urinating?	0	1	2	3	4	5
During the past month or so, how often did you have to stop and start the urinary stream several times while urinating?	0	1	2	3	4	5
During the past month or so, how often did you find it difficult to delay urination?	0	1	2	3	4	5
During the past month or so, how often did you find that your urinary stream was weak?	0	1	2	3	4	5
During the past month or so, how often did you have to push or strain to begin urinating?	0	1	2	3	4	5
	Frequency of Waking Up to Urinate					
During the past month or so, how often did you typically get up to urinate between going to bed in the evening and waking up in the morning?	0	1 (one time)	2 (two times)	3 (three times)	4 (four times)	5 (five times or more)

Score Key:

Mild = 0 to 7 points

Moderate = 8 to 9 points

Severe = 20 to 35 points

Adapted from Barry MJ, Fowler FJ, O'Leary MP, et al. The American Urological Association Symptom Index for benign prostatic hyperplasia. *Journal of Urology* 148:1549, 1992.