

## **GRAY SEAL (*Halichoerus grypus*): Western North Atlantic Stock**

### **STOCK DEFINITION AND GEOGRAPHIC RANGE**

The gray seal is found on both sides of the North Atlantic, with three major populations: eastern Canada, northwestern Europe and the Baltic Sea (Katona *et al.* 1993; R. DiGiovanni, pers. comm., The Riverhead Foundation, Riverhead, NY). The western North Atlantic stock is equivalent to the eastern Canada population, and ranges from New England to Labrador (Davies 1957; Mansfield 1966; Katona *et al.* 1993; Lesage and Hammill 2001). This stock is separated by geography, differences in the breeding season, and mitochondrial DNA variation from the northeastern Atlantic stock (Bonner 1981; Boskovic *et al.* 1996; Lesage and Hammill 2001). There are two breeding concentrations in eastern Canada; one at Sable Island, and one that breeds on the pack ice in the Gulf of St. Lawrence ((Laviguer and Hammill 1993)). Tagging studies indicate that there is little intermixing between the two breeding groups (Zwanenberg and Bowen 1990) and, for management purposes, they are treated by the Canadian DFO as separate stocks (Mohn and Bowen 1996). In the mid 1980s, small numbers of animals and pupping were observed on several isolated islands along the Maine coast and in Nantucket-Vineyard Sound, Massachusetts (Katona *et al.* 1993; J. R. Gilbert, pers. comm., University of Maine, Orono, ME; Rough 1995). In the late 1990s, a year-round breeding population of approximately 400+ animals was documented on outer Cape Cod and Muskeget Island (D. Murley, pers. comm., Mass. Audubon Society, Wellfleet, MA). In December 2001, NMFS initiated aerial surveys to monitor gray seal pup production on Muskeget Island and at the Monomoy National Wildlife Refuge. Gilbert (pers. comm.) has also documented resident colonies and pupping in Maine since 1994.

### **POPULATION SIZE**

Current estimates of the total western Atlantic gray seal population are not available; although estimates of portions of the stock are available for select time periods. The size of the Canadian population from 1993 to 2004 has been estimated from three surveys. A 1993 survey estimated the population at 144,000 animals (DFO 2003, Mohn and Bowen 1996), a 1997 survey estimated 195,000 (DFO [Dept. Of Fisheries and Oceans] 2003), and a 2004 survey obtained estimates ranging between 208,720 (SE=29,730) and 223,220 (SE=17,376) depending upon the model used (Trzcinski *et al.* 2005). The population at Sable Island had been increasing by approximately 13% per year for nearly 40 years (Bowen *et al.* 2003), but the most recent (2004) survey results indicate this population increase has declined to 7% (Trzcinski *et al.* 2005; Bowen *et al.* 2007). The non-Sable Island (Gulf of St Lawrence and Eastern Shore) abundance has increased from 20,900 (SE=200) in 1970 to 52,500 (SE=7,800) in 2004 (Hammill 2005).

The population in US waters is also increasing. Maine coast-wide surveys conducted during summer (all other surveys were conducted January-May) revealed 597 and 1,731 gray seals in 1993 and 2001, respectively (Gilbert *et al.* 2005). In 2002, the maximum counts of two breeding colonies in Maine, with number of pups in parentheses, were 193 (9) on Seal Island and 74 (31) on Green Island (S. Wood, pers. comm.). Gray seal numbers are increasing in Massachusetts at Muskeget Island off the coast of Nantucket, and at Monomoy Island, off the coast of Chatham, Cape Cod. Pup counts on Muskeget have increased from 0 in 1989 to 1,023 in 2002 (Rough 1995, S. Wood, pers. comm.). Gray seal numbers increase in this region in the spring (April-May) when molting occurs. In April-May 1994 a maximum count of 2,010 was obtained for Muskeget Island and Monomoy combined (Rough 1995). In March 1999 a maximum count of 5,611 was obtained in the region south of Maine (between Isles of Shoals, Maine and Woods Hole, Massachusetts) (Barlas 1999). No gray seals were recorded at haul out sites between Newport, Rhode Island and Montauk Pt., New York (Barlas 1999), although, more recently small numbers of gray seals have been recorded in this region (deHart 2002; R. DiGiovanni, pers. comm., The Riverhead Foundation, Riverhead, NY). Recently, a small number of gray seals have maintained a winter presence in the Woods Hole region (Vineyard Sound) (deHart 2002).

Table 1. Summary of abundance estimates for the western North Atlantic gray seal. Month, year, and area covered during each abundance survey, resulting abundance estimate ( $N_{best}$ ) and coefficient of variation (CV).			
Month/Year	Area	Nbest	CV
May 2001 <sup>a</sup>	Maine coast	1,731	NA
January 2004 <sup>b</sup>	Gulf of St Lawrence + Nova Scotia Eastern Shore	52,500	0.15
January 2004 <sup>b</sup>	Sable Island	208,720 216,490 223,220	0.14 0.11 0.08
<sup>a</sup> These counts pertain to animals seen in U.S. waters, and the stock relationship to animals in Canadian waters is unknown. <sup>b</sup> These are model based estimates derived from pup surveys.			

### Minimum Population Estimate

Depending on the model used, the  $N_{min}$  for the Canadian gray seal population was estimated to range between 125,541 and 169,064 (Trzcinski et al. 2005). Present data are insufficient to calculate the minimum population estimate for U.S. waters.

### Current Population Trend

Gray seal abundance is likely increasing in the U.S. Atlantic Exclusive Economic Zone (EEZ), but the rate of increase is unknown. The population in eastern Canada was greatly reduced by hunting and bounty programs, and in the 1950s the gray seal was considered rare (Lesage and Hammill 2001). The Sable Island population was less affected and has been increasing for several decades. Pup production on Sable Island, Nova Scotia, has increased exponentially at a rate of 12.8% annually for more than 40 years (Stobo and Zwanenburg 1990; Mohn and Bowen 1996; Bowen et al. 2003; Trzcinski et al. 2005; Bowen et al. 2007), but has declined to 7% in 2004 (Trzcinski et al. 2005; Bowen et al. 2007). The non Sable Island population increased from 6,900 in the mid-1980s to a peak of 11,100 (SE=1,300) animals in 1996 (Hammill and Gosselin 2005). Pup production declined to 6,100 (SE=900) in 2000, then increased to 15,900 (SE=1,200) in 2004 (Hammill and Gosselin 2005). Approximately 57% of the western North Atlantic population is from the Sable Island stock. In recent years pupping has been established on Hay Island, off the Cape Breton coast (Lesage and Hammill 2001).

Winter breeding colonies in Maine and on Muskeget Island may provide some measure of gray seal population trends and expansion in distribution. Sightings in New England increased during the 1980s as the gray seal population and range expanded in eastern Canada. Five pups were born at Muskeget in 1988. The number of pups increased to 12 in 1992, 30 in 1993, and 59 in 1994 (Rough 1995). In January 2002, between 900 and 1,000 pups were counted on Muskeget Island and surrounding shoals (S. Wood, pers. comm.). In recent years NMFS monitoring surveys have detected an occasional mother/pup (white coats) pair on both Monomoy Island and Nomans Land. These observations continue the increasing trend in pup production reported by Rough (1995). The change in gray seal counts at Muskeget and Monomoy from 2,010 in 1994 to 5,611 in 1999 represents an annual increase rate of 20.5%, however, it can not be determined what proportion of the increase represents growth or immigration.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. A recent study estimated the annual rate of increase at 7% on Sable Island (Trzcinski et al. 2005; Bowen et al. 2007), which represents a 45% decline from previous estimates (Mohn and Bowen 1996; Bowen et al. 2003). For purposes of this assessment, the maximum net productivity rate was assumed to be 0.12. This value is based on theoretical modeling showing that pinniped populations may not grow at rates much greater than 12% given the constraints of their reproductive life history (Barlow et al. 1995).

### POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a "recovery" factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum

population size is unknown. The maximum productivity rate is 0.12, the default value for pinnipeds. The recovery factor ( $F_R$ ) for this stock is 1.0, the value for stocks of unknown status, but is known to be increasing. PBR for the western North Atlantic gray seals in U.S. waters is unknown.

### **ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY**

For the period 2002-2006, the total estimated human caused mortality and serious injury to gray seals was 836 per year. The average was derived from three components: 1) 331 (CV=0.21) (Table 2) from the 2002-2006 U.S. observed fishery; 2) 2 from average 2002-2006 non-fishery related, human interaction stranding mortalities (NMFS unpublished data); and 3) 503 from average 2002-2006 kill in the Canadian hunt (DFO [Dept. Of Fisheries and Oceans] 2003; G. Stenson unpublished data; M. Hammill pers.comm., DFO, Mont-Joli, Quebec).

### **Fishery Information**

Detailed fishery information is given in Appendix III.

### **U.S.**

#### **Northeast Sink Gillnet**

Annual estimates of gray seal bycatch in the Northeast sink gillnet fishery reflect seasonal distribution of the species and of fishing effort. There were 105 gray seal mortalities observed in the Northeast sink gillnet fishery between 1993 and 2006. Estimated annual mortalities (CV in parentheses) from this fishery were 0 in 1990-1992, 18 in 1993 (1.00), 19 in 1994 (0.95), 117 in 1995 (0.42), 49 in 1996 (0.49), 131 in 1997 (0.50), 61 in 1998 (0.98), 155 in 1999 (0.51), 193 in 2000 (0.55), 117 in 2001 (0.59), 0 in 2002, 242 (0.47) in 2003, 504 (0.34) in 2004, 574 (0.44) in 2005, and 314 (0.22) in 2006 (Table 2). There were 2, 2, 9, 14, and 8 unidentified seals observed during 2002-2006, respectively. Since 1997 unidentified seals have not been prorated to a species. This is consistent with the treatment of other unidentified mammals that do not get prorated to a specific species. Average annual estimated fishery-related mortality and serious injury to this stock attributable to this fishery during 2002-2006 was 314 gray seals (CV=0.22) (Table 2). The stratification design used is the same as that for harbor porpoise (Bravington and Bisack 1996).

#### **Mid-Atlantic Coastal Gillnet**

No gray seals were taken in observed trips during 1998-2000, 2003, and 2006. One gray seal was observed taken in both 2001 and 2004 (Table 2). In 2001 the gray seal was taken in April off the coast of New Jersey near Hudson Canyon in 81 m of water. The 2004 take was off Virginia in April. Observed effort was scattered between New Jersey and North Carolina from 1 to 90 km off the beach. In 2002, 65% of sampling was concentrated in one area and not distributed proportionally across the fishery. Therefore, observed mortality is considered unknown in 2002. Average annual estimated fisher-related mortality and serious injury to this stock attributable to this fishery during 2002-2006 was 17 gray seals (CV=0.92) (Table 2).

#### **Gulf of Maine Atlantic Herring Purse Seine Fishery**

The Gulf of Maine Atlantic Herring Purse Seine Fishery is a Category III fishery. This fishery was not observed until 2003, and was not observed in 2006. No mortalities have been observed, but 15 gray seals were captured and released alive in 2004 and 19 in 2005. In addition, 5 seals of unknown species were captured and released alive in 2004 and 2 in 2005.

#### **Northeast Bottom Trawl**

Vessels in the North Atlantic bottom trawl fishery, a Category III fishery under MMPA, were observed in order to meet fishery management, rather than marine mammal management needs. No mortalities were observed prior to 2005, when four mortalities were attributed to this fishery. The estimated annual fishery-related mortality and serious injury attributable to this fishery was 0 between 2001 and 2004, and for 2006. Estimates have not been generated for 2005.

### **CANADA**

An unknown number of gray seals have been taken in Newfoundland and Labrador, Gulf of St. Lawrence, and Bay of Fundy groundfish gillnets, Atlantic Canada and Greenland salmon gillnets, Atlantic Canada cod traps, and in Bay of Fundy herring weirs ((Read 1994). In addition to incidental catches, some mortalities (e.g., seals trapped in herring weirs) were the result of direct shooting, and there were culls of about 1,700 animals annually during the

1970s and early 1980s on Sable Island (Anonymous 1986).

In 1996, observers recorded 3 gray seals (1 released alive) in Spanish deep-water trawl fishing on the southern edge of the Grand Banks (NAFO Areas 3) (Lens 1997). Seal bycatch occurred year-round, but interactions were highest during April-June. Many of the seals that died during fishing activities were unidentified. The proportion of sets with mortality (all seals) was 2.7 per 1,000 hauls (0.003).

Table 2. Summary of the incidental mortality of gray seal ( <i>Halichoerus grypus</i> ) by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the mortalities recorded by on-board observers (Observed Mortality), the estimated annual mortality (Estimated Mortality), the estimated CV of the annual mortality (Estimated CVs) and the mean annual mortality (CV in parentheses).								
Fishery	Years	Vessels	Data Type <sup>a</sup>	Observer Coverage <sup>b</sup>	Observed Mortality	Estimated Mortality	Estimated CVs	Mean Annual Mortality
Northeast Sink Gillnet <sup>c</sup>	02-06	unk	Obs. Data, Weighout, Logbooks	.02, .03, .06, .07, .04,	0, 5, 21, 33, 9	0, 242, 504, 574, 248	0, .47, .34, .44, .47	314 (0.22)
Mid-Atlantic Gillnet	02-06	unk	Obs. Data, Weighout	.01, .01, .02, .03, .04	unk <sup>e</sup> , 0, 1, 0, 0	unk <sup>d</sup> , 0, 69, 0, 0	unk <sup>d</sup> , 0, .92, 0, 0	17 <sup>d</sup> (0.92)
Northeast Bottom Trawl	02-06	unk	Obs. Data, Weighout	.03, .04, .05, .12, .06	0, 0, 0, 4, 0	0, 0, 0, unk <sup>f</sup> , 0	0, 0, 0, unk <sup>f</sup> , 0	unk <sup>f</sup>
TOTAL								331 (0.21)

a. Observer data (Obs. Data) are used to measure bycatch rates, and the data are collected within the Northeast Fisheries Observer Program. The Northeast Fisheries Observer Program collects landings data (Weighout), and total landings are used as a measure of total effort for the sink gillnet fishery. Mandatory logbook (Logbook) data are used to determine the spatial distribution of fishing effort in the Northeast multispecies sink gillnet fishery.

b. The observer coverages for the Northeast sink gillnet fishery and the mid-Atlantic gillnet fisheries are ratios based on tons of fish landed.

c. Since 1998, takes from pingered and non-pingered nets within a marine mammal time/area closure that required pingers, and takes from pingered and non-pingered nets not within a marine mammal time/area closure were pooled. The pooled bycatch rate was weighted by the total number of samples taken from the stratum and used to estimate the mortality. In 1998, 1 take was observed in a net without a pinger that was within a marine mammal closure that required pingers. In 2002 - 2006, respectively, 0, 1, 1, 1, and 1 takes were observed in nets with pingers. In 2002 - 2006, respectively, 2, 0, 4, 20, 32, and 8 takes were observed in nets without pingers.

d. Sixty-five percent of sampling in the mid-Atlantic coastal gillnet by the Northeast Fisheries Observer Program was concentrated in one area off the coast of Virginia. Because of the low level of sampling that was not distributed proportionately throughout the mid-Atlantic region observed mortality is considered unknown in 2002. The four year average (2003- 2006) estimated mortality was applied as the best representative estimate.

f. Analysis of bycatch mortality attributed to the Northeast bottom trawl fishery has not been generated.

### Other Mortality

**Canada:** In Canada, gray seals were hunted for several centuries by indigenous people and European settlers in the Gulf of St. Lawrence and along the Nova Scotia eastern shore, and were locally extirpated (Laviguier and Hammill 1993). Between 1999 and 2006 the annual kill of gray seals by hunters in Canada was: 1999 (98), 2000 (342), 2001 (76) 2002 (126), 2003 (6), 2004 (0), 2005 (579), and 2006 (1804). (DFO [Dept. Of Fisheries and Oceans] 2003; M. Hammill pers. comm.). The traditional hunt of a few hundred animals is expected to continue off the Magdalen Islands and in other areas, except Sable Island where commercial hunting is not permitted (DFO [Dept. Of Fisheries and Oceans] 2003). DFO established an annual (2006-2010) TAC of 2,100 gray seals in the Gulf of St. Lawrence, and 8,300 on the Scotian Shelf. The hunting of grey seals will continue to be prohibited on Sable Island ([http://www.dfo-mpo.gc.ca/seal-phoque/index\\_e.htm](http://www.dfo-mpo.gc.ca/seal-phoque/index_e.htm)).

Canada also issues personal hunting licenses which allow the holder to take six gray seals annually (Lesage and Hammill 2001). Hunting is not permitted during the breeding season and some additional seasonal/spatial restrictions are in effect (Lesage and Hammill 2001).

**U.S:** Gray seals, like harbor seals, were hunted for bounty in New England waters until the late 1960s. This hunt may have severely depleted this stock in U.S. waters (Rough 1995). Other sources of mortality include human interactions, storms, abandonment by the mother, disease, and predation. Mortalities caused by human interactions include boat strikes, fishing gear interactions, power plant entrainment, oil spill/exposure, harassment, and shooting. The Cape Cod stranding network has documented gray seals entangled in netting or plastic debris around the Cape Cod/Nantucket area, and in recent years have made successful disentanglement attempts.

From 2002-2006, 213 gray seal stranding mortalities were recorded, extending from Maine to North Carolina (Table 3; NMFS unpublished data). Most stranding mortalities were in Massachusetts. Thirty-seven (17.4%) of the total stranding mortalities showed signs of human interaction (6 in 2002, 7 in 2003, 16 in 2004, 3 in 2005, and 5 in 2006), with 25 having some indication of fishery interaction (3 in 2002, 5 in 2003, 11 in 2004, 1 in 2005, and 5 in 2006).

State	2002	2003	2004	2005	2006	Total
ME	2	2	3	4	3	14
NH		1				1
MA	32	58	33	26	29	178
RI	1	6	8	2	2	19
CT			2			2
NY	5	5	2	7	6	25
NJ		2		2	1	4
DE			1			1
MD			1	3		4
VA		1	2	1		4
NC	1				2	1
Total	41	75	52	45	43	213
Unspecified seals (all states)	35	27	33	59	46	200

a. Some of the data reported in this table differ from that reported in previous years. We have reviewed the records and made an effort to standardize reporting. Live releases and rehabbed animals have been eliminated. Mortalities include those which stranded dead, died at site, were euthanized, died during transport, or died soon after transfer to rehab.

## STATUS OF STOCK

The status of the gray seal population relative to OSP in U.S. Atlantic EEZ waters is unknown, but the stock's abundance appears to be increasing in Canadian and U.S. waters. The species is not listed as threatened or endangered under the Endangered Species Act. The total U.S. fishery-related mortality and serious injury for this stock is low relative to the stock size in Canadian and U.S. waters and can be considered insignificant and approaching zero mortality and serious injury rate. The level of human-caused mortality and serious injury in the U.S. Atlantic EEZ is unknown, but believed to be very low relative to the total stock size; therefore, this is not a strategic stock.

## REFERENCES CITED

- Anonymous 1986. Seals and sealing in Canada. Rep. of the Royal Commission on Seals and Sealing, Vol. 1. Canadian Government Publishing Centre, Ottawa, Ontario, Canada 65 pp.
- Barlas, M. E. 1999. The distribution and abundance of harbor seals (*Phoca vitulina concolor*) and gray seals (*Halichoerus grypus*) in southern New England, winter 1998-summer 1999. M.A. thesis. Graduate School of Arts and Sciences. Boston, MA, Boston University. 52 pp.

- Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade 1995. U.S. Marine Mammal Stock Assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- Bonner, W. N. 1981. Grey seal *Halichoerus grypus Fabricus*, 1791. Pages 111-144 in: Bonner, W. N., (ed.) Handbook of marine mammals, Vol. 2: Seals. Academic Press. London.
- Boskovic, R., K. M. Kovacs, M. O. Hammill and B. N. White 1996. Geographic distribution of mitochondrial DNA haplotypes in grey seals (*Halichoerus grypus*). Can. J. Zool. 74: 1787-1796.
- Bowen, W. D., J. McMillan and R. Mohn 2003. Sustained exponential population growth of grey seals at Sable Island, Nova Scotia. ICES J. Mar. Sci. 60: 1265-1274.
- Bowen, W. D., J. I. McMillan and W. Blanchard 2007. Reduced population growth of gray seals at Sable Island: Evidence from pup production and age of primiparity. Mar. Mamm. Sci. 23: 48-64.
- Bravington, M. V. and K. D. Bisack 1996. Estimates of harbour porpoise bycatch in the Gulf of Maine sink gillnet fishery, 1990-1993. Rep. Int. Whal. Comm. 46: 567-574.
- Davies, J. L. 1957. The geography of the gray seal. J. Mamm. 38: 297-310.
- deHart, P. A. P. 2002. The distribution and abundance of harbor seals (*Phoca vitulina concolor*) in the Woods Hole region. M.A. thesis. Graduate School of Arts and Sciences. Boston, MA, Boston University. 88 pp.
- DFO [Dept. Of Fisheries and Oceans] 2003. Atlantic Seal Hunt: 2003-2005 management plan. Department of Fisheries and Oceans, Ottawa, Ontario, Canada, K1A 0E6. Fisheries Resource Management – Atlantic 34 pp.
- Gilbert, J. R., G. T. Waring, K. M. Wynne and N. Guldager 2005. Changes in abundance and distribution of harbor seals in Maine, 1981-2001. Mar. Mamm. Sci. 21: 519-535.
- Hammill, M. O. 2005. Abundance of Northwest Atlantic grey seals in the Gulf of St. Lawrence and along the Nova Scotia Eastern Shore. C. D. o. F. a. Oceans, Ottawa, Ontario. DFO Research Document 2005/036. 11 pp.
- Hammill, M. O. and J. F. Gosselin 2005. Pup production of non-Sable Island grey seals in 2004. C. D. o. F. a. Oceans, Ottawa, Ontario. DFO Research Document 2005/036. 20 pp.
- Katona, S. K., V. Rough and D. T. Richardson 1993. A field guide to whales, porpoises, and seals from Cape Cod to Newfoundland. Washington, DC, Smithsonian Institution Press. 316 pp.
- Laviguer, L. and M. O. Hammill 1993. Distribution and seasonal movements of grey seals, *Halichoerus grypus*, born in the Gulf of St. Lawrence and eastern Nova Scotia shore. Can. Field-Nat. 107: 329-340.
- Lens, S. 1997. Interactions between marine mammals and deep water trawlers in the NAFO regulatory area. ICES [Int. Counc. Explor. Sea] C.M. 1997/Q:08: 10.
- Lesage, V. and M. O. Hammill 2001. The status of the grey seal, *Halichoerus grypus*, in the Northwest Atlantic. Can. Field-Nat. 115(4): 653-662.
- Mansfield, A. W. 1966. The grey seal in eastern Canadian waters. Can. Audubon Mag. 28: 161-166.
- Mohn, R. and W. D. Bowen 1996. Grey seal predation on the eastern Scotian Shelf: Modeling the impact on Atlantic cod. Can. J. Fish. Aquat. Sci 53: 2722-2738.
- Read, A. J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest Atlantic. Pages 133-147 in: Read, A. J., (ed.) Gillnets and cetaceans. Reports of the International Whaling Commission. (Special Issue) 15.
- Rough, V. 1995. Gray seals in Nantucket Sound, Massachusetts, winter and spring, 1994. Final report to Marine Mammal Commission. Contract T10155615: 28.
- Stobo, W. T. and K. C. T. Zwanenburg 1990. Grey seal (*Halichoerus grypus*) pup production on Sable Island and estimates of recent production in the northwest Atlantic. Pages 171-184 in: Stobo, W. T. and K. C. T. Zwanenburg, (eds.) Population biology of sealworm (*Pseudoterranova decipiens*) in relation to its intermediate and seal hosts. Can. Bull. Fish. and Aq. Sci. 222.
- Trzcinski, M. K., R. Mohn and W. D. Bowen 2005. Estimation of grey seal population size and trends at Sable Island. Canadian Department of Fisheries and Oceans, Ottawa, Ontario. DFO Research Document 2005/067. 10 pp.
- Wade, P. R. and R. P. Angliss 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Zwanenburg, K. C. T. and W. D. Bowen 1990. Population trends of the grey seal (*Halichoerus grypus*) in eastern Canada. Pages 185-197 in: Zwanenburg, K. C. T. and W. D. Bowen, (eds.) Population biology of sealworm (*Pseudoterranova decipiens*) in relation to its intermediate and seal hosts. Can. Bull. Fish. and Aq. Sci. 222.