

## RISSO'S DOLPHIN (*Grampus griseus*): Western North Atlantic Stock

### STOCK DEFINITION AND GEOGRAPHIC RANGE


Risso's dolphins are distributed worldwide in tropical and temperate seas, and in the Northwest Atlantic occur from Florida to eastern Newfoundland (Leatherwood *et al.* 1976; Baird and Stacey 1990). Off the northeastern U.S. coast, Risso's dolphins are distributed along the continental shelf edge from Cape Hatteras northward to Georges Bank during spring, summer, and autumn (CETAP 1982; Payne *et al.* 1984). In winter, the range is in the mid-Atlantic Bight and extends outward into oceanic waters (Payne *et al.* 1984). In general, the population occupies the mid-Atlantic continental shelf edge year round, and is rarely seen in the Gulf of Maine (Payne *et al.* 1984). During 1990, 1991 and 1993, spring/summer surveys conducted along the continental shelf edge and in deeper oceanic waters sighted Risso's dolphins associated with strong bathymetric features, Gulf Stream warm-core rings, and the Gulf Stream north wall (Waring *et al.* 1992; 1993; Hamazaki 2002). There is no information on stock structure of Risso's dolphin in the western North Atlantic, or to determine if separate stocks exist in the Gulf of Mexico and Atlantic. In 2006, a rehabilitated adult male Risso's dolphin stranded and released in the Gulf of Mexico off Florida was tracked via satellite to waters off Delaware (Wells *et al.* 2009). The Gulf of Mexico and Atlantic stocks are currently being treated as two separate stocks.

### POPULATION SIZE

Total numbers of Risso's dolphins off the U.S. or Canadian Atlantic coast are unknown, although eight abundance estimates are available from selected regions for select time periods. Sightings were almost exclusively in continental shelf edge and continental slope areas (Figure 1). The best abundance estimate for Risso's dolphins is the sum of the estimates from the two 2004 U.S. Atlantic surveys, 20,479 (CV=0.59), where the estimate from the northern U.S. Atlantic is 15,053 (CV=0.78), and from the southern U.S. Atlantic is 5,426 (CV=0.54). This joint estimate is considered best because these two surveys together have the most complete coverage of the population's habitat.

### Earlier abundance estimates

Please see appendix IV for earlier abundance estimates. As recommended in the GAMMS Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable, therefore should not be used for PBR determinations. Further, due to changes in survey methodology these data should not be used to make comparisons to more current estimates.



**Figure 1.** Distribution of Risso's dolphin sightings from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1998, 1999, 2002, 2004, 2006 and 2007. Isobaths are the 100-m, 1,000-m, and 4,000-m depth contours.

### Recent surveys and abundance estimates

An abundance estimate of 15,054 (CV=0.78) Risso's dolphins was obtained from a line-transect sighting survey conducted during 12 June to 4 August 2004 by a ship and plane that surveyed 10,761 km of track line in waters north of Maryland (38°N) to the Bay of Fundy (45°N) (Table 1; Palka 2006). Shipboard data were collected using the two-independent-team line-transect method and analyzed using the modified direct-duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and  $g(0)$ , the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line-transect method (Hiby 1999) and analyzed accounting for  $g(0)$  and biases due to school size and other potential covariates (Palka 2005).

A shipboard survey of the U.S. Atlantic outer continental shelf and continental slope (water depths >50 m) between Florida and Maryland (27.5-38°N latitude) was conducted during June-August 2004. The survey employed two independent visual teams searching with 25x bigeye binoculars. Survey effort was stratified to include increased effort along the continental shelf break and Gulf Stream front in the mid-Atlantic. The survey included 5,659 km of trackline, and recorded a total of 473 cetacean sightings. Sightings were most frequent in waters north of Cape Hatteras, North Carolina along the shelf break. Data were analyzed to correct for visibility bias ( $g(0)$ ) and group-size bias employing line-transect distance analysis and the direct-duplicate estimator (Palka 1995; Buckland *et al.* 2001). The resulting abundance estimate for Risso's dolphins between Florida and Maryland was 5,426 (CV=0.54).

An abundance estimate of 14,408 (CV=0.38) Risso's dolphins was obtained from an aerial survey conducted in August 2006 which covered 10,676 km of trackline in the region from the 2,000-m depth contour on the southern edge of Georges Bank to the upper Bay of Fundy and to the entrance of the Gulf of St. Lawrence (Table 1; Palka, pers. comm.). The value of  $g(0)$  used for this estimation was derived from the pooled 2002, 2004 and 2006 aerial survey data.

Table 1. Summary of abundance estimates for the western North Atlantic Risso's dolphin. Month, year, and area covered during each abundance survey, resulting abundance estimate ( $N_{best}$ ) and coefficient of variation (CV).			
Month/Year	Area	$N_{best}$	CV
Jun-Aug 2004	Maryland to Bay of Fundy	15,053	0.78
Jun-Aug 2004	Florida to Maryland	5,426	0.54
Jun-Aug 2004	Florida to Bay of Fundy (COMBINED)	20,479	0.59
Aug 2006	S. Gulf of Maine to upper Bay of Fundy to Gulf of St. Lawrence	14,408	0.38

### Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for Risso's dolphins is 20,479 (CV=0.59), obtained from the 2004 surveys. The minimum population estimate for the western North Atlantic Risso's dolphin is 12,920.

### Current Population Trend

There are insufficient data to determine population trends for this species.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

## POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 12,920. The maximum productivity rate is 0.04, the default value for cetaceans (Barlow *et al.* 1995). The “recovery” factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.48 because the CV of the average mortality estimate is between 0.3 and 0.6 (Wade and Angliss 1997). PBR for the western North Atlantic stock of Risso’s dolphin is 124.

## ANNUAL HUMAN-CAUSED MORTALITY

Total annual estimated average fishery-related mortality or serious injury to this stock during 2005-2009 was 18 Risso’s dolphins (CV=0.37; Table 2).

## Fishery Information

Detailed fishery information is reported in Appendix III.

## Earlier Interactions

Prior to 1977, there was no documentation of marine mammal bycatch in distant-water fleet (DWF) activities off the northeastern coast of the U.S. With implementation of the Fisheries Conservation and Management Act in that year, an observer program was established which recorded fishery data and information on incidental bycatch of marine mammals. NMFS foreign-fishery observers reported four deaths of Risso's dolphins incidental to squid and mackerel fishing activities in the continental shelf and continental slope waters between March 1977 and December 1991 (Waring *et al.* 1990; NMFS unpublished data).

In the pelagic drift gillnet fishery, 51 Risso's dolphin mortalities were observed between 1989 and 1998. One animal was entangled and released alive. Bycatch occurred during July, September and October along continental shelf edge canyons off the southern New England coast. Estimated annual mortality and serious injury (CV in parentheses) attributable to the drift gillnet fishery was 87 in 1989 (0.52), 144 in 1990 (0.46), 21 in 1991 (0.55), 31 in 1992 (0.27), 14 in 1993 (0.42), 1.5 in 1994 (0.16), 6 in 1995 (0), 0 in 1996, no fishery in 1997, and 9 in 1998 (0). This fishery was closed effective in 1999.

In the pelagic pair trawl fishery, one mortality was observed in 1992. Estimated annual fishery-related mortality (CV in parentheses) attributable to the pelagic pair trawl fishery was 0.6 dolphins in 1991 (1.0), 4.3 in 1992 (0.76), 3.2 in 1993 (1.0), 0 in 1994 and 3.7 in 1995 (0.45). This fishery ended as of 1996.

## Pelagic Longline

Pelagic longline bycatch estimates of Risso’s dolphins in 1998, 1999, and 2000 were obtained from Yeung (1999), Yeung *et al.* (2000), and Yeung (2001), respectively. Bycatch estimates for 2001 - 2009 were obtained from Garrison (2003), Garrison and Richards (2004), Garrison (2005), Fairfield Walsh and Garrison (2006), Fairfield Walsh and Garrison (2007), Fairfield and Garrison (2008), Garrison *et al.* (2009) and Garrison and Stokes (2010). Most of the estimated marine mammal bycatch was from U.S. Atlantic EEZ waters between South Carolina and Cape Cod. Excluding the Gulf of Mexico, from 1992 to 2000 one mortality was observed in both 1994 and 2000, and 0 in other years. The observed numbers of seriously-injured but released alive individuals from 1992 to 2009 were, respectively, 2, 0, 6, 4, 1, 0, 1, 1, 1, 6, 4, 2, 2, 0, 0, 1, 3, and 11 (Cramer 1994; Scott and Brown 1997; Johnson *et al.* 1999; Yeung 1999; Yeung *et al.* 2000; Yeung 2001; Garrison 2003; Garrison and Richards 2004; Garrison 2005; Fairfield Walsh and Garrison 2006; Fairfield Walsh and Garrison 2007; Fairfield and Garrison 2008 Garrison *et al.* (2009) and Garrison and Stokes (2010) (Table 2). Estimated annual fishery-related mortality (CV in parentheses) was 17 animals in 1994 (1.0), 41 in 2000 (1.0), 24 in 2001(1.0), 20 in 2002 (0.86), and 0 in 2003 to 2009 (Table 2). Seriously injured and released alive animals were estimated to be 54 dolphins (0.7) in 1992, 0 in 1993, 120 (0.57) in 1994, 103 (0.68) in 1995, 99 (1.0) in 1996, 0 in 1997, 57 (1.0) in 1998, 22 (1.0) in 1999, 23 (1.0) in 2000, 45 (0.7) in 2001, 8 (1.0) in 2002, 40 (0.63) in 2003 28(0.72) in 2004, 3(1.0), 0 in 2005, 0 in 2006, 9 in 2007, 17 in 2008, and 11 in 2009 (Table 2). There is a high likelihood that dolphins released alive with ingested gear or gear wrapped around appendages will not survive (Wells *et al.* 2008a). The annual average combined mortality and serious injury for 2005-2009 is 8 Risso’s dolphins (CV =0.40; Table 2).

## Northeast Sink Gillnet

Estimated annual mortalities (CV in parentheses) from this fishery are: 0 in 1999, 15 (1.06) in 2000, 0 in 2001-2004, 15 in 2005 (0.93), and 0 in 2006 through 2009 (Table 2). The 2005-2009 average mortality in this fishery is 3

Risso's dolphins (CV =0.93).

**Mid-Atlantic Gillnet**

A Risso's dolphin mortality was observed in this fishery for the first time in 2007. The resulting estimated annual mortality for 2007 was 34 (CV=0.73). The 2005-2009 average mortality in this fishery is 7 Risso's dolphins (CV=0.73).

**Mid-Atlantic Mid-water Trawl**

A Risso's dolphin mortality was observed in this fishery for the first time in 2008. No bycatch estimate has been generated.

Table 2. Summary of the incidental mortality of Risso's dolphin ( <i>Grampus griseus</i> ) by commercial fishery including the years sampled (Years), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the observed mortalities and serious injuries recorded by on-board observers, the estimated annual mortality and serious injury, the combined annual estimates of mortality and serious injury (Estimated Combined Mortality), the estimated CV of the combined estimates (Estimated CVs) and the mean of the combined estimates (CV in parentheses).										
Fishery	Years	Data Type <sup>a</sup>	Observer Coverage	Observed Serious Injury	Observed Mortality	Estimated Serious Injury	Estimated Mortality	Estimated Combined Mortality	Estimated CVs	Mean Annual Mortality
Pelagic Longline <sup>b</sup>	05-09	Obs. Data Logbook	.06, .07, .07, .07, .14	0, 0, 1, 2, 2	0, 0, 0, 0, 0	3, 0, 9, 17, 11	0, 0, 0, 0, 0	3, 0, 9, 17, 11	1, 0, .65, .73, .71	8 (0.40)
Northeast Sink Gillnet	05-09	Obs. Data Trip Logbook, Allocated Dealer Data	.04, .07, .05, .04	0, 0, 0, 0, 0	1, 0, 0, 0, 0	0, 0, 0, 0, 0	15, 0, 0, 0, 0	15, 0, 0, 0, 0	.93, 0, 0, 0, 0	3 (0.93)
Mid-Atlantic Gillnet	05-09	Obs. Data, Trip Logbook, Allocated Dealer Data	.03, .04, .04, .03, .03	0, 0, 0, 0, 0	0, 0, 1, 0, 0	0, 0, 0, 0, 0	0, 0, 34, 0, 0	0, 0, 33, 0, 0	0, 0, .73, 0, 0	7 (0.73)
Mid-Atlantic Midwater Trawl - Including Pair Trawl	05-09	Obs. Data Weighout Trip Logbook	.084, .089, .039, .133, .132	0,0,0,0,0	0,0,0,1,0	na	na	na	na	na
TOTAL										18 (0.37)
<sup>a</sup> Observer data (Obs. Data) are used to measure bycatch rates and the data are collected within the Northeast Fisheries Observer Program. The Observer Program collects landings data (Weighout), and total landings are used as a measure of total effort for the coastal gillnet fishery. <sup>b</sup> Estimates can include data pooled across years, so years without observed SI or Mortality may still have an estimated value.										

**Other Mortality**

From 2005 to 2009, 66 Risso's dolphin strandings were recorded along the U.S. Atlantic coast (NMFS unpublished data). Six animals during this time period had indications of human interaction, four of which were fishery interactions. Indications of human interaction are not necessarily the cause of death. In eastern Canada, one Risso's dolphin stranding was reported on Sable Island, Nova Scotia between 1970 and 1998 (Lucas and Hooker 2000).

A Virginia Coastal Small Cetacean Unusual Mortality Event (UME) occurred along the coast of Virginia from 1 May to 31 July 2004, when 66 small cetaceans, including one Risso's dolphin, stranded mostly along the outer (eastern) coast of Virginia's barrier islands.

A Mid-Atlantic Offshore Small Cetacean UME was declared when 33 small cetaceans stranded from Maryland to Georgia between July and September 2004. The species involved are generally found offshore and are not expected to strand along the coast. Three Risso's dolphins were involved in this UME.

Table 3. Risso's dolphin ( <i>Grampus griseus</i> ) reported strandings along the U.S. Atlantic coast, 2005-2009.						
STATE	2005	2006	2007	2008	2009	TOTALS
Maine		1		1	1	3
Massachusetts <sup>a,d</sup>	8	1	3	8	4	24
Rhode Island	1					1
New York	4	1				5
New Jersey	5		2			7
Delaware	1		1			2
Maryland	2	1		1	1	5
Virginia <sup>b</sup>	4	1	1		2	8
North Carolina <sup>c</sup>	2	1		1	3	7
Georgia					1	1
Florida <sup>e</sup>			1		2	3
TOTAL	27	6	8	11	14	66

a. One of the 2009 animals had propeller wounds.  
b. One of the 2005 animals showed signs of fishery interaction. One of the 2009 animals showed signs of human interaction.  
c. One animal in 2006 and 2 in 2009 showed signs of fishery interaction.  
d. 2008 includes 4 animals mass stranded in Massachusetts, 3 of which were released alive.  
e. The 2 animals in 2009 were considered a mass stranding.

Stranding data probably underestimate the extent of fishery-related mortality and serious injury because all of the marine mammals that die or are seriously injured may not wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interaction.

### STATUS OF STOCK

The status of Risso's dolphins relative to OSP in the U.S. Atlantic EEZ is unknown. The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine population trends for this species. The total U.S. fishery mortality and serious injury for this stock is not less than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant and approaching a zero mortality and serious injury rate. The 2005-2009 average annual human-related mortality does not exceed PBR; therefore, this is not a strategic stock.

### REFERENCES CITED

Baird, R. W. and P. J. Stacey 1990. Status of Risso's dolphin, *Grampus griseus*, in Canada. *Can. Field-Nat.* 105: 233-242.

Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.

Buckland, S. T., D. R. Anderson, K. P. Burnham, J. L. Laake, D. L. Borchers and L. Thomas 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press. 432 pp.

CETAP 1982. A characterization of marine mammals and turtles in the mid- and North Atlantic areas of the U.S. outer continental shelf, final report, Cetacean and Turtle Assessment Program, University of Rhode Island.

- Bureau of Land Management, Washington, DC. #AA551-CT8-48. 576 pp.
- Cramer, J. 1994. Large pelagic logbook newsletter - 1993. NOAA Tech. Memo. NMFS-SEFSC-352. 19 pp.
- Fairfield, C. P. and L. P. Garrison 2008. Estimated bycatch of marine mammals and sea turtles in the US Atlantic pelagic longline fleet during 2007. NOAA Tech. Memo. NMFS-SEFSC-572. 62 pp.
- Fairfield Walsh, C. and L. P. Garrison 2006. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2005. NOAA Tech. Memo. NMFS-SEFSC-539. 52 pp.
- Fairfield Walsh, C. and L. P. Garrison 2007. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2006. NOAA Tech. Memo. NMFS-SEFSC-560. 54 pp.
- Garrison, L. P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. NOAA Tech. Memo. NMFS-SEFSC-515. 52 pp.
- Garrison, L. P. 2005. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2004. NOAA Tech. Memo. NMFS-SEFSC-531. 57 pp.
- Garrison, L. P. and P. M. Richards 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. NOAA Tech. Memo. NMFS-SEFSC-527. 57 pp.
- Garrison, L. P., L. Stokes and C. Fairfield 2009. Estimated bycatch of marine mammals and sea turtles in the U.S. Atlantic pelagic longline fleet during 2008. NOAA Tech. Memo. NMFS-SEFSC-591. 63p.
- Garrison, L. P. and L. Stokes. 2010. Estimated bycatch of marine mammals and sea turtles in the U.S. Atlantic pelagic longline fleet during 2009. NOAA Tech. Memo. NMFS-SEFSC-607. 56p.
- Hamazaki, T. 2002. Spatiotemporal prediction models of cetacean habitats in the mid-western North Atlantic Ocean (from Cape Hatteras, No. Carolina, USA to Nova Scotia, Canada). *Mar. Mamm. Sci.* 18(4): 920-939.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. Pages 179-189 *in*: G.W. Garner, S.C. Amstrup, J.L. Laake, *et al.* (eds.). *Marine mammal survey and assessment methods*. Balkema, Rotterdam.
- Johnson, D. R., C. A. Brown and C. Yeung 1999. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1992-1997. NOAA Tech. Memo. NMFS- SEFSC-418. 70 pp.
- Leatherwood, S., D. K. Caldwell and H. E. Winn 1976. Whales, dolphins, and porpoises of the western North Atlantic. A guide to their identification. NOAA Tech. Rep. NMFS Circ. 396. 176 pp.
- Lucas, Z. N. and S. K. Hooker 2000. Cetacean strandings on Sable Island, Nova Scotia, 1970-1998. *Can. Field-Nat.* 114(1): 46-61.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. *Rep. Int. Whal. Comm.* (special issue) 16: 27-50.
- Palka, D. L. 2005. Aerial surveys in the northwest Atlantic: estimation of  $g(0)$ . Proceedings of a Workshop on Estimation of  $g(0)$  in Line-Transsect Surveys of Cetaceans, European Cetacean Society's 18th Annual Conference; Kolmården, Sweden; Mar. 28, 2004. pp12-17.
- Palka, D. L. and P. S. Hammond 2001. Accounting for responsive movement in line transect estimates of abundance. *Can. J. Fish. Aquat. Sci.* 58: 777-787.
- Payne, P. M., L. A. Selzer. and A. R. Knowlton. 1984. Distribution and density of cetaceans, marine turtles and seabirds in the shelf waters of the northeast U.S., June 1980 - Dec. 1983, based on shipboard observations. Contract report NA81FAC00023. National Marine Fisheries Service, Woods Hole. MA. 245 pp.
- Scott, G. P. and C. A. Brown 1997. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1994-1995. Miami Lab. Contribution MIA-96/97-28. National Marine Fisheries Service, Maimi, FL. 14 pp.
- Wade, P. R. and R. P. Angliss 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Waring, G. T., C. P. Fairfield, C. M. Ruhsam and M. Sano 1992. Cetaceans associated with Gulf Stream features off the northeastern USA Shelf. *ICES [Int. Coun. Explor. Sea] C.M.* 1992/N:12.
- Waring, G. T., C. P. Fairfield, C. M. Ruhsam and M. Sano 1993. Sperm whales associated with Gulf Stream features off the northeastern USA shelf. *Fish. Oceanogr.* 2(2): 101-105.
- Waring, G. T., P. Gerrior, P. M. Payne, B. L. Parry and J. R. Nicolas 1990. Incidental take of marine mammals in foreign fishery activities off the northeast United States, 1977-1988. *Fish. Bull.* 88(2): 347-360.
- Wells, R. S., J. B. Allen, S. Hofmann, K. Bassos-Hull, D. A. Fauquier, N. B. Barros, R. E. DeLynn, G. Sutton, V. Socha and M. D. Scott 2008. Consequences of injuries on survival and reproduction of common bottlenose dolphins (*Tursiops truncatus*) along the west coast of Florida. *Mar. Mamm. Sci.* 24: 774-794.
- Wells, R. S., C. A. Manire, L. Byrd, D. R. Smith, J. G. Gannon, D. Fauquier and K. D. Mullin 2009. Movements and dive patterns of a rehabilitated Risso's dolphin, *Grampus griseus*, in the Gulf of Mexico and Atlantic Ocean. *Mar. Mamm. Sci.* 25(2): 420-429.

- Yeung, C. 1999. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998. NOAA Tech. Memo. NMFS-SEFSC-430. 26 pp.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. NOAA Tech. Memo. NMFS-SEFSC-467. 43 pp.
- Yeung, C., S. Epperly and C. A. Brown 2000. Preliminary revised estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet, 1992-1999. PRD Contribution Number 99/00-13. National Marine Fisheries Service, Miami, FL 58 pp.