

Table 198. Selected Respiratory Diseases Among Persons 18 Years of Age and Over by Selected Characteristics: 2009

[In thousands (227,371 represents 227,371,000). Respondents were asked in two separate questions if they had ever been told by a doctor or other health professional that they had emphysema or asthma. Respondents who had been told they had asthma were asked if they still had asthma. Respondents were asked in three separate questions if they had been told by a doctor or other health professional in the past 12 months that they had hay fever, sinusitis, or bronchitis. Based on the National Health Interview Survey, a sample survey of the civilian noninstitutionalized population; see Appendix III]

Selected characteristic	Total persons	Selected respiratory conditions ¹					
		Emphysema	Asthma		Hay fever	Sinusitis	Chronic bronchitis
			Ever	Still			
Total ²	227,371	4,895	29,734	17,456	17,738	29,305	9,908
SEX							
Male	109,844	2,578	12,311	6,058	7,546	10,555	3,189
Female	117,527	2,317	17,423	11,398	10,192	18,750	6,718
AGE							
18 to 44 years	110,337	369	15,743	8,368	7,106	11,342	3,093
45 to 64 years	79,195	2,065	9,706	6,182	7,932	12,792	4,411
65 to 74 years	20,597	1,194	2,466	1,720	1,629	3,028	1,329
75 years and over	17,242	1,268	1,819	1,186	1,072	2,142	1,074
RACE							
Race alone ³	224,290	4,825	29,056	17,061	17,406	28,819	9,750
White	183,739	4,357	23,880	13,990	15,061	24,447	8,386
Black or African American	27,374	360	3,854	2,380	1,551	3,513	1,082
American Indian or Alaska Native	1,856	(B)	275	⁴ 1,177	⁴ 104	155	⁴ 78
Asian	10,763	⁴ 59	968	490	670	694	184
Native Hawaiian or other Pacific Islander	558	(B)	⁴ 25	(B)	(B)	(B)	(B)
Two or more races ⁵	3,082	⁴ 70	678	394	332	486	158
HISPANIC ORIGIN AND RACE ⁶							
Hispanic or Latino	31,312	200	3,225	1,721	1,648	2,584	820
Mexican or Mexican American	19,687	107	1,557	826	980	1,561	362
Not Hispanic or Latino	196,059	4,696	26,509	15,735	16,090	26,721	9,088
White, single race	155,185	4,173	21,035	12,484	13,532	22,123	7,647
Black or African American, single race	26,213	350	3,675	2,281	1,497	3,393	1,046

B Figure too small to meet statistical standards for reliability of a derived figure. ¹ A person may be represented in more than one column. ² Total includes other races not shown separately. ³ Refers to persons who indicated only a single race group. ⁴ Figure does not meet standard of reliability or precision. ⁵ Refers to all persons who indicated more than one race group. ⁶ Persons of Hispanic or Latino origin may be any race or combination of races.

Source: National Center for Health Statistics, *Summary Health Statistics for the U.S. Population: National Health Interview Survey, 2009*, Vital and Health Statistics, Series 10, Number 249, 2010. See also <http://www.cdc.gov/nchs/data/series/sr_10/sr10_249.pdf>.

Table 199. Persons 18 Years of Age and Over With Migraines and Pains in the Neck, Lower Back, Face, or Jaw by Selected Characteristics: 2009

[In thousands (227,371 represents 227,371,000). Based on National Health Interview Survey, a sample survey of the civilian noninstitutionalized population, Appendix III]

Selected characteristic	Total persons	Migraine and pain ¹			
		Migraine or severe headache ²	Pain in neck ³	Pain in lower back ³	Pain in face or jaw ³
Total ⁴	227,371	35,973	34,954	64,810	11,501
SEX					
Male	109,844	11,098	14,095	28,842	3,654
Female	117,527	24,875	20,859	35,968	7,847
AGE					
18 to 44 years	110,337	21,706	14,306	26,973	5,752
45 to 64 years	79,195	11,893	15,147	25,795	4,400
65 to 74 years	20,597	1,415	3,136	6,198	799
75 years and over	17,242	959	2,365	5,843	550
RACE					
Race alone ⁵	224,290	35,295	34,355	63,689	11,213
White	183,739	29,118	29,430	53,817	9,599
Black or African American	27,374	4,811	3,569	7,286	1,115
American Indian or Alaska Native	1,856	429	363	566	171
Asian	10,763	902	935	1,879	303
Native Hawaiian or other Pacific Islander	558	⁶ 33	⁶ 57	⁶ 141	(B)
Two or more races ⁷	3,082	678	598	1,121	288
HISPANIC ORIGIN AND RACE ⁸					
Hispanic or Latino	31,312	5,362	4,341	7,905	1,345
Mexican or Mexican American	19,687	3,297	2,446	4,319	818
Not Hispanic or Latino	196,059	30,611	30,613	56,905	10,156
White, single race	155,185	24,379	25,497	46,665	8,423
Black or African American, single race	26,213	4,546	3,426	6,982	1,065

B Figure to small to meet statistical standards for reliability of a derived figure. ¹ A person may be represented in more than one column. ² Respondents were asked, "During the past 3 months, did you have a severe headache or migraine?" Respondents were instructed to report pain that had lasted a whole day or more and, conversely, not to report fleeting or minor aches or pains. ³ Respondents were asked, "During the past 3 months, did you have a neck pain; or low back pain; or facial ache or pain in the jaw muscles or the joint in front of the ear?" Respondents were instructed to report pain that had lasted a whole day or more and, conversely, not to report fleeting or minor aches or pains. ⁴ Total includes other races not shown separately. ⁵ Refers to persons who indicated only a single race group. ⁶ Figure does not meet standard of reliability or precision. ⁷ Refers to all persons who indicated more than one race group. ⁸ Persons of Hispanic or Latino origin may be any race or combination of races.

Source: National Center for Health Statistics, *Summary Health Statistics for the U.S. Adults: National Health Interview Survey, 2009*, Vital and Health Statistics, Series 10, Number 249, 2010. See also <http://www.cdc.gov/nchs/data/series/sr_10/sr10_249.pdf>.