

Medication Guide

FORTEO® (for-TAY-o) teriparatide (rDNA origin) injection

Read this Medication Guide before you start taking FORTEO and each time you get a refill. There may be new information. Also, read the User Manual that comes with the FORTEO delivery device (pen) for information on how to use the device to inject your medicine the right way. This Medication Guide does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is the most important information I should know about FORTEO?

- During the drug testing process, the medicine in FORTEO caused some rats to develop a bone cancer called osteosarcoma. In people, osteosarcoma is a serious but rare cancer. Osteosarcoma has been reported rarely in people who took FORTEO. It is not known if people who take FORTEO have a higher chance of getting osteosarcoma.
- You should not take FORTEO for more than 2 years over your lifetime.
- There is a voluntary Patient Registry for people who take FORTEO. The purpose of the registry is to collect information about the possible risk of osteosarcoma in people who take FORTEO. For information about how to sign up for this patient registry, call 1-866-382-6813 or go to www.forteoregistry.rti.org.

What is FORTEO?

- FORTEO is a prescription medicine that is like a hormone made by the body called parathyroid hormone or PTH. FORTEO may help to form new bone, increase bone mineral density and bone strength.
- FORTEO can lessen the number of fractures of the spine and other bones in postmenopausal women with osteoporosis.
- The effect on fractures has not been studied in men.
- FORTEO is used in both men and postmenopausal women with osteoporosis who are at high risk for having fractures. FORTEO can be used by people who have had a fracture related to osteoporosis, or who have several risk factors for fracture, or who can not use other osteoporosis treatments.
- FORTEO is used in both men and women with osteoporosis due to use of glucocorticoid medicines, such as prednisone, for several months, who are at high risk for having broken bones (fractures). These include men and women with either a history of broken bones, who have several risk factors for fracture, or who can not use other osteoporosis treatments.

It is not known if FORTEO is safe and effective in children.

FORTEO should not be used in children and young adults whose bones are still growing.

Who should not use FORTEO?

Do not use FORTEO if you:

- are allergic to any of the ingredients in FORTEO. See the end of this Medication Guide for a complete list of the ingredients in FORTEO.

What should I tell my healthcare provider before taking FORTEO?

Before you take FORTEO, tell your healthcare provider if you:

- have the condition listed in the section “Who should not use FORTEO?”
- have Paget’s disease or other bone disease
- have cancer in your bones
- have trouble injecting yourself and do not have someone who can help you
- are a child or young adult whose bones are still growing
- have or have had kidney stones
- have had radiation therapy
- have or had too much calcium in your blood

- 52
- have any other medical conditions
 - are pregnant or thinking about becoming pregnant. It is not known if FORTEO will harm your unborn baby.
 - are breast-feeding or plan to breast-feed. It is not known if FORTEO passes into your breast milk. You and your doctor should decide if you will take FORTEO or breast feed. You should not do both.

57

58 **Tell your healthcare provider about all the medicines you take** including prescription and non-prescription
59 medicines, vitamins, and herbal supplements. Your healthcare provider needs this information to help keep you from
60 taking FORTEO with other medicines that may harm you.

- Especially tell your doctor if you take medicines that contain digoxin (Digoxin, Lanoxicaps, Lanoxin).

62 **How should I use FORTEO?**

- Inject FORTEO one time each day in your thigh or abdomen (lower stomach area). Talk to a healthcare provider about how to rotate injection sites.
- Before you try to inject FORTEO yourself, a healthcare provider should teach you how to use the FORTEO delivery device to give your injection the right way.
- Read the detailed User Manual at the end of this Medication Guide.
- You can take FORTEO with or without food or drink.
- The FORTEO delivery device has enough medicine for 28 days. It is set to give a 20 microgram dose of medicine each day. Do not inject all the medicine in the FORTEO delivery device at any one time.
- Do not transfer the medicine from the FORTEO delivery device to a syringe. This can result in taking the wrong dose of FORTEO. If you do not have pen needles to use with your FORTEO delivery device, talk with your healthcare provider.
- FORTEO should look clear and colorless. Do not use FORTEO if it has particles in it, or if it is cloudy or colored.
- Inject FORTEO right away after you take the delivery device out of the refrigerator.
- After each use, safely remove the needle, recap the delivery device, and put it back in the refrigerator right away.
- You can take FORTEO at any time of the day. To help you remember to take FORTEO, take it at about the same time each day.
- If you forget or can not take FORTEO at your usual time, take it as soon as you can on that day. Do not take more than one injection in the same day.
- If you take more FORTEO than prescribed, call your healthcare provider. If you take too much FORTEO, you may have nausea, vomiting, weakness or dizziness.

86 Follow your healthcare provider's instructions about other ways you can help your osteoporosis, such as exercise,
87 diet, and reducing or stopping your use of tobacco and alcohol. If your healthcare provider recommends calcium and
88 vitamin D supplements, you can take them at the same time you take FORTEO.

89 **What are the possible side effects of FORTEO?**

90

91 FORTEO can cause serious side effects including:

- See "What is the most important information I should know about FORTEO?"
- **Decrease in blood pressure when you change positions.** Some people feel dizzy, get a fast heartbeat, or feel faint right after the first few doses. This usually happens within 4 hours of taking FORTEO and goes away within a few hours. For the first few doses, take your injections of FORTEO in a place where you can sit or lie down right away if you get these symptoms. If your symptoms get worse or do not go away, stop taking FORTEO and call your healthcare provider.
- **Increased calcium in your blood.** Tell your healthcare provider if you have nausea, vomiting, constipation, low energy, or muscle weakness. These may be signs there is too much calcium in your blood.

100

101 Common side effects of FORTEO include:

- nausea
- joint aches
- pain

105

106 Your healthcare provider may take samples of blood and urine during treatment to check your response to FORTEO.
107 Also, your healthcare provider may ask you to have follow-up tests of bone mineral density.

108
109 Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

110
111 These are not all the possible side effects of FORTEO. For more information, ask your doctor or pharmacist.

112
113 Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

114 **How should I store FORTEO?**

- 115 • Keep your FORTEO delivery device in the refrigerator between 36° to 46°F (2° to 8°C).
- 116 • Do not freeze the FORTEO delivery device. Do not use FORTEO if it has been frozen.
- 117 • Do not use FORTEO after the expiration date printed on the delivery device and packaging.
- 118 • Throw away the FORTEO delivery device after 28 days even if it has medicine in it (see the User
119 Manual).

120 **Keep FORTEO and all medicines out of the reach of children.**

121 **General information about FORTEO**

122 Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use
123 FORTEO for a condition for which it was not prescribed. Do not give FORTEO to other people, even if they have
124 the same condition you have.

125
126 This Medication Guide summarizes the most important information about FORTEO. If you would like more
127 information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information
128 about FORTEO that is written for healthcare professionals. For more information, go to www.FORTEO.com or call
129 Lilly at 1-866-436-7836.

130 **What are the ingredients in FORTEO?**

131
132 Active ingredient: teriparatide

133
134 Inactive ingredients: glacial acetic acid, sodium acetate (anhydrous), mannitol, metacresol, and water for injection.
135 In addition, hydrochloric acid solution 10% and/or sodium hydroxide solution 10% may have been added to adjust
136 the product to pH 4.

137 **What is Osteoporosis?**

138 Osteoporosis is a disease in which the bones become thin and weak, increasing the chance of having a broken bone.
139 Osteoporosis usually causes no symptoms until a fracture happens. The most common fractures are in the spine
140 (backbone). They can shorten height, even without causing pain. Over time, the spine can become curved or
141 deformed and the body bent over. Fractures from osteoporosis can also happen in almost any bone in the body, for
142 example, the wrist, rib, or hip. Once you have had a fracture, the chance for more fractures greatly increases.

143
144 The following risk factors increase your chance of getting fractures from osteoporosis:

- 145 • past broken bones from osteoporosis.
- 146 • very low bone mineral density (BMD).
- 147 • frequent falls.
- 148 • limited movement, such as using a wheelchair.
- 149 • medical conditions likely to cause bone loss, such as some kinds of arthritis.
- 150 • taking steroid medicines called glucocorticoids, such as prednisone.
- 151 • other medicines that may cause bone loss, for example: seizure medicines (such as phenytoin), blood
152 thinners (such as heparin), high doses of vitamin A.

153
154 *This Medication Guide has been approved by the U.S. Food and Drug Administration.*

155 Medication Guide revised Month XX, YYYY

156

157 Manufactured by Lilly France - F-67640 Fegersheim, France
158 for Eli Lilly and Company - Indianapolis, IN 46285, USA
159
160 Copyright © 2002, 200X, Eli Lilly and Company. All rights reserved.
161
162 PA 9411 FSAMP