

HIV and AIDS

Use this guide to help you talk to your doctor, pharmacist, or nurse about your HIV medicines. The guide lists all of the FDA-approved products now available to treat this condition. You will also find some general information to help you use your medicines wisely.



HIV stands for **H**uman **I**mmunodeficiency **V**irus. HIV is the virus that causes AIDS. A person with HIV is called HIV positive (HIV+).

HIV makes it hard for your body to fight off sickness. There are cells in your blood called “CD4 cells” or “T cells.” These cells help protect your body from disease. HIV kills these cells. A person with HIV does not have as many of these cells as a healthy person.

There is hope.

There are drugs that can treat HIV and help people live longer. These medicines help to keep the virus from building up in your body. These drugs do not stop you from spreading HIV. You can still give the disease to someone else.

People with HIV may need to take 3 or more different medicines every day. You and your doctor will decide which ones are right for you. It is important that you take your HIV medicines everyday. Do not stop taking your medicines without talking to your doctor. Over time, you can get very sick if you do not take your medicines.

Use this guide to help you talk to your doctor, pharmacist, or nurse about the HIV medicines you are taking. This guide provides some basic facts about the HIV medicines that have been approved by the FDA.

There are many things you can do to help improve your health.

- Work closely with your doctor to monitor your health.
- Take your medicine.
- Eat a healthy, balanced diet.
- Quit smoking and using illegal drugs.
- Get regular exercise.

Women and HIV

Women from all backgrounds and cultures can get HIV. However, increasing numbers of African American and Latino women have HIV.

Learn the facts about women and HIV. Educate yourself to help you live longer. Educate yourself so that you can teach other women how to prevent HIV and AIDS.

Did you know?

- Most women get HIV from having unprotected sex with men.
- A woman can pass HIV to her baby during pregnancy, labor, or delivery.
- A woman can also pass HIV to her baby during breastfeeding.
- A pregnant woman can take medicine to lower the chance of giving her baby HIV. Pregnant women should talk to their doctor about the pros and cons of taking medicines for HIV. Some anti-HIV medicines should not be taken during pregnancy because they can cause birth defects.
- Babies born to women with HIV may need to take anti-HIV medicines after birth to lower the chance that they will get HIV. Talk to your doctor about the best way to treat your baby.
- Women who are HIV positive should get regular Pap smears and exams to test them for other health problems.

Women with HIV are more likely to have:

- o Vaginal yeast infections
- o Other infections such as bacterial vaginosis
- o Pelvic Inflammatory Disease (PID)
- o Sexually Transmitted Infections like gonorrhea or HPV
- o Changes in the cervix that may lead to cervical cancer

Anti-HIV Medicines

The main drug treatment for people with HIV is **Highly Active Antiretroviral Therapy** (also called **HAART**). HAART drugs help to slow the growth of HIV in your body.

HAART is made up of different kinds of medicines:

1. **Nucleoside Reverse Transcriptase Inhibitors** (NRTIs)
2. **Nonnucleoside Reverse Transcriptase Inhibitors** (NNRTIs)
3. **Protease Inhibitors**
4. **Fusion Inhibitors**
5. **Integrase Inhibitors**
6. **Entry Inhibitors**
7. **Combination Drugs**

The seven different groups of HAART drugs are listed on the next few pages. The brand names and generic names are listed for each drug.

Risks and Side Effects

The drugs used to treat HIV can sometimes cause side effects. Side effects may be different depending on the person and the kind of medicine. Some people have no side effects. Others can have very bad side effects.

Tell your doctor about any side effects you are having. Do not stop taking your medicine without talking to your doctor. Your doctor may tell you tips to help you cope with the side effects. The doctor may also decide to have you take different drugs.

This guide does not give the specific side effects or warnings for each HAART drug. **Check the drug label and ask your doctor for the side effects and warnings for the HIV medicines you are taking.**

Information about specific drugs can also be found on the FDA website at: <http://www.fda.gov/cder/drug/DrugSafety/DrugIndex.htm>

Nucleoside Reverse Transcriptase Inhibitors (NRTIs)

Brand Name	Other Names
Combivir	Lamivudine and Zidovudine
Emtriva	Emtricitabine FTC
Epivir	Lamivudine 3TC
Epzicom	Abacavir and Lamivudine
Hivid	Zalcitabine Dideoxycytidine, ddC
Retrovir	Zidovudine, AZT, Azidothymidine, ZDV
Trizivir	Abacavir, Zidovudine, and Lamivudine
Truvada	Tenofovir Disoproxil and Emtricitabine
Videx	Didanosine, ddl, Dideoxyinosine
Videx EC	Enteric Coated Didanosine
Viread	Tenofovir Disoproxil Fumarate, TDF
Zerit	Stavudine d4T
Ziagen	Abacavir Sulfate, ABC

For up-to-date information about the risks and side effects for each drug
Check <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/>

Nucleoside Reverse Transcriptase Inhibitors: What You Should Know

This guide does not give the specific side effects or warnings for each drug. **Check the drug label and ask your doctor for the side effects and warnings for the HIV medicines you are taking.**

Warnings

- Women should not breastfeed while taking these medicines.
- These medicines may cause lactic acidosis (too much acid in the blood).
- These medicines may cause serious liver or pancreas problems.
- People with liver problems including hepatitis and people with kidney problems should talk to their doctor before taking these medicines.
- In some cases, people taking HIV medicines notice changes in body fat (like extra fat in the neck or upper back or loss of fat in the face or arms).

Warning Signs

Call your doctor right away if you have any of these signs:

- Upset Stomach
- Vomiting
- Feeling Very Weak or Tired
- Problems Breathing
- Weakness in Arms and Legs
- Tingling, Numbness, or Pain in Feet or Hands
- Jaundice (Skin or Eyes Look Yellow)
- Pain in the Upper Stomach Area

Common Side Effects

- Trouble Sleeping
- Headache
- Feeling Tired
- Upset Stomach
- Mild Nausea
- Vomiting
- Diarrhea
- Dizziness
- Do Not Feel Like Eating

Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs)

Brand Name	Other Names
Edurant	Rilpivirine
Intelence	Etravirine
Rescriptor	Delavirdine DLV
Sustiva	Efavirenz EFV
Viramune	Nevirapine NVP
Viramune XR (extended release)	Nevirapine NVP

For up-to-date information about the risks and side effects for each drug
Check <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/>

Non-Nucleoside Reverse Transcriptase Inhibitors: What You Should Know

This guide does not give the specific side effects or warnings for each drug. **Check the drug label and ask your doctor for the side effects and warnings for the HIV medicines you are taking.**

Warnings

- Women should not breastfeed while taking these medicines.
- These medicines may cause serious liver problems or severe skin rashes.
- People with liver problems including hepatitis and people with kidney problems should talk to their doctor before taking these medicines.
- Women with CD4 counts higher than 250 should talk to their doctor about the risks of taking Viramune (Nevirapine).
- In some cases, people taking HIV medicines notice changes in body fat (like extra fat in the neck or upper back or loss of fat in the face or arms).

Warning Signs

Call your doctor right away if you have any of these signs:

- Flu-Like Symptoms
- Feeling Tired
- Do Not Feel Like Eating
- Dark Urine (Looks Like Tea)
- Pale Stools
- Upset Stomach/Nausea
- Jaundice (Skin or Eyes Look Yellow)
- Pain, Aches, or Sensitivity to Touch on Right Side below Your Ribs

Also call your doctor right away if you have a severe rash along with blisters, swelling, pink eye, fever, muscle/joint pain, or mouth sores.

Common Side Effects

- Skin Rash
- Upset Stomach
- Dizziness
- Problems Concentrating
- Feeling Tired
- Vomiting
- Trouble Sleeping
- Headache
- Diarrhea
- Strange Dreams

Protease Inhibitors

Brand Name	Other Names
Agenerase	Amprenavir APV
Aptivus	Tipranavir TPV
Crixivan	Indinavir IDV
Fortovase	Saquinavir (No Longer Marketed)
Invirase	Saquinavir Mesylate SQV
Kaletra	Lopinavir and Ritonavir LPV/RTV
Lexiva	Fosamprenavir Calcium FOS-APV
Norvir	Ritonavir RTV
Prezista	Darunavir
Reyataz	Atazanavir Sulfate ATV
Viracept	Nelfinavir Mesylate NFV

For up-to-date information about the risks and side effects for each drug
Check <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/>

Protease Inhibitors: What You Should Know

This guide does not give the specific side effects or warnings for each drug. **Check the drug label and ask your doctor for the side effects and warnings for the HIV medicines you are taking.**

Warnings

- Women taking birth control pills need to use another birth control method.
- Tell your doctor about all medicines that you are taking. Protease Inhibitors may cause serious health problems or death if mixed with other medicines.
- These medicines may cause serious liver problems.
- These medicines may cause increased bleeding in people with hemophilia.
- These medicines may make diabetes worse or cause people to get diabetes.
- Some people taking protease inhibitors notice large increases in their cholesterol.
- In some cases, people taking HIV medicines notice changes in body fat (like extra fat in the neck or upper back or loss of fat in the face or arms).
- Women should not breastfeed while taking these medicines.

Warning Signs

Call your doctor if you have any of these signs:

- Serious Skin Rash
- Feeling Very Weak or Tired
- Unusual Muscle Pain
- Trouble Breathing
- Stomach Pain with Nausea and Vomiting

Common Side Effects

- Rash
- Diarrhea
- Nausea (Upset Stomach)
- Vomiting
- Feeling Tired
- Headache

Fusion Inhibitors

Brand Name	Other Names
<p data-bbox="301 261 417 289">Fuzeon</p> <p data-bbox="176 349 542 376">This medicine is a shot.</p>	<p data-bbox="895 261 1069 289">Enfuvirtide</p> <p data-bbox="946 306 1018 334">T-20</p>

**For up-to-date information about the risks and side effects for each drug
Check <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/>**

Fusion Inhibitors: What You Should Know

This guide does not give the specific side effects or warnings for each drug. **Check the drug label and ask your doctor for the side effects and warnings for the HIV medicines you are taking.**

Warnings

- People taking Fuzeon (Enfuvirtide) with other HIV medicines may be more likely to get pneumonia. Tell your doctor if you have a cough, fever, or trouble breathing.
- Women should not breastfeed while taking Fuzeon (Enfuvirtide).
- In some cases, people taking HIV medicines notice changes in body fat (like extra fat in the neck or upper back or loss of fat in the face or arms).

Warning Signs

Fuzeon may cause serious allergic reactions. Call your doctor right away if you have any of these signs:

- Trouble Breathing
- Fever with Vomiting and a Skin Rash
- Blood in Urine
- Swelling of the Feet

Common Side Effects

- Itching, Redness, Pain, Bumps, or Swelling Where the Shot is Given
- Pain and Numbness in Feet or Legs
- Problems Sleeping
- Depression
- Weakness or Loss of Strength
- Muscle Pain
- Constipation

Integrase Inhibitors

Brand Name	Other Names
Isentress	Raltegravir

For up-to-date information about the risks and side effects for each drug
Check <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/>

Integrase Inhibitors: What You Should Know

This guide does not give all of the specific side effects or warnings for each drug. **Check the drug label and ask your doctor for the side effects and warnings for the HIV medicines you are taking.**

Warnings

- Women who are pregnant or plan to become pregnant should talk to their doctor before taking Isentress (Raltegravir).
- Women should not breastfeed while taking Isentress (Raltegravir).
- Tell your doctor if you have any allergies before you start taking Isentress (Raltegravir).
- In some cases, people taking HIV medicines notice changes in body fat (like extra fat in the neck or upper back or loss of fat in the face or arms).

Warning Signs

You should also see your doctor right away if you have any of these signs:

- Unexplained Muscle Pain, Tenderness or Weakness
- Signs of an Infection

Common Side Effects

- Diarrhea
- Nausea
- Headache

Entry Inhibitors

Brand Name	Other Names
Selzentry	Maraviroc

For up-to-date information about the risks and side effects for each drug
Check <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/>

Entry Inhibitors: What You Should Know

This guide does not give all of the specific side effects or warnings for each drug. **Check the drug label and ask your doctor for the side effects and warnings for the HIV medicines you are taking.**

- People with liver problems including Hepatitis B or C should talk to their doctor before taking Selzentry (Maraviroc).
- People with kidney problems or heart problems should talk to their doctor before taking Selzentry (Maraviroc).
- Tell your doctor if you are taking high blood pressure medicines or if you have low blood pressure.
- Do not drive a car or use heavy machinery if you feel dizzy while taking Selzentry (Maraviroc).
- Tell your doctor if you have any allergies before you start taking Selzentry (Maraviroc).
- Women should not breastfeed while taking Selzentry (Maraviroc).
- In some cases, people taking HIV medicines notice changes in body fat (like extra fat in the neck or upper back or loss of fat in the face or arms).

Warning Signs

Selzentry may cause serious liver problems. Call your doctor right away if you have any of these signs:

- Itchy Rash on Your Body
- Skin or Eyes Look Yellow
- Dark (Tea-Colored) Urine
- Vomiting and Stomach Pain

You should also see your doctor right away if you have any of these signs:

- Nausea
- Flu-Like Symptoms
- Fatigue

Serious Side Effects

- Possible Chance of Infection or Cancer

Common Side Effects

- Cough
- Fever
- Colds
- Dizziness
- Rash
- Muscle and Joint Pain
- Stomach Pain

Multi-Class Combination Drugs

Brand Name	Other Names
Atripla	Efavirenz, Emtricitabine, and Tenofovir Disoproxil Fumarate
Complera	Emtricitabine, Rilpivirine, and Tenofovir Disoproxil Fumarate

For up-to-date information about the risks and side effects for each drug
Check <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/>

Combination Drugs: What You Should Know

This guide does not give the specific side effects or warnings for each drug. **Check the drug label and ask your doctor for the side effects and warnings for the HIV medicines you are taking.**

Warnings

- Atripla may cause lactic acidosis (too much acid in the blood).
- Atripla may cause serious liver problems.
- People who have kidney or liver problems, including Hepatitis B, should talk to their doctor before using this medicine.
- People who have ever had seizures and people taking medicines for seizures should talk to their doctor before using this medicine.
- People who have ever had mental illness and people with bone problems should talk to their doctor before using this medicine.
- Women should not get pregnant or breastfeed while taking Atripla.
- Women taking birth control pills need to use another birth control method.
- In some cases, people taking HIV medicines notice changes in body fat (like extra fat in the neck or upper back or loss of fat in the face or arms).

Warning Signs

Call your doctor right away if you have any of these signs:

- Feeling Very Weak or Tired
- Unusual Muscle Pain
- Trouble Breathing
- Stomach Pain with Nausea and Vomiting
- Feeling Dizzy or Lightheaded
- Dark Urine
- Light Colored Stools
- Loss of Appetite for Several Days
- Upset Stomach (Nausea)
- Stomach Pain
- Jaundice (Skin or Eyes Look Yellow)

Common Side Effects

- Dizziness
- Headache
- Trouble Sleeping
- Feeling Drowsy
- Problems Concentrating
- Unusual Dreams

Important Questions to Ask Your Doctor

- What drugs am I taking?
-
-

- What are the side effects of taking these drugs?
-
-

- What do I do if I start having bad side effects?
-
-

- What other prescription drugs should I avoid while taking my HIV medicines?
-
-

- What herbs (like St. John's Wort) or over-the-counter medicines should I avoid?
-
-

- When should I take each drug?
-
-

- Should I take my medicines with food?
-
-

- How should I keep my medicines when I go out of town?
-
-

- How long can I stay on this regimen?
-
-

**Ask your doctor to tell you what you should know about your HIV medicines.
Write down the important facts in the space below.**

My Regimen

It is important that you take your HIV medicines exactly as your doctor tells you. **Do not skip a pill.** The medicines may not work correctly if you skip a pill.



Here are some tips to help you remember when to take your HIV medicines.

- Use a schedule or planner.
- Set the alarm on your watch or clock.
- Find a friend to remind you.
- Use a pillbox to help you organize your pills.

My Medicine Schedule

Time	Drug Name	Dose (How many pills)	Notes
6:30	XXX (example)	1 Pill	Take with food



FDA

Office of
Women's
Health

www.fda.gov/womens



This document was developed by the FDA Office of Women's Health (OWH).

To Learn More:

FDA Office of Special Health Issues

FDA HIV/AIDS Info

<http://www.fda.gov/oashi/aids/virals.html>

AIDSinfo

US Department of Health
and Human Services

<http://www.aidsinfo.nih.gov>

Phone: 1 800 448 0440

TTY/TTD: 888 480 3739

US Centers for Disease Control and Prevention (CDC)

Department of Health and Human Services

<http://www.cdc.gov/hiv/>

Phone: 1 800 232 4636

TTY/TTD: 1 888 232 6348

U.S. Government HIV/AIDS Website

<http://aids.gov/>

This guide should not be used in place of talking to your doctor or reading the label on your medicine bottle. The drug and risk information in this guide may change. Check <http://www.accessdata.fda.gov/scripts/cder/drugsatfda/> for the latest facts on each product listed in this guide.