

## References

- Franken RJ, Hik DS (2004) Influence of habitat quality, patch size, and connectivity on colonization and extinction dynamics of collared pikas (*Ochotona collaris*). *Journal of Animal Ecology*, **73**, 889–896.
- Hamilton MB, Pincus EL, Di Fiore A, Fleischer RC (1999) Universal linker and ligation procedures for construction of genomic DNA libraries enriched for microsatellites. *BioTechniques*, **27**, 500–507.
- Morrison SF, Hik DS (2007) Demographic analysis of a declining pika *Ochotona collaris* population: linking survival to broad-scale climate patterns via spring snowmelt patterns. *Journal of Animal Ecology*, **76**, 899–907.
- Park SDE (2001) *Trypanotolerance in West African cattle and the population genetic effects of selection*. PhD Thesis. University of Dublin.
- Peacock MM, Kirchoff VS, Merideth SJ (2002) Identification and characterization of nine polymorphic microsatellite loci in the North American pika, *Ochotona princeps*. *Molecular Ecology Notes*, **2**, 360–362.
- Rice WR (1989) Analyzing tables of statistical tests. *Evolution*, **43**, 223–225.
- Rousset F (2008) genepop'007: a complete re-implementation of the genepop software for Windows and Linux. *Molecular Ecology Resources*, **8**, 103–106.
- Rozen S, Skaletsky HJ (2000) Primer 3 on the WWW for general users and for biologist programmers. In: *Bioinformatics Methods and Protocols: Methods in Molecular Biology* (eds Krawetz S, Misener S), pp. 365–386. Humana Press, Totowa, New Jersey.
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: A Laboratory Manual*, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) micro-checker: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, **4**, 535–538.

doi: 10.1111/j.1755-0998.2008.02397.x

© 2009 The Authors

Journal compilation © 2009 Blackwell Publishing Ltd

# Twelve microsatellite loci for lake trout (*Salvelinus namaycush*)

MARYCLARE F. ROLLINS,\* NINH V. VU,\* INGRID B. SPIES† and STEVEN T. KALINOWSKI\*

\*Department of Ecology, 310 Lewis Hall, Montana State University, Bozeman, MT 59717, USA, †NOAA Fisheries, Alaska Fisheries Science Center, 7600 Sand Point Way N.E., Seattle, WA 98115, USA

## Abstract

We describe 12 microsatellite loci isolated from lake trout (*Salvelinus namaycush*). The number of alleles at these loci ranged from two to 11 with an average of 5.3 alleles per locus. The expected heterozygosity ranged from 0.29 to 0.76, with an average of 0.68. Accidental (or illegal) introductions of lake trout into watersheds are decimating native trout populations in the northern Rocky Mountains, and these loci will be useful for identifying the source of these introductions and for estimating the number of founding individuals.

*Keywords:* Lake trout, microsatellite, primer, *Salvelinus namaycush*

Received 13 May 2008; revision accepted 7 June 2008

## Introduction

Lake trout (*Salvelinus namaycush*) are large, long-lived fish predators native to deep lakes of Canada and northern parts of the USA, including the Great Lakes (Benke 2002). During the late 19th and early 20th century, lake trout were

extensively introduced into lakes outside their native ranges (Crossman 1995), and the species is currently expanding its range through dispersal into adjacent waters. Illegal (or accidental) introductions are also common. These introductions and population expansions have decimated native trout species, including populations of cutthroat trout in Yellowstone National Park (Ruzycki *et al.* 2003; Koel *et al.* 2005) and bull trout in Glacier National Park (Fredenberg 2002). Recently, lake trout have been discovered in Swan

Correspondence: Steven Kalinowski, Fax: (406) 994-3190;

E-mail: skalinowski@montana.edu

**Table 1** Summary information for 12 microsatellite loci in lake trout genotyped in 96 individuals: locus names, primer sequences, repeat motifs, fragment size, annealing temperature ( $T_a$ ), number of alleles ( $N_a$ ), observed heterozygosity ( $H_o$ ), expected heterozygosity ( $H_e$ ), Hardy–Weinberg  $P$  value, and GenBank Accession number

Locus	Primer sequence (5'–3')	Repeat motifs	Fragment size (bp)	$T_a$ (°C)	$N_a$	$H_o/H_e$	$P$	GenBank Accession no.
SnaMSU01	F: 6FAM-TCACACACCCATTTCGTTTCAT R: AGCATGGGATAACCACAACC	(GACA) <sub>28</sub>	246–282	57	6	0.68/0.75	0.42	EU331427
SnaMSU02	F: VIC-GCCCTTCATTGAGGAACAGA R: CTCACACACACGCACAACAA	(GTCT) <sub>2</sub> , GTCAAT, (GT) <sub>5</sub> , (GTCT) <sub>11</sub> , GT, (CCGT) <sub>15</sub>	221–257	57	4	0.67/0.75	0.63	EU331428
SnaMSU03	F: NED-TGGGCAAATTTATTGAAGACAAA R: CAGTATACGTCTCTGCCTGTCTG	(GACA) <sub>3</sub> , TATACATTCCTCT (GACA) <sub>22</sub>	197–256	57	4	0.77/0.75	0.66	EU331429
SnaMSU05	F: VIC-TTGAGGGCAAATTCACAATG R: GAGGCAGACAGTGGTAGCTTG	(GT) <sub>23</sub> , (GTCT) <sub>8</sub>	172–204	66	5	0.84/0.75	0.15	EU331431
SnaMSU06	F: 6FAM-GCTGGTGAGGGAGAGATGAC R: CAGCCATGAGAATGGGATTT	(CA) <sub>12</sub> , (GACA) <sub>22</sub>	263–292	60	7	0.88/0.74	0.06	EU331433
SnaMSU07	F: PET-TGCCTATTCTATTCAATACAGG R: CCCCCACAGTCTGAATCAAT	(GACA) <sub>33</sub> , (GA) <sub>34</sub> , (GACA) <sub>20</sub>	351–374	60	2	0.80/0.75	0.20	EU331435
SnaMSU08	F: NED-AGAGCAGTTCGATTGCGATGAC R: ACTGCCCTCACTGATGGTG	(GTATGTCT) <sub>10</sub> , (GTAT) <sub>5</sub>	157–177	66	11	0.29/0.29	1.00	EU331432
SnaMSU09	F: NED-TGATCAGAGATGGCAGTTTCA R: CATTATCCTGGCAACATGGA	(GACA) <sub>25</sub>	251–329	66	7	0.83/0.75	0.10	EU331439
SnaMSU10	F: NED-GCACCTCACCACTCACCTTT R: TTATACAGCAGGGCTGAGCA	(GTCT) <sub>36</sub>	196–243	63	4	0.51/0.60	0.19	EU331434
SnaMSU11	F: 6FAM-TGATGATGGAAAGGCAGAGG R: CCATTTGGGATGCACATACA	(GACA) <sub>36</sub>	226–251	63	4	0.67/0.66	1.00	EU331436
SnaMSU12	F: VIC-ATTTTCCACATGCTGCGTCT R: TGAAATAGCTTGGAGCAGTAGC	(GTCT) <sub>27</sub>	198–216	63	3	0.68/0.60	0.12	EU331437
SnaMSU13	F: PET-AGTTTCCAAGGCAGCACTGT R: TGCTACACAGCAAATGTGTCA	(GACA) <sub>36</sub>	237–256	63	6	0.80/0.76	0.65	EU331438

Lake, Montana, and the population has been growing rapidly. Neither the source of this introduction nor the number of founders is known. Microsatellite genetic variation could help answer these questions, but no microsatellite loci have been specifically developed for lake trout, and loci identified in other salmonids show modest levels of genetic variation (N. Vu, unpublished). In this study, we describe 12 microsatellite loci isolated from lake trout in Montana.

We used a variation of the method of Hamilton *et al.* (1999) to develop a lake trout microsatellite library in which the restriction of genomic DNA and the ligation of a universal DNA linker were combined into one reaction (DIG/LIG). The DIG/LIG reaction consisted of 2.0  $\mu$ M double-stranded SNX linker, 6  $\mu$ g Bovine Serum Albumin, 20 U restriction enzyme, 40 U *Xmn*I, 800 U T4 DNA Ligase with rATP, 1 $\times$  NEBuffer2, 250 ng genomic DNA, in a total volume of 60  $\mu$ L (New England BioLabs). To maximize diversity of DNA fragments, we performed two separate DIG/LIG reactions, each with a different restriction enzyme (*Hinc*II and *Pvu*II). The thermal profile for the DIG/LIG reaction was 22 cycles of (37 °C for 10 min and 16 °C for 30 min), followed by a 20-min enzyme denatura-

tion step at 65 °C. Products were visualized on an agarose gel stained with ethidium bromide, and resulting genomic DNA fragments ranged in size from 300 to 1500 base pairs. We used tetra-oligonucleotide probes (GATA)<sub>8</sub> and (GACA)<sub>4</sub> for the subtractive hybridization procedure; both were performed at 48 °C. To isolate DNA fragments from the enriched libraries, we used the TOPO TA cloning kit (Invitrogen). From the two libraries, we purified and amplified a total of 1344 clones via a Templphi reaction (GE Healthcare). Cloned DNA fragments from the Templphi reaction were ready for DNA sequencing without any modification. We used BigDye Terminator version 3.1 kit, and for sequencing cleanup, we used the Ethanol/EDTA/Sodium Acetate Precipitation method (Applied Biosystems). Sequencing products were visualized with a 3100-*Avant* Genetic Analyser (Applied Biosystems).

We cloned and isolated 1344 DNA fragments from *Hinc*II and *Pvu*II libraries, and of these, 711 DNA fragments contained microsatellite repeat motifs, a 53% enrichment efficiency. We selected 128 DNA fragments that contained more than 12 microsatellite repeat motifs for primer design and subsequent screening (Rozen & Skaletsky 2000). To screen for potentially informative loci, we performed

polymerase chain reaction (PCR) and visualized the PCR products using agarose gel-electrophoresis. For loci that amplified and appeared variable, we fluorescently labelled the left primer at the 5' end, and performed PCR on a larger sample size. A typical PCR volume consisted of 1 µM of each primer, 5 µL 2× QIAGEN Multiplex PCR Kit, 50 ng DNA, and enough water for a final volume of 10 µL. The thermo profile consisted of one activation step at 95 °C for 15 min followed by 30 cycles (94 °C for 30 s, 60 °C for 90 s and 72 °C for 30 s), with a final extension step at 72 °C for 30 min (QIAGEN, catalogue no. 206143). For all PCR thermocycling, we used the DNA Engine DYAD thermocycler (Bio-Rad Laboratories). Fluorescently labelled PCR products were visualized on a 3100-*Avant* Genetic Analyser. From the screening processes, 12 loci were shown to be polymorphic. For each of these loci, we genotyped 96 individuals to screen for variability and assess Hardy-Weinberg (HW) equilibrium.

We summarized the genetic variation at these 12 loci with a few summary statistics and statistical tests (Table 1). The number of alleles ranged from two to 11, with an average of 5.3 alleles per locus. The expected heterozygosity for each locus was calculated using HW-QuickCheck (Kalinowski 2006) and ranged from 0.29 to 0.76, with an average of 0.68. Deviations from HW expectations were tested using the exact probability test of Guo & Thompson (1992), as implemented by the web version of GenePop (<http://genepop.curtin.edu.au/>; Raymond & Rousset 1995). None of the loci showed deviations from HW expectations at the 0.05 level of significance. We tested for genotypic equilibrium between all pairs of loci using a Markov chain version of Fisher's exact test as implemented by the web version of GenePop (<http://genepop.curtin.edu.au/>; Raymond & Rousset 1995). Two pairs of loci showed significant association of genotypes at the 0.001 level of significance: Sna\_MSU02 and Sna\_MSU10, and Sna\_MSU07 and Sna\_MSU12. These results show that the microsatellite loci here possess high levels of polymorphism and should be useful for identifying the source of lake trout introductions and for estimating the number of founders.

## Acknowledgements

This work was primarily funded by Montana Fish Wildlife and Parks, with some support provided by the National Science Foundation (DEB 071756). We thank Clint Muhlfeld (US Geological Survey, Flathead, MT) and Ben Cox (Montana Cooperative Fishery Research Unit, Bozeman, MT) for providing tissue samples. Lastly, we thank Bob Sharrock (Montana State University, Plant Science and Plant Pathology, Bozeman, MT) for providing facilities and equipment needed to perform the microsatellite enrichment.

## References

- Benke RJ (2002) *Trout and Salmon of North America*. The Free Press, New York.
- Crossman E (1995) Introduction of the lake trout in areas outside its native distribution: a review. *Journal of Great Lakes Research*, **21** (Suppl. 1), 17–29.
- Fredenberg W (2002) Further evidence that lake trout displace bull trout in mountain lakes. *Intermountain Journal of Sciences*, **8**, 143–152.
- Guo SW, Thompson EA (1992) Performing the exact test of Hardy-Weinberg proportion for multiple alleles. *Biometrics*, **48**, 359.
- Hamilton MB, Pincus EL, DiFiore A, Fleischer RC (1999) Microsatellite-enhanced genomic library construction streptavidin-biotin bench version SNX linker. *BioTechniques*, **27**, 500–507.
- Kalinowski ST (2006) HW-QuickCheck: a computer program for checking genotypes for agreement with Hardy-Weinberg expectations. *Molecular Ecology Notes*, **6**, 974–979.
- Koel TM, Bigelow PE, Doepke PD, Ertel BD, Mahony DL (2005) Nonnative lake trout result in Yellowstone cutthroat trout decline and impacts to bears and anglers. *Fisheries*, **30**(11), 10–19.
- Raymond M, Rousset F (1995) GenePop (version 1.2): population genetics software for exact tests and ecumenicism. *The Journal of Heredity*, **86**, 248–249.
- Rozen S, Skaletsky HS (2000) Primer 3 on the WWW for general users and for biologist programmers. In: *Bioinformatics Methods and Protocols: Methods in Molecular Biology* (eds Krawetz S, Misener S), pp. 365–386. Humana Press, Totowa, New Jersey.
- Ruzycki JR, Beauchamp DA, Yule DL (2003) Effects of introduced lake trout on native cutthroat trout in Yellowstone Lake. *Ecological Applications*, **31**, 23–37.

doi: 10.1111/j.1755-0998.2008.02403.x

© 2009 The Authors

Journal compilation © 2009 Blackwell Publishing Ltd