


# Trends in SEER Incidence & US Death Rates by Primary Cancer Site 1996-2005

## Trends in SEER Incidence Rates



## Trends in US Cancer Death Rates




Figure 1-4

Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia) and US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

For sex-specific cancer sites, the population was limited to the population of the appropriate sex.

Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

\* The Annual Percent Change is significantly different from zero ( $p < .05$ ).

<sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

# Trends in SEER Incidence Rates by Primary Cancer Site 1996-2005




Figure 1-5

Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). For sex-specific cancer sites, the population was limited to the population of the appropriate sex. Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

\* The Annual Percent Change is significantly different from zero ( $p < .05$ ).

<sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

# Trends in US Death Rates by Primary Cancer Site 1996-2005




Figure 1-6

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.  
 For sex-specific cancer sites, the population was limited to the population of the appropriate sex.  
 Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).  
 \* The Annual Percent Change is significantly different from zero ( $p < .05$ ).

# Trends in SEER Incidence Rates by Primary Cancer Site 1996-2005




Figure 1-7

Source: SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).

Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

\* The Annual Percent Change is significantly different from zero ( $p < .05$ ).

<sup>a</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

# Trends in US Death Rates by Primary Cancer Site 1996-2005





Figure 1-8

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.  
Underlying rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).  
\* The Annual Percent Change is significantly different from zero (p<.05).

Figure I-10

# Trends in US Death Rates, 1996-2005 by Primary Cancer Site All Ages, by Race



Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. The APC is the Annual Percent Change over the time interval. Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

# US Death Rates 1975-2005

## Ages Less Than 65

## Ages 65 and Over





Figure 1-3

Source: US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).