

Table I-21
US PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2005^a
USING DIFFERENT TUMOR INCLUSION CRITERIA^b

Site/Sex	5-Year Limited Duration			30-year Limited Duration	
	1st Invasive Tumor Ever ^c	1st Per Site in Previous 30 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 30 Years ^d
All Sites	4,047,630	4,124,002	4,471,606	10,505,741	10,700,724
Male	2,069,338	2,097,658	2,265,792	4,896,353	4,954,654
Female	1,978,292	2,026,344	2,205,814	5,609,388	5,746,070
Oral Cavity & Pharynx	86,675	99,393	102,699	223,620	245,611
Male	58,447	66,437	68,409	144,387	157,204
Female	28,228	32,956	34,290	79,233	88,407
Esophagus	18,953	22,591	22,591	27,187	31,941
Male	14,462	17,126	17,126	20,510	23,821
Female	4,491	5,465	5,465	6,677	8,120
Stomach	31,518	37,554	37,666	60,885	69,650
Male	18,834	22,619	22,647	34,789	40,033
Female	12,684	14,935	15,019	26,096	29,617
Colon & Rectum	414,431	477,379	485,167	1,050,283	1,168,268
Male	209,554	241,797	245,238	515,575	570,013
Female	204,877	235,582	239,929	534,708	598,255
Liver & Intrahep	15,981	18,086	18,086	21,259	23,745
Male	11,073	12,473	12,473	14,023	15,605
Female	4,908	5,613	5,613	7,236	8,140
Pancreas	21,905	26,051	26,051	29,026	33,751
Male	10,950	13,152	13,152	14,178	16,577
Female	10,955	12,899	12,899	14,848	17,174
Larynx	30,668	35,999	36,272	88,989	98,327
Male	24,542	28,792	29,019	71,335	78,509
Female	6,126	7,207	7,253	17,654	19,818
Lung & Bronchus	206,013	255,746	260,876	348,273	417,553
Male	99,962	125,160	127,386	165,661	198,810
Female	106,051	130,586	133,490	182,612	218,743
Melanoma of the Skin	221,975	247,168	256,223	677,773	725,006
Male	116,685	132,017	137,768	333,977	360,601
Female	105,290	115,151	118,455	343,796	364,405
Breast					
Female	808,737	867,285	909,008	2,377,471	2,521,460
Cervix					
Female	40,655	42,514	42,620	189,952	195,487
Corpus & Uterus					
Female	145,463	163,808	163,843	511,115	554,123
Ovary ^f					
Female	53,959	63,067	63,111	151,232	170,276

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER 9 registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census.

^{bcd} (b) Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion: (c) First invasive tumor ever; (d) First invasive tumor for each cancer site diagnosed during the previous 30 years (1975-2004); (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2000-2004). For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2000; Melanoma in 2001.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 30-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2005.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 30-year limited duration prevalence. The 2000 breast cancer is counted for the breast 5-year and 30-year limited duration prevalence.

In method (e) the 2000 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2001 melanoma is counted for 5-year limited duration prevalence for melanoma.

^f Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table I-21 - continued
US PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2005^a
USING DIFFERENT TUMOR INCLUSION CRITERIA^b

Site/Sex	5-Year Limited Duration			30-year Limited Duration	
	1st Invasive Tumor Ever ^c	1st Per Site in Previous 30 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 30 Years ^d
Prostate					
Male	939,384	1,013,473	1,013,521	2,102,958	2,244,449
Testis					
Male	39,561	40,196	40,664	165,843	167,969
Urinary Bladder					
Male	187,611	229,342	232,628	500,529	575,400
Female	140,466	171,744	174,234	370,894	424,832
Female	47,145	57,598	58,394	129,635	150,568
Kidney & Renal Pelvis					
Male	105,967	127,975	128,978	242,021	279,591
Female	63,208	77,412	78,092	142,504	165,759
Female	42,759	50,563	50,886	99,517	113,832
Brain & Nervous System					
Male	40,067	41,895	42,078	105,744	108,660
Female	21,582	22,580	22,690	56,701	58,150
Female	18,485	19,315	19,388	49,043	50,510
Thyroid					
Male	111,720	120,840	121,123	344,182	362,308
Female	24,952	28,074	28,096	76,616	81,765
Female	86,768	92,766	93,027	267,566	280,543
Hodgkin Lymphoma					
Male	34,567	36,114	36,138	140,716	143,814
Female	18,082	18,848	18,859	72,571	74,112
Female	16,485	17,266	17,279	68,145	69,702
Non-Hodgkin Lymphoma					
Male	173,017	198,936	200,239	389,438	430,711
Female	89,995	104,311	104,889	201,376	222,211
Female	83,022	94,625	95,350	188,062	208,500
Myeloma					
Male	37,951	43,567	43,601	55,888	63,084
Female	20,764	24,306	24,340	30,945	35,402
Female	17,187	19,261	19,261	24,943	27,682
Leukemia					
Male	91,507	103,055	103,168	213,808	231,461
Female	52,959	59,875	59,932	120,855	131,096
Female	38,548	43,180	43,236	92,953	100,365
Acute Lymphocytic Leuk					
Male	14,454	14,746	14,746	52,633	53,040
Female	8,169	8,268	8,268	28,919	29,056
Female	6,285	6,478	6,478	23,714	23,984
Childhood (0-19)					
Male	59,044	59,122	59,392	248,912	249,391
Female	31,593	31,637	31,734	127,804	128,005
Female	27,451	27,485	27,658	121,108	121,386
Kaposi Sarcoma					
Male	6,456	6,920	6,920	20,669	21,698
Female	5,958	6,317	6,317	19,259	20,131
Female	498	603	603	1,410	1,567
Mesothelioma					
Male	2,847	3,600	3,600	4,361	5,219
Female	2,040	2,599	2,599	2,646	3,276
Female	807	1,001	1,001	1,715	1,943

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER 9 registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census.

^{bcde} (b) Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion: (c) First invasive tumor ever; (d) First invasive tumor for each cancer site diagnosed during the previous 30 years (1975-2004); (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (2000-2004). For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 2000; Melanoma in 2001.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 30-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred more than 5 years prior to 1/1/2005.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 30-year limited duration prevalence. The 2000 breast cancer is counted for the breast 5-year and 30-year limited duration prevalence.

In method (e) the 2000 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 2001 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table I-22
US COMPLETE PREVALENCE COUNTS, INVASIVE CANCERS ONLY, January 1, 2005^a
BY AGE AT PREVALENCE

Age at Prevalence	Age Specific								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
<u>Site/Sex</u>									
All Sites									
Males	5,017,159	16,399	39,131	72,187	144,424	325,302	684,471	1,158,366	2,576,879
Females	6,081,291	14,148	32,907	78,183	213,178	591,903	1,084,176	1,295,883	2,770,913
Oral Cavity & Pharynx									
Males	154,159	22	457	1,378	3,285	14,388	34,829	39,166	60,632
Females	86,017	94	445	1,590	3,294	8,286	15,001	18,032	39,275
Esophagus									
Males	20,618	0	0	14	190	993	4,149	6,451	8,820
Females	6,790	0	12	24	39	245	870	1,475	4,126
Stomach									
Males	36,118	0	17	64	601	2,306	5,417	8,526	19,186
Females	27,093	17	23	133	452	1,835	3,582	4,955	16,095
Colon & Rectum									
Males	531,875	11	58	908	4,952	21,683	65,986	117,180	321,098
Females	563,408	0	92	1,021	4,786	19,725	55,223	95,601	386,961
Liver & Intrahep									
Males	14,098	318	426	302	385	1,286	4,581	3,125	3,676
Females	7,544	406	317	308	348	739	1,283	1,519	2,624
Pancreas									
Males	14,355	0	35	69	243	1,227	3,204	3,844	5,732
Females	15,090	0	62	115	368	1,335	2,493	3,533	7,183
Larynx									
Males	74,702	0	0	46	211	2,653	10,886	19,951	40,955
Females	18,394	0	0	46	186	1,184	2,730	5,061	9,186
Lung & Bronchus									
Males	172,426	22	88	267	1,144	6,635	23,556	50,032	90,682
Females	187,655	0	44	360	1,473	8,522	25,008	50,522	101,726
Melanoma of the Skin									
Males	349,414	70	796	4,912	16,062	46,512	78,727	81,518	120,817
Females	374,002	78	890	10,158	31,346	67,262	88,525	69,337	106,407

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

Table I-22 - continued
US COMPLETE PREVALENCE COUNTS, INVASIVE CANCERS ONLY, January 1, 2005^a
 BY AGE AT PREVALENCE

Age at Prevalence	Age Specific								
	All Ages ^c	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70+
<u>Site/Sex</u>									
Breast									
Males	12,864	0	0	23	86	500	2,112	3,338	6,804
Females	2,477,847	0	70	2,367	35,542	205,803	485,718	597,083	1,151,266
Cervix									
Females	249,756	23	58	2,406	18,587	44,912	56,492	49,040	78,238
Corpus & Uterus, NOS									
Females	572,626	0	43	598	4,635	23,170	73,904	120,851	349,424
Ovary ^d									
Females	174,236	44	985	3,228	7,229	20,668	36,444	39,782	65,855
Prostate									
Males	2,106,499	22	69	93	286	16,233	179,722	532,480	1,377,594
Urinary Bladder									
Males	384,294	39	139	619	2,498	12,873	40,736	85,424	241,966
Females	136,858	11	56	425	1,212	4,625	12,496	26,832	91,201
Kidney & Renal Pelvis									
Males	148,837	1,493	2,452	2,268	4,063	13,380	29,131	38,394	57,657
Females	104,665	1,495	2,574	2,621	3,821	9,794	16,463	22,752	45,145
Hodgkin Lymphoma									
Males	80,386	127	2,071	8,803	16,282	22,142	17,052	8,546	5,364
Females	75,786	66	1,541	9,602	16,859	20,963	14,940	6,428	5,387
Non-Hodgkin Lymphoma									
Males	207,821	585	3,464	6,304	11,912	25,726	40,865	46,986	71,979
Females	193,876	313	1,515	3,644	7,917	18,360	33,094	41,373	87,660
Myeloma									
Males	31,066	0	0	40	550	2,132	6,299	9,490	12,555
Females	25,134	0	0	23	283	1,525	4,520	6,533	12,251
Leukemia									
Males	123,731	6,018	11,203	9,962	8,677	10,545	16,300	21,846	39,180
Females	96,332	4,782	9,585	9,056	7,170	7,633	10,616	13,737	33,754
Acute Lymphocytic Leuk									
Males	30,765	4,979	9,818	7,749	4,881	1,772	722	528	315
Females	25,452	4,032	8,130	6,555	3,892	1,532	651	356	304

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER 9 registries (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta) and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^b Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^c Due to rounding, the sum of the age specific estimates may not equal the all ages estimate.

^d Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Table II-16
ALL CANCER SITES (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	4,047,630	2,669,631	1,766,113	1,019,963	599,763	402,640	8,483,375	10,505,741	592,709	11,098,450
Males	2,069,338	1,307,711	836,905	374,429	197,838	110,132	4,213,955	4,896,353	120,806	5,017,159
Females	1,978,292	1,361,920	929,208	645,534	401,925	292,508	4,269,420	5,609,388	471,903	6,081,291
White ^b	3,505,535	2,345,934	1,582,582	927,604	548,404	368,182	7,434,050	9,278,240	524,364	9,802,604
Males	1,783,465	1,140,227	749,465	342,954	182,989	101,749	3,673,157	4,300,849	112,370	4,413,219
Females	1,722,070	1,205,707	833,117	584,650	365,415	266,433	3,760,893	4,977,391	411,994	5,389,385
Black ^b	376,663	221,875	126,932	61,263	34,507	22,272	725,470	843,511	32,984	876,495
Males	204,157	119,998	63,513	21,191	9,722	4,883	387,668	423,463	3,508	426,971
Females	172,506	101,877	63,419	40,072	24,785	17,389	337,802	420,048	29,476	449,524
Asian/Pacific Islander ^c	101,457	57,985	33,530	+	+	+	192,972	+	+	+
Males	44,546	23,346	12,527	+	+	+	80,419	+	+	+
Females	56,911	34,639	21,003	+	+	+	112,553	+	+	+
Hispanic ^d	225,763	129,311	80,093	+	+	+	435,168	+	+	+
Males	109,189	60,160	33,879	+	+	+	203,229	+	+	+
Females	116,574	69,151	46,214	+	+	+	231,939	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	2.5665%	0.0751%	0.1545%	0.2626%	0.6521%	1.5479%	3.7013%	8.1540%	13.5257%	14.5615%	2.6954%
Males	2.5517%	0.0794%	0.1635%	0.2445%	0.5114%	1.0541%	2.9999%	8.9457%	18.1752%	21.4303%	3.0858%
Females	2.5810%	0.0705%	0.1450%	0.2818%	0.7972%	2.0393%	4.3674%	7.4419%	9.9672%	10.7713%	2.4751%
White ^c	2.8091%	0.0807%	0.1690%	0.2917%	0.7220%	1.6386%	3.8364%	8.5408%	14.1986%	14.9919%	2.8235%
Males	2.7614%	0.0860%	0.1795%	0.2717%	0.5769%	1.1295%	3.0751%	9.1701%	18.6971%	21.7580%	3.1798%
Females	2.8568%	0.0752%	0.1581%	0.3137%	0.8780%	2.1616%	4.5817%	7.9592%	10.6784%	11.3062%	2.6334%
Black ^c	1.8442%	0.0523%	0.1040%	0.1706%	0.4569%	1.2391%	3.6484%	8.4041%	13.3074%	12.4859%	2.5325%
Males	2.0672%	0.0498%	0.1067%	0.1540%	0.3212%	0.8620%	3.6986%	11.4989%	21.3769%	21.4479%	3.4670%
Females	1.6420%	0.0548%	0.1012%	0.1863%	0.5761%	1.5699%	3.6067%	5.9835%	7.9668%	8.1636%	1.9573%
Asian/Pacific Islander ^c	1.6210%	0.0598%	0.1181%	0.1624%	0.3852%	1.1320%	2.5941%	4.8656%	8.2375%	10.3676%	1.7546%
Males	1.4420%	0.0646%	0.1277%	0.1453%	0.2430%	0.5881%	1.5462%	4.5856%	10.6326%	14.9282%	1.8100%
Females	1.7887%	0.0548%	0.1078%	0.1793%	0.5197%	1.6296%	3.5117%	5.1007%	6.4987%	7.3349%	1.7650%
Hispanic ^d	1.0253%	0.0713%	0.1408%	0.1805%	0.4458%	1.1046%	2.5056%	5.4872%	9.4444%	10.2114%	1.8788%
Males	0.9155%	0.0757%	0.1478%	0.1678%	0.3329%	0.6242%	1.7701%	5.8656%	13.2222%	15.5695%	2.1411%
Females	1.1415%	0.0667%	0.1335%	0.1958%	0.5764%	1.6095%	3.1963%	5.1701%	6.7139%	7.1273%	1.7625%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^{f h} Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table III-16
BRAIN AND OTHER NERVOUS SYSTEM (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	40,067	21,818	17,952	12,639	7,531	5,737	79,837	105,744	14,507	120,251
Males	21,582	11,766	9,954	6,516	3,998	2,885	43,302	56,701	6,626	63,327
Females	18,485	10,052	7,998	6,123	3,533	2,852	36,535	49,043	7,881	56,924
White ^b	35,522	19,272	15,765	11,339	6,716	5,186	70,558	93,799	7,819	101,618
Males	19,364	10,493	8,685	5,736	3,588	2,637	38,541	50,503	4,129 ⁱ	54,632
Females	16,158	8,779	7,080	5,603	3,128	2,549	32,017	43,296	3,690 ⁱ	46,986
Black ^b	2,970	1,561	1,515	952	599	416	6,046	8,014	+	+
Males	1,429	834	823	559	311	180	3,086	4,137	+	+
Females	1,541	727	692	393	288	236	2,960	3,877	+	+
Asian/Pacific Islander ^c	1,008	572	385	+	+	+	1,965	+	+	+
Males	501	268	232	+	+	+	1,001	+	+	+
Females	507	304	153	+	+	+	964	+	+	+
Hispanic ^d	3,797	1,985	1,433	+	+	+	7,215	+	+	+
Males	2,086	986	740	+	+	+	3,812	+	+	+
Females	1,711	999	693	+	+	+	3,403	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0247%	0.0134%	0.0279%	0.0233%	0.0243%	0.0300%	0.0300%	0.0286%	0.0236%	0.0151%	0.0247%
Males	0.0270%	0.0138%	0.0303%	0.0250%	0.0259%	0.0323%	0.0342%	0.0329%	0.0284%	0.0160%	0.0270%
Females	0.0225%	0.0129%	0.0254%	0.0216%	0.0227%	0.0278%	0.0260%	0.0248%	0.0198%	0.0146%	0.0225%
White ^c	0.0277%	0.0146%	0.0310%	0.0254%	0.0282%	0.0341%	0.0339%	0.0321%	0.0259%	0.0148%	0.0276%
Males	0.0303%	0.0152%	0.0338%	0.0271%	0.0298%	0.0363%	0.0386%	0.0371%	0.0307%	0.0170%	0.0301%
Females	0.0251%	0.0140%	0.0281%	0.0235%	0.0264%	0.0319%	0.0294%	0.0275%	0.0221%	0.0136%	0.0252%
Black ^c	0.0157%	0.0096%	0.0191%	0.0175%	0.0140%	0.0160%	0.0184%	0.0177%	0.0138%	0.0183%	0.0158%
Males	0.0165%	0.0100%	0.0199%	0.0189%	0.0142%	0.0174%	0.0202%	0.0175%	0.0231%	-	0.0164%
Females	0.0150%	0.0091%	0.0182%	0.0162%	0.0138%	0.0147%	0.0169%	0.0178%	0.0076%	0.0272%	0.0150%
Asian/Pacific Islander ^c	0.0141%	0.0085%	0.0183%	0.0158%	0.0121%	0.0169%	0.0137%	0.0129%	0.0144%	0.0124%	0.0141%
Males	0.0148%	0.0079%	0.0199%	0.0163%	0.0124%	0.0180%	0.0153%	0.0121%	0.0163%	0.0137%	0.0148%
Females	0.0134%	0.0091%	0.0165%	0.0153%	0.0117%	0.0160%	0.0122%	0.0137%	0.0130%	0.0116%	0.0135%
Hispanic ^d	0.0172%	0.0108%	0.0197%	0.0128%	0.0160%	0.0244%	0.0245%	0.0274%	0.0208%	0.0120%	0.0187%
Males	0.0176%	0.0110%	0.0196%	0.0130%	0.0163%	0.0248%	0.0265%	0.0311%	0.0250%	-	0.0195%
Females	0.0169%	0.0106%	0.0198%	0.0125%	0.0156%	0.0240%	0.0227%	0.0242%	0.0177%	0.0128%	0.0180%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table IV-21
BREAST CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	814,237	629,517	423,556	279,066	147,527	96,146	1,867,310	2,390,048	100,663	2,490,711
Males	5,500	3,095	1,898	1,018	729	337	10,493	12,577	287	12,864
Females	808,737	626,422	421,658	278,048	146,798	95,809	1,856,817	2,377,471	100,376	2,477,847
White ^b	709,422	558,831	381,068	254,704	134,348	88,312	1,649,320	2,126,685	89,590	2,216,275
Males	4,624	2,736	1,654	890	701	324	9,014	10,929	166 [†]	11,095
Females	704,798	556,095	379,414	253,814	133,647	87,988	1,640,306	2,115,756	89,424	2,205,180
Black ^b	72,473	48,161	29,705	17,448	9,670	5,483	150,338	182,939	+	+
Males	706	252	172	110	25	13	1,130	1,278	+	+
Females	71,767	47,909	29,533	17,338	9,645	5,470	149,208	181,661	7,001	188,662
Asian/Pacific Islander ^c	24,107	16,145	9,249	+	+	+	49,502	+	+	+
Males	77	55	43	+	+	+	176	+	+	+
Females	24,030	16,090	9,206	+	+	+	49,326	+	+	+
Hispanic ^d	43,674	27,804	17,423	+	+	+	88,901	+	+	+
Males	179	70	44	+	+	+	293	+	+	+
Females	43,495	27,734	17,379	+	+	+	88,608	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.5704%	-	0.0002%	0.0056%	0.0807%	0.4250%	1.1343%	1.9165%	2.5023%	2.9250%	0.5883%
Males	0.0063%	-	-	-	0.0004%	0.0023%	0.0097%	0.0251%	0.0396%	0.0563%	0.0076%
Females	1.1236%	-	0.0003%	0.0114%	0.1634%	0.8456%	2.2024%	3.6179%	4.3872%	4.5080%	1.0717%
White ^c	0.6272%	-	0.0001%	0.0053%	0.0797%	0.4304%	1.1734%	2.0381%	2.6905%	3.1255%	0.6195%
Males	0.0068%	-	-	-	0.0003%	0.0020%	0.0098%	0.0248%	0.0419%	0.0601%	0.0078%
Females	1.2473%	-	0.0002%	0.0111%	0.1650%	0.8706%	2.3125%	3.8984%	4.7632%	4.7953%	1.1396%
Black ^c	0.3885%	-	-	0.0075%	0.0967%	0.4080%	1.0318%	1.5911%	2.0442%	2.3373%	0.5082%
Males	0.0063%	-	-	-	-	0.0040%	0.0143%	0.0436%	0.0407%	0.0564%	0.0098%
Females	0.7350%	-	-	0.0143%	0.1811%	0.7624%	1.8754%	2.8014%	3.3702%	3.4373%	0.8732%
Asian/Pacific Islander ^c	0.4054%	-	-	0.0051%	0.0727%	0.4045%	0.9608%	1.3614%	1.5556%	1.5646%	0.4236%
Males	0.0031%	-	-	-	-	-	0.0044%	0.0116%	0.0216%	0.0267%	0.0038%
Females	0.7822%	-	-	0.0102%	0.1409%	0.7733%	1.7982%	2.4946%	2.6692%	2.5872%	0.7660%
Hispanic ^d	0.2125%	-	-	0.0043%	0.0550%	0.2973%	0.7711%	1.2670%	1.6036%	1.7517%	0.3875%
Males	0.0013%	-	-	-	-	0.0012%	0.0038%	0.0097%	0.0157%	0.0216%	0.0031%
Females	0.4359%	-	-	0.0094%	0.1181%	0.6086%	1.4916%	2.3208%	2.7513%	2.7476%	0.7040%

- ^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- ^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
- ^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- ^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
- ^{g h i} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table V-12
CERVIX UTERI CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race</u>										
All Races ^b	40,655	39,359	35,168	29,369	23,160	22,241	115,182	189,952	59,804	249,756
White ^b	31,965	31,027	29,121	24,595	19,114	18,103	92,113	153,925	49,832	203,757
Black ^b	6,045	5,475	4,026	3,232	3,093	3,258	15,546	25,129	+	+
Asian/Pacific Islander ^c	1,960	1,772	1,409	+	+	+	5,141	+	+	+
Hispanic ^d	8,273	6,351	5,650	+	+	+	20,274	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	0-9	10-19	20-29	Age Specific (Crude)						Age-Adjusted ^f
					30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.0837%	-	0.0003%	0.0106%	0.0847%	0.1657%	0.1750%	0.1550%	0.1374%	0.0852%	0.0817%
White ^c	0.0849%	-	0.0003%	0.0115%	0.0963%	0.1758%	0.1722%	0.1427%	0.1207%	0.0643%	0.0823%
Black ^c	0.0763%	-	-	0.0104%	0.0680%	0.1433%	0.1713%	0.1864%	0.1962%	0.1855%	0.0841%
Asian/Pacific Islander ^c	0.0791%	-	-	0.0032%	0.0324%	0.1173%	0.1821%	0.2021%	0.2002%	0.1900%	0.0764%
Hispanic ^d	0.1004%	-	-	0.0115%	0.0929%	0.2617%	0.3186%	0.3077%	0.3045%	0.1828%	0.1387%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table VI-23
COLON AND RECTUM CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
Race/Sex										
All Races ^b	414,431	254,745	166,882	113,245	64,365	36,616	836,057	1,050,283	45,000	1,095,283
Males	209,554	126,290	81,920	55,117	28,251	14,443	417,764	515,575	16,300	531,875
Females	204,877	128,455	84,962	58,128	36,114	22,173	418,293	534,708	28,700	563,408
White ^b	353,567	222,365	147,039	100,310	57,833	32,917	722,972	914,032	40,137	954,169
Males	180,274	111,254	72,595	49,312	25,614	13,030	364,124	452,080	14,680	466,760
Females	173,293	111,111	74,444	50,998	32,219	19,887	358,848	461,952	25,457	487,409
Black ^b	41,826	21,999	13,724	9,403	4,640	2,471	77,549	94,063	3,145	97,208
Males	19,169	9,449	6,118	3,940	1,657	846	34,736	41,179	1,052	42,231
Females	22,657	12,550	7,606	5,463	2,983	1,625	42,813	52,884	2,093	54,977
Asian/Pacific Islander ^c	13,422	7,484	4,338	+	+	+	25,245	+	+	+
Males	6,984	3,951	2,196	+	+	+	13,132	+	+	+
Females	6,438	3,533	2,142	+	+	+	12,113	+	+	+
Hispanic ^d	21,478	11,024	6,580	+	+	+	39,082	+	+	+
Males	11,505	5,926	3,291	+	+	+	20,722	+	+	+
Females	9,973	5,098	3,289	+	+	+	18,360	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.2532%	-	0.0003%	0.0040%	0.0214%	0.0843%	0.2945%	0.7579%	1.5313%	2.2292%	0.2696%
Males	0.2574%	-	0.0002%	0.0040%	0.0218%	0.0881%	0.3317%	0.9023%	1.8701%	2.5412%	0.3163%
Females	0.2491%	-	0.0004%	0.0041%	0.0209%	0.0805%	0.2591%	0.6279%	1.2719%	2.0570%	0.2337%
White ^c	0.2675%	-	0.0004%	0.0041%	0.0211%	0.0817%	0.2816%	0.7495%	1.5457%	2.2752%	0.2693%
Males	0.2708%	-	0.0003%	0.0041%	0.0214%	0.0856%	0.3209%	0.8972%	1.8844%	2.5918%	0.3167%
Females	0.2642%	-	0.0005%	0.0041%	0.0207%	0.0778%	0.2430%	0.6130%	1.2807%	2.1028%	0.2322%
Black ^c	0.1971%	-	-	0.0031%	0.0215%	0.0957%	0.3678%	0.9130%	1.5638%	1.9501%	0.2845%
Males	0.1893%	-	-	0.0028%	0.0213%	0.1000%	0.4009%	1.0187%	1.8080%	2.1170%	0.3170%
Females	0.2042%	-	-	0.0035%	0.0217%	0.0919%	0.3405%	0.8304%	1.4022%	1.8696%	0.2622%
Asian/Pacific Islander ^c	0.2216%	-	-	0.0038%	0.0208%	0.0859%	0.3022%	0.6740%	1.3787%	2.0007%	0.2480%
Males	0.2369%	-	-	0.0041%	0.0226%	0.0889%	0.3282%	0.8173%	1.7612%	2.3637%	0.2975%
Females	0.2073%	-	-	0.0034%	0.0192%	0.0832%	0.2795%	0.5537%	1.1010%	1.7593%	0.2112%
Hispanic ^d	0.0915%	-	-	0.0030%	0.0161%	0.0636%	0.2147%	0.5509%	1.1124%	1.4555%	0.1922%
Males	0.0931%	-	-	0.0034%	0.0156%	0.0630%	0.2405%	0.6781%	1.4755%	1.6414%	0.2327%
Females	0.0899%	-	-	0.0026%	0.0167%	0.0643%	0.1904%	0.4442%	0.8500%	1.3485%	0.1621%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table VII-12
CORPUS AND UTERUS, NOS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race</u>										
All Races ^b	145,463	113,728	85,844	64,851	50,894	50,336	345,035	511,115	61,511	572,626
White ^b	129,156	103,694	80,149	60,781	48,301	47,993	312,999	470,075	54,349	524,424
Black ^b	9,616	5,388	3,094	2,382	1,289	1,176	18,097	22,944	1,309	24,253
Asian/Pacific Islander ^c	4,227	2,777	1,662	+	+	+	8,665	+	+	+
Hispanic ^d	8,104	4,966	3,562	+	+	+	16,632	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	0-9	10-19	20-29	Age Specific (Crude)						Age-Adjusted ^f
					30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.2043%	-	-	0.0029%	0.0233%	0.0975%	0.3384%	0.7521%	0.9500%	0.8676%	0.1962%
White ^c	0.2342%	-	-	0.0027%	0.0233%	0.0974%	0.3600%	0.8335%	1.0696%	0.9611%	0.2146%
Black ^c	0.0882%	-	-	0.0032%	0.0190%	0.0590%	0.1638%	0.4363%	0.5727%	0.3929%	0.1097%
Asian/Pacific Islander ^c	0.1364%	-	-	0.0026%	0.0248%	0.1261%	0.3426%	0.4651%	0.4328%	0.3596%	0.1327%
Hispanic ^d	0.0816%	-	-	0.0037%	0.0309%	0.0954%	0.2513%	0.5095%	0.5305%	0.4599%	0.1322%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table VIII-16
ESOPHAGUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	18,953	4,963	2,127	700	346	99	26,042	27,187	221	27,408
Males	14,462	3,732	1,564	456	238	60	19,757	20,510	108	20,618
Females	4,491	1,231	563	244	108	39	6,285	6,677	113	6,790
White ^b	16,776	4,378	1,774	585	291	73	22,928	23,878	182	24,060
Males	13,062	3,335	1,299	382	209	60	17,696	18,347	101	18,448
Females	3,714	1,043	475	203	82	13	5,232	5,531	81	5,612
Black ^b	1,707	438	293	92	54	26	2,438	2,610	55	2,665
Males	1,026	279	218	62	28	0	1,523	1,613	16	1,629
Females	681	159	75	30	26	26	915	997	39	1,036
Asian/Pacific Islander ^c	330	90	54	+	+	+	474	+	+	+
Males	240	70	42	+	+	+	351	+	+	+
Females	90	20	12	+	+	+	123	+	+	+
Hispanic ^d	716	166	106	+	+	+	987	+	+	+
Males	589	136	86	+	+	+	811	+	+	+
Females	127	30	20	+	+	+	176	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0073%	-	-	0.0001%	0.0005%	0.0024%	0.0114%	0.0295%	0.0445%	0.0364%	0.0077%
Males	0.0111%	-	-	-	0.0009%	0.0038%	0.0191%	0.0503%	0.0724%	0.0656%	0.0130%
Females	0.0036%	-	-	-	-	0.0009%	0.0040%	0.0109%	0.0232%	0.0203%	0.0035%
White ^c	0.0081%	-	-	-	0.0006%	0.0025%	0.0124%	0.0307%	0.0482%	0.0374%	0.0082%
Males	0.0125%	-	-	-	0.0011%	0.0041%	0.0213%	0.0528%	0.0790%	0.0679%	0.0139%
Females	0.0038%	-	-	-	-	0.0008%	0.0037%	0.0104%	0.0241%	0.0208%	0.0035%
Black ^c	0.0056%	-	-	-	-	0.0031%	0.0120%	0.0358%	0.0371%	0.0314%	0.0077%
Males	0.0072%	-	-	-	-	0.0040%	0.0148%	0.0544%	0.0535%	0.0535%	0.0112%
Females	0.0042%	-	-	-	-	0.0024%	0.0097%	0.0212%	0.0263%	0.0207%	0.0053%
Asian/Pacific Islander ^c	0.0041%	-	-	-	-	0.0008%	0.0049%	0.0172%	0.0255%	0.0330%	0.0046%
Males	0.0063%	-	-	-	-	0.0015%	0.0082%	0.0301%	0.0418%	0.0572%	0.0078%
Females	0.0021%	-	-	-	-	-	0.0019%	0.0065%	0.0136%	0.0168%	0.0022%
Hispanic ^d	0.0023%	-	-	-	0.0006%	0.0012%	0.0064%	0.0182%	0.0246%	0.0237%	0.0046%
Males	0.0036%	-	-	-	0.0009%	0.0019%	0.0107%	0.0350%	0.0475%	0.0460%	0.0086%
Females	0.0009%	-	-	-	-	-	0.0024%	0.0042%	0.0081%	0.0109%	0.0015%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table IX-16
HODGKIN LYMPHOMA (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
Race/Sex										
All Races ^b	34,567	29,785	26,194	21,942	16,553	11,673	90,547	140,716	15,456	156,172
Males	18,082	15,406	13,403	10,953	8,916	5,810	46,892	72,571	7,815	80,386
Females	16,485	14,379	12,791	10,989	7,637	5,863	43,655	68,145	7,641	75,786
White ^b	29,524	26,293	23,582	20,337	15,379	11,017	79,400	126,133	17,264	143,397
Males	15,566	13,586	12,191	10,089	8,160	5,369	41,343	64,961	8,878	73,839
Females	13,958	12,707	11,391	10,248	7,219	5,648	38,057	61,172	8,386	69,558
Black ^b	3,741	2,805	2,210	1,244	963	542	8,757	11,506	1,176	12,682
Males	1,782	1,429	1,039	680	644	358	4,251	5,932	808	6,740
Females	1,959	1,376	1,171	564	319	184	4,506	5,574	368	5,942
Asian/Pacific Islander ^c	746	370	259	+	+	+	1,373	+	+	+
Males	392	210	110	+	+	+	711	+	+	+
Females	354	160	149	+	+	+	662	+	+	+
Hispanic ^d	2,924	2,216	1,677	+	+	+	6,817	+	+	+
Males	1,741	1,219	938	+	+	+	3,898	+	+	+
Females	1,183	997	739	+	+	+	2,919	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0279%	0.0006%	0.0089%	0.0371%	0.0519%	0.0420%	0.0296%	0.0253%	0.0257%	0.0160%	0.0278%
Males	0.0296%	0.0008%	0.0103%	0.0341%	0.0511%	0.0458%	0.0347%	0.0311%	0.0318%	0.0202%	0.0296%
Females	0.0262%	0.0003%	0.0074%	0.0404%	0.0527%	0.0383%	0.0247%	0.0200%	0.0210%	0.0138%	0.0263%
White ^c	0.0315%	0.0006%	0.0099%	0.0421%	0.0604%	0.0472%	0.0327%	0.0278%	0.0291%	0.0174%	0.0315%
Males	0.0334%	0.0009%	0.0114%	0.0380%	0.0591%	0.0509%	0.0384%	0.0340%	0.0356%	0.0219%	0.0332%
Females	0.0297%	0.0003%	0.0083%	0.0467%	0.0618%	0.0434%	0.0271%	0.0221%	0.0240%	0.0150%	0.0301%
Black ^c	0.0233%	0.0007%	0.0064%	0.0306%	0.0411%	0.0393%	0.0289%	0.0269%	0.0184%	0.0115%	0.0240%
Males	0.0238%	-	0.0063%	0.0288%	0.0391%	0.0436%	0.0319%	0.0332%	0.0220%	0.0211%	0.0254%
Females	0.0229%	-	0.0065%	0.0323%	0.0428%	0.0356%	0.0265%	0.0219%	0.0160%	-	0.0229%
Asian/Pacific Islander ^c	0.0097%	-	0.0049%	0.0155%	0.0155%	0.0123%	0.0088%	0.0083%	0.0079%	0.0061%	0.0093%
Males	0.0104%	-	0.0074%	0.0156%	0.0140%	0.0135%	0.0092%	0.0117%	0.0116%	-	0.0102%
Females	0.0090%	-	0.0023%	0.0154%	0.0170%	0.0111%	0.0083%	0.0054%	0.0052%	-	0.0085%
Hispanic ^d	0.0162%	0.0009%	0.0089%	0.0211%	0.0253%	0.0230%	0.0210%	0.0286%	0.0242%	0.0163%	0.0180%
Males	0.0179%	0.0015%	0.0119%	0.0193%	0.0258%	0.0259%	0.0275%	0.0349%	0.0359%	0.0243%	0.0210%
Females	0.0143%	-	0.0057%	0.0233%	0.0248%	0.0200%	0.0149%	0.0233%	0.0157%	0.0117%	0.0154%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table X-13
KAPOSI SARCOMAEstimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	6,456	8,047	4,059	1,425	410	271	18,563	20,669	+	+
Males	5,958	7,622	3,772	1,387	387	132	17,353	19,259	+	+
Females	498	425	287	38	23	139	1,210	1,410	+	+
White ^b	4,544	6,310	3,404	1,197	323	224	14,258	16,002	+	+
Males	4,132	6,027	3,175	1,159	311	108	13,334	14,912	+	+
Females	412	283	229	38	12	116	924	1,090	+	+
Black ^b	1,586	1,276	403	103	40	12	3,266	3,421	+	+
Males	1,505	1,179	368	103	40	12	3,052	3,207	+	+
Females	81	97	35	0	0	0	214	214	+	+
Asian/Pacific Islander ^c	84	102	23	+	+	+	208	+	+	+
Males	79	99	23	+	+	+	201	+	+	+
Females	5	3	0	+	+	+	7	+	+	+
Hispanic ^d	1,109	1,364	590	+	+	+	3,063	+	+	+
Males	1,039	1,321	556	+	+	+	2,916	+	+	+
Females	70	43	34	+	+	+	147	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0073%	-	-	0.0009%	0.0084%	0.0196%	0.0126%	0.0071%	0.0070%	0.0150%	0.0073%
Males	0.0140%	-	-	0.0015%	0.0163%	0.0388%	0.0254%	0.0137%	0.0133%	0.0281%	0.0143%
Females	0.0007%	-	-	-	0.0003%	0.0005%	0.0005%	0.0013%	0.0021%	0.0078%	0.0007%
White ^c	0.0074%	-	-	0.0006%	0.0077%	0.0199%	0.0127%	0.0073%	0.0077%	0.0155%	0.0073%
Males	0.0140%	-	-	0.0011%	0.0144%	0.0389%	0.0252%	0.0138%	0.0144%	0.0293%	0.0140%
Females	0.0008%	-	-	-	0.0004%	0.0004%	0.0004%	0.0014%	0.0024%	0.0080%	0.0007%
Black ^c	0.0094%	-	-	0.0025%	0.0178%	0.0247%	0.0154%	0.0093%	0.0046%	0.0115%	0.0099%
Males	0.0188%	-	-	0.0046%	0.0377%	0.0520%	0.0325%	0.0195%	0.0082%	-	0.0203%
Females	0.0008%	-	-	-	-	-	-	-	-	0.0119%	0.0010%
Asian/Pacific Islander ^c	0.0016%	-	-	-	0.0012%	0.0033%	0.0041%	0.0021%	0.0022%	0.0039%	0.0016%
Males	0.0032%	-	-	-	0.0025%	0.0065%	0.0088%	0.0039%	0.0052%	-	0.0033%
Females	-	-	-	-	-	-	-	-	-	-	-
Hispanic ^d	0.0073%	-	-	0.0008%	0.0113%	0.0229%	0.0142%	0.0127%	0.0115%	0.0413%	0.0100%
Males	0.0135%	-	-	0.0014%	0.0208%	0.0443%	0.0287%	0.0239%	0.0239%	0.0755%	0.0192%
Females	0.0007%	-	-	-	-	-	-	0.0033%	-	0.0217%	0.0013%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XI-16
KIDNEY AND RENAL PELVIS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	105,967	58,131	37,454	21,035	12,406	7,030	201,552	242,021	11,481	253,502
Males	63,208	34,461	22,036	11,751	7,299	3,750	119,705	142,504	6,333	148,837
Females	42,759	23,670	15,418	9,284	5,107	3,280	81,847	99,517	5,148	104,665
White ^b	90,865	49,645	33,011	18,876	11,094	6,342	173,522	209,834	13,090	222,924
Males	54,469	29,869	19,625	10,677	6,755	3,426	103,963	124,821	6,103	130,924
Females	36,396	19,776	13,386	8,199	4,339	2,916	69,559	85,013	6,987	92,000
Black ^b	11,109	6,423	3,492	1,785	1,057	556	21,022	24,422	1,954	26,376
Males	6,308	3,349	1,859	854	421	234	11,515	13,025	698	13,723
Females	4,801	3,074	1,633	931	636	322	9,507	11,397	1,256	12,653
Asian/Pacific Islander ^c	2,199	1,087	473	+	+	+	3,759	+	+	+
Males	1,374	662	288	+	+	+	2,324	+	+	+
Females	825	425	185	+	+	+	1,435	+	+	+
Hispanic ^d	8,039	3,879	2,227	+	+	+	14,145	+	+	+
Males	4,713	2,157	1,226	+	+	+	8,096	+	+	+
Females	3,326	1,722	1,001	+	+	+	6,049	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0601%	0.0065%	0.0088%	0.0024%	0.0096%	0.0372%	0.1041%	0.2116%	0.2893%	0.2642%	0.0626%
Males	0.0729%	0.0063%	0.0078%	0.0022%	0.0103%	0.0449%	0.1379%	0.2874%	0.3967%	0.3847%	0.0833%
Females	0.0475%	0.0067%	0.0098%	0.0026%	0.0090%	0.0296%	0.0720%	0.1434%	0.2071%	0.1977%	0.0459%
White ^c	0.0657%	0.0066%	0.0096%	0.0023%	0.0102%	0.0388%	0.1085%	0.2244%	0.3038%	0.2758%	0.0657%
Males	0.0796%	0.0068%	0.0087%	0.0021%	0.0110%	0.0460%	0.1422%	0.3016%	0.4164%	0.4092%	0.0872%
Females	0.0519%	0.0065%	0.0105%	0.0024%	0.0093%	0.0313%	0.0756%	0.1531%	0.2156%	0.2031%	0.0481%
Black ^c	0.0516%	0.0074%	0.0097%	0.0041%	0.0113%	0.0413%	0.1207%	0.2193%	0.3055%	0.2587%	0.0671%
Males	0.0607%	0.0058%	0.0079%	0.0039%	0.0115%	0.0508%	0.1687%	0.2991%	0.4148%	0.3367%	0.0883%
Females	0.0433%	0.0090%	0.0117%	0.0044%	0.0111%	0.0330%	0.0808%	0.1569%	0.2332%	0.2211%	0.0519%
Asian/Pacific Islander ^c	0.0315%	0.0032%	0.0025%	-	0.0049%	0.0205%	0.0554%	0.1068%	0.1689%	0.1603%	0.0340%
Males	0.0400%	0.0025%	0.0021%	-	0.0047%	0.0290%	0.0768%	0.1558%	0.2354%	0.2180%	0.0468%
Females	0.0235%	0.0039%	0.0030%	-	0.0051%	0.0127%	0.0366%	0.0656%	0.1206%	0.1219%	0.0238%
Hispanic ^d	0.0333%	0.0055%	0.0066%	0.0016%	0.0082%	0.0327%	0.0952%	0.1989%	0.3130%	0.2676%	0.0609%
Males	0.0367%	0.0054%	0.0046%	0.0013%	0.0083%	0.0361%	0.1193%	0.2600%	0.4318%	0.3604%	0.0783%
Females	0.0297%	0.0056%	0.0087%	0.0020%	0.0080%	0.0291%	0.0726%	0.1476%	0.2271%	0.2142%	0.0476%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XII-16
LARYNX CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	30,668	21,764	15,514	10,707	6,913	3,422	67,947	88,989	4,107	93,096
Males	24,542	17,315	12,404	8,755	5,470	2,849	54,262	71,335	3,367	74,702
Females	6,126	4,449	3,110	1,952	1,443	573	13,685	17,654	740	18,394
White ^b	25,720	18,695	13,416	9,297	6,282	3,079	57,830	76,488	3,323	79,811
Males	20,593	14,947	10,715	7,724	5,035	2,563	46,254	61,576	2,713	64,289
Females	5,127	3,748	2,701	1,573	1,247	516	11,576	14,912	610	15,522
Black ^b	4,058	2,563	1,810	1,199	527	270	8,431	10,429	283	10,712
Males	3,162	1,938	1,469	909	351	221	6,569	8,051	242	8,293
Females	896	625	341	290	176	49	1,862	2,378	41	2,419
Asian/Pacific Islander ^c	527	328	171	+	+	+	1,026	+	+	+
Males	465	283	148	+	+	+	896	+	+	+
Females	62	45	23	+	+	+	130	+	+	+
Hispanic ^d	1,449	853	619	+	+	+	2,919	+	+	+
Males	1,201	741	509	+	+	+	2,450	+	+	+
Females	248	112	110	+	+	+	469	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0195%	-	-	0.0002%	0.0009%	0.0066%	0.0299%	0.0834%	0.1211%	0.0890%	0.0207%
Males	0.0317%	-	-	0.0002%	0.0010%	0.0094%	0.0501%	0.1429%	0.2250%	0.2042%	0.0378%
Females	0.0076%	-	-	0.0003%	0.0008%	0.0037%	0.0107%	0.0299%	0.0415%	0.0255%	0.0074%
White ^c	0.0211%	-	-	0.0003%	0.0011%	0.0071%	0.0302%	0.0859%	0.1244%	0.0914%	0.0213%
Males	0.0338%	-	-	-	0.0011%	0.0100%	0.0501%	0.1455%	0.2265%	0.2098%	0.0383%
Females	0.0084%	-	-	0.0004%	0.0010%	0.0042%	0.0108%	0.0309%	0.0445%	0.0269%	0.0078%
Black ^c	0.0209%	-	-	-	0.0010%	0.0077%	0.0499%	0.1261%	0.1655%	0.0917%	0.0290%
Males	0.0346%	-	-	-	-	0.0110%	0.0859%	0.2232%	0.3418%	0.2290%	0.0555%
Females	0.0086%	-	-	-	-	0.0048%	0.0200%	0.0501%	0.0488%	0.0255%	0.0105%
Asian/Pacific Islander ^c	0.0090%	-	-	-	-	0.0023%	0.0102%	0.0340%	0.0664%	0.0676%	0.0102%
Males	0.0163%	-	-	-	-	0.0044%	0.0192%	0.0654%	0.1346%	0.1484%	0.0206%
Females	0.0022%	-	-	-	-	-	0.0023%	0.0077%	0.0169%	0.0139%	0.0023%
Hispanic ^d	0.0068%	-	-	0.0003%	0.0008%	0.0039%	0.0148%	0.0532%	0.0920%	0.0658%	0.0140%
Males	0.0110%	-	-	-	0.0009%	0.0051%	0.0258%	0.1036%	0.1858%	0.1487%	0.0275%
Females	0.0023%	-	-	-	-	0.0026%	0.0045%	0.0110%	0.0243%	0.0180%	0.0039%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIII-24
LEUKEMIA (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	91,507	51,541	32,197	20,082	11,660	6,822	175,244	213,808	6,255	220,063
Males	52,959	29,090	18,065	11,265	6,219	3,257	100,114	120,855	2,876	123,731
Females	38,548	22,451	14,132	8,817	5,441	3,565	75,130	92,953	3,379	96,332
White ^b	80,801	46,082	29,268	18,225	10,618	6,317	156,151	191,311	5,309	196,620
Males	47,284	25,862	16,671	10,344	5,686	3,079	89,817	108,926	2,099	111,025
Females	33,517	20,220	12,597	7,881	4,932	3,238	66,334	82,385	3,210	85,595
Black ^b	6,378	3,104	1,618	974	610	341	11,100	13,025	651	13,676
Males	3,214	1,681	715	475	325	110	5,609	6,520	304	6,824
Females	3,164	1,423	903	499	285	231	5,491	6,505	347	6,852
Asian/Pacific Islander ^c	2,003	1,188	768	+	+	+	3,959	+	+	+
Males	1,144	684	401	+	+	+	2,229	+	+	+
Females	859	504	367	+	+	+	1,730	+	+	+
Hispanic ^d	7,572	4,391	2,957	+	+	+	14,920	+	+	+
Males	4,208	2,501	1,705	+	+	+	8,414	+	+	+
Females	3,364	1,890	1,252	+	+	+	6,506	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0542%	0.0276%	0.0491%	0.0208%	0.0200%	0.0296%	0.0578%	0.1265%	0.1983%	0.2209%	0.0562%
Males	0.0629%	0.0298%	0.0522%	0.0231%	0.0227%	0.0354%	0.0739%	0.1664%	0.2689%	0.2975%	0.0700%
Females	0.0456%	0.0254%	0.0460%	0.0184%	0.0171%	0.0238%	0.0426%	0.0906%	0.1444%	0.1787%	0.0448%
White ^c	0.0610%	0.0307%	0.0550%	0.0233%	0.0212%	0.0311%	0.0616%	0.1398%	0.2206%	0.2421%	0.0616%
Males	0.0705%	0.0335%	0.0582%	0.0259%	0.0240%	0.0371%	0.0782%	0.1829%	0.2977%	0.3273%	0.0766%
Females	0.0514%	0.0277%	0.0516%	0.0204%	0.0182%	0.0250%	0.0454%	0.0999%	0.1603%	0.1957%	0.0491%
Black ^c	0.0286%	0.0123%	0.0244%	0.0136%	0.0154%	0.0208%	0.0434%	0.0704%	0.1271%	0.1135%	0.0339%
Males	0.0319%	0.0113%	0.0240%	0.0156%	0.0177%	0.0258%	0.0562%	0.0897%	0.1587%	0.1488%	0.0409%
Females	0.0257%	0.0133%	0.0248%	0.0117%	0.0134%	0.0164%	0.0329%	0.0553%	0.1061%	0.0965%	0.0288%
Asian/Pacific Islander ^c	0.0296%	0.0251%	0.0413%	0.0137%	0.0174%	0.0236%	0.0314%	0.0561%	0.0601%	0.0615%	0.0310%
Males	0.0346%	0.0257%	0.0466%	0.0144%	0.0203%	0.0279%	0.0377%	0.0714%	0.0811%	0.0923%	0.0372%
Females	0.0250%	0.0246%	0.0356%	0.0131%	0.0146%	0.0196%	0.0259%	0.0433%	0.0449%	0.0410%	0.0257%
Hispanic ^d	0.0356%	0.0293%	0.0565%	0.0214%	0.0197%	0.0259%	0.0369%	0.0755%	0.1011%	0.1141%	0.0419%
Males	0.0388%	0.0316%	0.0591%	0.0256%	0.0220%	0.0289%	0.0455%	0.0796%	0.1376%	0.1492%	0.0487%
Females	0.0322%	0.0270%	0.0538%	0.0163%	0.0170%	0.0228%	0.0289%	0.0722%	0.0748%	0.0938%	0.0361%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIII-25
ACUTE LYMPHOCYTIC LEUKEMIA (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
Race/Sex										
All Races ^b	14,454	11,565	9,002	7,886	5,919	3,808	35,021	52,633	3,584	56,217
Males	8,169	6,463	4,953	4,421	3,013	1,901	19,585	28,919	1,846	30,765
Females	6,285	5,102	4,049	3,465	2,906	1,907	15,436	23,714	1,738	25,452
White ^b	12,583	10,035	7,887	6,935	5,341	3,573	30,506	46,356	+	+
Males	7,271	5,560	4,475	3,938	2,732	1,824	17,306	25,801	+	+
Females	5,312	4,475	3,412	2,997	2,609	1,749	13,200	20,555	+	+
Black ^b	969	817	599	462	309	131	2,386	3,289	+	+
Males	422	471	234	228	161	41	1,127	1,557	+	+
Females	547	346	365	234	148	90	1,259	1,732	+	+
Asian/Pacific Islander ^c	599	468	370	+	+	+	1,436	+	+	+
Males	327	294	173	+	+	+	794	+	+	+
Females	272	174	197	+	+	+	642	+	+	+
Hispanic ^d	3,486	2,412	1,726	+	+	+	7,623	+	+	+
Males	1,922	1,361	1,014	+	+	+	4,297	+	+	+
Females	1,564	1,051	712	+	+	+	3,326	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0128%	0.0234%	0.0424%	0.0123%	0.0046%	0.0029%	0.0025%	0.0030%	0.0018%	0.0014%	0.0130%
Males	0.0145%	0.0251%	0.0456%	0.0147%	0.0056%	0.0032%	0.0029%	0.0033%	0.0024%	0.0013%	0.0143%
Females	0.0111%	0.0217%	0.0391%	0.0097%	0.0035%	0.0026%	0.0021%	0.0028%	0.0014%	0.0015%	0.0116%
White ^c	0.0142%	0.0265%	0.0481%	0.0144%	0.0051%	0.0032%	0.0026%	0.0030%	0.0019%	0.0014%	0.0147%
Males	0.0161%	0.0288%	0.0518%	0.0171%	0.0063%	0.0034%	0.0031%	0.0035%	0.0024%	-	0.0162%
Females	0.0123%	0.0241%	0.0444%	0.0114%	0.0037%	0.0029%	0.0021%	0.0026%	0.0016%	0.0016%	0.0130%
Black ^c	0.0062%	0.0079%	0.0183%	0.0063%	0.0023%	0.0013%	-	0.0019%	-	-	0.0055%
Males	0.0064%	0.0071%	0.0177%	0.0077%	0.0018%	0.0021%	-	-	-	-	0.0055%
Females	0.0060%	0.0087%	0.0190%	0.0050%	0.0026%	-	-	-	-	-	0.0055%
Asian/Pacific Islander ^c	0.0100%	0.0207%	0.0343%	0.0066%	0.0040%	0.0028%	0.0029%	0.0036%	0.0022%	-	0.0107%
Males	0.0114%	0.0202%	0.0395%	0.0079%	0.0045%	0.0028%	0.0037%	-	-	-	0.0117%
Females	0.0086%	0.0213%	0.0288%	0.0054%	0.0034%	0.0028%	0.0023%	0.0044%	-	-	0.0096%
Hispanic ^d	0.0184%	0.0257%	0.0502%	0.0132%	0.0054%	0.0045%	0.0039%	0.0049%	-	-	0.0153%
Males	0.0201%	0.0276%	0.0531%	0.0164%	0.0055%	0.0058%	0.0037%	0.0039%	-	-	0.0166%
Females	0.0166%	0.0237%	0.0472%	0.0093%	0.0052%	0.0032%	0.0041%	0.0058%	-	-	0.0139%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIV-16
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	15,981	2,875	1,140	765	334	164	19,996	21,259	383	21,642
Males	11,073	1,734	687	296	146	87	13,494	14,023	75	14,098
Females	4,908	1,141	453	469	188	77	6,502	7,236	308	7,544
White ^b	11,560	2,156	866	625	277	98	14,582	15,582	339	15,921
Males	7,953	1,272	515	209	122	58	9,740	10,128	118 ⁱ	10,246
Females	3,607	884	351	416	155	40	4,842	5,454	221 ⁱ	5,675
Black ^b	2,210	228	117	72	35	44	2,556	2,707	+	+
Males	1,529	141	70	37	13	21	1,740	1,811	+	+
Females	681	87	47	35	22	23	816	896	+	+
Asian/Pacific Islander ^c	1,976	407	129	+	+	+	2,511	+	+	+
Males	1,385	290	83	+	+	+	1,758	+	+	+
Females	591	117	46	+	+	+	753	+	+	+
Hispanic ^d	2,119	345	197	+	+	+	2,661	+	+	+
Males	1,456	193	145	+	+	+	1,794	+	+	+
Females	663	152	52	+	+	+	867	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0082%	0.0019%	0.0012%	0.0007%	0.0015%	0.0048%	0.0182%	0.0277%	0.0374%	0.0216%	0.0085%
Males	0.0113%	0.0018%	0.0014%	0.0006%	0.0020%	0.0069%	0.0292%	0.0405%	0.0547%	0.0326%	0.0124%
Females	0.0052%	0.0020%	0.0011%	0.0007%	0.0009%	0.0027%	0.0077%	0.0161%	0.0241%	0.0155%	0.0051%
White ^c	0.0064%	0.0020%	0.0013%	0.0006%	0.0011%	0.0037%	0.0151%	0.0192%	0.0247%	0.0152%	0.0063%
Males	0.0087%	0.0018%	0.0015%	0.0006%	0.0012%	0.0049%	0.0244%	0.0278%	0.0356%	0.0219%	0.0090%
Females	0.0041%	0.0022%	0.0012%	0.0006%	0.0009%	0.0024%	0.0061%	0.0113%	0.0162%	0.0115%	0.0039%
Black ^c	0.0066%	-	0.0008%	0.0013%	0.0021%	0.0039%	0.0217%	0.0283%	0.0327%	0.0175%	0.0082%
Males	0.0097%	-	-	-	0.0033%	0.0055%	0.0368%	0.0464%	0.0505%	0.0290%	0.0131%
Females	0.0038%	-	-	0.0017%	-	0.0025%	0.0093%	0.0142%	0.0210%	0.0119%	0.0045%
Asian/Pacific Islander ^c	0.0209%	0.0029%	-	-	0.0031%	0.0125%	0.0341%	0.0817%	0.1222%	0.0778%	0.0225%
Males	0.0299%	0.0028%	-	-	0.0054%	0.0208%	0.0552%	0.1253%	0.1868%	0.1160%	0.0348%
Females	0.0125%	0.0030%	-	-	-	0.0050%	0.0157%	0.0451%	0.0753%	0.0524%	0.0127%
Hispanic ^d	0.0062%	0.0018%	0.0014%	0.0006%	0.0009%	0.0052%	0.0249%	0.0388%	0.0443%	0.0277%	0.0106%
Males	0.0082%	0.0018%	0.0017%	0.0007%	0.0012%	0.0079%	0.0403%	0.0556%	0.0563%	0.0398%	0.0152%
Females	0.0042%	0.0018%	0.0010%	-	-	0.0024%	0.0104%	0.0247%	0.0357%	0.0208%	0.0067%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XV-22
LUNG AND BRONCHUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
Race/Sex										
All Races ^b	206,013	69,132	37,536	19,609	10,036	5,948	312,681	348,273	11,808	360,081
Males	99,962	31,236	17,208	9,534	4,735	2,986	148,406	165,661	6,765	172,426
Females	106,051	37,896	20,328	10,075	5,301	2,962	164,275	182,612	5,043	187,655
White ^b	177,913	60,655	33,309	17,471	8,821	5,266	271,877	303,433	5,759	309,192
Males	85,424	26,799	15,029	8,320	4,135	2,604	127,252	142,310	2,817	145,127
Females	92,489	33,856	18,280	9,151	4,686	2,662	144,625	161,123	2,942	164,065
Black ^b	20,971	6,356	3,320	1,677	950	517	30,646	33,791	541	34,332
Males	10,688	3,231	1,679	944	473	309	15,598	17,324	302	17,626
Females	10,283	3,125	1,641	733	477	208	15,048	16,467	239	16,706
Asian/Pacific Islander ^c	5,287	1,477	620	+	+	+	7,384	+	+	+
Males	2,887	837	357	+	+	+	4,081	+	+	+
Females	2,400	640	263	+	+	+	3,303	+	+	+
Hispanic ^d	6,338	1,679	940	+	+	+	8,956	+	+	+
Males	3,127	871	429	+	+	+	4,427	+	+	+
Females	3,211	808	511	+	+	+	4,529	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0900%	-	0.0003%	0.0014%	0.0051%	0.0272%	0.1104%	0.3550%	0.6240%	0.4906%	0.0967%
Males	0.0870%	-	0.0003%	0.0012%	0.0045%	0.0241%	0.1122%	0.3748%	0.6954%	0.5974%	0.1054%
Females	0.0930%	-	0.0002%	0.0015%	0.0057%	0.0302%	0.1087%	0.3372%	0.5693%	0.4317%	0.0905%
White ^c	0.0970%	-	0.0003%	0.0015%	0.0054%	0.0266%	0.1078%	0.3688%	0.6505%	0.4981%	0.0992%
Males	0.0907%	-	0.0004%	0.0013%	0.0048%	0.0235%	0.1047%	0.3730%	0.7043%	0.5921%	0.1048%
Females	0.1032%	-	-	0.0018%	0.0060%	0.0298%	0.1107%	0.3649%	0.6084%	0.4468%	0.0957%
Black ^c	0.0762%	-	-	0.0009%	0.0047%	0.0386%	0.1667%	0.4080%	0.6350%	0.4155%	0.1064%
Males	0.0811%	-	-	-	0.0037%	0.0341%	0.1963%	0.5083%	0.7877%	0.5571%	0.1299%
Females	0.0718%	-	-	-	0.0055%	0.0426%	0.1421%	0.3295%	0.5339%	0.3472%	0.0904%
Asian/Pacific Islander ^c	0.0645%	-	-	-	0.0042%	0.0205%	0.0819%	0.2336%	0.4479%	0.4696%	0.0720%
Males	0.0736%	-	-	-	0.0039%	0.0201%	0.0958%	0.2991%	0.5798%	0.6511%	0.0920%
Females	0.0560%	-	-	-	0.0045%	0.0208%	0.0698%	0.1786%	0.3520%	0.3490%	0.0570%
Hispanic ^d	0.0209%	-	-	0.0013%	0.0029%	0.0125%	0.0423%	0.1445%	0.2846%	0.2750%	0.0440%
Males	0.0198%	-	-	0.0008%	0.0032%	0.0114%	0.0439%	0.1504%	0.3331%	0.3663%	0.0504%
Females	0.0221%	-	-	0.0020%	0.0025%	0.0136%	0.0408%	0.1395%	0.2495%	0.2224%	0.0399%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XVI-16
MELANOMA OF THE SKIN (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	221,975	164,459	113,357	86,036	56,675	35,271	499,791	677,773	45,643	723,416
Males	116,685	83,923	56,297	38,871	24,087	14,113	256,905	333,977	15,437	349,414
Females	105,290	80,536	57,060	47,165	32,588	21,158	242,886	343,796	30,206	374,002
White ^b	214,382	157,948	108,815	81,798	54,869	33,652	481,145	651,464	43,173	694,637
Males	112,902	80,546	54,250	37,170	23,376	13,404	247,698	321,648	14,735	336,383
Females	101,480	77,402	54,565	44,628	31,493	20,248	233,447	329,816	28,438	358,254
Black ^b	758	681	323	307	110	122	1,763	2,304	120	2,424
Males	357	284	150	81	24	37	792	935	45	980
Females	401	397	173	226	86	85	971	1,369	75	1,444
Asian/Pacific Islander ^c	657	347	178	+	+	+	1,182	+	+	+
Males	286	149	70	+	+	+	505	+	+	+
Females	371	198	108	+	+	+	677	+	+	+
Hispanic ^d	3,901	2,715	1,850	+	+	+	8,465	+	+	+
Males	1,457	1,004	678	+	+	+	3,138	+	+	+
Females	2,444	1,711	1,172	+	+	+	5,327	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.1426%	0.0004%	0.0033%	0.0285%	0.0864%	0.1711%	0.2746%	0.3769%	0.4778%	0.4631%	0.1453%
Males	0.1498%	0.0004%	0.0031%	0.0188%	0.0606%	0.1510%	0.2950%	0.4804%	0.6739%	0.7216%	0.1671%
Females	0.1356%	0.0004%	0.0036%	0.0387%	0.1129%	0.1911%	0.2552%	0.2838%	0.3278%	0.3205%	0.1312%
White ^c	0.1784%	0.0004%	0.0042%	0.0360%	0.1088%	0.2116%	0.3351%	0.4545%	0.5712%	0.5336%	0.1762%
Males	0.1861%	0.0004%	0.0038%	0.0230%	0.0742%	0.1836%	0.3557%	0.5730%	0.7970%	0.8421%	0.1995%
Females	0.1707%	0.0004%	0.0046%	0.0502%	0.1460%	0.2404%	0.3151%	0.3450%	0.3945%	0.3655%	0.1620%
Black ^c	0.0045%	-	-	0.0010%	0.0032%	0.0044%	0.0077%	0.0198%	0.0211%	0.0274%	0.0059%
Males	0.0042%	-	-	-	0.0024%	0.0040%	0.0072%	0.0232%	0.0280%	0.0335%	0.0065%
Females	0.0048%	-	-	-	0.0039%	0.0048%	0.0082%	0.0171%	0.0166%	0.0244%	0.0056%
Asian/Pacific Islander ^c	0.0093%	-	0.0009%	0.0018%	0.0079%	0.0095%	0.0163%	0.0243%	0.0290%	0.0362%	0.0096%
Males	0.0083%	-	-	0.0021%	0.0044%	0.0072%	0.0159%	0.0241%	0.0385%	0.0349%	0.0093%
Females	0.0103%	-	-	0.0015%	0.0112%	0.0117%	0.0167%	0.0245%	0.0221%	0.0371%	0.0100%
Hispanic ^d	0.0202%	-	0.0007%	0.0049%	0.0170%	0.0374%	0.0584%	0.0734%	0.1080%	0.1513%	0.0326%
Males	0.0143%	-	-	0.0024%	0.0070%	0.0235%	0.0439%	0.0720%	0.1190%	0.1973%	0.0291%
Females	0.0264%	-	0.0008%	0.0079%	0.0284%	0.0520%	0.0721%	0.0745%	0.1000%	0.1248%	0.0372%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XVII-11
MESOTHELIOMAEstimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	2,847	646	368	214	178	110	3,860	4,361	+	+
Males	2,040	312	104	74	93	24	2,456	2,646	+	+
Females	807	334	264	140	85	86	1,404	1,715	+	+
White ^b	2,583	565	313	199	163	86	3,460	3,908	+	+
Males	1,868	279	80	59	78	11	2,226	2,374	+	+
Females	715	286	233	140	85	75	1,234	1,534	+	+
Black ^b	177	49	36	11	11	24	263	308	+	+
Males	118	12	24	11	11	13	154	189	+	+
Females	59	37	12	0	0	11	109	119	+	+
Asian/Pacific Islander ^c	49	15	4	+	+	+	68	+	+	+
Males	31	12	0	+	+	+	43	+	+	+
Females	18	3	4	+	+	+	25	+	+	+
Hispanic ^d	206	48	15	+	+	+	270	+	+	+
Males	139	34	0	+	+	+	173	+	+	+
Females	67	14	15	+	+	+	97	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0011%	-	-	-	0.0001%	0.0006%	0.0017%	0.0039%	0.0074%	0.0054%	0.0012%
Males	0.0015%	-	-	-	-	0.0005%	0.0019%	0.0057%	0.0132%	0.0103%	0.0018%
Females	0.0008%	-	-	-	0.0002%	0.0007%	0.0015%	0.0022%	0.0030%	0.0027%	0.0007%
White ^c	0.0013%	-	-	-	0.0002%	0.0006%	0.0019%	0.0041%	0.0083%	0.0060%	0.0013%
Males	0.0017%	-	-	-	-	0.0005%	0.0021%	0.0058%	0.0142%	0.0118%	0.0020%
Females	0.0009%	-	-	-	-	0.0008%	0.0016%	0.0024%	0.0036%	0.0029%	0.0008%
Black ^c	0.0007%	-	-	-	-	-	0.0016%	0.0050%	0.0039%	-	0.0009%
Males	0.0009%	-	-	-	-	-	-	0.0070%	0.0082%	-	0.0014%
Females	0.0005%	-	-	-	-	-	-	0.0034%	-	-	0.0006%
Asian/Pacific Islander ^c	0.0006%	-	-	-	-	-	-	0.0015%	0.0035%	-	0.0006%
Males	0.0008%	-	-	-	-	-	-	-	0.0082%	-	0.0010%
Females	0.0004%	-	-	-	-	-	-	-	-	-	0.0003%
Hispanic ^d	0.0006%	-	-	-	-	0.0007%	0.0022%	0.0022%	0.0106%	-	0.0013%
Males	0.0008%	-	-	-	-	-	0.0027%	0.0046%	0.0161%	-	0.0018%
Females	0.0005%	-	-	-	-	-	0.0018%	-	0.0067%	-	0.0008%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XVIII-16
MYELOMA (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
Race/Sex										
All Races ^b	37,951	11,508	4,195	1,517	505	209	53,656	55,888	312	56,200
Males	20,764	6,583	2,371	891	272	63	29,719	30,945	121	31,066
Females	17,187	4,925	1,824	626	233	146	23,937	24,943	191	25,134
White ^b	29,837	9,005	3,188	1,101	316	171	42,030	43,618	77	43,695
Males	16,767	5,284	1,838	665	162	60	23,889	24,775	0 ⁱ	24,775
Females	13,070	3,721	1,350	436	154	111	18,141	18,843	77 ⁱ	18,920
Black ^b	6,614	2,103	851	343	171	26	9,567	10,107	+	+
Males	3,264	1,062	470	203	101	0	4,796	5,100	+	+
Females	3,350	1,041	381	140	70	26	4,771	5,007	+	+
Asian/Pacific Islander ^c	832	240	97	+	+	+	1,169	+	+	+
Males	402	141	50	+	+	+	593	+	+	+
Females	430	99	47	+	+	+	576	+	+	+
Hispanic ^d	2,506	779	233	+	+	+	3,519	+	+	+
Males	1,337	404	140	+	+	+	1,881	+	+	+
Females	1,169	375	93	+	+	+	1,638	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0165%	-	-	0.0002%	0.0019%	0.0080%	0.0277%	0.0637%	0.0910%	0.0808%	0.0173%
Males	0.0183%	-	-	0.0003%	0.0025%	0.0092%	0.0324%	0.0796%	0.1117%	0.1027%	0.0212%
Females	0.0147%	-	-	-	0.0012%	0.0069%	0.0232%	0.0494%	0.0751%	0.0687%	0.0142%
White ^c	0.0161%	-	-	0.0001%	0.0017%	0.0074%	0.0249%	0.0600%	0.0860%	0.0750%	0.0161%
Males	0.0181%	-	-	-	0.0024%	0.0084%	0.0298%	0.0762%	0.1077%	0.0978%	0.0201%
Females	0.0140%	-	-	-	0.0009%	0.0064%	0.0202%	0.0450%	0.0690%	0.0626%	0.0128%
Black ^c	0.0249%	-	-	-	0.0047%	0.0149%	0.0588%	0.1245%	0.1735%	0.1514%	0.0340%
Males	0.0265%	-	-	-	0.0051%	0.0178%	0.0676%	0.1590%	0.2009%	0.1744%	0.0404%
Females	0.0235%	-	-	-	0.0042%	0.0124%	0.0514%	0.0975%	0.1554%	0.1402%	0.0292%
Asian/Pacific Islander ^c	0.0100%	-	-	-	-	0.0048%	0.0180%	0.0365%	0.0547%	0.0665%	0.0109%
Males	0.0104%	-	-	-	-	0.0052%	0.0207%	0.0379%	0.0641%	0.0833%	0.0126%
Females	0.0096%	-	-	-	-	0.0043%	0.0156%	0.0354%	0.0478%	0.0553%	0.0096%
Hispanic ^d	0.0083%	-	-	-	0.0022%	0.0075%	0.0234%	0.0595%	0.0870%	0.0712%	0.0161%
Males	0.0085%	-	-	-	0.0032%	0.0083%	0.0247%	0.0706%	0.0964%	0.1025%	0.0189%
Females	0.0080%	-	-	-	0.0010%	0.0066%	0.0221%	0.0502%	0.0801%	0.0533%	0.0139%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h i} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIX-20
NON-HODGKIN LYMPHOMA (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	173,017	100,265	56,408	32,341	17,996	9,412	329,689	389,438	12,259	401,697
Males	89,995	51,648	29,367	16,592	9,217	4,558	171,009	201,376	6,445	207,821
Females	83,022	48,617	27,041	15,749	8,779	4,854	158,680	188,062	5,814	193,876
White ^b	153,057	89,140	50,726	29,470	16,353	8,734	292,923	347,479	12,455	359,934
Males	79,876	45,756	26,361	15,197	8,400	4,257	151,993	179,846	6,944	186,790
Females	73,181	43,384	24,365	14,273	7,953	4,477	140,930	167,633	5,511	173,144
Black ^b	12,833	7,065	3,581	1,858	1,067	498	23,479	26,901	921	27,822
Males	6,443	3,773	1,853	912	509	207	12,069	13,697	407	14,104
Females	6,390	3,292	1,728	946	558	291	11,410	13,204	514	13,718
Asian/Pacific Islander ^c	4,285	2,295	1,254	+	+	+	7,836	+	+	+
Males	2,186	1,197	621	+	+	+	4,005	+	+	+
Females	2,099	1,098	633	+	+	+	3,831	+	+	+
Hispanic ^d	10,689	5,477	2,903	+	+	+	19,068	+	+	+
Males	5,761	2,836	1,424	+	+	+	10,020	+	+	+
Females	4,928	2,641	1,479	+	+	+	9,048	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.1006%	0.0021%	0.0112%	0.0187%	0.0364%	0.0787%	0.1604%	0.3012%	0.4474%	0.4835%	0.1043%
Males	0.1059%	0.0028%	0.0153%	0.0230%	0.0418%	0.0925%	0.1828%	0.3385%	0.4973%	0.5542%	0.1190%
Females	0.0953%	0.0014%	0.0069%	0.0140%	0.0308%	0.0650%	0.1392%	0.2676%	0.4092%	0.4445%	0.0917%
White ^c	0.1115%	0.0022%	0.0116%	0.0193%	0.0379%	0.0826%	0.1706%	0.3249%	0.4853%	0.5131%	0.1114%
Males	0.1167%	0.0028%	0.0159%	0.0238%	0.0440%	0.0964%	0.1939%	0.3638%	0.5367%	0.5863%	0.1265%
Females	0.1063%	0.0016%	0.0072%	0.0145%	0.0315%	0.0685%	0.1477%	0.2889%	0.4451%	0.4732%	0.0980%
Black ^c	0.0598%	0.0019%	0.0082%	0.0187%	0.0360%	0.0745%	0.1346%	0.2137%	0.2478%	0.2167%	0.0733%
Males	0.0654%	0.0021%	0.0126%	0.0228%	0.0410%	0.0916%	0.1608%	0.2442%	0.2654%	0.2509%	0.0851%
Females	0.0548%	0.0016%	0.0038%	0.0149%	0.0317%	0.0594%	0.1130%	0.1898%	0.2362%	0.2003%	0.0638%
Asian/Pacific Islander ^c	0.0643%	0.0018%	0.0123%	0.0146%	0.0242%	0.0511%	0.0996%	0.1784%	0.2942%	0.3498%	0.0685%
Males	0.0671%	0.0028%	0.0152%	0.0192%	0.0263%	0.0604%	0.1036%	0.1919%	0.3350%	0.4053%	0.0773%
Females	0.0616%	-	0.0092%	0.0101%	0.0223%	0.0425%	0.0961%	0.1670%	0.2647%	0.3129%	0.0614%
Hispanic ^d	0.0451%	0.0020%	0.0088%	0.0123%	0.0273%	0.0558%	0.1188%	0.2122%	0.3499%	0.3409%	0.0771%
Males	0.0456%	0.0024%	0.0124%	0.0153%	0.0313%	0.0627%	0.1307%	0.2136%	0.3696%	0.3658%	0.0831%
Females	0.0445%	0.0015%	0.0050%	0.0087%	0.0226%	0.0485%	0.1077%	0.2110%	0.3357%	0.3266%	0.0714%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XX-16
 CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
Race/Sex										
All Races ^b	86,675	53,639	34,936	23,293	14,733	10,344	175,251	223,620	16,556	240,176
Males	58,447	34,569	22,553	14,085	8,536	6,197	115,569	144,387	9,772	154,159
Females	28,228	19,070	12,383	9,208	6,197	4,147	59,682	79,233	6,784	86,017
White ^b	74,568	46,949	30,741	20,893	13,144	9,147	152,259	195,443	14,063	209,506
Males	50,805	30,734	20,145	12,834	7,822	5,644	101,684	127,984	8,385	136,369
Females	23,763	16,215	10,596	8,059	5,322	3,503	50,575	67,459	5,678	73,137
Black ^b	7,446	3,675	2,426	1,442	936	683	13,546	16,606	824	17,430
Males	4,649	2,054	1,326	758	379	252	8,028	9,417	325	9,742
Females	2,797	1,621	1,100	684	557	431	5,518	7,189	499	7,688
Asian/Pacific Islander ^c	3,074	1,909	1,124	+	+	+	6,106	+	+	+
Males	1,933	1,154	678	+	+	+	3,764	+	+	+
Females	1,141	755	446	+	+	+	2,342	+	+	+
Hispanic ^d	3,721	2,315	1,397	+	+	+	7,433	+	+	+
Males	2,395	1,468	765	+	+	+	4,629	+	+	+
Females	1,326	847	632	+	+	+	2,804	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0542%	0.0003%	0.0019%	0.0058%	0.0137%	0.0423%	0.1109%	0.1844%	0.2284%	0.2059%	0.0557%
Males	0.0719%	0.0002%	0.0017%	0.0052%	0.0141%	0.0556%	0.1641%	0.2755%	0.3382%	0.3056%	0.0800%
Females	0.0368%	0.0004%	0.0021%	0.0064%	0.0134%	0.0291%	0.0605%	0.1025%	0.1444%	0.1509%	0.0354%
White ^c	0.0575%	0.0003%	0.0019%	0.0056%	0.0121%	0.0403%	0.1120%	0.1928%	0.2423%	0.2146%	0.0570%
Males	0.0769%	-	0.0015%	0.0047%	0.0125%	0.0537%	0.1679%	0.2871%	0.3594%	0.3253%	0.0825%
Females	0.0381%	0.0003%	0.0023%	0.0066%	0.0117%	0.0266%	0.0573%	0.1058%	0.1506%	0.1543%	0.0352%
Black ^c	0.0356%	-	0.0022%	0.0056%	0.0135%	0.0367%	0.1035%	0.1498%	0.1592%	0.1076%	0.0442%
Males	0.0455%	-	0.0034%	0.0066%	0.0123%	0.0419%	0.1508%	0.2360%	0.2361%	0.1351%	0.0620%
Females	0.0266%	-	-	0.0047%	0.0146%	0.0321%	0.0643%	0.0823%	0.1084%	0.0943%	0.0308%
Asian/Pacific Islander ^c	0.0489%	-	0.0016%	0.0066%	0.0203%	0.0568%	0.1018%	0.1458%	0.1676%	0.1754%	0.0505%
Males	0.0620%	-	-	0.0071%	0.0224%	0.0764%	0.1392%	0.2130%	0.2299%	0.2295%	0.0683%
Females	0.0366%	-	0.0020%	0.0061%	0.0184%	0.0388%	0.0692%	0.0893%	0.1224%	0.1395%	0.0359%
Hispanic ^d	0.0174%	-	0.0010%	0.0029%	0.0072%	0.0207%	0.0546%	0.1013%	0.1408%	0.1360%	0.0312%
Males	0.0209%	-	0.0009%	0.0022%	0.0070%	0.0241%	0.0768%	0.1459%	0.2063%	0.2069%	0.0436%
Females	0.0138%	-	0.0012%	0.0037%	0.0074%	0.0172%	0.0337%	0.0638%	0.0934%	0.0951%	0.0216%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXI-12
OVARY CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
 By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race</u>										
All Races ^b	53,959	29,988	22,631	18,687	14,732	11,235	106,578	151,232	23,004	174,236
White ^b	47,957	26,541	20,135	16,969	13,263	9,994	94,633	134,859	19,944	154,803
Black ^b	3,431	1,927	1,493	1,122	947	912	6,850	9,831	1,987	11,818
Asian/Pacific Islander ^c	1,899	1,060	750	+	+	+	3,710	+	+	+
Hispanic ^d	3,936	1,956	1,356	+	+	+	7,248	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
 By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>												
All Races ^c	0.0671%	0.0004%	0.0053%	0.0160%	0.0268%	0.0636%	0.1349%	0.2043%	0.2159%	0.1635%	0.0648%	
White ^c	0.0741%	0.0005%	0.0054%	0.0162%	0.0268%	0.0670%	0.1446%	0.2245%	0.2345%	0.1740%	0.0693%	
Black ^c	0.0346%	-	0.0048%	0.0105%	0.0233%	0.0419%	0.0747%	0.1054%	0.1249%	0.1174%	0.0394%	
Asian/Pacific Islander ^c	0.0563%	-	0.0055%	0.0205%	0.0286%	0.0614%	0.1242%	0.1507%	0.1564%	0.0949%	0.0543%	
Hispanic ^d	0.0356%	0.0007%	0.0057%	0.0168%	0.0218%	0.0473%	0.1043%	0.1580%	0.1794%	0.1310%	0.0519%	

- ^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
 US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- ^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.
- ^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- ^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.
- ^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table XXII-16
PANCREAS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	21,905	3,471	1,875	930	569	276	27,250	29,026	419	29,445
Males	10,950	1,575	849	401	231	171	13,374	14,178	177	14,355
Females	10,955	1,896	1,026	529	338	105	13,876	14,848	242	15,090
White ^b	18,683	2,711	1,549	607	463	245	22,943	24,259	341	24,600
Males	9,623	1,333	710	282	202	140	11,666	12,290	158	12,448
Females	9,060	1,378	839	325	261	105	11,277	11,969	183	12,152
Black ^b	2,372	444	203	272	70	27	3,019	3,387	187	3,574
Males	962	114	75	116	13	27	1,151	1,307	70	1,377
Females	1,410	330	128	156	57	0	1,868	2,080	117	2,197
Asian/Pacific Islander ^c	679	160	52	+	+	+	892	+	+	+
Males	315	58	24	+	+	+	398	+	+	+
Females	364	102	28	+	+	+	494	+	+	+
Hispanic ^d	1,272	217	167	+	+	+	1,656	+	+	+
Males	629	63	76	+	+	+	769	+	+	+
Females	643	154	91	+	+	+	887	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0085%	-	0.0002%	0.0004%	0.0013%	0.0051%	0.0142%	0.0295%	0.0448%	0.0426%	0.0089%
Males	0.0084%	-	-	0.0003%	0.0011%	0.0050%	0.0164%	0.0327%	0.0506%	0.0458%	0.0097%
Females	0.0085%	-	0.0003%	0.0006%	0.0016%	0.0053%	0.0121%	0.0266%	0.0404%	0.0409%	0.0082%
White ^c	0.0088%	-	0.0002%	0.0005%	0.0014%	0.0048%	0.0143%	0.0298%	0.0446%	0.0407%	0.0088%
Males	0.0090%	-	-	0.0003%	0.0010%	0.0050%	0.0170%	0.0339%	0.0517%	0.0445%	0.0099%
Females	0.0086%	-	0.0003%	0.0006%	0.0018%	0.0046%	0.0116%	0.0260%	0.0390%	0.0387%	0.0079%
Black ^c	0.0075%	-	-	-	0.0013%	0.0061%	0.0174%	0.0297%	0.0540%	0.0517%	0.0102%
Males	0.0061%	-	-	-	0.0015%	0.0050%	0.0193%	0.0297%	0.0431%	0.0247%	0.0088%
Females	0.0088%	-	-	-	-	0.0070%	0.0158%	0.0297%	0.0612%	0.0647%	0.0110%
Asian/Pacific Islander ^c	0.0076%	-	-	-	0.0009%	0.0058%	0.0107%	0.0271%	0.0380%	0.0527%	0.0083%
Males	0.0070%	-	-	-	-	0.0042%	0.0097%	0.0272%	0.0435%	0.0656%	0.0086%
Females	0.0081%	-	-	-	-	0.0073%	0.0116%	0.0271%	0.0341%	0.0442%	0.0081%
Hispanic ^d	0.0039%	-	-	0.0003%	0.0012%	0.0034%	0.0106%	0.0279%	0.0358%	0.0382%	0.0074%
Males	0.0035%	-	-	-	-	0.0029%	0.0108%	0.0306%	0.0414%	0.0410%	0.0080%
Females	0.0043%	-	-	-	0.0018%	0.0039%	0.0105%	0.0257%	0.0318%	0.0366%	0.0072%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXIII-12
PROSTATE CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
 By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race</u>										
All Races ^b	939,384	648,082	398,529	87,363	22,685	6,915	1,985,995	2,102,958	3,541	2,106,499
White ^b	783,783	548,161	351,255	79,693	20,616	6,271	1,683,200	1,789,780	3,856	1,793,636
Black ^b	120,582	79,242	38,475	6,415	1,814	560	238,299	247,089	311	247,400
Asian/Pacific Islander ^c	16,643	9,136	4,383	+	+	+	30,162	+	+	+
Hispanic ^d	47,271	27,667	13,419	+	+	+	88,358	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
 By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	1.1821%	-	0.0002%	0.0002%	0.0012%	0.0652%	0.9189%	4.5830%	10.7558%	12.6607%	1.5073%
White ^c	1.2449%	-	0.0002%	-	0.0010%	0.0582%	0.8855%	4.5615%	10.8543%	12.5270%	1.5017%
Black ^c	1.2525%	-	-	-	0.0030%	0.1606%	1.8297%	7.7507%	15.7410%	15.9066%	2.2660%
Asian/Pacific Islander ^c	0.5771%	-	-	-	-	0.0181%	0.2794%	1.8531%	5.5462%	8.4108%	0.7795%
Hispanic ^d	0.3934%	-	-	-	0.0009%	0.0326%	0.5447%	3.1912%	8.3546%	10.1096%	1.1377%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXIV-16
STOMACH CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	31,518	12,494	7,436	5,263	2,692	1,482	51,447	60,885	2,326	63,211
Males	18,834	7,223	3,806	2,668	1,407	851	29,862	34,789	1,329	36,118
Females	12,684	5,271	3,630	2,595	1,285	631	21,585	26,096	997	27,093
White ^b	24,038	9,323	5,610	4,072	2,022	1,233	38,972	46,299	1,886	48,185
Males	14,804	5,403	2,840	2,069	1,098	697	23,047	26,911	1,027	27,938
Females	9,234	3,920	2,770	2,003	924	536	15,925	19,388	859	20,247
Black ^b	4,559	1,829	920	699	415	132	7,308	8,554	303	8,857
Males	2,357	959	512	358	215	93	3,828	4,494	176	4,670
Females	2,202	870	408	341	200	39	3,480	4,060	127	4,187
Asian/Pacific Islander ^c	2,669	1,215	803	+	+	+	4,687	+	+	+
Males	1,462	713	439	+	+	+	2,614	+	+	+
Females	1,207	502	364	+	+	+	2,073	+	+	+
Hispanic ^d	3,749	1,450	786	+	+	+	5,985	+	+	+
Males	2,038	687	400	+	+	+	3,126	+	+	+
Females	1,711	763	386	+	+	+	2,859	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0196%	-	0.0001%	0.0005%	0.0028%	0.0100%	0.0258%	0.0575%	0.1123%	0.1488%	0.0207%
Males	0.0224%	-	-	0.0003%	0.0029%	0.0108%	0.0313%	0.0787%	0.1511%	0.2001%	0.0270%
Females	0.0168%	-	-	0.0006%	0.0027%	0.0092%	0.0205%	0.0383%	0.0827%	0.1205%	0.0159%
White ^c	0.0163%	-	0.0001%	0.0004%	0.0026%	0.0081%	0.0208%	0.0476%	0.0860%	0.1138%	0.0164%
Males	0.0189%	-	-	0.0003%	0.0026%	0.0090%	0.0257%	0.0654%	0.1178%	0.1534%	0.0216%
Females	0.0138%	-	-	0.0005%	0.0026%	0.0072%	0.0160%	0.0311%	0.0611%	0.0922%	0.0123%
Black ^c	0.0183%	-	-	-	0.0026%	0.0147%	0.0367%	0.0672%	0.1353%	0.1838%	0.0259%
Males	0.0202%	-	-	-	0.0042%	0.0166%	0.0430%	0.0940%	0.1802%	0.2185%	0.0329%
Females	0.0166%	-	-	-	-	0.0131%	0.0315%	0.0462%	0.1055%	0.1671%	0.0209%
Asian/Pacific Islander ^c	0.0415%	-	-	-	0.0042%	0.0178%	0.0475%	0.1148%	0.2701%	0.4139%	0.0469%
Males	0.0476%	-	-	-	0.0037%	0.0178%	0.0573%	0.1586%	0.3629%	0.5289%	0.0605%
Females	0.0358%	-	-	-	0.0046%	0.0177%	0.0389%	0.0779%	0.2027%	0.3375%	0.0368%
Hispanic ^d	0.0141%	-	-	0.0006%	0.0043%	0.0133%	0.0353%	0.0787%	0.1457%	0.2066%	0.0282%
Males	0.0141%	-	-	-	0.0041%	0.0116%	0.0371%	0.1062%	0.1846%	0.2439%	0.0338%
Females	0.0140%	-	-	-	0.0047%	0.0151%	0.0336%	0.0556%	0.1176%	0.1852%	0.0242%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXV-12
TESTIS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
 By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race</u>										
All Races ^b	39,561	34,456	31,670	27,089	20,164	12,903	105,687	165,843	18,231	184,074
White ^b	37,315	32,470	30,249	26,111	19,493	12,393	100,034	158,031	17,493	175,524
Black ^b	970	895	477	436	307	229	2,342	3,314	181	3,495
Asian/Pacific Islander ^c	636	509	454	+	+	+	1,600	+	+	+
Hispanic ^d	4,103	2,979	2,385	+	+	+	9,467	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
 By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.0683%	0.0020%	0.0058%	0.0580%	0.1406%	0.1535%	0.0824%	0.0341%	0.0167%	0.0097%	0.0669%
White ^c	0.0827%	0.0020%	0.0065%	0.0708%	0.1744%	0.1841%	0.0973%	0.0401%	0.0200%	0.0110%	0.0809%
Black ^c	0.0138%	-	0.0034%	0.0090%	0.0274%	0.0339%	0.0216%	0.0070%	-	-	0.0141%
Asian/Pacific Islander ^c	0.0233%	0.0037%	0.0043%	0.0209%	0.0379%	0.0538%	0.0292%	0.0065%	0.0052%	-	0.0221%
Hispanic ^d	0.0437%	0.0025%	0.0064%	0.0502%	0.0961%	0.0917%	0.0433%	0.0189%	0.0079%	-	0.0433%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXVI-16
 THYROID CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	111,720	75,355	54,707	41,816	31,647	28,938	241,781	344,182	44,204	388,386
Males	24,952	16,619	12,751	9,163	6,611	6,520	54,322	76,616	10,304	86,920
Females	86,768	58,736	41,956	32,653	25,036	22,418	187,459	267,566	33,900	301,466
White ^b	97,772	66,163	48,807	37,480	28,642	26,087	212,741	304,950	40,017	344,967
Males	22,349	15,081	11,638	8,451	5,899	6,057	49,068	69,475	8,737	78,212
Females	75,423	51,082	37,169	29,029	22,743	20,030	163,673	235,475	31,280	266,755
Black ^b	6,893	4,203	2,917	1,867	1,447	1,408	14,014	18,734	1,822	20,556
Males	1,350	625	615	308	330	137	2,591	3,365	202	3,567
Females	5,543	3,578	2,302	1,559	1,117	1,271	11,423	15,369	1,620	16,989
Asian/Pacific Islander ^c	4,965	3,328	2,238	+	+	+	10,531	+	+	+
Males	927	589	377	+	+	+	1,893	+	+	+
Females	4,038	2,739	1,861	+	+	+	8,638	+	+	+
Hispanic ^d	9,996	6,145	4,401	+	+	+	20,542	+	+	+
Males	1,773	1,084	765	+	+	+	3,622	+	+	+
Females	8,223	5,061	3,636	+	+	+	16,920	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0785%	0.0001%	0.0038%	0.0332%	0.0921%	0.1340%	0.1511%	0.1535%	0.1293%	0.0928%	0.0783%
Males	0.0349%	-	0.0012%	0.0091%	0.0313%	0.0514%	0.0738%	0.0951%	0.0848%	0.0677%	0.0364%
Females	0.1212%	-	0.0065%	0.0588%	0.1547%	0.2163%	0.2246%	0.2061%	0.1634%	0.1067%	0.1187%
White ^c	0.0845%	0.0001%	0.0041%	0.0367%	0.1018%	0.1464%	0.1578%	0.1561%	0.1302%	0.0914%	0.0831%
Males	0.0386%	-	0.0014%	0.0097%	0.0349%	0.0570%	0.0796%	0.0995%	0.0895%	0.0693%	0.0392%
Females	0.1304%	-	0.0069%	0.0665%	0.1738%	0.2381%	0.2342%	0.2085%	0.1621%	0.1035%	0.1272%
Black ^c	0.0357%	-	0.0015%	0.0129%	0.0405%	0.0569%	0.0835%	0.1093%	0.0807%	0.0558%	0.0408%
Males	0.0143%	-	-	0.0043%	0.0129%	0.0180%	0.0368%	0.0603%	0.0544%	0.0350%	0.0184%
Females	0.0551%	-	0.0020%	0.0210%	0.0648%	0.0910%	0.1222%	0.1475%	0.0981%	0.0658%	0.0593%
Asian/Pacific Islander ^c	0.0798%	-	0.0040%	0.0319%	0.0797%	0.1250%	0.1571%	0.1656%	0.1501%	0.1263%	0.0787%
Males	0.0299%	-	-	0.0103%	0.0249%	0.0433%	0.0599%	0.0877%	0.0687%	0.0719%	0.0313%
Females	0.1265%	-	0.0079%	0.0532%	0.1315%	0.1997%	0.2421%	0.2310%	0.2092%	0.1625%	0.1203%
Hispanic ^d	0.0492%	-	0.0039%	0.0249%	0.0638%	0.1071%	0.1305%	0.1447%	0.1270%	0.1056%	0.0664%
Males	0.0166%	-	0.0009%	0.0051%	0.0200%	0.0283%	0.0525%	0.0833%	0.0618%	0.0687%	0.0262%
Females	0.0837%	-	0.0070%	0.0487%	0.1144%	0.1900%	0.2036%	0.1962%	0.1740%	0.1269%	0.1058%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXVII-16
URINARY BLADDER CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
Race/Sex										
All Races ^b	187,611	126,491	84,612	52,902	32,195	16,718	398,715	500,529	20,623	521,152
Males	140,466	93,789	62,886	39,033	23,367	11,352	297,142	370,894	13,400	384,294
Females	47,145	32,702	21,726	13,869	8,828	5,366	101,573	129,635	7,223	136,858
White ^b	174,922	118,854	80,014	50,515	30,891	15,903	373,790	471,099	16,204	487,303
Males	131,715	88,351	59,508	37,286	22,473	10,846	279,574	350,179	10,350	360,529
Females	43,207	30,503	20,506	13,229	8,418	5,057	94,216	120,920	5,854	126,774
Black ^b	7,779	4,423	2,940	1,538	843	414	15,143	17,937	331	18,268
Males	4,884	3,064	2,119	1,136	555	265	10,067	12,022	265	12,287
Females	2,895	1,359	821	402	288	149	5,076	5,915	66	5,981
Asian/Pacific Islander ^c	2,588	1,518	972	+	+	+	5,078	+	+	+
Males	1,999	1,099	737	+	+	+	3,835	+	+	+
Females	589	419	235	+	+	+	1,243	+	+	+
Hispanic ^d	5,767	3,355	2,273	+	+	+	11,395	+	+	+
Males	4,326	2,542	1,698	+	+	+	8,566	+	+	+
Females	1,441	813	575	+	+	+	2,829	+	+	+

Estimated prevalence percent^a on January 1, 2005, of the SEER 11 population diagnosed in the previous 15 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.1143%	0.0001%	0.0003%	0.0019%	0.0069%	0.0294%	0.1119%	0.3610%	0.7631%	0.9913%	0.1228%
Males	0.1734%	0.0002%	0.0005%	0.0022%	0.0097%	0.0446%	0.1776%	0.5900%	1.3519%	1.9651%	0.2175%
Females	0.0564%	-	0.0002%	0.0015%	0.0040%	0.0144%	0.0494%	0.1549%	0.3125%	0.4539%	0.0533%
White ^c	0.1367%	0.0001%	0.0003%	0.0023%	0.0082%	0.0338%	0.1279%	0.4169%	0.8689%	1.0806%	0.1388%
Males	0.2061%	0.0002%	0.0004%	0.0027%	0.0113%	0.0501%	0.1994%	0.6729%	1.5294%	2.1762%	0.2446%
Females	0.0673%	-	-	0.0018%	0.0049%	0.0170%	0.0579%	0.1803%	0.3520%	0.4838%	0.0600%
Black ^c	0.0394%	-	-	0.0009%	0.0030%	0.0163%	0.0653%	0.1744%	0.3296%	0.4576%	0.0582%
Males	0.0563%	-	-	-	0.0042%	0.0271%	0.1117%	0.2846%	0.5574%	0.7743%	0.0982%
Females	0.0241%	-	-	-	0.0019%	0.0067%	0.0268%	0.0883%	0.1788%	0.3049%	0.0318%
Asian/Pacific Islander ^c	0.0462%	-	-	0.0007%	0.0021%	0.0122%	0.0379%	0.1218%	0.3185%	0.5692%	0.0533%
Males	0.0716%	-	-	-	0.0034%	0.0211%	0.0646%	0.2118%	0.5636%	1.0041%	0.0937%
Females	0.0224%	-	-	-	-	0.0041%	0.0145%	0.0462%	0.1405%	0.2800%	0.0234%
Hispanic ^d	0.0265%	-	-	0.0008%	0.0042%	0.0149%	0.0492%	0.1525%	0.3517%	0.5126%	0.0579%
Males	0.0384%	-	-	0.0008%	0.0050%	0.0203%	0.0764%	0.2580%	0.6487%	0.9954%	0.1039%
Females	0.0139%	-	-	0.0009%	0.0032%	0.0092%	0.0236%	0.0641%	0.1370%	0.2346%	0.0253%

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^{g h} (g) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXVIII-12
 CHILDHOOD CANCER (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts^a on January 1, 2005
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	25 to <30	0 to <15 ^e	0 to <30 ^e	>=30 ^g	Complete ^h
<u>Race/Sex</u>										
All Races ^b	59,044	49,156	44,096	37,983	31,355	27,278	152,296	248,912	+	+
Males	31,593	25,700	23,188	19,236	15,639	12,448	80,482	127,804	+	+
Females	27,451	23,456	20,908	18,747	15,716	14,830	71,814	121,108	+	+
White ^b	49,623	41,351	37,330	32,907	27,558	24,164	128,302	212,932	+	+
Males	26,872	21,488	19,892	16,696	13,913	11,156	68,251	110,016	+	+
Females	22,751	19,863	17,438	16,211	13,645	13,008	60,051	102,916	+	+
Black ^b	6,262	4,910	4,400	3,359	2,656	2,209	15,572	23,796	+	+
Males	3,043	2,620	2,113	1,628	1,205	883	7,776	11,492	+	+
Females	3,219	2,290	2,287	1,731	1,451	1,326	7,796	12,304	+	+
Asian/Pacific Islander ^c	1,944	1,699	1,410	+	+	+	5,053	+	+	+
Males	1,043	928	698	+	+	+	2,669	+	+	+
Females	901	771	712	+	+	+	2,384	+	+	+
Hispanic ^d	9,728	7,000	5,800	+	+	+	22,527	+	+	+
Males	5,288	3,619	3,250	+	+	+	12,157	+	+	+
Females	4,440	3,381	2,550	+	+	+	10,370	+	+	+

^a US 2005 cancer prevalence counts are based on 2005 cancer prevalence proportions from the SEER registries and 1/1/2005 US population estimates based on the average of 2004 and 2005 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 30 years for 1975-2005 SEER 9 data; 15 years for 1990-2005 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^{f g} (f) Cases diagnosed more than 30 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <30 and >=30.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.