

Table 8.51A Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Age Group and Demographic Characteristics: 2000 and 2001

Demographic Characteristic	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2000	2001	2000	2001	2000	2001	2000	2001
TOTAL	3,386 ^b	4,302	1,064 ^b	1,449	1,227 ^b	1,495	1,096 ^b	1,358
GENDER								
Male	1,558 ^b	2,034	565 ^b	782	562 ^b	710	432 ^b	543
Female	1,828 ^b	2,267	499 ^b	667	665 ^b	785	664 ^b	815
HISPANIC ORIGIN AND RACE								
Not Hispanic	2,922 ^b	3,717	915 ^b	1,263	1,068 ^b	1,294	939 ^b	1,160
White Only	2,312 ^b	2,884	698 ^b	969	868 ^b	996	746 ^b	919
Black Only	464 ^b	619	177 ^a	238	149 ^b	226	138	155
American Indian or Alaska Native Only	26	38	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian Only	69	82	19	24	27	23	23	35
More Than One Race	43 ^b	88	6 ^a	16	14 ^a	35	*	36
Hispanic	465 ^b	585	148	186	159	201	157	198

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis. See Table 8.55 for a list of the 10 specific sources of treatment.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.51B Percentages of Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Age Group and Demographic Characteristics: 2000 and 2001

Demographic Characteristic	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2000	2001	2000	2001	2000	2001	2000	2001
TOTAL	14.6 ^b	18.4	13.8 ^b	18.9	15.3 ^b	18.4	14.6 ^b	17.7
GENDER								
Male	13.1 ^b	17.0	14.2 ^b	19.9	13.5 ^b	17.2	11.5 ^a	13.9
Female	16.1 ^b	19.7	13.4 ^b	17.9	17.2 ^a	19.6	17.7 ^b	21.7
HISPANIC ORIGIN AND RACE								
Not Hispanic	14.7 ^b	18.5	13.9 ^b	19.3	15.4 ^b	18.6	14.6 ^b	17.8
White Only	15.2 ^b	18.9	13.9 ^b	19.5	16.3 ^b	18.8	15.3 ^b	18.5
Black Only	14.0 ^b	18.5	15.4 ^b	20.8	13.8 ^b	19.3	12.9	15.1
American Indian or Alaska Native Only	17.9	21.7	*	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*	*	*
Asian Only	8.3	9.8	7.7	9.9	9.1	8.0	8.1	11.4
More Than One Race	11.9 ^b	22.4	5.1 ^a	14.4	10.1 ^a	23.5	*	28.3
Hispanic	14.1 ^b	17.2	13.4	16.9	14.5	17.2	14.3	17.5

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis. See Table 8.55 for a list of the 10 specific sources of treatment.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.52A Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Age Group, Geographic and Socioeconomic Characteristics: 2000 and 2001

Geographic/Socioeconomic Characteristic	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2000	2001	2000	2001	2000	2001	2000	2001
TOTAL	3,386 ^b	4,302	1,064 ^b	1,449	1,227 ^b	1,495	1,096 ^b	1,358
GEOGRAPHIC REGION								
Northeast	692 ^b	894	212 ^b	300	246 ^b	327	234	267
Midwest	775 ^b	987	230 ^b	329	308	343	238 ^b	315
South	1,158 ^b	1,389	378 ^a	472	427	480	353 ^a	437
West	761 ^b	1,032	243 ^b	349	246 ^b	345	272	339
COUNTY TYPE								
Large Metro	1,692 ^b	2,144	517 ^b	723	611 ^b	740	565 ^a	680
Small Metro	1,046 ^b	1,359	353 ^b	459	379 ^b	480	314 ^b	421
250K - 1 Mil. Pop.	771 ^b	1,039	267 ^b	344	290 ^b	381	214 ^b	314
<250K Pop.	275	320	86	114	89	99	101	107
Nonmetro	649 ^b	799	194 ^b	267	238	275	217	257
Urbanized	187	233	59	73	69	80	58	80
Less Urbanized	377 ^b	483	115 ^b	174	142	165	120	145
Completely Rural	85	83	20	20	26	30	39	32
FAMILY INCOME								
Less Than \$20,000	697 ^b	828	213 ^b	296	256	282	228	250
\$20,000 - \$49,999	1,351 ^b	1,581	448 ^b	550	471 ^a	565	432	466
\$50,000 - \$74,999	616 ^b	811	188 ^b	274	235	249	193 ^b	288
\$75,000 or More	721 ^b	1,081	214 ^b	328	265 ^b	400	243 ^b	354
GOVERNMENT ASSISTANCE¹								
Yes	650 ^b	791	232	266	220 ^a	273	198 ^a	251
No	2,737 ^b	3,511	832 ^b	1,183	1,007 ^b	1,222	898 ^b	1,107

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis. See Table 8.55 for a list of the 10 specific sources of treatment.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

¹ At least one family member received Supplemental Security Income (SSI), cash assistance or non-cash assistance (Temporary Assistance for Needy Families, TANF), or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.52B Percentages of Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Age Group, Geographic and Socioeconomic Characteristics: 2000 and 2001

Geographic/Socioeconomic Characteristic	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2000	2001	2000	2001	2000	2001	2000	2001
TOTAL	14.6 ^b	18.4	13.8 ^b	18.9	15.3 ^b	18.4	14.6 ^b	17.7
GEOGRAPHIC REGION								
Northeast	16.6 ^b	21.2	15.2 ^b	21.2	17.5 ^a	21.8	17.1	20.3
Midwest	14.3 ^b	17.6	12.9 ^b	17.5	16.0	17.4	13.8 ^b	17.9
South	14.0 ^b	16.8	13.8 ^b	17.7	14.9	17.1	13.3	15.7
West	14.2 ^b	19.3	13.7 ^b	20.5	13.6 ^b	18.6	15.3 ^a	18.9
COUNTY TYPE								
Large Metro	15.0 ^b	19.2	13.7 ^b	19.6	15.8 ^b	19.2	15.4 ^b	18.8
Small Metro	14.6 ^b	18.6	14.7 ^b	19.4	15.4 ^a	18.5	13.7 ^b	18.0
250K - 1 Mil. Pop.	14.4 ^b	19.1	14.7 ^b	19.4	15.8 ^b	19.7	12.6 ^b	18.2
<250K Pop.	15.3	17.3	14.8	19.7	14.3	15.0	16.8	17.4
Nonmetro	13.6 ^b	16.0	12.6 ^b	16.5	14.1	16.3	14.0	15.2
Urbanized	14.2	16.9	13.8	16.3	15.3	17.2	13.4	17.2
Less Urbanized	13.1 ^a	15.6	12.3 ^b	17.2	14.0	15.9	13.0	13.9
Completely Rural	14.4	16.0	11.1	12.7	12.4	16.8	19.6	18.3
FAMILY INCOME								
Less Than \$20,000	16.6 ^b	20.6	15.2 ^b	21.7	17.7	20.9	16.9	19.3
\$20,000 - \$49,999	15.0 ^b	18.3	14.8 ^b	19.2	15.2 ^b	18.8	14.9	16.8
\$50,000 - \$74,999	13.2 ^b	17.0	12.2 ^b	17.6	14.5	15.0	12.8 ^b	18.4
\$75,000 or More	13.5 ^b	18.0	12.4 ^b	17.5	14.3 ^b	18.8	13.8 ^a	17.5
GOVERNMENT ASSISTANCE¹								
Yes	18.8 ^b	23.5	19.1 ^a	24.3	18.9 ^a	23.0	18.4 ^a	23.3
No	13.8 ^b	17.5	12.8 ^b	18.0	14.7 ^b	17.6	13.9 ^b	16.8

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis. See Table 8.55 for a list of the 10 specific sources of treatment.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

¹ At least one family member received Supplemental Security Income (SSI), cash assistance or non-cash assistance (Temporary Assistance for Needy Families, TANF), or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.53A Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Past Year Any Illicit Drug Use and Demographic Characteristics: 2000 and 2001

Demographic Characteristic	Total ¹		ANY ILLICIT DRUG USE ²			
			Yes		No	
	2000	2001	2000	2001	2000	2001
TOTAL	3,386 ^b	4,302	1,106 ^b	1,272	2,280 ^b	3,030
GENDER						
Male	1,558 ^b	2,034	452 ^a	533	1,106 ^b	1,501
Female	1,828 ^b	2,267	654	739	1,174 ^b	1,529
HISPANIC ORIGIN AND RACE						
Not Hispanic	2,922 ^b	3,717	959 ^a	1,095	1,963 ^b	2,622
White Only	2,312 ^b	2,884	790	880	1,522 ^b	2,005
Black Only	464 ^b	619	121	146	343 ^b	473
American Indian or Alaska Native Only	26	38	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	69	82	*	*	50	64
Other or More Than One Race	43 ^b	88	*	*	30	55
Hispanic	465 ^b	585	147	177	318 ^b	408

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis. See Table 8.55 for a list of the 10 specific sources of treatment.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

¹ Estimates in the total column represent all persons aged 12 to 17 who received mental health treatment/counseling in the past year.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.53B Percentages of Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Past Year Any Illicit Drug Use and Demographic Characteristics: 2000 and 2001

Demographic Characteristic	Total ¹		ANY ILLICIT DRUG USE ²			
			Yes		No	
	2000	2001	2000	2001	2000	2001
TOTAL	14.6 ^b	18.4	25.6	26.2	12.1 ^b	16.3
GENDER						
Male	13.1 ^b	17.0	20.7	21.3	11.4 ^b	15.9
Female	16.1 ^b	19.7	30.6	31.4	12.7 ^b	16.7
HISPANIC ORIGIN AND RACE						
Not Hispanic	14.7 ^b	18.5	25.8	26.3	12.1 ^b	16.5
White Only	15.2 ^b	18.9	26.4	26.8	12.4 ^b	16.8
Black Only	14.0 ^b	18.5	24.1	24.5	12.2 ^b	17.2
American Indian or Alaska Native Only	17.9	21.7	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	8.3	9.8	*	*	6.9	8.9
Other or More Than One Race	11.9 ^b	22.4	*	*	10.1 ^a	18.3
Hispanic	14.1 ^b	17.2	24.5	25.4	11.8 ^b	15.1

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis. See Table 8.55 for a list of the 10 specific sources of treatment.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

¹ Estimates in the total column represent all persons aged 12 to 17 who received mental health treatment/counseling in the past year.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.54A Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Past Year Any Illicit Drug Use, Geographic and Socioeconomic Characteristics: 2000 and 2001

Geographic/Socioeconomic Characteristic	Total ¹		ANY ILLICIT DRUG USE ²			
			Yes		No	
	2000	2001	2000	2001	2000	2001
TOTAL	3,386 ^b	4,302	1,106 ^b	1,272	2,280 ^b	3,030
GEOGRAPHIC REGION						
Northeast	692 ^b	894	245	260	447 ^b	634
Midwest	775 ^b	987	252	303	524 ^b	683
South	1,158 ^b	1,389	366	396	792 ^b	993
West	761 ^b	1,032	243 ^a	312	518 ^b	720
COUNTY TYPE						
Large Metro	1,692 ^b	2,144	540	618	1,152 ^b	1,525
Small Metro	1,046 ^b	1,359	339 ^a	413	707 ^b	946
250K - 1 Mil. Pop.	771 ^b	1,039	240 ^b	314	531 ^b	725
<250K Pop.	275	320	100	99	176	221
Nonmetro	649 ^b	799	227	241	422 ^b	558
Urbanized	187	233	76	81	111 ^a	152
Less Urbanized	377 ^b	483	120	135	257 ^b	348
Completely Rural	85	83	31	25	55	57
FAMILY INCOME						
Less Than \$20,000	697 ^b	828	226	223	472 ^b	605
\$20,000 - \$49,999	1,351 ^b	1,581	444	489	907 ^b	1,092
\$50,000 - \$74,999	616 ^b	811	209	208	407 ^b	603
\$75,000 or More	721 ^b	1,081	227 ^b	351	494 ^b	730
GOVERNMENT ASSISTANCE³						
Yes	650 ^b	791	213	250	437 ^b	541
No	2,737 ^b	3,511	893 ^a	1,022	1,843 ^b	2,489

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis. See Table 8.55 for a list of the 10 specific sources of treatment.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

¹ Estimates in the total column represent all persons aged 12 to 17 who received mental health treatment/counseling in the past year.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ At least one family member received Supplemental Security Income (SSI), cash assistance or non-cash assistance (Temporary Assistance for Needy Families, TANF), or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.54B Percentages of Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Past Year Any Illicit Drug Use, Geographic and Socioeconomic Characteristics: 2000 and 2001

Geographic/Socioeconomic Characteristic	Total ¹		ANY ILLICIT DRUG USE ²			
			Yes		No	
	2000	2001	2000	2001	2000	2001
TOTAL	14.6 ^b	18.4	25.6	26.2	12.1 ^b	16.3
GEOGRAPHIC REGION						
Northeast	16.6 ^b	21.2	30.8	29.9	13.3 ^b	18.9
Midwest	14.3 ^b	17.6	25.9	25.7	11.7 ^b	15.4
South	14.0 ^b	16.8	24.3	24.5	11.7 ^b	14.9
West	14.2 ^b	19.3	23.3	26.3	12.0 ^b	17.3
COUNTY TYPE						
Large Metro	15.0 ^b	19.2	26.7	27.6	12.4 ^b	17.1
Small Metro	14.6 ^b	18.6	23.8	26.0	12.4 ^b	16.6
250K - 1 Mil. Pop.	14.4 ^b	19.1	22.8	26.4	12.4 ^b	17.1
<250K Pop.	15.3	17.3	26.7	24.8	12.3 ^a	15.2
Nonmetro	13.6 ^b	16.0	26.1	23.6	10.8 ^b	14.1
Urbanized	14.2	16.9	27.2	26.3	10.7 ^a	14.2
Less Urbanized	13.1 ^a	15.6	24.9	22.2	10.8 ^b	14.0
Completely Rural	14.4	16.0	28.7	23.2	11.2	14.1
FAMILY INCOME						
Less Than \$20,000	16.6 ^b	20.6	28.2	26.5	13.9 ^b	19.1
\$20,000 - \$49,999	15.0 ^b	18.3	25.3	26.3	12.5 ^b	16.1
\$50,000 - \$74,999	13.2 ^b	17.0	25.8	22.5	10.5 ^b	15.6
\$75,000 or More	13.5 ^b	18.0	23.9 ^a	28.6	11.3 ^b	15.2
GOVERNMENT ASSISTANCE³						
Yes	18.8 ^b	23.5	29.7	32.6	16.0 ^b	20.8
No	13.8 ^b	17.5	24.8	25.0	11.4 ^b	15.6

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis. See Table 8.55 for a list of the 10 specific sources of treatment.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

¹ Estimates in the total column represent all persons aged 12 to 17 who received mental health treatment/counseling in the past year.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ At least one family member received Supplemental Security Income (SSI), cash assistance or non-cash assistance (Temporary Assistance for Needy Families, TANF), or the family received food stamps.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.55A Estimated Numbers (in Thousands) Reporting Source of Mental Health Treatment/Counseling in the Past Year Among Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Age Group: 2000 and 2001

Source of Treatment/Counseling	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2000	2001	2000	2001	2000	2001	2000	2001
Received treatment/counseling from any of following 10 sources ¹	3,386 ^b	4,302	1,064 ^b	1,449	1,227 ^b	1,495	1,096 ^b	1,358
Private therapist, psychologist, psychiatrist, social worker, or counselor	1,577 ^b	1,881	429 ^b	573	604	671	544 ^a	638
School counselors, school psychologists, or having regular meetings with teachers ²	1,351 ^b	1,977	440 ^b	728	508 ^b	660	403 ^b	590
Mental health clinic or center	407 ^a	481	110	113	160	177	138 ^a	192
In-home therapist, counselor, or family preservation worker	457 ^b	581	142	178	179 ^a	231	136	171
Pediatrician or other family doctor	443 ^b	610	139 ^b	191	165	191	140 ^b	228
Overnight or longer stay in any type of hospital	308	332	102	87	117	126	90	120
Special education services while in a regular classroom or in a special classroom, a special program, or in a special school ²	570 ^b	837	186 ^b	324	221 ^a	287	162 ^b	227
Partial day hospital or day treatment program	261 ^a	328	96	111	110	120	56 ^b	97
Overnight or longer stay in a residential treatment center	172	206	53	61	70	77	49	69
Overnight or longer stay in foster care or in a therapeutic foster care home	115	127	31	27	39	57	46	43

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

¹ Respondents were asked if they had received mental health treatment/counseling from each of these sources; thus, the response categories are not mutually exclusive.

² Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.55B Percentages Reporting Source of Mental Health Treatment/Counseling in the Past Year Among Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Age Group: 2000 and 2001

Source of Treatment/Counseling	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2000	2001	2000	2001	2000	2001	2000	2001
Received treatment/counseling from any of following 10 sources ¹	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Private therapist, psychologist, psychiatrist, social worker, or counselor	46.9 ^a	44.1	40.7	39.7	49.6	45.5	49.7	47.2
School counselors, school psychologists, or having regular meetings with teachers ²	40.2 ^b	46.5	41.8 ^b	51.0	41.8	44.7	36.9 ^b	43.7
Mental health clinic or center	12.1	11.3	10.4 ^a	7.9	13.1	11.9	12.6	14.2
In-home therapist, counselor, or family preservation worker	13.6	13.6	13.4	12.4	14.7	15.7	12.4	12.7
Pediatrician or other family doctor	13.2	14.3	13.1	13.3	13.5	12.9	12.8 ^a	16.9
Overnight or longer stay in any type of hospital	9.1	7.8	9.6 ^b	6.0	9.6	8.4	8.2	8.8
Special education services while in a regular classroom or in a special classroom, a special program, or in a special school ²	16.9 ^a	19.7	17.7 ^a	22.7	18.2	19.4	14.8	16.8
Partial day hospital or day treatment program	7.7	7.7	9.0	7.7	9.1	8.1	5.1	7.2
Overnight or longer stay in a residential treatment center	5.1	4.8	5.0	4.3	5.8	5.1	4.5	5.1
Overnight or longer stay in foster care or in a therapeutic foster care home	3.4	3.0	2.9	1.9	3.2	3.9	4.2	3.2

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

¹ Respondents were asked if they had received mental health treatment/counseling from each of these sources; thus, the response categories are not mutually exclusive.

² Respondents who did not report their school enrollment status or who reported not being enrolled in school in the past 12 months were not asked about receipt of treatment/counseling from this source.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.56A Estimated Numbers (in Thousands) Reporting Reason for Mental Health Treatment/Counseling in the Past Year Among Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Age Group: 2000 and 2001

	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2000	2001	2000	2001	2000	2001	2000	2001
Received treatment/counseling from any of 9 sources	3,159 ^b	3,915	993 ^b	1,299	1,141 ^b	1,353	1,026 ^b	1,263
Reason for Treatment/Counseling¹								
Thought about killing self or tried to kill self	458 ^b	607	93 ^b	155	186	230	179	222
Felt depressed	1,277 ^b	1,643	328 ^b	467	482 ^a	574	467 ^b	601
Felt very afraid or tense	392 ^b	544	97 ^b	190	174	174	121 ^b	180
Breaking rules or "acting out" ²	669 ^b	819	241	286	260	303	167 ^a	229
Eating problems	185 ^b	279	47	63	73	100	65 ^b	115
Family/home problems ³	381 ^b	473	120 ^a	160	132 ^a	174	129	139
Social/friend problems ³	138 ^b	230	50 ^b	101	56 ^a	84	32	45
School-related issues ³	235 ^b	387	74 ^b	142	89 ^b	137	72 ^a	108
Other reason/reasons	957 ^b	1,207	297 ^a	371	330 ^a	406	331 ^b	429

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis. Youths who reported they received treatment/counseling from one specific source (special education services while in a regular or special classroom, as special program, or a special school) were not asked for the reasons they received these services and are therefore not included in this table. See Table 8.55 for a list of the 10 specific sources of treatment.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

¹ Respondents were asked the reasons for the last time they received mental health treatment/counseling from each of their reported sources of treatment; thus, the response categories are not mutually exclusive.

² Includes other specify responses of "difficulty controlling anger or fighting."

³ Respondents were permitted to specify other reasons for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.

Table 8.56B Percentages Reporting Reason for Mental Health Treatment/Counseling in the Past Year Among Persons Aged 12 to 17 Receiving Mental Health Treatment/Counseling in the Past Year, by Age Group: 2000 and 2001

	Total		AGE GROUP (Years)					
			12-13		14-15		16-17	
	2000	2001	2000	2001	2000	2001	2000	2001
Received treatment/counseling from any of 9 sources	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Reason for Treatment/Counseling¹								
Thought about killing self or tried to kill self	15.6	16.6	10.4	13.0	17.4	18.1	18.3	18.5
Felt depressed	43.4	44.9	36.8	39.3	45.2	45.2	47.6	50.2
Felt very afraid or tense	13.3	14.9	10.9 ^a	16.0	16.3	13.7	12.4	15.0
Breaking rules or "acting out" ²	22.8	22.4	27.1	24.1	24.4	23.9	17.0	19.1
Eating problems	6.3	7.6	5.2	5.3	6.9	7.9	6.6 ^a	9.6
Family/home problems ³	12.9	12.9	13.5	13.5	12.4	13.7	13.1	11.6
Social/friend problems ³	4.7 ^a	6.3	5.6 ^a	8.5	5.2	6.6	3.2	3.7
School-related issues ³	8.0 ^b	10.6	8.3 ^a	12.0	8.4	10.8	7.3	9.0
Other reason/reasons	32.6	33.0	33.3	31.3	30.9	32.0	33.7	35.8

*Low precision; no estimate reported.

NOTE: Mental health treatment/counseling for youths is defined as having received treatment or counseling from any of 10 specific sources for emotional or behavioral problems NOT caused by alcohol or drug use. Youths who did not answer four or more of the source of treatment questions with a "yes" or "no" response were excluded from this analysis. Youths who reported they received treatment/counseling from one specific source (special education services while in a regular or special classroom, as special program, or a special school) were not asked for the reasons they received these services and are therefore not included in this table. See Table 8.55 for a list of the 10 specific sources of treatment.

^aDifference between estimate and 2001 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2001 estimate is statistically significant at the .01 level.

¹ Respondents were asked the reasons for the last time they received mental health treatment/counseling from each of their reported sources of treatment; thus, the response categories are not mutually exclusive.

² Includes other specify responses of "difficulty controlling anger or fighting."

³ Respondents were permitted to specify other reasons for having received mental health treatment/counseling. This reason is one of the most commonly reported other reasons for having received treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000 and 2001.