

**Table 6.19A Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Detailed Age Categories: Numbers in Thousands, 2002**

Age Category	Total	SERIOUS MENTAL ILLNESS <sup>1</sup>	
		Yes	No
Total 18 or Older	27,322	8,355	18,968
18	431	153	277
19	407	170	237
20	377	137	240
21	469	209	260
22	398	168	230
23	427	189	238
24	396	155	240
25	380	213	166
26-29	1,738	595	1,142
30-34	2,811	1,051	1,760
35-39	3,006	1,052	1,954
40-44	3,668	1,312	2,356
45-49	3,333	1,163	2,171
50-54	2,900	*	2,044
55-59	2,408	*	1,876
60-64	1,188	*	1,082
65 or Older	2,988	*	2,694

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for alcohol or drug use. Respondents with missing treatment/counseling information are excluded.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.6 of Appendix B of the *Results from the 2002 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 6.19B Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Detailed Age Categories: Percentages, 2002**

Age Category	Total	SERIOUS MENTAL ILLNESS <sup>1</sup>	
		Yes	No
Total 18 or Older	13.0	47.9	9.9
18	9.9	28.4	7.3
19	10.0	33.1	6.6
20	9.5	26.8	6.9
21	11.6	36.6	7.5
22	10.3	31.5	6.9
23	11.2	36.6	7.2
24	11.2	36.4	7.8
25	11.4	45.4	5.8
26-29	11.4	41.0	8.3
30-34	14.2	49.3	9.9
35-39	13.7	52.7	9.8
40-44	16.8	58.7	12.0
45-49	15.8	68.5	11.2
50-54	15.4	*	11.7
55-59	16.4	*	13.6
60-64	10.3	*	9.7
65 or Older	8.8	*	8.2

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for alcohol or drug use. Respondents with missing treatment/counseling information are excluded.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.6 of Appendix B of the *Results from the 2002 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 6.20A Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Demographic Characteristics: Numbers in Thousands, 2002**

Demographic Characteristic	Total	SERIOUS MENTAL ILLNESS <sup>1</sup>	
		Yes	No
<b>TOTAL</b>	27,322	8,355	18,968
<b>AGE</b>			
18-25	3,283	1,394	1,888
26-49	14,556	5,173	9,383
50 or Older	9,483	1,787	7,696
<b>GENDER</b>			
Male	8,784	2,376	6,408
Female	18,538	5,978	12,560
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	25,277	7,703	17,574
White	22,014	6,547	15,467
Black or African American	1,973	750	1,223
American Indian or Alaska Native	228	*	134
Native Hawaiian or Other Pacific Islander	29	*	24
Asian	682	*	486
Two or More Races	352	*	240
Hispanic or Latino	2,045	651	1,394
<b>EDUCATION</b>			
< High School	4,311	1,404	2,907
High School Graduate	7,912	2,631	5,281
Some College	7,528	2,576	4,952
College Graduate	7,571	1,743	5,828
<b>CURRENT EMPLOYMENT</b>			
Full-Time	13,090	3,796	9,295
Part-Time	3,901	1,346	2,554
Unemployed	1,077	457	620
Other <sup>2</sup>	9,255	2,756	6,499

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for alcohol or drug use. Respondents with missing treatment/counseling information are excluded.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.6 of Appendix B of the *Results from the 2002 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 6.20B Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Demographic Characteristics: Percentages, 2002**

Demographic Characteristic	Total	SERIOUS MENTAL ILLNESS <sup>1</sup>	
		Yes	No
<b>TOTAL</b>	13.0	47.9	9.9
<b>AGE</b>			
18-25	10.6	34.2	7.0
26-49	14.6	54.4	10.4
50 or Older	12.0	46.4	10.3
<b>GENDER</b>			
Male	8.7	39.5	6.8
Female	17.0	52.3	12.8
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	13.7	49.0	10.4
White	14.7	51.9	11.3
Black or African American	8.5	36.9	5.8
American Indian or Alaska Native	17.8	*	11.9
Native Hawaiian or Other Pacific Islander	3.9	*	3.5
Asian	8.5	*	6.6
Two or More Races	16.6	*	13.1
Hispanic or Latino	8.2	37.8	6.0
<b>EDUCATION</b>			
< High School	11.7	39.3	8.7
High School Graduate	11.7	45.1	8.5
Some College	14.3	51.7	10.4
College Graduate	14.4	57.1	11.8
<b>CURRENT EMPLOYMENT</b>			
Full-Time	11.3	45.0	8.6
Part-Time	14.2	50.7	10.3
Unemployed	14.2	42.4	9.6
Other <sup>2</sup>	15.8	52.2	12.2

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for alcohol or drug use. Respondents with missing treatment/counseling information are excluded.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.6 of Appendix B of the *Results from the 2002 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 6.21A Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Geographic Characteristics: Numbers in Thousands, 2002**

Geographic Characteristic	Total	SERIOUS MENTAL ILLNESS <sup>1</sup>	
		Yes	No
<b>TOTAL</b>	27,322	8,355	18,968
<b>GEOGRAPHIC DIVISION</b>			
Northeast	5,597	1,745	3,852
New England	1,828	653	1,175
Middle Atlantic	3,769	1,092	2,677
Midwest	6,208	1,912	4,297
East North Central	4,192	1,330	2,861
West North Central	2,017	581	1,435
South	9,239	2,999	6,239
South Atlantic	4,846	1,594	3,252
East South Central	1,747	636	1,110
West South Central	2,646	769	1,877
West	6,278	1,699	4,580
Mountain	1,810	507	1,303
Pacific	4,468	1,191	3,276
<b>COUNTY TYPE</b>			
Large Metro	12,885	3,681	9,204
Small Metro	8,953	2,879	6,074
250K - 1 Mil. Pop.	6,628	2,027	4,601
<250K Pop.	2,325	852	1,473
Nonmetro	5,485	1,795	3,690
Urbanized	2,051	625	1,426
Less Urbanized	2,990	1,045	1,945
Completely Rural	443	*	319

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for alcohol or drug use. Respondents with missing treatment/counseling information are excluded.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.6 of Appendix B of the *Results from the 2002 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 6.21B Received Mental Health Treatment/Counseling in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Geographic Characteristics: Percentages, 2002**

Geographic Characteristic	Total	SERIOUS MENTAL ILLNESS <sup>1</sup>	
		Yes	No
<b>TOTAL</b>	13.0	47.9	9.9
<b>GEOGRAPHIC DIVISION</b>			
Northeast	13.8	50.7	10.4
New England	17.2	64.9	12.2
Middle Atlantic	12.6	44.9	9.8
Midwest	13.0	47.5	9.9
East North Central	12.6	47.4	9.4
West North Central	14.1	47.8	11.0
South	12.3	47.5	9.1
South Atlantic	12.3	47.6	9.0
East South Central	13.8	53.3	9.7
West South Central	11.6	43.5	8.9
West	13.4	46.3	10.6
Mountain	13.3	45.4	10.4
Pacific	13.4	46.7	10.6
<b>COUNTY TYPE</b>			
Large Metro	12.3	46.1	9.5
Small Metro	14.1	52.0	10.5
250K - 1 Mil. Pop.	14.4	51.1	10.9
<250K Pop.	13.5	54.2	9.4
Nonmetro	13.1	45.7	9.7
Urbanized	15.4	56.1	11.7
Less Urbanized	12.4	43.5	9.0
Completely Rural	9.6	*	7.5

\*Low precision; no estimate reported.

NOTE: Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for alcohol or drug use. Respondents with missing treatment/counseling information are excluded.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.6 of Appendix B of the *Results from the 2002 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.