

WOOD STRAWBERRY

Fragaria vesca L.

Plant Symbol = FRVE

Contributed By: USDA NRCS National Plant Data Center



Carl Lindman
©Gerhard Keuck from Bilder ur Nordens
@ Project Runeburg

Alternate Names

Woodland strawberry; California strawberry

Uses

Ethnobotanic: The fruit was gathered by native peoples throughout the United States and Canada. Such cultural groups include the Micmac, Huron, Potawatomi, Creek, Blackfoot, Iroquois, and many other groups. The fruit was eaten raw and not preserved by California Indian tribes including the Coast Yuki and the Karok. Furthermore, a tea was made from the leaves by the Upriver Halkomelem and Sechelt of British Columbia, the Cowlitz of Washington and the Micmac of the maritimes.

Wildlife: The Portola woodrat and the valley quail eat the fruit and leaves of wild strawberries.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

Description

General: Rose Family (Rosaceae). This herbaceous perennial plant spreads by seed, short rhizomes and leafless stolons. The toothed leaves are thin and basal with a petiole generally 3-12 cm. They appear in leaflets of 3 and are sparsely hairy above. The flowers have 5 white petals with numerous pistils and 20-35 stamens. The five bractlets are often 2-lobed. The red fleshy fruit is covered with achenes.

Distribution

For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site. It is found in northwestern California, the Cascade Ranges, the Sierra Nevada, central-western California, San Bernardino Mountains, Peninsular Ranges, to eastern North America, and south to Baja California and also Europe.

Establishment

Adaptation: This plant is found below 2000 m in partial shade of closed-cone pine, evergreen, mixed conifer forests, and chaparral and has a very wide distribution.

Planting: Dig up plantlets or runners and plant them in pots in summer, make sure to cover the stems and roots in soil. Place the pots in a hothouse to establish good, strong roots. Water the plants or runners and keep them moist. Plant the seedlings outdoors in the ground in the fall or winter after the rains have started. They should be planted in full sun in a light, loose soil, about ten inches apart. It will not take long for the plants to make a complete ground cover. Lightly fertilize the plants during the growing season. Note that those plants that have bigger flowers usually have less fruit while those with smaller flowers have more fruit. Protect the plants from gophers, deer, squirrels, raccoons, and other wildlife.

Management

Keep the runners pruned back because they can be invasive. It is necessary to divide the patch every 3 to 4 years and start a new patch for increased vigor. Younger plants are more vigorous and produce more berries.

Cultivars, Improved and Selected Materials (and area of origin)

FRVE is somewhat available through native plant nurseries within its range. Contact your local Natural Resources Conservation Service (formerly Soil Conservation Service) office for more information. Look in the phone book under "United States Government." The Natural Resources Conservation Service will be listed under the subheading "Department of Agriculture."

References

- Chesnut, V.K. 1902. *Plants used by the Indians of Mendocino County, California*. Contributions from the U.S. National Herbarium Vol. VII. Reprinted by the Mendocino County Historical Society in 1974.
- Dixon, R.B. 1905. *The Huntington California expedition: the Northern Maidu*. Bulletin of the American Museum of Natural History 17(3):119-346.
- Ertter, B. 1993. *Fragaria*. Page 952 IN: The Jepson Manual: Higher Plants of California. J.C. Hickman (ed.). University of California Press, Berkeley, California.
- Gifford, E.W. 1965. *The Coast Yuki*. Sacramento Anthropological Society Paper 2:1-97. Sacramento, California.
- Kuhnlein, H.V. & N.J. Turner 1991. *Traditional plant foods of Canadian indigenous peoples: Nutrition, botany, and use*. Gordon and Breach Science Publishers, Philadelphia, Pennsylvania.
- Martin, A.C., H.S. Zim, & A.L. Nelson 1951. *American wildlife and plants: A guide to wildlife food habits*. Dover Publications, Inc., New York, New York.
- Schenck, S.M. & E.W. Gifford 1952. *Karok ethnobotany*. Anthropological Records 13(6):377-392.
- USDA, NRCS 1999. *The PLANTS database*. National Plant Data Center, Baton Rouge, Louisiana. <<http://plants.usda.gov>>. Version: 990412.

Prepared By & Species Coordinator

M. Kat Anderson

USDA, NRCS, National Plant Data Center
c/o Plant Sciences Department, University of California, Davis, California

Wayne Roderick

Former Director of the East Bay Regional Parks
Botanic Garden, Berkeley, California

Edited: 05dec00 jsp; 19may03 ahv; 07jun06 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Read about [Civil Rights at the Natural Resources Conservation Service](#).