

WOOLGRASS

Scirpus cyperinus (L.) Kunth

Plant Symbol = SCCY

Contributed by: USDA NRCS National Plant Data Center



R. Mohlenbrock
USDA, NRCS, Wetlands Institute
@ PLANTS

Alternative Names

cotton grass bulrush, common wool-grass

Uses

Ethnobotanic: Woolgrass stems were woven to make matting and ropes. The fruiting tops of the plant were used as a resilient material for stuffing and making pillows (Moerman 1998). The small rushes were used in making woven mats and storage bags (Ibid.).

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

Description

General: Sedge family (Cyperaceae). Woolgrass (*Scirpus cyperinus*) is a tall perennial with slender culms. This species is an erect grasslike plant that commonly grows four to five feet (Tiner 1987). The leaves are smooth, flat, elongated, and up to ½ inch wide. The flowers occur in dense rounded clusters of greenish-brown spiklets arising from the top of the culm. The fruits are yellow-gray to white achenes surpassed by long red-brown bristles at maturity.

Distribution: *Scirpus cyperinus* ranges from New England and New York westward across Ohio to

Iowa, and southward to North Carolina and Oklahoma. It is also found from Newfoundland to Minnesota south to Florida and Louisiana (Tiner 1987). For current distribution, please consult the Plant profile page for this species on the PLANTS Web site.

Adaptation

Woolgrass is found in irregularly flooded tidal fresh marshes, inland marshes, wet meadows, and swamps. This species grows best in areas with wet soil moisture content and is seldom found in more than a few inches of water (Voss 1972). It prefers peat or sandy soil types in full to partially sunny locations.

Establishment

Propagation by Seed: *Scirpus cyperinus* seeds should be sown in a cold frame as soon as they are ripe in a pot standing in three centimeters of water. The seeds germinate quickly. When they are large enough to handle, plant them into their permanent positions in early summer.

Large divisions can be planted directly into their permanent positions. It is best to pot smaller divisions and grow them in a cold frame, out-planting when they are well established in the summer.

Management

After seed planting, water level over *Scirpus cyperinus* seeds should be maintained at one foot for two weeks. Periodic flooding up to three feet should occur until the seeds are established.

Cultivars, Improved and Selected Materials (and area of origin)

Available through wetland plant nurseries. Contact your local Natural Resources Conservation Service office for more information. Look in the phone book under "United States Government." The Natural Resources Conservation Service will be listed under the subheading "Department of Agriculture."

References

Braun, L.E. 1967. *The monocotyledoneae from cat-tails to orchids*. The Ohio State University Press, Columbus, Ohio.

Britton, N.L. & A. Brown 1970. *An illustrated flora of the northern United States and Canada*. Dover Publications, New York, New York.

Plant Materials <<http://plant-materials.nrcs.usda.gov/>>

Plant Fact Sheet/Guide Coordination Page <<http://plant-materials.nrcs.usda.gov/intranet/pfs.html>>

National Plant Data Center <<http://npdc.usda.gov>>

Bruggen, T. V. 1976. *The vascular plants of South Dakota*. The Iowa State University Press, Ames, Iowa.

Gleason, H. A. & A. Cronquist 1993. *Manual of vascular plants of northeastern United States and adjacent Canada*. 2nd ed. The New York Botanical Garden, Bronx, New York.

Moerman, D. 1998. *Native American ethnobotany*. Timber Press, Oregon.

Radford, A.E., H.E. Ahles, & C.R. Bell 1968. *Manual of the vascular flora of the Carolinas*. The University of North Carolina Press, Chapel Hill, North Carolina.

Straughbaugh, P. D. & E. L. Core 1977. *Flora of West Virginia*. 2nd ed. Seneca Books, Inc., Morgantown, West Virginia.

The Great Plains Flora Association 1986. *Flora of the Great Plains*. University Press of Kansas, Lawrence, Kansas.

Tiner, R.W. Jr. 1987. *A field guide to coastal wetland plants of the northeastern United States*. The University of Massachusetts Press, Amherst, Massachusetts.

Voss, E.G. 1972. *Michigan flora*. Cranbrook Institute of Science, Bloomfield Hills, Michigan, and University of Michigan Herbarium, Ann Arbor, Michigan.

Prepared By

Jammie Favorite

formerly USDA, NRCS, National Plant Data Center
Baton Rouge, Louisiana

Species Coordinator

M. Kat Anderson

USDA, NRCS, National Plant Data Center, c/o Plant
Sciences Dept., Davis, California

Edited: 19jun02 jsp; 03jun03 ahv; 060816 jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's [TARGET Center](#) at 202-720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Read about [Civil Rights at the Natural Resources Conservation Service](#).

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities