

NEW ENGLAND ASTER

Symphyotrichum novae-angliae

(L.) Nesom

Plant Symbol = SYNO2

Contributed by: USDA NRCS National Plant Data Center



Robert Mohlenbrock
USDA, NRCS, Wetland Sciences Institute
@ PLANTS

Alternative Names

Aster novae-angliae

Uses

Ethnobotanic: A decoction of the plant has been used in the treatment of weak skin (Moerman 1998). A poultice has been used in the treatment of pain, fevers, and diarrhea.

Wildlife: New England aster is known for attracting butterflies and moths to areas where it is found growing. This is a good bee plant providing nectar in the autumn. Most species in this genus seem to be immune to the predations of rabbits (Thomas 1990).

Status

Please consult the Plants Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

Description

General: Aster family (Asteraceae). New England aster (*Aster novae-angliae*) is a perennial with several

erect stems in clumps, stems 0.8 to 2 meters, densely spreading pubescent, thirty or more nodes below the branches of the inflorescence (Radford, Ahles & Bell 1968). The leaves are lanceolate to elliptic lanceolate, two to nine centimeters long and one to two centimeters wide. The disc flowers are yellow, hermaphrodite (having both male and female organs) and are pollinated by bees, butterflies, flies, beetles and moths.

Distribution: *Aster novae-angliae* ranges from Quebec to Alberta, south to North Carolina, Alabama, Arkansas, Kansas, Colorado, Wyoming and New Mexico (Steyermark 1963). For current distribution, please consult the Plant profile page for this species on the PLANTS Web site.

Adaptation

This plant is commonly found in moist prairies, meadows, roadsides and streams (Dension 1998). It requires well-drained soil and prefers sandy, loamy and clay soils. This species can grow on nutritionally poor soil but prefers rich soil. New England aster grows well in a sunny location and can succeed in partial shade.

Establishment

Propagation by Seed: *Aster novae-angliae* seeds should be sown fresh in the fall or spring (Heuser 1997). Pre-chill spring sown seeds to improve germination (Ibid.). When the seedlings are large enough to handle, place them into individual pots and plant them out in the summer.

Division of this species should be done in the spring or autumn. Large divisions can be planted into their permanent positions whereas smaller clumps should be kept in a cold frame until they are growing well.

Management

Divisions of *Aster novae-angliae* species should be done in the spring every three years to maintain vigor (Heuser 1997).

Cultivars, Improved and Selected Materials (and area of origin)

Available somewhat through native plant seed sources. Contact your local Natural Resources Conservation Service (formerly Soil Conservation Service) office for more information. Look in the phone book under "United States Government." The Natural Resources Conservation Service will be

listed under the subheading "Department of Agriculture."

References

Bruggen, T.V. 1976. *The vascular plants of South Dakota*. The Iowa State University Press, Ames, Iowa.

Dension, E. 1998. *Missouri wildflowers*. Missouri Department of Conservation, Jefferson City, Missouri.

Gleason, H.A. & A. Cronquist 1991. *Manual of the vascular plants of northeastern United States and adjacent Canada*. 2nd ed. The New York Botanical Garden, Bronx, New York.

Heuser, C.W. 1997. *The complete book of plant propagation*. The Taunton Press, Newtown, Connecticut.

Huxley, A. 1992. *The new RHS dictionary of gardening*. MacMillan Press, New York, New York.

Moerman, D. 1998. *Native American ethnobotany*. Timber Press, Portland, Oregon.

Radford, A.E., H.E. Ahles, & C. Bell 1968. *Manual of the vascular flora of the Carolinas*. The University of North Carolina Press, Chapel Hill, North Carolina.

Steyermark, J.A. 1963. *Flora of Missouri*. The Iowa State University Press, Ames, Iowa.

Strausbaugh, P.D. & E.L. Core 1971. *Flora of West Virginia*. 2nd ed. Seneca Books, Inc., Morgantown, West Virginia.

The Great Plains Flora Association 1986. *Flora of the Great Plains*. University Press of Kansas, Lawrence, Kansas.

Prepared By

Lincoln M. Moore

USDA, NRCS, National Plant Data Center, Baton Rouge, Louisiana

Species Coordinator

M. Kat Anderson

USDA, NRCS, National Plant Data Center, c/o Plant Sciences Dept., Davis, California

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

Read about [Civil Rights at the Natural Resources Conservation Service](#).