

EFFECT OF FARM MILK ON DEVELOPMENT OF ASTHMA AND ALLERGIES

Georg Loss, *Swiss Tropical and Public Health Institute, Basel, Switzerland and University of Basel, Switzerland*

Silvia Apprich, *BOKU, University of Natural Resources and Life Sciences, Vienna, Austria*

Marco Waser, *Swiss Tropical and Public Health Institute, Basel, Switzerland and University of Basel, Switzerland*

Wolfgang Kneifel, *BOKU Vienna, University of Natural Resources and Life Sciences, Vienna, Austria*

Erika von Mutius, *LMU Munich, University Children's Hospital, Munich, Germany*

Jon Genuneit, *Ulm University, Germany*

Gisela Büchele, *Ulm University, Germany*

Barbara Sozanska, *Wroclaw Medical University, Poland*

Charlotte Braun-Fahrlander, *Swiss Tropical and Public Health Institute, Basel, Switzerland and University of Basel, Switzerland*

Background and Aims: Several epidemiological studies found consumption of farm milk to be inversely associated with the occurrence of hay fever, pollen sensitization. GABRIELA, a large cross-sectional multi-centre study, offered for the first time the opportunity to relate analytically measured milk components to asthma and atopy outcomes in children and possibly explain the previously reported protective farm milk effect.

Methods: In four rural regions of Germany, Austria, and Switzerland, farm and milk exposure of 7606 5–13 year old children was assessed by a detailed questionnaire and they also provided blood samples to determine specific IgE levels. Viable bacterial counts, whey protein levels, and total fat content were analysed in 800 milk samples that were collected at the participant's homes. Associations of milk consumption and asthma, atopy, hay fever, and atopic dermatitis were computed by stratified weighted logistic regression analyses adjusted for potential confounders.

Results: Raw milk consumption, that was reported in questionnaires, was inversely associated with asthma (aOR: 0.59, 95%-CI: 0.46–0.74), atopy (0.74, 0.61–0.90), and hay fever (0.51, 0.37–0.69) independent of other farm exposures. Consumption of exclusively boiled farm did not show a protective effect. The heating status of consumed milk as reported by parents was in very good agreement with objective measurements. Total viable bacterial counts or total fat content of milk were not significantly related to asthma or atopy. Yet, increased levels of the whey proteins were inversely associated with asthma but not with atopy. Significant inverse dose-response relations were observed for lactoferrin (p -value for linear trend = 0.02), TGF-beta2 (0.02), alpha-lactalbumin (0.02), and beta-lactoglobulin (0.04).

Conclusions: The protective effect of raw milk consumption on asthma seems to be associated with the whey protein fraction of unprocessed farm milk. Relevant immunomodulatory compounds of farm milk and underlying mechanisms need to be explored in more detail.