

METHODS TO ASSESS SELENIUM EXPOSURE: EVALUATION OF DIETARY INTAKE AND BIOMARKERS IN MELANOMA PATIENTS AND POPULATION CONTROLS

Simone Storani, *Department of Public Health Sciences, University of Modena and Reggio Emilia, Italy*
Carlotta Malagoli, *Department of Public Health Sciences, University of Modena and Reggio Emilia, Italy*
Ilaria Bottecchi, *Department of Public Health Sciences, University of Modena and Reggio Emilia, Italy*
Angela Ferrari, *Department of Public Health Sciences, University of Modena and Reggio Emilia, Italy*
Marco Vinceti, *Department of Public Health Sciences, University of Modena and Reggio Emilia, Italy*

Background and Aims: Selenium is a metalloid of strong nutritional and toxicological importance, whose relation with risk of chronic diseases, including cancer, is still uncertain and currently under active investigation. Different methods to estimate selenium exposure in the human have been suggested, but their strengths and limitations are still controversial. We assessed selenium status in a group of melanoma and cancer patients, by evaluating their average dietary intake of the metalloid and its concentrations in plasma and toenails.

Methods: We investigated selenium status of 54 patients with newly-diagnosed cutaneous melanoma and 56 controls, originally enrolled in a population-based case-control study of environmental and life-style risk factors of cutaneous melanoma in Modena, northern Italy. We sampled subjects' blood and toenails and we administered a food frequency questionnaire including 248 questions on frequency and quantity of consumption of 188 food items, developed within the EPIC (European Prospective Investigation on Cancer and Nutrition) project in northern Italy. Plasma and toenail selenium concentrations were determined by atomic absorption spectrometry.

Results: In the overall study population, plasma selenium levels correlated with toenail concentrations (Spearman rank correlation coefficient (ρ) = 0.22, $p=0.02$), but these biomarkers were not associated with dietary intake of the metalloid. In subgroup analysis according to disease status, plasma selenium levels were associated with dietary intake only among cases ($\rho=0.25$, $p=0.07$), while plasma and toenail concentrations correlated both in cases and in controls.

Conclusions: Study results may be explained by intake of different selenium species in cases compared with controls, or more likely by differences in absorption and metabolism of selenium compounds between the two study groups. These findings suggest that the validity of methods assessing selenium exposure may vary according to disease status.