SOCIOECONOMIC STATUS AND ENVIRONMENTAL TOBACCO SMOKE EXPOSURE OF CHILDREN IN KOREA.

Okhee Yi, Department of Preventive Medicine, Dankook University College of Medicine, Cheonan, Republic of Korea Ho-Jang Kwon, Department of Preventive Medicine, Dankook University College of Medicine, Cheonan, Republic of Korea Dongjin Kim, Health Promotion Research Division, Korea institute for health and social affairs, Seoul, Republic of Korea Ho Kim, Graduate School of Public Health, Seoul National University, Seoul, Republic of Korea

Mina Ha, Department of Preventive Medicine, Dankook University College of Medicine, Cheonan, Republic of Korea Soo-Jong Hong, Department of Pediatrics, Asan Medical Center, Ulsan University, College of Medicine, Seoul, Republic of Korea

Yun-Chul Hong, Institute of Environmental Medicine, Seoul National University College of Medicine, Seoul, Republic of Korea Jong-Han Leem, Department of Occupational and Environmental Medicine, Inha University, Incheon, Republic of Korea Joon Sakong, Department of Preventive Medicine and Public Health, College of Medicine, Yeungnam University, Republic of Korea

Chul Gab Lee, Department of Occupational Environmental Medicine, Chosun University, Kwangju, Republic of Korea Su-Young Kim, Department of Preventive Medicine College of Medicine Cheju National University, Cheju, Republic of Korea Dongmug Kang, Department of Preventive and Occupational Medicine, Pusan National University, Pusan, Republic of Korea

Background and Aims: Exposure to environment tobacco smoke (ETS) can have serious health consequences, especially for children. It was reported that most of ETS exposure occurred at home and lower parental socioeconomic status (SES) increased a risk of ETS exposure at home. This study aims to investigate the relationship between individual level SES of parents and children's ETS exposure, to examine the effects of individual level SES depending on the characteristic of residential areas, and to evaluate the interactive effects of area level and individual level SES in Korea.

Methods Whether ETS exposure was associated with parental SES was evaluated using information on 7,059 children aged 6–13 years from the Children's Health and Environment Research (CHEER) study in Korea. First, bivariate and multivariate associations between parental SES and children's ETS exposure were examined using logistic regression analysis. Secondly, to evaluate the effects of parental SES on children's ETS exposure by deprivation level of residential areas, we estimated the model including the interaction term of deprivation level of residential areas with parental SES.

Results: After adjustment for possible confounding variables, paternal education was found to be highly correlated with children's ETS exposure, especially among children whose father had less than 12 year education living in the most deprived area (Odds ratio=2.82, 95% CI=(2.02-3.94)).

Conclusions: Our findings suggest that characteristics of residential areas may influence ETS exposure of children directly and interactively with parental SES in Korea. Findings from this study could inform policy decision making of evidences regarding the associations of community as well as individual level socioeconomic factors with ETS.

References:

Bolte G, Hermann Fromme for the GME Study Group. Socioeconomic determinants of children's environmental tobacco smoke exposure and family's home smoking policy. European Journal of Public Health. 2008;19:52–58.

Cook DG, Whincup PH, Jarvis MJ, Strachan DP, Papacosta O, Bryant A. Passive exposure to tobacco smoke in children aged 5–7 years: individual, family, and community factors. BMJ. 1994;308:384–389.

Huisman M, Kunst A.E, Mackenbach. Inequlities in the prevalence of smoking in the European Union: comparing education and income. Preventive Medicine 2005;40:756–764

Laaksonen M, Rahkonen O, Karvonen S, Lahelma E. Socioeconomic status and smoking. European Journal of Public Health. 2005:15:262–269.