


B O C E P R E V I R

Treatment for Hepatitis C | A Patient's Workbook

What You Need to Know When Taking Triple Therapy with Boceprevir (Victrelis™)



CONTACT INFORMATION

Prescriber contact: _____

Contact for questions about medications: _____

Number to call for refills: _____

Order your medications 2 weeks ahead of time or as far in advance as possible.

Appointments: _____

Other important contacts: _____

CALL YOUR PROVIDER IF:

Your temperature is above 101 F. for more than 24-48 hours.

You miss more than one day of the drug boceprevir.

You miss more than 1-2 days of ribavirin.

You miss any dose of pegylated interferon.

Table of Contents

What is Triple Therapy?	2
Preparing for Treatment with Boceprevir	3
How to Take Your Medications	4
When to Take Your Medications	4
Sample Medication Schedules	5
If You Miss a Dose	6
How Long Does Treatment Last?	7
After Treatment is Completed	7
If Treatment Doesn't Work for You	7
Which Treatment Track Should YOU Follow?	8
Treatment Tracks:	9
• Tracks A & B.....	10
• Track C.....	14
• Track D.....	18
Dealing with Side Effects	22
Notes to Bring to Your Provider	28
My Viral Load Chart	30
My Treatment Visit Schedule	31

What is Triple Therapy?

Until recently, the standard treatment for chronic hepatitis C was a combination of peginterferon and ribavirin.

In May 2011 the Food and Drug Administration approved two new drugs for the treatment of hepatitis C: **boceprevir** (brand name Victrelis™) and **telaprevir** (brand name Incivek™). These drugs must be taken with peginterferon and ribavirin. Boceprevir and telaprevir cannot be used together; a patient would take one or the other.

This combination of three drugs—peginterferon, ribavirin, and EITHER boceprevir or telaprevir—is called **triple therapy**. Triple therapy is used only in patients with a strain of hepatitis C called genotype 1, the most common strain in the U.S.

This booklet is for people taking **boceprevir**.



RIBAVIRIN

Ribavirin is a medicine that fights certain viruses. Ribavirin works against hepatitis C when taken with peginterferon and boceprevir.



PEGINTERFERON

Interferon is a protein that your body makes to fight infections. When used as a treatment, interferon increases the amount of interferon in your body. Long-acting interferon, given as a shot, once a week, is called pegylated interferon, or peginterferon.



BOCEPREVIR

Boceprevir directly attacks the hepatitis C virus to keep it from growing. It can't get rid of the virus if just taken by itself, but it greatly increases the chances of getting rid of the virus (a cure) when used together with peginterferon and ribavirin.

Preparing for Treatment with Boceprevir

- Before starting hepatitis C medications, you should check with your regular doctor and your hepatitis doctor on any existing problems, such as diabetes mellitus, gout, hypertension, COPD, thyroid disorder, depression, coronary artery disease, skin disease and high cholesterol. Your doctors may recommend you have other tests done—an eye exam, for example, or an X ray--to be sure you are in the best condition possible before starting treatment. Tell your doctor if you are taking any herbal supplements.
- Be sure you and your sexual partner are using 2 forms of contraception—such as condoms plus IUD or condoms plus a diaphragm. Birth control pills may not be as effective when they are used with the new medicines. Ribavirin can cause serious birth defects, and you and your partner should be extremely cautious not to become pregnant while either of you is taking treatment and for 6 months after treatment ends.
- Expect that you may be extremely tired and run down while taking treatment. Consider discussing with family, friends and coworkers ahead of time.
- Sleep well, exercise regularly, eat a healthy diet, and drink plenty of water.
- It is best if you do not drink any alcohol during your course of treatment.
- Plan a routine for home and work that will ensure:
 - You take your boceprevir every 8 hours, with food
 - You take ribavirin twice daily.
 - You have a set day and time of the week for your peginterferon injection.
 - You plan on having blood draws approximately every 4 weeks or as directed by your provider. Blood tests will show how you are responding to treatment and can flag side effects, such as low white or red blood cells.
 - You plan on having office visits with your provider, approximately every 4 weeks. These may be on different dates than the blood draws. (See Treatment Visit Schedule on p. 31.)

How to Take Your Medications

RIBAVIRIN



- Take 2 or 3 capsules or tablets, 2 times per day, as prescribed. Each pill is 200 mg.
- Your dose will be based on your weight and may change during the course of treatment because of side effects.
- Take with food to minimize stomach upset.
- Store at room temperature between 59-86° F.

PEGINTERFERON



- You will be given prefilled syringes or pens of either Pegasys® or Peg-Intron®.
- Keep in the refrigerator.
- Inject under the skin once weekly.

BOCEPREVIR



- Take 4 capsules, every 8 hours. Each capsule is 200 mg.
- You **MUST** take within 30 minutes of either a meal or light snack at all 3 times for the medication to be absorbed into the bloodstream.
- Start this medication 4 weeks **after** starting ribavirin and peginterferon.
- Refrigerate if room temperature is greater than 77° F.

When to Take Your Medications




















Boceprevir pills must be taken with food and on a strict schedule of every 8 hours to work properly, so make sure the schedule you and your provider pick will work for you. A common schedule is 6 am–2 pm–10 pm. You can choose whether to take ribavirin and boceprevir together, or at separate times of day. Because **ribavirin** can interfere with sleep, take your last dose before 6 pm.

Think about your personal routine when deciding which day to take your **peginterferon** shot. Many patients feel more fatigued on the first day or two after each shot. If you work Monday through Friday, for example, consider taking your shot on Friday night so you have the weekend to recover from the worst of the fatigue.

SAMPLE MEDICATION SCHEDULES


















































The charts below show how you might fill in times for taking your medications. Some weeks you'll take 2 drugs, other weeks you'll take 3 drugs. (Follow the medication schedule for your particular Treatment Track.)

2 Drugs

Time of Day	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
 Morning 	 2 # ribavirin	 2 # ribavirin	 2 # ribavirin	 2 # ribavirin	 2 # ribavirin	 2 # ribavirin	 2 # ribavirin
 Afternoon 	 2 # ribavirin	 2 # ribavirin	 2 # ribavirin	 2 # ribavirin	 2 # ribavirin	 2 # ribavirin	 2 # ribavirin
 Peginterferon Mark day & time						7 PM	

3 Drugs

Patients can choose whether to take ribavirin and boceprevir together, or at separate times of day. However, boceprevir *must* be taken every 8 hours with a light meal or snack.

Time of Day	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
 Morning 	 +  2 # ribavirin	 +  2 # ribavirin	 +  2 # ribavirin	 +  2 # ribavirin	 +  2 # ribavirin	 +  2 # ribavirin	 +  2 # ribavirin
 Afternoon 	 +  2 # ribavirin	 +  2 # ribavirin	 +  2 # ribavirin	 +  2 # ribavirin	 +  2 # ribavirin	 +  2 # ribavirin	 +  2 # ribavirin
 Night 	 	 	 	 	 	 	 
 Peginterferon Mark day & time						7 PM	

If You Miss a Dose

BOCEPREVIR:

It is very important that you **do not miss any doses**. The hepatitis C virus can become resistant to boceprevir if you miss even a single dose. Future drugs also may be ineffective if resistance develops.

- If you take your pills within an hour of your scheduled time, it is not considered a missed dose.
- If your next dose is due more than 2 hours from now, **TAKE THE MISSED DOSE NOW** with food and continue with your normal schedule.
- If your next dose is due less than 2 hours from now, **SKIP THE MISSED DOSE** and continue with your normal schedule.
- If you miss multiple doses, call your prescriber contact, listed in the inside cover of this booklet.

RIBAVIRIN:

If you miss more than 1-2 days of ribavirin, call your prescriber contact listed in the inside cover of this booklet.

PEGINTERFERON:

If you miss any dose of peginterferon, call your prescriber contact listed in the inside cover of this booklet.

TIP: USE A MEDICINE TIMER!



The best way to ensure that you do not miss a dose is to remind yourself of your pill times by setting a timer. Perhaps use the alarm on your cell phone if you have one.

How Long Does Treatment Last?

Your treatment with the combination of boceprevir, peginterferon and ribavirin may be prescribed for as short as 28 weeks, or as long as 48 weeks, depending on how the virus responds to treatment.

If you are not responding to therapy, treatment may be stopped as early as 12 weeks.

After Treatment is Completed

Once you have finished your treatment, you will continue to have blood draws. One of the blood draws will be done 6 months after treatment is complete. If the virus is still undetectable at that time, then you have successfully cleared the virus and are cured.

If Treatment Doesn't Work for You

Treatments for hepatitis C are changing, and new drugs are always being tested. Some are currently in clinical trials and may be available within a few years. Keep in mind that even if the treatment didn't get rid of the virus, it may have improved the health of your liver.

It's important to continue to see your provider for follow-up. Your provider will set up a schedule for appointments.

Which Treatment Track Should YOU Follow?

Often patients who were treated before but not cured are not sure what happened during their course of treatment. It is completely fine if you are not sure. But before you start treatment with boceprevir, you need to talk over your history carefully with your doctor so that you both understand what happened in earlier treatments.

Also, if you are not sure whether you have cirrhosis (a scarred liver), please specifically ask your doctor.

With this information, you and your doctor can find your treatment track below:

	Tracks A & B	Track C	Track D
I have cirrhosis			X
I have never been treated before I do not have cirrhosis	X		
I have been treated before and the virus came down somewhat I do not have cirrhosis		X	
I have been treated before and the virus cleared but then returned I do not have cirrhosis		X	
I have been treated before and the virus did not respond at all I do not have cirrhosis			X

Treatment Tracks

■ TRACKS A & B

Treatment Roadmap.....	10
My Medication Schedule.....	12

■ TRACK C


Treatment Roadmap.....	14
My Medication Schedule.....	16

■ TRACK D

Treatment Roadmap.....	18
My Medication Schedule.....	20

Tracks A & B Treatment Roadmap

If it's your first time being treated, AND
you do not have cirrhosis



TRACK A1: 2 drugs for 4 weeks, 3 drugs for 24 weeks. Total: 28 weeks.

TRACK A2: 2 drugs for 4 weeks, 3 drugs for 32 weeks, 2 drugs for 12 weeks.
Total: 48 weeks.

TRACK B: 2 drugs for 4 weeks, 3 drugs for 44 weeks. Total: 48 weeks.

*IU stands for “international unit”—a standard measure of how much hepatitis C virus is in the blood.

Here are the forks in your road:

- **AT WEEK 4:** My viral load _____
How far your viral load has dropped will determine if you go onto Track A or Track B.

- **AT WEEK 8:** My viral load _____
Track A: If viral load undetectable: Go on Track A1 for a shortened, 28-week course.
If viral load detectable: Go on Track A2 for a full 48-week course.
Track B: Continue treatment.

- **AT WEEK 12:** My viral load _____
All Tracks: If viral load less than 100 IU: Continue treatment.
If viral load greater or equal to 100 IU: Treatment isn't working and will be stopped.

- **AT WEEK 24:** My viral load _____
All Tracks: If viral load undetectable: Continue treatment.
If viral load detectable: Treatment isn't working and will be stopped.

- **AT WEEK 28:**
Track A1: See "After Treatment is Completed," p. 7.

- **AT WEEK 36:** My viral load _____
Track A2: If viral load undetectable: Stop boceprevir. Continue to take peginterferon and ribavirin.
If viral load detectable: Treatment isn't working and will be stopped.
Track B: If viral load undetectable: Continue treatment with all 3 drugs.
If viral load detectable: Treatment isn't working and will be stopped.

- **AT WEEK 48:**
Tracks A2 & B: See "After Treatment is Completed," p. 7.

Tracks A & B (cont.)

My Medication Schedule



Ribavirin:

Take ___ pills every morning and ___ pills every afternoon or evening, as prescribed. You can take ribavirin with boceprevir, or at separate times of day.



Peginterferon:

Pegasys® ___ mcg/week or Peg-Intron® ___ ml/week
Inject weekly--same day, same time



Boceprevir:

Take 4 capsules, every 8 hours, with a light meal or snack




















Take with food

2 DRUGS

All Tracks: Weeks 1-4















































Start date: _____ Take until: _____

Time of Day	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Morning  ___ : ___ am	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin
Afternoon  ___ : ___ pm	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin
 Peginterferon Mark day & time							

3 DRUGS


















Track A1 & B: Weeks 5 to end Start date:* _____ Take until: _____

Track A2: Weeks 5-36 Start date:* _____ Take until: _____

Time of Day	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Morning  ____ : ____ am	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin
Afternoon  ____ : ____ pm	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin
Night  ____ : ____ pm	 	 	 	 	 	 	 
 Peginterferon Mark day & time							

2 DRUGS


Track A2: Weeks 37-48 Start date:* _____ Take until: _____

Time of Day	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Morning  ____ : ____ am	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin
Afternoon  ____ : ____ pm	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin
 Peginterferon Mark day & time							

*It can take time to receive lab results. Confirm with your provider the exact date to begin.

Track C Treatment Roadmap

You were treated before and the virus came down somewhat or
the virus cleared but came back AND
You do not have cirrhosis



TRACK C1: 2 drugs for 4 weeks, 3 drugs for 32 weeks. Total: 36 weeks.

TRACK C2: 2 drugs for 4 weeks, 3 drugs for 32 weeks, 2 drugs for 12 weeks.
Total: 48 weeks.

*IU stands for “international unit”—a standard measure of how much hepatitis C virus is in the blood.

HERE ARE THE FORKS IN YOUR ROAD:

- **AT WEEK 8:** My viral load _____
If viral load undetectable: Go on Track C1 for a shortened, 36-week course.
If viral load detectable: Go on Track C2 for a full 48-week course.

- **AT WEEK 12:** My viral load _____
Both Tracks: If viral load less than 100 IU: Continue treatment.
If viral load greater or equal to 100 IU: Treatment isn't working and will be stopped.

- **AT WEEK 24:** My viral load _____
Both Tracks: If viral load undetectable: Continue treatment.
If viral load detectable: Treatment isn't working and will be stopped.

- **AT WEEK 36:** My viral load _____
Track C1: See "After Treatment is Completed," p. 7.
Track C2: If viral load undetectable: Stop boceprevir. Take peginterferon and ribavirin only.
Track C2: If viral load detectable: Treatment isn't working and will be stopped

- **AT WEEK 48:**
Tracks C2: See "After Treatment is Completed," p. 7.

Track C (cont.) My Medication Schedule



Ribavirin:

Take ___ pills every morning and ___ pills every afternoon or evening, as prescribed. You can take ribavirin with boceprevir, or at separate times of day.



Peginterferon:

Pegasys® ___ mcg/week or Peg-Intron® ___ ml/week
Inject weekly--same day, same time



Boceprevir:

Take 4 capsules, every 8 hours, with a light meal or snack



Take with Food

2 DRUGS








































Weeks 1-4

Start date: _____ Take until: _____

Time of Day	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Morning ___:___ am	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin
Afternoon ___:___ pm	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin
 Peginterferon Mark day & time							


















3 DRUGS

Track C1 and C2: Weeks 5-36 Start date*: _____ Take until: _____

Time of Day	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Morning  ____ : ____ am ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin
Afternoon  ____ : ____ pm ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin
Night  ____ : ____ pm ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin
 Peginterferon Mark day & time							

2 DRUGS


Track C2: Weeks 37-48 Start date*: _____ Take until: _____

Time of Day	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Morning  ____ : ____ am ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin
Afternoon  ____ : ____ pm ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin	 ____ # ribavirin
 Peginterferon Mark day & time							

*It can take time to receive lab results. Confirm with your provider the exact date to begin.

Track D Treatment Roadmap

You have cirrhosis OR
You were treated before and had no viral response



TRACK D: Take 2 drugs for 4 weeks, then 3 drugs for 36 weeks. Total: 48 weeks.

*IU stands for “international unit”—a standard measure of how much hepatitis C virus is in the blood.

HERE ARE THE FORKS IN YOUR ROAD:

- **AT WEEK 12:** My viral load _____

If viral load less than 100 IU: Continue treatment.

If viral load greater or equal to 100 IU: Treatment isn't working and will be stopped.

- **AT WEEK 24:** My viral load _____

If viral load undetectable: Continue treatment.

If viral load detectable: Treatment isn't working and will be stopped.

- **AT WEEK 36:** My viral load _____

If viral load undetectable: Continue treatment.

If viral load detectable: Treatment isn't working and will be stopped.

- **AT WEEK 48:**

See "After Treatment is Completed," p. 7.

Track D (cont.)

My Medication Schedule



Ribavirin:

Take ___ pills every morning and ___ pills every afternoon or evening, as prescribed. You can take ribavirin with boceprevir, or at separate times of day.



Peginterferon:

Pegasys® ___ mcg/week or Peg-Intron® ___ ml/week
Inject weekly--same day, same time



Boceprevir:

Take 4 capsules, every 8 hours, with a light meal or snack



Take with food















































2 DRUGS

Weeks 1-4 Start date: _____ Take until: _____

Time of Day	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Morning ___:___ am ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin
Afternoon ___:___ pm ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin	 ___ # ribavirin
 Peginterferon Mark day & time							

3 DRUGS

Weeks 5-48 Start date*: _____ Take until: _____

Time of Day	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Morning  ____ : ____ am	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin
Afternoon  ____ : ____ pm	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin
Night  ____ : ____ pm	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin	 +  ____ # ribavirin
 Peginterferon Mark day & time							

*It can take time to receive lab results. Confirm with your provider the exact date to begin.

Dealing with Side Effects

*indicates brand name

SIDE EFFECT	WHAT TO DO
<p>FEVER/CHILLS</p>	<p>Try injecting interferon at bedtime if fever/chills develop 1-3 hours after the interferon injection</p> <p>Take acetaminophen (Tylenol[*]) about 30-60 minutes before weekly interferon injection</p> <p>Take acetaminophen (Tylenol[*]) 1-2 tablets prior to interferon injection and repeat 4-6 hours later if needed up to a maximum of 2,000 mg/day. Thus, do not exceed 6 tablets/day of 325 mg or 4 tablets/day of 500 mg acetaminophen (Tylenol[*]).</p> <p>Use ibuprofen [Advil[*], Motrin[*]] or naproxen [Aleve[*]] if approved by your provider and you do not have cirrhosis</p> <p>Try a cool sponge bath, ice pack or cold pack when you have a fever</p> <p>Use extra blankets and clothes when you have the chills</p> <p>Notify your provider if your temperature is above 101°F for more than 24-48 hours</p>
<p>MUSCLE AND BODY ACHES</p>	<p>Try injecting interferon at bedtime if body aches develop 1-3 hours after the interferon injection</p> <p>Take acetaminophen (Tylenol[*]) about 30-60 minutes before weekly interferon injection</p> <p>Take acetaminophen (Tylenol[*]) 1-2 tablets prior to interferon injection and repeat 4-6 hours later if needed up to a maximum of 2,000 mg/day. Thus, do not exceed 6 tablets/day of 325 mg or 4 tablets/day of 500 mg of acetaminophen (Tylenol[*]).</p> <p>Use ibuprofen [Advil[*], Motrin[*]] or naproxen [Aleve[*]] if approved by your provider and you do not have cirrhosis</p> <p>Try low-impact exercise such as walking or low-impact aerobics as directed by your provider</p> <p>Maintain adequate fluid intake (at least six to eight non-caffeinated 8-oz glasses/day)</p> <p>Apply warm moist heat or massage areas</p>

SIDE EFFECT	WHAT TO DO
HEADACHES	<p>Maintain adequate fluid intake (at least six to eight non-caffeinated 8-oz glasses/day)</p> <p>Take acetaminophen (Tylenol*) or ibuprofen (Motrin*, Advil*) as directed by your provider</p> <p>Keep lights dim, wear sunglasses or stay in darkened rooms</p> <p>Try to get plenty of rest</p>
FATIGUE	<p>Try low-impact exercise such as walking or low-impact aerobics as directed by your provider</p> <p>Maintain adequate fluid intake (at least six to eight non-caffeinated 8-oz glasses/day; you may have a caffeinated beverage in the morning)</p> <p>Take a short nap during the day</p> <p>Lessen your work schedule if possible</p> <p>Eat well-balanced meals every day</p>
DEPRESSION	<p>Talk to your provider about your symptoms and medications that might help</p> <p>Always discuss thoughts of harming yourself or someone else with your provider</p> <p>Go to the nearest emergency room if you are in danger of harming yourself or others</p> <p>Do mild to moderate exercises at least 3 times/week as directed by your provider</p> <p>Join a hepatitis C support group</p>

SIDE EFFECT	WHAT TO DO
<p>ANXIETY AND IRRITABILITY</p>	<p>Talk to your provider about your symptoms and medications that might help</p> <p>Always discuss thoughts of harming yourself or someone else with your provider</p> <p>Go to the nearest emergency room if you are in danger of harming yourself or others</p> <p>Do mild to moderate exercises at least 3 times/week as directed by your provider</p> <p>Try relaxation techniques such as deep breathing, taped exercises, yoga, Tai Chi or meditation</p> <p>Avoid stimulants like caffeine and maintain adequate fluid intake</p> <p>Join a hepatitis C support group</p>
<p>INSOMNIA</p>	<p>Go to sleep and wake up at the same time every day</p> <p>Do not read or watch TV in bed</p> <p>Limit daytime naps</p> <p>If ribavirin makes you jittery, take it before 6 pm instead of right before bedtime</p> <p>Limit fluid intake for 2 hours before bedtime to avoid having to get up to go to the bathroom</p> <p>Avoid caffeinated products, especially in the afternoon and at night</p> <p>Avoid heavy meals close to bedtime</p> <p>Take warm baths, read or listen to music, get a massage</p> <p>Try a glass of warm milk (contains tryptophan, a natural sleep agent)</p> <p>Diphenhydramine (Benadryl*) or other medications may be recommended by your provider</p>

SIDE EFFECT	WHAT TO DO
<p>DRY MOUTH OR MOUTH ULCERS</p>	<p>Brush teeth frequently, especially after eating</p> <p>Avoid mouthwash containing alcohol</p> <p>Drink plenty of water or use ice chips or sugar-free lemon drops</p> <p>Other medications may be needed for mouth sores/ulcers</p> <p>Ask your provider about medications that may help</p>
<p>BAD TASTE IN MOUTH</p>	<p>Use sugar-free lemon drops or real lemon wedges</p> <p>Eat a small amount of yogurt, 1/2 hour before meals or eat dark chocolate (monitor your glucose levels if you have diabetes)</p> <p>Drink lemonade or cranberry juice (monitor your glucose levels if you have diabetes)</p> <p>Eat food cold or at room temperature</p> <p>Brush teeth frequently, especially after eating, to eliminate metallic taste</p> <p>Use plastic utensils if experiencing metallic taste</p>
<p>POOR APPETITE</p>	<p>Eat smaller, more frequent (4-6) meals throughout the day</p> <p>Drink protein drinks (Carnation Instant Breakfast*, Ensure* or Boost*)</p> <p>Eat snacks with protein (cheese, peanut butter, eggs)</p> <p>Eat whatever appeals to you even if you're not hungry. Eat a variety of foods</p> <p>Walk before a meal</p>

SIDE EFFECT	WHAT TO DO
<p>NAUSEA AND VOMITING</p>	<p>Take ribavirin with food</p> <p>Eat small meals</p> <p>Avoid foods or smells that trigger nausea. Try eating cold foods and avoiding cooking smells</p> <p>Eat healthy foods. Avoid greasy, spicy, acidic or sweet foods</p> <p>Try eating ginger in ginger tea, ginger ale or gingersnaps</p> <p>Eat some crackers or dry white toast if you feel sick in the morning</p> <p>Over-the-counter antacids or other medications may be recommended by your provider</p>
<p>DIARRHEA</p>	<p>Eat more soluble fiber like Bananas, white Rice, Applesauce and white Toast (the “BRAT” diet)</p> <p>Avoid foods that are spicy or acidic (like citrus)</p> <p>Avoid dairy products up to several days after diarrhea resolves</p> <p>Maintain adequate fluid intake (at least six to eight 8-oz glasses/day)</p> <p>Your provider may recommend loperamide (Imodium*) or methylcellulose (Citrucel*) or psyllium (Metamucil*)</p>
<p>DEHYDRATION</p>	<p>Increase intake of water or non-caffeinated beverages; try to drink at least 8-10 glasses of water each day.</p> <p>Water down juices, Gatorade* or Crystal Light*</p> <p>Avoid caffeinated beverages</p>
<p>COUGH</p>	<p>Increase intake of water or non-caffeinated beverages</p> <p>Use a humidifier</p> <p>Try sugar-free hard candy or cough drops</p>

SIDE EFFECT	WHAT TO DO
<p>DRY SKIN/ RASHES</p>	<p>Avoid long, hot showers or baths</p> <p>Use moisturizing soaps (Dove*, Oil of Olay*, Cetaphil*)</p> <p>Use moisturizing lotion after showers (Aquaphor*, Absorbace*, Lac-Hydrin*)</p> <p>Use mild unscented laundry detergents and avoid fabric softeners</p> <p>Use sunscreen</p> <p>Try rubbing or pressing on the itchy areas rather than scratching</p> <p>Use petroleum jelly on dry, itchy areas. Your provider may recommend other agents to help</p>
<p>HAIR THINNING OR HAIR LOSS</p>	<p>Avoid harsh hair products such as dyes, perms, gels, sprays and mousses</p> <p>Use a mild shampoo such as baby shampoo</p> <p>Avoid braiding hair; use a wide-tooth comb or soft brush</p> <p>Wear a cap, scarf, turban or wig</p>
<p>INJECTION SITE REACTIONS</p>	<p>Before injecting, warm your medicine by gently rolling the syringe in your hands for a minute</p> <p>Rotate/alternate your injection site—thigh, upper arm and abdomen. If you are thin, use the thigh area</p> <p>Do not inject into an area that is irritated, bruised or red</p> <p>Do not rub injection site</p> <p>Apply a cold pack</p> <p>Your provider may recommend applying hydrocortisone cream or other medications to help</p>

My Viral Load Chart

As you progress through treatment, you may find it helpful to mark down the viral load results of your blood work. (Viral load tests are required before you start treatment; for treatment weeks 4, 8, 12, 24, 36, and 48; and 6 months after treatment.)

Date:	Week:	Results:
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

OFFICE OF PUBLIC HEALTH

U.S. DEPARTMENT OF VETERANS ADMINISTRATION

JULY 2012

WWW.HEPATITIS.VA.GOV