

TABLE 11—FINAL 2012 AND 2013 LISTED BSAI AMERICAN FISHERIES ACT CATCHER/PROCESSOR GROUND FISH  
 SIDEBOARD LIMITS

[Amounts are in metric tons]

Target species	Area/season	1995–1997			2012 ITAC available to trawl C/Ps <sup>1</sup>	2012 AFA C/P side-board limit	2013 ITAC available to trawl C/Ps <sup>1</sup>	2013 AFA C/P side-board limit
		Retained catch	Total catch	Ratio of retained catch to total catch				
Sablefish trawl .....	BS .....	8	497	0.016	948	15	935	15
	AI .....	0	145	0	436	0	429	0
Atka mackerel .....	Central AI A season <sup>2</sup> .	n/a	n/a	0.115	4,806	553	3,966	456
	Central AI B season <sup>2</sup> .	n/a	n/a	0.115	4,806	553	3,966	456
	Western AI A season <sup>2</sup> .	n/a	n/a	0.2	670	134	670	134
	Western AI B season <sup>2</sup> .	n/a	n/a	0.2	670	134	670	134
Rock sole .....	BSAI .....	6,317	169,362	0.037	77,691	2,875	77,691	2,875

Greenland turbot ...	BS	121	17,305	0.007	5,296	37	5,109	36
	AI	23	4,987	0.005	2,066	10	1,717	9
Arrowtooth flounder	BSAI	76	33,987	0.002	21,250	43	21,250	43
Kamchatka flounder.	BSAI	76	33,987	0.002	15,045	30	15,045	30
Flathead sole .....	BSAI	1,925	52,755	0.036	30,482	1,097	30,482	1,097
Alaska plaice .....	BSAI	14	9,438	0.001	20,400	20	20,400	20
Other flatfish .....	BSAI	3,058	52,298	0.058	2,720	158	2,720	158
Pacific ocean perch.	BS	12	4,879	0.002	4,854	10	5,559	11
	Eastern AI	125	6,179	0.02	5,019	100	5,751	115
	Central AI	3	5,698	0.001	4,456	4	5,099	5
	Western AI	54	13,598	0.004	7,483	30	8,582	34
Northern rockfish ..	BSAI	91	13,040	0.007	4,700	33	4,700	33
Shortraker rockfish	BSAI	50	2,811	0.018	393	7	393	7
Rougheye rockfish	EBS/EAI	50	2,811	0.018	231	4	241	4
	CAI/WAI	50	2,811	0.018	244	4	258	5
Other rockfish .....	BS	18	621	0.029	500	15	500	15
	AI	22	806	0.027	570	15	570	15
Squids .....	BSAI	73	3,328	0.022	361	8	361	8
Skates .....	BSAI	553	68,672	0.008	24,700	198	24,746	198
Sharks .....	BSAI	553	68,672	0.008	200	2	200	2
Octopuses .....	BSAI	553	68,672	0.008	900	7	900	7
Sculpins .....	BSAI	553	68,672	0.008	5,200	42	5,200	42

<sup>1</sup> Aleutian Islands Pacific ocean perch, and BSAI Atka mackerel, flathead sole, rock sole, yellowfin sole are multiplied by the remainder of the TAC after the subtraction of the CDQ reserve under § 679.20(b)(1)(ii)(C).

<sup>2</sup> The seasonal apportionment of Atka mackerel in the open access fishery is 50 percent in the A season and 50 percent in the B season. Listed AFA catcher/processors are limited to harvesting no more than zero in the Eastern Aleutian District and Bering Sea subarea, 20 percent of the annual ITAC specified for the Western Aleutian District, and 11.5 percent of the annual ITAC specified for the Central Aleutian District.