



## Health Disparities Calculator

<http://seer.cancer.gov/hdcalc/>

### Overview

The Health Disparities Calculator (HD\*Calc) is designed to generate multiple summary measures to evaluate and monitor health disparities. The HD\*Calc statistical software can be used either as an extension of SEER\*Stat—allowing users to import Surveillance, Epidemiology, and End Results (SEER) data—or with other population-based health data, including data from the National Health Interview Survey (NHIS), California Health Interview Survey (CHIS), Tobacco Use Supplement to the Current Population Survey (TUS-CPS), and National Health and Nutrition Examination Survey (NHANES).

HD\*Calc can be used to produce 11 summary measures of disparity. Cancer rates and other data categorized by disparity groups such as race/ethnicity, socioeconomic status (SES), and geographic area can be used with HD\*Calc to generate output that can be presented in both tabular and graphic formats. HD\*Calc program users are able to specify a range of conditions and formats.

HD\*Calc can be used to apply the measures of health disparities described in the NCI monograph, *Methods for Measuring Cancer Disparities: A Review Using Data Relevant to Healthy People 2010 Cancer-Related Objectives* (NCI, 2005; <http://seer.cancer.gov/publications/disparities>). This monograph evaluates major

issues that may affect the choice of summary measures of disparity, and it is recommended for those who are unfamiliar with the measures available in HD\*Calc.

For those interested in a comparative look at measures of health disparities, the monograph *Selected Comparisons of Measures of Health Disparities: A Review Using Databases Relevant to Healthy People 2010 Cancer-Related Objectives* (NCI, 2007; <http://seer.cancer.gov/publications/disparities2>) uses case studies to analyze the performance and appropriateness of various health disparity measures.

These monographs are recommended to anyone involved in health disparities research.

### Use HD\*Calc to:

- Quickly and accurately compare different measures of disparities by race/ethnicity, SES, and geographic area;
- Evaluate multiple summary measures for monitoring and presenting health disparities;
- Graphically explore underlying trends in data; and
- Easily export tables and graphs.

## HD\*Calc Tutorials

Exercises on the following topics are available at <http://seer.cancer.gov/hdcalc/tutorials>:

- Preparing SEER cancer incidence data using SEER\*Stat (<http://seer.cancer.gov/seerstat>);
- Importing the data into HD\*Calc for analysis;
- Using data that did not come from SEER\*Stat to calculate health disparities in HD\*Calc.

The tutorials demonstrate how to measure health disparities:

- By race using SEER incidence data.
- By SES group using SEER incidence data.
- By race using SEER incidence data showing a 3-year moving average.
- By income level using CHIS colorectal cancer screening data.

## References

Harper S, Lynch J. Methods for measuring cancer disparities: using data relevant to Healthy People 2010 cancer-related objectives. NCI Cancer Surveillance Monograph Series, Number 6. Bethesda, MD: National Cancer Institute, 2005. NIH Publication No. 05-5777.

Harper S, Lynch J. Selected comparisons of measures of health disparities: a review using databases relevant to Healthy People 2010 cancer-related objectives. NCI Cancer Surveillance Monograph Series, Number 7. National Cancer Institute. NIH Pub. No. 07-6281, Bethesda, MD, 2007.

Harper S, Lynch J, Meersman, SC, Breen N, Davis WW, Reichman ME. An overview of methods for monitoring social disparities in cancer with an example using trends in lung cancer incidence by area-socioeconomic position and race-ethnicity, 1992-2004. Am J Epidemiol. 2008 Apr 15;167(8):889-99. Epub 2008 Mar 15. (<http://aje.oxfordjournals.org/cgi/content/abstract/167/8/889>)

Harper S, Lynch J, Meersman, SC, Breen N, Davis WW, Reichman ME. Trends in area-socioeconomic and race-ethnic disparities in breast cancer incidence, stage at diagnosis, screening, mortality, and survival among women ages 50 years and over (1987-2005). Cancer Epidemiol Biomarkers Prev. 2009; 18(1):121-31. (<http://cebp.aacrjournals.org/cgi/content/abstract/18/1/121>)

Harper S, King NB, Meersman SC, Reichman ME, Breen N, Lynch J. Implicit value judgments in the measurement of health inequalities. Milbank Q. 2010 Mar;88(1):4-29.

**For more information, contact [hdcalculator@imsweb.com](mailto:hdcalculator@imsweb.com).**