

## NEW EVIDENCE FOR THE ROLE OF EMERGING FLUX IN A SOLAR FILAMENT'S SLOW RISE PRECEDING ITS CME-PRODUCING FAST ERUPTION

ALPHONSE C. STERLING<sup>1</sup>

NASA Marshall Space Flight Center, Space Science Office, VP62, Huntsville, AL 35805; alphonse.sterling@nasa.gov

LOUISE K. HARRA

Mullard Space Science Laboratory, University College London, Holmbury St. Mary, Dorking, Surrey RH5 6NT, UK; lkh@mssl.ucl.ac.uk

AND

RONALD L. MOORE

NASA Marshall Space Flight Center, Space Science Office, VP62, Huntsville, AL 35805; ron.moore@nasa.gov

Received 2007 January 28; accepted 2007 June 12

### ABSTRACT

We observe the eruption of a large-scale ( $\approx 300,000$  km) quiet-region solar filament leading to an Earth-directed “halo” coronal mass ejection (CME), using data from EIT, CDS, MDI, and LASCO on *SOHO* and from SXT on *Yohkoh*. Initially the filament shows a slow ( $\sim 1$  km s<sup>-1</sup> projected against the solar disk) and approximately constant velocity rise for about 6 hr, before erupting rapidly, reaching a velocity of  $\sim 8$  km s<sup>-1</sup> over the next  $\approx 25$  minutes. CDS Doppler data show Earth-directed filament velocities ranging from  $< 20$  km s<sup>-1</sup> (the noise limit) during the slow-rise phase, to  $\sim 100$  km s<sup>-1</sup> early in the eruption. Beginning within 10 hr prior to the start of the slow rise, localized new magnetic flux emerged near one end of the filament. Near the start of and during the slow-rise phase, soft X-ray (SXR) microflaring occurred repeatedly at the flux-emergence site, and the magnetic arcade over the filament progressively brightened in a fan of illumination in SXR. These observations are consistent with “tether-weakening” reconnection occurring between the newly emerging flux and the overlying arcade field containing the filament, and apparently this reconnection is the cause of the filament’s slow rise. We cannot, however, discern whether the transition from slow rise to fast eruption was caused by a final episode of tether-weakening reconnection, or by one or some combination of other possible mechanisms allowed by the observations. Intensity “dimmings” and “brightenings” occurring both near to and relatively far from the location of the filament are possible signatures of the expansion (“opening”) of the erupting field and its reconnection with overarching field during the eruption.

*Subject headings:* Sun: coronal mass ejections (CMEs) — Sun: filaments — Sun: flares — Sun: UV radiation — Sun: X-rays, gamma rays

*Online material:* mpeg animations

### 1. INTRODUCTION

Solar eruptions result in the rapid expulsion from the Sun of magnetic field, material, and radiations in the form of coronal mass ejections (CMEs) and solar flares. Why eruptions occur is a major outstanding question of solar physics. Research over the past several decades, and in particular that based on observations from space over the last three decades, has vastly improved our understanding of these events and led to the formation of several ideas for the eruption mechanism.

We know that eruptions occur at locations on the Sun where there is a buildup of a large amount of magnetic energy, and eruptions occur when this energy is released via some process. Many of the eruption models deal with how this energy release is accomplished. For example, Forbes & Isenberg (1991) and Lin et al. (1998) discuss how convergence of magnetic flux below a flux tube situated on a magnetic-polarity inversion line can lead to enhancement of the flux tube and its eventual disruption due to an instability. Chen & Shibata (2000) present a model where eruption results when tension in a sheared arcade or flux rope field is weakened via reconnection between that main field and an emerging flux; this is similar to the tether-weakening and tether-cutting

schemes presented by Moore & Roumeliotis (1992) and by Moore & LaBonte (1980), Sturrock (1989), and Moore et al. (2001). Antiochos (1998) and Antiochos et al. (1999) discuss instead disruption of a sheared core field from above, when tension between the core field and an overlying restraining field (of polarity opposite that of the core field) is reduced via reconnection between the two field systems. Another theoretical possibility for the eruption cause is kinking (kink instability) of an emerging flux rope (e.g., Sturrock et al. 2001; Linker et al. 2001; Fan & Gibson 2004; Rust & LaBonte 2005; Gibson & Fan 2006). Some reviews describing basic aspects of several eruption ideas are Forbes (2000), Klimchuk (2001), Lin et al. (2003), and Moore & Sterling (2006).

To determine which (if any) of these mechanisms is predominantly responsible for eruptions, it is necessary to observe the evolution of eruptions and compare with expectations from the models. In particular, we would like to know the behavior of the magnetic field around the core of the erupting region prior to and during the explosive eruption. We have been attempting to study this for several events through detailed analysis using filaments as tracers of the otherwise-invisible evolving field, and for some of these also examining intensity variations (dimmings and brightenings) in EUV and soft X-rays (SXR) that can be indicative of evolution of the field outside of the filament (Sterling & Moore 2003, 2004a, 2004b, 2005). The purpose of the present

<sup>1</sup> Current address: JAXA, Institute of Space and Astronautical Science, Hinode Group, 3-1-1 Yoshinodai, Sagami-hara, Kanagawa 229-8510, Japan.

paper is to continue these investigations by studying an eruption that occurred near solar disk center on 2001 February 28. Similar to the previous eruptions we have reported on, this eruption also involves a filament whose early movement is a slow rise (as often occurs; e.g., Tandberg-Hanssen et al. 1980; Kahler et al. 1988; Feynman & Ruzmaikin 2004), indicative of a slow preeruption expansion of the magnetic fields involved in the eruption. We also investigate spectroscopic Doppler motions of the filament and associated features from around the time of eruption onset for this event. We show that for this event there is strong evidence that the cause of the slow rise of the filament was magnetic reconnection between the main filament field and newly emerging flux, but we could not isolate a single cause of the onset of the rapid eruption (similar to the collective results of several of our previous investigations; Moore & Sterling 2006).

This eruption occurred in a quiet region, and therefore the evolution toward eruption was relatively slow compared to typical active-region eruptions; this is a consequence of the relatively weak field strength in quiet regions, and correspondingly lower Alfvén velocities. Because of this slow evolution, we are able to follow closely the timings of filament motions compared with various intensity changes, even with the relatively low or non-uniform cadence of our observations.

## 2. INSTRUMENTATION AND DATA

We use imaging, magnetic, and spectroscopic data for our analysis. Imaging data include those in EUV from the EUV Imaging Telescope (EIT; Delaboudinière et al. 1995) on the *Solar and Heliospheric Observatory* (*SOHO*) satellite, and in SXR from the Soft X-ray Telescope (SXT; Tsuneta et al. 1991) on the *Yohkoh* satellite. Magnetic data are from the Michelson Doppler Imager (MDI; Scherrer et al. 1995) and spectroscopic data, including rastered images, are from the Coronal Diagnostic Spectrometer (CDS; Harrison et al. 1995), both on *SOHO*.

Our EIT data are primarily from its 195 Å Fe XII filter at a 12 minute cadence and spatial pixel resolution of 2.6". EIT's full-disk coverage is ideal for observing larger scale eruptions, such as the one we study here. We use mainly full-disk AlMg-filter images from SXT, with cadence varying from about 2 minutes, to over an hour during times of spacecraft night or data gaps. Filaments are visible in the EUV images (e.g., Mein et al. 2001; Schmieder et al. 2004; Anzer & Heinzel 2005), and we use these images to track the filament movements.

CDS provides spectral information in various lines, and for this study we generally used O V data. CDS has a slit length of 240", and a spatial resolution of 4.06" × 1.68". We used CDS images generated by rastering across an area of the Sun from west to east (right to left in the images). Each raster consisted of 260 steps and took ≈10 minutes, and so that is the cadence of our CDS raster images. For the current set of observations the CDS field of view (FOV) was 250" × 160", and it was in a mode where the rastering occurred repeatedly over the same fixed heliocentric location, with solar features rotating with time into its FOV from the east and out of its FOV in the west. In computing velocities in the region of the filament eruption, we took zero velocity to be the average of the rasters before the filament appeared in the CDS field of view. Hence the Doppler velocities measured are relative to the quiet background velocity, which is the most accurate way to deal with this issue. The data were calibrated using the standard CDS software, which corrects for geometrical distortions as well as making the standard instrumental corrections.

The eruption occurred on 2001 February 28, beginning near 12:00 UT, with the filament that erupted located at longitude between about 5° and 15° west of Sun center and latitude between




FIG. 1.—LASCO difference image at 16:50 UT from the C2 coronagraph, showing the halo CME resulting from the filament eruption of this paper. The insert is an EIT difference image from 16:48 UT. This image is from the LASCO online catalog (Yashiro et al. 2004).

5° north and 20° south. It occurred in a weak magnetic region, with only small spots present. Evolution toward eruption was relatively slow, with a slow-rise preeruption phase lasting several hours (compared to typically only a few minutes for strong-field active-region eruptions) and the SXR output relatively weak; SXR flux from the 1–8 Å channel of the *GOES* satellite showed a weak increase above background to the mid-B level near 13:30 UT that likely resulted from this eruption, and several even weaker increases at earlier times that may have been from imaged SXR enhancements that we observed during the preeruption phase (see Fig. 5). White-light coronal images from the Large Angle and Spectrometric Coronagraph Experiment (LASCO; Brueckner et al. 1995) on *SOHO* recorded a partial halo CME from 14:50 UT, with a possible precursor visible from as early as 13:50 UT (Fig. 1). Its projected velocity against the plane of the sky was ≈300 km s<sup>-1</sup>, according to the LASCO CME catalog (Yashiro et al. 2004).

## 3. OBSERVATIONS AND ANALYSIS

### 3.1. Filament Motions

Figure 2 shows EIT images of the erupting region, where arrows point out the filament. Prior to eruption the filament undergoes slow movement beginning about 6 hr prior to eruption; it shows substantial change between Figures 2a and 2b, particularly the southern portion of the filament (compare the two southernmost arrows in the two panels). By the time of Figure 2c the filament has erupted, with a bright flare loop visible in box 5 and regions around boxes 2 and 3 having undergone intensity dimming compared to the previous panel. Figure 2d shows the situation several hours after eruption, where additional dimmings have occurred in boxes 6 and 7. The accompanying video is an EIT movie of the erupting filament.

Figure 3a (and also Fig. 5a) shows the movement of the filament along the fiducials of Figure 2a. Our error bars are 1  $\sigma$  uncertainties based on three independent measurements of the filament motion. Some of the motions prior to 6:00 UT may be real, indicating that filament is undulating prior to eruption,




FIG. 2.—EUV images of the erupting region from EIT, taken with its 195 Å filter. Arrows in (a) and (b) point to the filament, which is starting to erupt in (b) and has erupted and by the time of (c), and therefore the filament is not visible in (c) or (d). The nearly horizontal lines in (a) are fiducials along which the height of the filament projected on the disk is measured in Figs. 3a and 5a. Boxes, labeled in (d), are regions for which light curves are generated in Fig. 3b. North is up and west is to the right here and in all solar images in this paper. [This figure is available as an mpeg animation in the electronic edition of the Journal.]




FIG. 3.—(a) Trajectory of the filament as a function of time along fiducials of Fig. 2a, with the bottom (top) trajectory from the southern (northern) fiducial. Initial heights are arbitrarily set at approximately 0 and 10,000 km, and motion upward (downward) reflects motions projected on the disk toward the west (east). Error bars are  $1\sigma$  uncertainties based on three independent measurements of the filament motion. The curve labeled “B flux” is from Fig. 8 and discussed further below. Vertical bars show times of the peaks of SXR microflare bursts in Fig. 5. (b) Light curves from correspondingly numbered boxed regions in Fig. 2. Curve 10 is from a region showing little change and represents background variations. Different line thicknesses, and dashed and solid lines, are for clarity only, and all plots are shifted arbitrarily on the vertical axis for clarity.

perhaps due to unresolved counterstreaming or twist (e.g., Wiik et al. 1997). At about 5:30 UT the filament begins moving out along the southern fiducial. This motion continues until about 11:30 UT, after which time the filament rapidly accelerates and erupts. Thus, for about 6 hr the filament undergoes a slow-rise phase prior to eruption, similar to what we have seen in earlier studies. Motion along the northern fiducial of Figure 2 becomes unambiguously pronounced after about 11:00 UT, although more subtle motions of the filament are apparent at earlier times (see the video accompanying Fig. 2), and these are reflected in the plot in Figure 3a. Motions of the northern portion of the filament are less obvious than in its southern portion, perhaps because the motions are directed along the line of sight, or because this portion of the filament is more stationary than the southern portion during the slow rise. Figure 3a shows the north filament’s curve apparently decreasing in height with time; actually it is merely moving in a direction opposite to that of the southern portion of the filament.

It is difficult to detect the northern portion of the filament at the time of eruption in the EIT images (just past 12:00 UT), and so the curve in the plot just terminates.

From Figure 3a we can measure the velocity of the filament. If we take the motion during the slow-rise phase to be constant, a least-square fit between 07:24:57 and 11:11:30 UT gives a speed of  $0.8\text{ km s}^{-1}$  for the slow rise along the southern fiducial. After eruption starts a linear fit does not seem appropriate for a very extended time period. Between 11:35:32 and 12:47:33 UT the motion is roughly linear, and a linear fit gives a speed of about  $7.5\text{ km s}^{-1}$ . These values are for motion projected against the disk; as the eruption is near disk center and appears to be roughly vertical, the radial velocity away the Sun will be substantially larger. We can estimate the outward velocity for the southern fiducial, assuming vertical motions. At the time of eruption the filament along that fiducial was between about  $8^\circ$  and  $12^\circ$  away from disk center. This gives the radial velocities to be between about 5 and 7 times the projected values; that is, the slow-rise velocity is  $5 \pm 1\text{ km s}^{-1}$ , and early in the eruption the velocity is  $45 \pm 8\text{ km s}^{-1}$ . Motions of the filament projected on the disk along the northern fiducial is initially to the east, opposite that of the southern part of the filament. This indicates that as the filament erupts it either twists as it rises, or it expands and appears to twist because different portions of it move toward opposite sides of the line of sight between the Sun and the *SOHO* satellite.

### 3.2. Intensity Variations during Slow-Rise Phase and Eruption

Figure 3b shows the EIT intensities as functions of time (light curves) integrated over the boxes of Figure 2. Box 5 is at the location of the main flaring loops, and its light curve shows that the primary flare starts in EUV near 13:00 UT. Weaker brightenings are discernible at earlier times at and around the box 5 location, and these brightenings are reflected in some of the fluctuations in the box 5 light curve from at least 10:00 UT. Box 1 is of a region that shows substantial low-level activity during the event; this is one of our most important boxes since, as we discuss in § 4, activity at this location could have played a primary role in the initiation of the eruption of the filament and CME. Boxes 2 and 3 are locations of a pair of dimmings (“twin dimmings”; § 3.3) that form on either end of the flare arcade. Boxes 6, 7, and 9 are at regions where we have identified dimmings that are farther removed from the main erupting region than those of boxes 2 and 3 (“remote dimmings”; § 3.3). Box 8 is a location where we also see intensity changes, but primarily in SXRs.

Figure 4 shows AIMg-filter SXR images from SXT of the same region as that in Figure 2. Loops in the region prior to eruption are consistent with a sheared magnetic geometry, with a roughly sigmoid shape (Sterling & Hudson 1997; Canfield et al. 1999). The filament is not visible in the SXR images, but arrows in Figure 4b point out a bright SXR loop; this is the last loop of a fan of loops that shows up prominently in the SXR images from about the time of the start of the filament’s slow-rise phase (see Fig. 3a), and persists until the time of eruption, as we discuss in § 4. This SXR feature has no obvious significant manifestation in the EUV images.

Figure 5 is analogous to Figure 3, but for SXRs instead of EUV. As in Figure 3, the box 5 light curve represents the flux of the flare arcade and the background light curve is again that of box 10. We use box 4 in place of box 3, however, because box 4 better captures the region that dims in SXRs than does box 3, while box 3 better captures the EUV dimming. Boxes 3 and 4 are separated by about  $60''$  (both are plotted in Fig. 6b); this is larger than the uncertainty in the alignment between the SXT and EIT images, which we estimate to be about  $10''$  based on the co-alignment of features




FIG. 4.—Similar to Fig. 2, but from SXR images from SXT, taken with its AIMg filter. Boxes, labeled in (d), are regions for which light curves are generated in Fig. 5b. Box 5 is omitted from (a) to show the central features more clearly. Arrows in (b) point to a loop that brightens in SXR near the start of the filament’s fast-rise phase. A coronal hole is visible in the northeast of the frames, east of  $-450$  in the abscissa, and between approximately  $-200$  and  $+400$  in the ordinate. [This figure is available as an mpeg animation in the electronic edition of the Journal.]




FIG. 5.—(a) Same as in Fig. 3a, but with the filament trajectory plotted on a logarithmic scale; error bars are given in Fig. 3a but omitted here for clarity. The curve labeled “B flux” is from Fig. 8 and discussed below. (b) Light curves from correspondingly numbered boxed regions in Fig. 4. Curve 10 is from a region showing little change and represents background variations. Curve 2 has been magnified by a factor of 5 to show better its variations. Different line thicknesses, and dashed and solid lines, are for clarity only, and all plots are shifted arbitrarily on the vertical axis for clarity. Stars on curve 10 are plotted at times of SXT images, showing the cadence of the observations. In (a), vertical bars show times of the peaks of SXR microflare bursts from the box 1 light curve of (b).

in the other boxes between Figures 2 and 4. Thus the shift in position between the EUV and SXR images for this dimming region may be due to features in the respective lines of sight in the two wave bands, or due to evolution of various dynamic flare-associated surrounding features over the time required for visibility in the respective wave bands. The video accompanying Figure 4 shows an SXT movie of the same region as the video accompanying Figure 2.

Figure 6 shows difference images in EUV from EIT (Figs. 6a–6c) and in SXR from SXT (Figs. 6d–6f). These are “fixed frame” difference images, in which the same EIT or SXT pre-erupt image was subtracted from the subsequent images. In addition, the EIT images are normalized to the image that was subtracted off; that is, these are “percentage difference images” (Wills-Davey & Thompson 1999).

Taken together, Figures 2–6 reveal coronal-evolution characteristics of this eruption; in particular they show the intensity

variations in the two wavelength regimes as the magnetic field evolves toward eruption, where we take the filament motions to be indicative of the field evolution. While the box 1 feature shows a nearly steady increase in intensity in EUV (Fig. 3b), it shows much more dramatic fluctuations in SXR (Fig. 5b). The video accompanying Figure 4 shows that this feature undergoes several episodes of microflaring, the last of which is occurring in Figure 4b. This microflaring is temporally related to brightening of the loop feature indicated by the arrows in Figure 4b; brightening of this loop is partially captured by the box 4 light curves in Figure 5b, between about 7:00 and 12:00 UT. After 12:00 UT there is a sharp intensity drop in box 4, coinciding with onset of the main flare in box 5, both of which are close in time to the filament’s fast-eruption onset. There are three particularly strong SXR microflares at the box 1 location, and these correlate well with the movement of the filament. Vertical bars in Figures 3a and 5a are at times of the microflare intensity peaks: the earliest peak is near 5:00 UT and corresponds to about the time of the start of the earliest monotonic slow rise of the filament; the second peak is near 8:30 UT and corresponds with a fluctuation in the slow-rise trajectory; and the third peak is near 11:45 UT, which is near the time of the end of the slow rise and the start of the rapid ejection of the filament.

### 3.3. Dimmings

EUV light curves 2 and 3 in Figure 3 and SXR light curves 2 and 4 in Figure 5 are from the pair of dimming regions closest to the core eruption region; light curve 2 is from the southern region and light curves 3 and 4 are from the northern region. We call these “twin dimmings,” and similar dimming pairs are often seen in eruptions (e.g., Sterling & Hudson 1997; Hudson et al. 1999; Zarro et al. 1999; Thompson et al. 1998, 2000; Kahler & Hudson 2001), although more complicated dimmings are also not uncommon (e.g., de Toma et al. 2005). Twin dimmings apparently result when the magnetic field containing a filament or filament channel and the magnetic fields bulging out from this channel leaves the Sun; either this field is a flux rope prior to eruption, or a flux rope forms in the process of erupting. As it lifts off the Sun, the flux rope leaves below it a relatively long and narrow channel of reduced density, and therefore reduced emission. Near the middle of this evacuated channel, however, flare loops rapidly develop, leaving the two end regions as the dimming pair. Figures 2c and 4c show these dimmings best.

Light curves from EIT (Fig. 3b) have adequate time cadence to see some details in the evolution of these dimmings. There is a gradual downturn in the intensity beginning at about 07:00 UT in the southern region and 11:00 UT for the northern region. Both locations show an increased rate of dimming from about 12:00 UT, with the maximum dimming occurring at about 13:00 UT. Overall, the southern dimming’s decrease roughly tracks the rise of the filament, and the increased rate of intensity decrease in both dimmings coincides closely with the onset of rapid filament rise. At later times there is an upturn in the intensity at both locations due to the intrusion of post-eruption-arcade emission into the dimmed regions. Prior to the onset of either of the twin dimmings, both regions show an increase in intensity, beginning between 4 and 6 UT in the southern region and between 6 and 9 UT in the northern region. In either case, this seems to be due to brightening of isolated looplike features; it is not certain whether these brightenings are directly related to the subsequent dimmings, which occur over a more extended region than the brightening features alone.

Light curves from SXT also show intensity decreases due to these twin dimmings from about 12:00 UT. Details of intensity






FIG. 6.—Fixed-frame intensity difference images of the erupting region, all from 2001 February 28. Panels *a–c* are percentage difference images from EIT, with the image used in forming the percentage difference from 06:00 UT. Panels *d–f* are SXT difference images, with the subtracted image from 06:36 UT. Arrows in (*e*) point out a ridge next to a coronal hole that brightens in SXT images and dims in both EIT (*c*) and SXT (*f*) images.

changes with time, however, are not as easy to determine as with the EUV light curves due to the less-regular cadence of SXT compared to EIT.

Remote dimmings and brightenings would be expected to precede the onset of fast filament eruption if the breakout mechanism is the trigger for the eruption (Moore & Sterling 2006). Light curves 6, 7, and 8 in both EUV (Fig. 3*b*) and SXR (Fig. 5*b*) all show dimming at locations that are remote, in the sense that they are farther removed from the erupting filament than are the twin dimmings; more specifically, they are associated with magnetic elements outside the bipolar region that erupts. (Light curve 9 is from a location of a set of loops that show changes consistent with temperature variation in a set of loops; see Sterling & Moore [2004*a*].) After a careful analysis of these dimmings using methods similar to those of Sterling & Moore (2004*a*), we find that we cannot say whether onset of the primary intensity changes (dimmings and brightenings) or onset of the fast rise begins first, and the data allow that they are concurrent. That is, we cannot determine whether the breakout reconnection was the trigger of, or a consequence of, the eruption. This is similar to our previous findings, as discussed in Moore & Sterling (2006; also see Williams et al. 2005).

#### 3.4. Magnetic Environment

Figure 7 shows the magnetic environment in the vicinity of the eruption. Overall, at this time the southern hemisphere of the Sun consisted of multiple polarities. We have identified in Figure 7*b* four alternating-polarity regions that form a quadruple, and from the coronal images from EIT and SXT we suspect that all four of these regions are involved in or affected by the eruption. The filament that erupts resides on a magnetic neutral line between regions R2 and R3. Figure 7 also shows that the box 1 region, where microflaring activity prior to eruption occurred, is a location of stronger mixed field, including at least three spotlike negative-field concentrations.

Figure 8 shows in more detail the field concentrations of and around box 1. Flux emerges over about 16 hr. As late as 20:48 UT on February 27, box 1 contained no black flux at all, and Figures 8*a–c* show that emergence occurred over the next several hours. Figure 8*d* shows the unsigned magnetic flux integrated over box 1 as a function of time; this curve is reproduced in Figures 3*a* and 5*a*, showing that the largest growth of the flux coincides with the start of the filament's slow rise. We have already pointed out that the SXR microflaring coincides closely with the filament's slow rise, and since the microflares emanate from the location of the growing flux, it is likely that the emerging flux is responsible for both the generation of the microflares and the instigation of the filament's slow rise (see § 4).

#### 3.5. CDS Spectra

Figure 9 shows the CDS FOV overlaid onto selected EIT and SXT images. As the northern portion of the filament crosses the CDS FOV, two significant events occur within the CDS FOV. The first is a microflaring of the feature at the location of box 1 in Figures 2 and 4, and the second is the eruption of the northern part of the filament. Arrows in Figures 9*a*, 9*b*, and 9*d* show the microflaring feature; it is in the CDS FOV only in Figure 9*a*, but Figure 9*d* shows that there is a bright SXR loop emanating from this microflaring region and extending to the southeast. This is the early flare arcade, also visible in Figure 4*c* and represented by the light curve for box 5 in Figure 5*b*. This feature becomes visible in SXT images between 12:10 and 12:55 UT, and therefore forms after the start of the fast eruption of the filament.

Figures 10*a* and 10*b* show CDS O v velocity rasters at times when the box 1 emerging-flux region is in the CDS FOV and when it is undergoing microflaring. From the raster, this microflaring results in motions along or near the magnetic field traced by the bright SXR loop emanating from this region in Figure 9*d*. Figure 9*b* also overlays onto the CDS raster of Figure 10*a* the magnetogram of Figure 8. This overlay shows that the microflaring




FIG. 7.—Magnetic setting in the vicinity of the eruption. (a) MDI magnetogram at the time indicated in the heading. (b) Same as (a), but with red and blue contours representing positive and negative flux regions, respectively. Labels R1–R4 denote alternating positive and negative regions that form a quadrupole involved in the eruption. (c) Magnetogram contours of (b) overlaid on an EIT image prior to eruption. (d) Magnetogram contours of (b) overlaid on an SXT image from after the eruption. Contour levels are 25, 40, and 300 G. Boxes in (b–d) are those of Figs. 2 and 4.






FIG. 8.—Close-up of the magnetic region around box 1. Panels *a–c* show MDI magnetograms at three different times, where the black rectangle is the same as box 1 in Figs. 2 and 4. Panel *d* shows the unsigned total magnetic flux in box 1; this curve is also plotted in Figs. 3*a* and 5*a*.

feature occurs at the location of the newly emerged flux concentration, and this match supports our conclusion that the emerging flux was responsible for the microflare.

Figures 10*c–10f* show where motions associated with the erupting filament are clearly apparent in CDS. At times just prior to these, from 12:01 to 12:23 UT, it is barely possible for us to pick out the filament in the velocity movies, indicating filament velocities  $< 20 \text{ km s}^{-1}$ , which is the approximate noise level of the velocity images. Figure 3*a* shows that there may be subtle motions (of magnitude similar to the error bars) in this northern portion of the filament projected on the solar disk from about 08:00 UT. From Figure 10, the filament has Earthward-directed velocities of up to a few tens of  $\text{km s}^{-1}$  from about 12:23 UT, reaching  $100 \text{ km s}^{-1}$  or more from around 12:44 UT. Since the lowest detectable CDS velocities are around  $20 \text{ km s}^{-1}$ , we cannot resolve possible slow-rise filament motions of this portion of the filament, which are likely to be much smaller than  $20 \text{ km s}^{-1}$  based on our estimates in § 3.1 for the velocities of the southern portion of the filament. Figures 10*d–10f* do tell us, however, that between 12:34 and about 13:00 UT the northern portion of the filament does have a line-of-sight velocity comparable to our estimated velocity of about  $45 \text{ km s}^{-1}$  for the southern portion of the filament at similar times.

#### 4. IMPLICATIONS FOR THE SLOW-RISE MECHANISM AND FOR THE FAST-RISE-ONSET MECHANISM

First we consider what may have driven the slow-rise of the filament. A strong hint comes from the activity around the box 1 location in Figures 2 and 4. Figure 8 shows that there is flux emergence at this location, and Figures 3*a* and 5*a* show that the emergence coincides approximately with the filament’s slow rise and the start of the SXR microflaring.

Figure 11 shows our interpretation of these events, in the immediate vicinity of the filament eruption. The emerging flux of box 1 interacts with the preexisting magnetic arcade that arches

over and ties down the filament field, and “tether-weakening” reconnection occurs between the emerging and the overlying flux systems. In contrast to “tether-cutting” reconnection, which reduces the filament field’s connection to the photosphere and builds a flux rope, tether-weakening reconnection, in the present setup, results in longer, higher arching loops in the overlying arcade (Moore & Roumeliotis 1992). This reconnection heats the plasma on the reconnected field lines, producing the fanlike SXR structure together with the SXR microflare brightenings. As the reconnection progresses the fan field expands, allowing the filament-carrying field to expand, and the filament rises slowly in step with the fan-field expansion.

As box 1 is spatially separated from the location where we measure the strongest filament motion (southern fiducial of Fig. 2*a*), we must consider how long it would take for a microflare disturbance generated by an emerging magnetic element of box 1 to be transmitted to our measurement point on the filament. From Figure 2*a* this distance is of order  $300''$ , or about  $2 \times 10^5 \text{ km}$ . We measured the average magnetic field strength of the region around the filament (spanning heliocentric longitude  $0''$  to  $200''$ , and latitude  $-400''$  to  $0''$ ; see Fig. 2*b*) to be 12 G. Taking  $\rho \sim 10^{-15} \text{ g cm}^{-3}$  to represent a quiet-region loop density (e.g., Priest 1982, Table 6.2) gives an Alfvén velocity of  $v_A = B/(4\pi\rho)^{1/2} \sim 1000 \text{ km s}^{-1}$ , with a corresponding Alfvén travel time of about 3.5 minutes between the emerging flux and the filament-measurement location. Given the roughness of our estimates and the poor cadence of the SXT images (resulting in poor cadence for the SXR microflare curve in Fig. 5), this travel time is consistent with the changes in filament motion occurring in response to the SXR microflares.

Eventually, the slow rise switches to fast rise as the filament erupts, producing the CME and strong flare heating. But as in our previous studies (Moore & Sterling 2006), we are not able to determine the exact mechanism for the onset of this transition from slow to fast rise. A possibility for our event is that continued tether-weakening reconnection could result in the overall system




FIG. 9.—FOV of CDS (*rectangular box*) overlaid onto (a–c) EIT images and (d) an SXT image; since CDS was pointing at the same fixed heliocentric location (i.e., CDS was not tracking solar rotation), the solar images rotate with respect to the CDS FOV over time. Arrows in (a), (b), and (d) indicate a location showing microflaring during the preeruption phase; this is at the location of box 1 in Figs. 2 and 4 and is in the CDS FOV in (a).




FIG. 10.—CDS O  $v$  velocity rasters (a) when a microflare is occurring (indicated by the arrow) at the location of box 1 in Figs. 2 and 4 and at the location indicated by the arrows in Figs. 9a, 9b, and 9d. Panel b is the same as panel a with the MDI magnetogram of 2001 February 28, 01:35 UT overlaid, with contours the same level as those in Fig. 7. Panels c–f show the filament when it is most visible in the velocity diagrams. Prior to the time of (a), it is barely detectable, with velocities near the noise level (about  $-20 \text{ km s}^{-1}$ ) in movies constructed from the CDS velocity rasters. Negative values indicate blueshifted motions. Values in (f) have higher uncertainties (and are likely stronger than they appear), since the signal is weak and therefore the profiles cannot be fit as well.

evolving to an unstable state, resulting in the fast eruption. In fact, the third and final SXR microflare could be due to the tether-weakening reconnection episode that finally allows the system to “fall off the table” and become explosively unstable. If instead, tether cutting is the mechanism triggering fast eruption, then we would expect a brightening increase at the location of box 5 at the time of the onset of the filament’s fast rise. From Figure 5, the time of flare brightening and fast filament rise onset are very close, but it is not possible to make a definitive conclusion of whether the brightening begins early enough to be consistent with a tether-cutting mechanism. Similarly, as discussed in § 3.3, from our data we also cannot determine whether breakout was the trigger of the fast-rise onset for this event.

## 5. SUMMARY AND DISCUSSION

We observed a large-scale ( $\approx 300,000 \text{ km}$ ) filament that erupted from a quiet solar region, a case in which the filament and asso-

ciated ejected fields produced a CME. By following the motions of the filament we can infer how the otherwise-invisible enveloping field structures evolve toward eruption. The earliest motion that we can detect of the filament is a nearly constant slow rise of about  $1 \text{ km s}^{-1}$  projected against the disk, which we estimate to be about  $5 \text{ km s}^{-1}$  in the vertical direction. After about 6 hr the filament undergoes acceleration and erupts from the Sun, leaving a postflare arcade in its wake. Weak-intensity dimmings began near the filament during the slow-rise phase, and these dimmings became pronounced twin dimmings at the time of eruption. Dimmings also occur at more remote locations, which could be indicative of breakout-type reconnection. New magnetic flux emerges adjacent to the filament over a 15 hr period that includes the slow-rise and eruption of the filament, with the fastest rate of emergence coinciding with the onset of the slow rise. Several SXR microflares occur at the location of emergence, and the three strongest ones peak respectively near the times of filament slow-rise onset, a kink in the slow-rise trajectory, and the time of transition from slow rise to fast eruption of the filament. Spectra from CDS of the first microflare show that it has rapid jetlike upflows. These spectra also show that the northern portion of the filament erupts with velocities similar to that of the southern portion of the filament, a fact that we could not obtain without the spectra since the northern part of the filament erupts nearly directly along the Earth-Sun line of sight.

The coronal hole in the vicinity of the filament (see Figs. 4 and 6) undergoes intensity changes during eruption, and it could have some implication in the large-scale evolution of the magnetic system involving the filament. In Figure 12 we present such a possibility, including a modified breakout-type scenario similar to that of Sterling & Moore (2001). Our interpretation of these events and our 2D cartoon sketches, however, are speculative and necessarily simplistic, and a full understanding of the observations would require nonlinear force-free 3D MHD modeling.

It has long been known that localized newly emerging flux frequently plays an important role in the onset of solar eruptions (e.g., Rust 1976; Moore et al. 1984; Parker 1987), and this has resulted in emerging-flux-based theoretical models since at least Heyvaerts et al. (1977). Wang et al. (2004) have observed rapid flux emergence near the time and place of flare onset, although their event was not associated with a filament eruption. SXR microflares have also been seen in association with emerging flux (Tang et al. 2000; Shimizu et al. 2002). Also, Feynman & Ruzmaikin (2004) found an association between emerging flux and the onset of the slow rise in a filament; our work supports their findings, and we give new evidence that the agent by which the emerging flux leads to the slow rise is tether-weakening reconnection.

Our strongest evidence for tether-weakening reconnection causing the slow rise of the filament field (which is part of a more general eruption field) is the bright SXR fan structure stemming from the SXR microflares, the connection of these with the emerging flux, and their coordination with the filament trajectory. Both the fan and the microflares, however, are prominent only in SXRs, not EUV. Previous filament slow-rise studies may have missed signatures for tether weakening because they generally either did not include SXRs, or used SXT SXR images of insufficient cadence. An active-region filament eruption studied by Sterling & Moore (2005) also showed a connection between localized EUV and SXR brightenings, recently emerged flux, and the slow-rise phase of a filament, although the data quality were not sufficient to conclude whether tether weakening was operating in that case. Similarly, Williams et al. (2005) present another




FIG. 11.—Schematic drawings of the erupting region in the vicinity of the filament during the slow-rise phase. (a) Three-dimensional view, with black dashed line representing the magnetic neutral line, black wavy line the filament, arched blue lines the magnetic field prior to a tether-weakening-reconnection episode, and adjacent red lines the same field slightly displaced by the reconnection episode; the red cross shows the location of the reconnection. These red field lines form a fanlike structure that arches over the filament, and is illuminated in SXR. (b) Cross-sectional cut of (a) at the position of the emerging flux. Blue and red lines are the same as in (a), and the filament is in the dip of a twisted field line; the tether-weakening reconnection lengthens field lines of the confining arcade, allowing the sheared core field to expand upward, resulting in the filament's slow rise.




FIG. 12.—Schematic drawings of the larger scale region, illustrating possible interactions between the erupting field and the coronal hole noted in Figs. 4 and 6. Panel *a* shows the setup near the time of the start of the filament's slow rise. The bipole associated with the emerging flux is adjacent to R3, and the blue field line above it is one of the loops arching over the filament in Fig. 11*a*. Black loops of still higher altitude are oriented so that reconnection with open-coronal-hole fields emanating from R1 is possible at the location of the red X. In (b), reconnection between these fields has started, with the reconnection-product fields dashed. The new loops formed between R1 and R2 lead to the remote SXT brightening of Fig. 6*e*, and to a reduction of the coronal hole. Meanwhile, the filament-arcade blue field continues to expand upward. In (c) the erupting filament-arcade blue field has “broken out” through the external reconnection site and the legs of the erupting blue field internally reconnect at the location of the red X, causing the main flare and unleashing the CME flux rope plasmoid. After the CME plasmoid pinches off and escapes, the most recently created R1–R2 loop will reconnect with the most recently created open field (both dashed). In (d) the dashed field lines of (c) have reconnected to produce new dashed reconnection-product fields, and a new set of fields are coming together and will reconnect. Eventually, the entire system relaxes via these reconnections until a state similar to (a) is reinstated, although in this final state the R2–R3 arcade is more relaxed and less inflated than that of (a) because the eruption has occurred. This could result in the dimming apparent in Fig. 6*f*.

observational example of possible tether weakening due to the intrusion of localized newly emerging flux into a magnetically stressed region; they did not see brightening by the inferred reconnection, however, probably because they observed in EUV without SXR. Schmieder et al. (2000) also present an example of an X-ray-bright point in spatial and temporal proximity to a destabilized filament, which they suggested could be related to emerging flux. These additional examples suggest that tether weakening causes the slow-rise phase of erupting filaments in at least a class of events. It still remains to be seen, however, whether tether weakening by localized emerging flux, leading to a CME explosion, is rare or common. In addition, more

observations are needed to determine what triggers the fast eruption.

A. C. S. and R. L. M. were supported by funding from NASA's Office of Space Science through the Solar Physics Supporting Research and Technology Program and the Sun-Earth Connection Guest Investigator Program. We thank the anonymous referee for many suggestions that improved the paper. *Yohkoh* is a mission of the Institute of Space and Astronautical Sciences (Japan), with participation from the US and UK, and *SOHO* is a project of international cooperation between ESA and NASA.

## REFERENCES

- Antiochos, S. K. 1998, *ApJ*, 502, L181  
 Antiochos, S. K., DeVore, C. R., & Klimchuk, J. A. 1999, *ApJ*, 510, 485  
 Anzer, U., & Heinzel, P. 2005, *ApJ*, 622, 714  
 Brueckner, G. E., et al. 1995, *Sol. Phys.*, 162, 357  
 Canfield, R. C., Hudson, H. S., & McKenzie, D. E. 1999, *Geophys. Res. Lett.*, 26, 627  
 Chen, P., & Shibata, K. 2000, *ApJ*, 545, 524  
 Delaboudinière, J.-P., et al. 1995, *Sol. Phys.*, 162, 291  
 de Toma, G., Holzer, T. E., Burckpile, J. T., & Gilbert, H. R. 2005, *ApJ*, 621, 1109  
 Fan, Y., & Gibson, S. E. 2004, *ApJ*, 609, 1123  
 Feynman, J., & Ruzmaikin, A. 2004, *Sol. Phys.*, 219, 301  
 Forbes, T. G. 2000, *J. Geophys. Res.*, 105, 23153  
 Forbes, T. G., & Isenberg, P. A. 1991, *ApJ*, 373, 294  
 Gibson, S. E., & Fan, Y. 2006, *J. Geophys. Res.*, 111, A12103  
 Harrison, R. A., et al. 1995, *Sol. Phys.*, 162, 233  
 Heyvaerts, J., Priest, E., & Rust, D. M. 1977, *Sol. Phys.*, 53, 255  
 Hudson, H. S., Acton, L. W., Harvey, K. L., & McKenzie, D. E. 1999, *ApJ*, 513, L83  
 Kahler, S. W., & Hudson, H. S. 2001, *J. Geophys. Res.*, 106, 29239  
 Kahler, S. W., Moore, R. L., Kane, S. R., & Zirin, H. 1988, *Sol. Phys.*, 328, 824  
 Klimchuk, J. A. 2001, in *Space Weather*, ed. P. Song, H. J. Singer, & G. L. Siscoe (Geophys. Monogr. 125; Washington: AGU), 143  
 Lin, J., Forbes, T. G., Isenberg, P. A., & Démoulin, P. 1998, *ApJ*, 504, 1006  
 Lin, J., Soon, W., & Baliunas, S. L. 2003, *NewA Rev.*, 47, 53  
 Linker, J., Lionello, R., Mikic, Z., & Amari, T. 2001, *J. Geophys. Res.*, 106, 25165  
 Mein, N., Schmieder, B., DeLuca, E. E., Heinzel, P., Mein, P., Malherbe, J.-M., & Staiger, J. 2001, *ApJ*, 556, 438  
 Moore, R. L., Hurford, G. J., Hones, H. P., & Kane, S. R. 1984, *ApJ*, 276, 379  
 Moore, R. L., & LaBonte, B. 1980, in *IAU Symp. 91, Solar and Interplanetary Dynamics*, ed. M. Dryer & E. Tandberg-Hanssen (Dordrecht: Reidel), 207  
 Moore, R. L., & Roumeliotis, G. 1992, in *IAU Colloq. 133, Eruptive Solar Flares*, ed. Z. Svestka, B. V. Jackson, & M. E. Machado (Berlin: Springer), 69  
 Moore, R. L., & Sterling, A. C. 2006, in *Solar Eruptions and Energetic Particles*, ed. N. Gopalswamy, R. A. Mewaldt, & J. Torsti (Geophys. Monogr. 165; Washington: AGU), 43  
 Moore, R. L., Sterling, A. C., Hudson, H. S., & Lemen, J. R. 2001, *ApJ*, 552, 833  
 Parker, E. N. 1987, *ApJ*, 312, 868  
 Priest, E. R. 1982, *Solar Magnetohydrodynamics* (Dordrecht: Reidel)  
 Rust, D. M. 1976, *Sol. Phys.*, 47, 21  
 Rust, D. M., & LaBonte, B. J. 2005, *ApJ*, 622, L69  
 Scherrer, P. H., et al. 1995, *Sol. Phys.*, 162, 129  
 Schmieder, B., Delannée, C., Yong, D. Y., Vial, J. C., & Madjarska, M. 2000, *A&A*, 358, 728  
 Schmieder, B., Lin, Y., Heinzel, P., & Schwartz, P. 2004, *Sol. Phys.*, 221, 297  
 Shimizu, T., Shine, R. A., Title, A. M., Tarbell, T. D., & Frank, Z. 2002, *ApJ*, 574, 1074  
 Sterling, A. C., & Hudson, H. S. 1997, *ApJ*, 491, L55  
 Sterling, A. C., & Moore, R. L. 2001, *J. Geophys. Res.*, 106, 25227  
 ———. 2003, *ApJ*, 599, 1418  
 ———. 2004a, *ApJ*, 602, 1024  
 ———. 2004b, *ApJ*, 613, 1221  
 ———. 2005, *ApJ*, 630, 1148  
 Sturrock, P. A. 1989, *Sol. Phys.*, 121, 387  
 Sturrock, P. A., Weber, M., Wheatland, M. S., & Wolfson, R. 2001, *ApJ*, 548, 492  
 Tandberg-Hanssen, E., Martin, S. F., & Hansen, R. T. 1980, *Sol. Phys.*, 65, 357  
 Tang, Y. H., Li, Y. N., Fang, C., Aulanier, G., Schmieder, B., Démoulin, P., & Sakurai, T. 2000, *ApJ*, 534, 482  
 Thompson, B. J., Cliver, E. W., Nitta, N., Delannée, C., & Delaboudinière, J. P. 2000, *Geophys. Res. Lett.*, 27, 1431  
 Thompson, B. J., Plunkett, S. P., Gurman, J. B., Newmark, J. S., St. Cyr, O. C., & Michels, D. J. 1998, *Geophys. Res. Lett.*, 25, 2465  
 Tsuneta, S., et al. 1991, *Sol. Phys.*, 136, 37  
 Wang, H., et al. 2004, *ApJ*, 605, 931  
 Wiik, J. E., Schmieder, B., Kucera, T., Poland, A., Brekke, P., & Simnett, G. 1997, *Sol. Phys.*, 175, 411  
 Williams, D. R., Török, T., Démoulin, P., van Driel-Gesztelyi, L., & Kliem, B. 2005, *ApJ*, 628, L163  
 Wills-Davey, M. J., & Thompson, B. J. 1999, *Sol. Phys.*, 190, 467  
 Yashiro, S., Gopalswamy, N., Michalek, G., St. Cyr, O. C., Plunkett, S. P., Rich, N. B., & Howard, R. A. 2004, *J. Geophys. Res.*, 109, A07105  
 Zarro, D. M., Sterling, A. C., Thompson, B. J., Hudson, H. S., & Nitta, N. 1999, *ApJ*, 520, L139