

## Furan

### CAS No. 110-00-9

Reasonably anticipated to be a human carcinogen

First listed in the *Eighth Report on Carcinogens* (1998)



### Carcinogenicity

Furan is *reasonably anticipated to be a human carcinogen* based on sufficient evidence of carcinogenicity from studies in experimental animals.

#### Cancer Studies in Experimental Animals

Oral exposure to furan caused tumors at several different tissue sites in mice and rats. Administration of furan by stomach tube for up to 2 years caused benign and/or malignant liver tumors (hepatocellular adenoma or carcinoma) in mice and rats of both sexes. It also caused bile-duct cancer (cholangiocarcinoma) and mononuclear-cell leukemia in rats of both sexes and benign adrenal-gland tumors (pheochromocytoma) in mice of both sexes (NTP 1993). Similar administration of furan to male rats for 9 to 13 weeks caused bile-duct cancer (cholangiocarcinoma) by 16 months after the end of exposure (Maronpot *et al.* 1991, Elmore and Sirica 1993). Since furan was listed in the *Eighth Report on Carcinogens*, an additional study in mice has been identified. Intraperitoneal injection of furan caused benign or malignant liver tumors (hepatocellular adenoma or carcinoma) in newborn male mice (Johansson *et al.* 1997).

#### Cancer Studies in Humans

No epidemiological studies were identified that evaluated the relationship between human cancer and exposure specifically to furan.

#### Studies on Mechanisms of Carcinogenesis

In bacteria, furan caused gene mutations in *Salmonella typhimurium* strain TA100 (Lee *et al.* 1994) and in *Escherichia coli* containing bacteriophage T7 (Ronto *et al.* 1992), but not in *S. typhimurium* strains TA98 (Lee *et al.* 1994), TA1535, or TA1537 (Mortelmans *et al.* 1986). It did not cause gene mutations in *Drosophila melanogaster* (Fourman *et al.* 1994). In mammalian *in vitro* systems, it caused gene mutations in mouse lymphoma cells (McGregor *et al.* 1988), DNA damage in Chinese hamster ovary (CHO) cells (NTP 1993), and chromosomal damage in CHO cells with mammalian metabolic activation (NTP 1993, IARC 1995), but it did not cause DNA damage in mouse or rat hepatocytes (Wilson *et al.* 1992, NTP 1993). In mammalian *in vivo* systems, furan caused chromosomal aberrations in bone marrow of mice (NTP 1993, Johansson 1997), but did not cause DNA damage in mouse bone marrow or hepatocytes or rat hepatocytes (Wilson *et al.* 1992, NTP 1993).

A current hypothesis for the mechanism of furan-induced carcinogenesis is metabolic activation of furan by cytochrome P450 to a reactive and cytotoxic intermediate that stimulates cell replication, increasing the likelihood of tumor induction (Kedderis *et al.* 1993, Chen *et al.* 1995). The postulated reactive metabolite is *cis*-2-butene-1,4-dial, which was characterized as a furan metabolite by Chen *et al.* (1995). This reactive metabolite probably explains furan's binding reactivity with proteins both *in vitro* (in uninduced and induced male rat liver microsomes) and *in vivo* (with male rat liver protein) (Burka *et al.* 1991, Parmar and Burka 1993). Furan metabolites may

react with DNA, but no radiotracer was detected in DNA from livers of rats administered [<sup>14</sup>C]furan (Burka *et al.* 1991).

### Properties

Furan is a cyclic dienic ether that is a clear, colorless liquid with an ethereal odor (Akron 2009, HSDB 2009). It can turn brown upon standing (HSDB 2009). Furan is slightly soluble in water and is soluble at greater than 10% in acetone, benzene, ether, and ethanol. It is extremely flammable and may form explosive peroxides in the absence of inhibitors (Akron 2009). Physical and chemical properties of furan are listed in the following table.

Property	Information
Molecular weight	68.1 <sup>a</sup>
Specific gravity	0.9371 at 19.4°C/4°C <sup>a</sup>
Melting point	-85.6°C <sup>a</sup>
Boiling point	31.4°C at 760 mm Hg <sup>a</sup>
Log <i>K</i> <sub>ow</sub>	1.34 <sup>a</sup>
Water solubility	10 g/L at 25°C <sup>b</sup>
Vapor pressure	600 mm Hg at 25°C <sup>b</sup>
Vapor density relative to air	2.3 <sup>a</sup>

Sources: <sup>a</sup>HSDB 2009, <sup>b</sup>ChemIDplus 2009.

### Use

Furan is used primarily as an intermediate in the synthesis and production of tetrahydrofuran, pyrrole, and thiophene. Hydrogenation of furan over a nickel catalyst produces high yields of tetrahydrofuran and is a source of commercial tetrahydrofuran (NTP 1993, IARC 1995). Furan is also used in the formation of lacquers, as a solvent for resins, and in the production of agricultural chemicals, stabilizers, and pharmaceuticals (IARC 1995, HSDB 2009).

### Production

Commercial production of furan involves decarbonylation of furfural over a palladium-charcoal catalyst. The commercial product is at least 99% pure (IARC 1995). In 2009, furan was produced by one manufacturer worldwide, in the United States (SRI 2009), and was available from 20 suppliers, including 11 U.S. suppliers (ChemSources 2009). U.S. imports of furan resins totaled about 9.7 million pounds in 1986 (HSDB 2009). Reports filed from 1986 to 1998 under the U.S. Environmental Protection Agency's Toxic Substances Control Act Inventory Update Rule indicated that U.S. production plus imports of furan totaled 10 million to 50 million pounds (EPA 2004); no reports were filed in 2002.

### Exposure

The routes of potential human exposure to furan are inhalation, ingestion, and dermal contact. The pattern of commercial use suggests that minimal exposure to the general population would be expected through contact with products contaminated with furan (NTP 1993). However, furan can be formed in foods during processing. Furan has been detected in the breath of both smokers and nonsmokers and in indoor air, foods, and human milk samples (IARC 1995, NTP 1999, FDA 2005). Furan also occurs naturally in pine rosin and in volatile emissions from sorb trees (HSDB 2009).

Furan was measured by the U.S. Food and Drug Administration in various foods and beverages, including infant formulas, baby foods, soups and sauces, fruits and vegetables, bread, and meat products. The maximum concentration found was 125 ppb (µg/kg) in canned soup (FDA 2005). A second study confirmed that heat-treated foods, such as canned and jarred foods, contained measurable quantities of furan (up to 240 µg/kg in canned chili) (Becalski *et al.* 2005). Furan was measured in fruit juice at concentrations near 1 µg/kg (Goldmann

## Report on Carcinogens, Twelfth Edition (2011)

*et al.* 2005). In several brands of brewed coffee, the highest furan concentration found was 84.2 ppb (FDA 2005, Ho *et al.* 2005). Furan was also identified as a component of coffee aroma that has antioxidant activity (Fuster *et al.* 2000). Furan was also detected at a concentration of 110 µg/kg in jarred baby food containing cooked vegetables (Goldmann *et al.* 2005). However, furan concentrations decreased after the jar was opened and the contents were heated. When food is heated in a container, furan concentrations increase if the container remains closed, but not if it is open (Hasnip *et al.* 2006). Furan does not appear to be transferred from the packaging or gasket of the can or jar. Furan is formed from ascorbic acid, fructose, sucrose, and glucose when foods are heated or irradiated (Fan 2005). Furan production increases greatly with decreasing pH of the medium; 1,600 times as much furan is formed at pH 3 as is formed at pH 8. Furan was detected in 1 of 11 breast-milk samples from women in four different urban areas (HSDB 2009).

In one study in Texas, furan was detected in the exhaled breath of two of three male smokers and four of five male nonsmokers (HSDB 2009). Smokers exhaled between 0.25 and 98 µg of furan per hour, and nonsmokers exhaled between 0.33 and 28 µg/h. In a study in Chicago, 15 of 387 breath samples collected from 54 male and female nonsmokers had detectable levels of furan, with a mean concentration of 0.55 ng/L. Furan was also detected in the indoor air of homes in the Chicago, Illinois, and Washington, D.C., metropolitan areas (NTP 1999).

If furan is released to air, it will exist almost entirely in the vapor phase (Howard 1989). In daylight, it will react with hydroxyl radicals, with a half-life of 9.5 hours. Furan is resistant to hydrolysis. Its estimated half-life in a shallow model river is 2.5 hours. If released to surface water, it will volatilize rapidly and will not adsorb to sediment or suspended solids or bioaccumulate in aquatic organisms. If released to soil, it will volatilize or leach rapidly. Furan has been detected in industrial effluents, ambient air, wood smoke, and automobile exhaust and in surface water. However, the frequency of detection and concentration generally were low. For example, furan was detected in 1 of 63 industrial effluents at concentrations of less than 10 µg/L and in aqueous condensate samples from low-temperature gasification of rosebud coal at 7 µg/L (IARC 1995, HSDB 2009).

The primary route of occupational exposure to furan is inhalation. The industrial processes in which furan is used are conducted in closed systems, and its volatility requires that furan be handled in closed containers; therefore, occupational exposure is limited (NTP 1993). The National Occupational Hazard Survey (conducted from 1972 to 1974) estimated that 244 workers potentially were exposed to furan (NIOSH 1976). The National Occupational Exposure Survey (conducted from 1981 to 1983) estimated that 35 workers (mostly in the Business Services industry), including 7 women, potentially were exposed to furan (NIOSH 1990).

## Regulations

### Department of Transportation (DOT)

Furan is considered a hazardous material, and special requirements have been set for marking, labeling, and transporting this material.

### Environmental Protection Agency (EPA)

#### Clean Air Act

*Prevention of Accidental Release:* Threshold quantity (TQ) = 5,000 lb.

*Comprehensive Environmental Response, Compensation, and Liability Act*  
Reportable quantity (RQ) = 100 lb.

#### Emergency Planning and Community Right-To-Know Act

Reportable quantity (RQ) = 100 lb.

Threshold planning quantity (TPQ) = 500 lb.

### Resource Conservation and Recovery Act

*Listed Hazardous Waste:* Waste code for which the listing is based wholly or partly on the presence of furan = U124.

### Occupational Safety and Health Administration (OSHA)

Considered a highly hazardous chemical; threshold quantity (TQ) = 500 lb.

## References

- Akron. 2009. *The Chemical Database*. The Department of Chemistry at the University of Akron. <http://ull.chemistry.uakron.edu/erd> and search on CAS number. Last accessed: 5/09.
- Becalski A, Forsyth D, Casey V, Lau BP-Y, Pepper K, Seaman S. 2005. Development and validation of a headspace method for determination of furan in food. *Food Addit Contam* 22(6): 535-540.
- Burka LT, Washburn KD, Irwin RD. 1991. Disposition of [<sup>14</sup>C]furan in the male F344 rat. *J Toxicol Environ Health* 34(2): 245-257.
- ChemIDplus. 2009. *ChemIDplus Advanced*. National Library of Medicine. <http://chem.sis.nlm.nih.gov/chemidplus/chemidheavy.jsp> and select Registry Number and search on CAS number. Last accessed: 5/09.
- ChemSources. 2009. *Chem Sources - Chemical Search*. Chemical Sources International. <http://www.chemsources.com/chemonline.html> and search on furan. Last accessed: 5/09.
- Chen LJ, Hecht SS, Peterson LA. 1995. Identification of *cis*-2-butene-1,4-dial as a microsomal metabolite of furan. *Chem Res Toxicol* 8(7): 903-906.
- Elmore LR, Sirica AE. 1993. "Intestinal-type" of adenocarcinoma preferentially induced in right/caudate liver lobes of rats treated with furan. *Cancer Res* 53(2): 254-259.
- EPA. 2004. *Non-confidential IUR Production Volume Information*. U.S. Environmental Protection Agency. <http://www.epa.gov/oppt/iur/tools/data/2002-vol.html> and search on CAS number.
- Fan X. 2005. Formation of furan from carbohydrates and ascorbic acid following exposure to ionizing radiation and thermal processing. *J Agric Food Chem* 53(20): 7826-7831.
- FDA. 2005. *Exploratory Data on Furan in Food: Individual Food Products*. Last updated: 6/15/05. <http://www.fda.gov/Food/FoodSafety/FoodContaminantsAdulteration/ChemicalContaminants/Furan/UCM078439>.
- Fourman P, Mason JM, Valencia R, Zimmering S. 1994. Chemical mutagenesis testing in *Drosophila*. IX. Results of 50 coded compounds tested for the National Toxicology Program. *Environ Mol Mutagen* 23(1): 51-63.
- Fuster MD, Mitchell AE, Ochi H, Shibamoto T. 2000. Antioxidative activities of heterocyclic compounds formed in brewed coffee. *J Agric Food Chem* 48(11): 5600-5603.
- Goldmann T, Perisset A, Scanlan F, Stadler RH. 2005. Rapid determination of furan in heated foodstuffs by isotope dilution solid phase micro-extraction-gas chromatography-mass spectrometry (SPME-GC-MS). *Analyst* 130(6): 878-883.
- Hasnip S, Crews C, Castle L. 2006. Some factors affecting the formation of furan in heated foods. *Food Addit Contam* 23(3): 219-227.
- Ho IP, Yoo SJ, Tefera S. 2005. Determination of furan levels in coffee using automated solid-phase microextraction and gas chromatography/mass spectrometry. *J AOAC Int* 88(2): 574-576.
- Howard PH. 1989. Furan. In *Handbook of Environmental Fate and Exposure Data for Organic Chemicals*. vol. 1. Chelsea, MI: Lewis Publishers. pp. 342-350.
- HSDB. 2009. *Hazardous Substances Data Bank*. National Library of Medicine. <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB> and search on CAS number. Last accessed: 3/22/09.
- IARC. 1995. Furan. In *Dry Cleaning, Some Chlorinated Solvents and Other Industrial Chemicals*. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 63. Lyon, France: International Agency for Research on Cancer. pp. 393-407.
- Johansson E, Reynolds S, Anderson M, Maronpot R. 1997. Frequency of Ha-ras-1 gene mutations inversely correlated with furan dose in mouse liver tumors. *Mol Carcinog* 18(4): 199-205.
- Kedderis GL, Carfagna MA, Held SD, Batra R, Murphy JE, Gargas ML. 1993. Kinetic analysis of furan biotransformation by F-344 rats *in vivo* and *in vitro*. *Toxicol Appl Pharmacol* 123(2): 274-282.
- Lee H, Bian SS, Chen YL. 1994. Genotoxicity of 1,3-dithiane and 1,4-dithiane in the CHO/SCE assay and the Salmonella/microsomal test. *Mutat Res* 321(4): 213-218.
- Maronpot RR, Giles HD, Dykes DJ, Irwin RD. 1991. Furan-induced hepatic cholangiocarcinomas in Fischer 344 rats. *Toxicol Pathol* 19(4 Pt 2): 561-570.
- McGregor DB, Brown A, Cattanach P, Edwards I, McBride D, Riach C, Caspary WJ. 1988. Responses of the L5178Y tk+/tk- mouse lymphoma cell forward mutation assay: III. 72 coded chemicals. *Environ Mol Mutagen* 12(1): 85-154.
- Mortelmans K, Haworth S, Lawlor T, Speck W, Tainer B, Zeiger E. 1986. Salmonella mutagenicity tests: II. Results from the testing of 270 chemicals. *Environ Mutagen* 8(Suppl 7): 1-119.
- NIOSH. 1976. *National Occupational Hazard Survey (1972-74)*. DHEW (NIOSH) Publication No. 78-114. Cincinnati, OH: National Institute for Occupational Safety and Health.
- NIOSH. 1990. *National Occupational Exposure Survey (1981-83)*. National Institute for Occupational Safety and Health. Last updated: 7/1/90. <http://www.cdc.gov/noes/noes1/34185sic.html>.
- NTP. 1993. *Toxicology and Carcinogenesis Studies of Furan (CAS No. 110-00-9) in F344 Rats and B6C3F<sub>1</sub> Mice (Gavage Studies)*. NTP Technical Report Series no. 402. Research Triangle Park, NC: National Toxicology Program. 286 pp.
- NTP. 1999. *NTP Report on Carcinogens Background Document for Furan*. Research Triangle Park: Integrated Laboratory Systems. 60 pp.

## Report on Carcinogens, Twelfth Edition (2011)

Parmar D, Burka LT. 1993. Studies on the interaction of furan with hepatic cytochrome P-450. *J Biochem Toxicol* 8(1): 1-9.

Ronto G, Grof P, Buisson JP, Einhorn J, Demerseman P. 1992. Genotoxic effectivity—comparison of 36 nitrated furan and arenofuran derivatives on a quantitative scale. Statistical comparison of T7 and other short-term tests. *Mutagenesis* 7(4): 243-249.

SRI. 2009. *Directory of Chemical Producers*. Menlo Park, CA: SRI Consulting. Database edition. Last accessed: 5/09.

Wilson DM, Goldsworthy TL, Popp JA, Butterworth BE. 1992. Evaluation of genotoxicity, pathological lesions, and cell proliferation in livers of rats and mice treated with furan. *Environ Mol Mutagen* 19(3): 209-222.