Coronary Artery Bypass Graft (CABG) Rate

Inpatient Quality Indicators #26 Technical Specifications

Area-Level Indicator
Procedure Utilization Indicator
AHRQ Quality Indicators, Version 4.3, August 2011

Numerator

Number of discharges with a CABG procedure age 40 years and older.

ICD-9-CM CABG procedure codes:

o o, 120 p. occuu. o couco.		
AORTOCORONARY BYPASS NOS	3615	1 INT MAM-COR ART BYPASS
AORTOCOR BYPAS-1 COR ART	3616	2 INT MAM-COR ART BYPASS
AORTOCOR BYPAS-2 COR ART	3617	ABD-CORON ART BYPASS (OCT96)
AORTOCOR BYPAS-3 COR ART	3619	HRT REVAS BYPS ANAS NEC
AORTCOR BYPAS-4+ COR ART		
	AORTOCORONARY BYPASS NOS AORTOCOR BYPAS-1 COR ART AORTOCOR BYPAS-2 COR ART AORTOCOR BYPAS-3 COR ART	AORTOCORONARY BYPASS NOS 3615 AORTOCOR BYPAS-1 COR ART 3616 AORTOCOR BYPAS-2 COR ART 3617 AORTOCOR BYPAS-3 COR ART 3619

Exclude cases:

• MDC 14 (pregnancy, childbirth, and puerperium)

 with missing discharge disposition (DISP=missing), gender (SEX=missing), age (AGE=missing), quarter (DQTR=missing), year (YEAR=missing) or principal diagnosis (DX1=missing)

Denominator

Discharges in the numerator are assigned to the denominator based on the Metro Area¹ or county of the patient residence, not the Metro Area or county of the hospital where the discharge occurred.

¹ The term "metropolitan area" (MA) was adopted by the U.S. Census in 1990 and referred collectively to metropolitan statistical areas (MSAs), consolidated metropolitan statistical areas (CMSAs), and primary metropolitan statistical areas (PMSAs). In addition, "area" could refer to either 1) FIPS county, 2) modified FIPS county, 3) 1999 OMB Metropolitan Statistical Area, or 4) 2003 OMB Metropolitan Statistical Area. Micropolitan Statistical Areas are not used in the QI software.