


Table II-13
ALL CANCER SITES (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	189.8	0.11		
High Five States				
Kentucky	222.7 ^b	1.03	(01)	17.3 ^c
Louisiana	221.0 ^b	1.02	(02)	16.4 ^c
West Virginia	213.4 ^b	1.40	(03)	12.4 ^c
Mississippi	211.9 ^b	1.23	(04)	11.6 ^c
Tennessee	211.2 ^b	0.84	(05)	11.3 ^c
Low Five States				
Arizona	167.5 ^b	0.77	(47)	-11.7 ^c
New Mexico	167.3 ^b	1.36	(48)	-11.9 ^c
Colorado	166.4 ^b	0.95	(49)	-12.3 ^c
Hawaii	152.2 ^b	1.50	(50)	-19.8 ^c
Utah	140.9 ^b	1.29	(51)	-25.8 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	206.2 ^b	0.93	(09)	8.6 ^c	Montana	186.6	1.92	(32)	-1.7
Alaska	186.5	3.36	(33)	-1.7	Nebraska	179.6 ^b	1.39	(42)	-5.4 ^c
Arizona	167.5 ^b	0.77	(47)	-11.7 ^c	Nevada	197.1 ^b	1.41	(18)	3.8
Arkansas	207.3 ^b	1.18	(07)	9.2 ^c	New Hampshire	193.0	1.73	(27)	1.7
California	173.1 ^b	0.33	(46)	-8.8 ^c	New Jersey	193.4 ^b	0.65	(25)	1.9
Colorado	166.4 ^b	0.95	(49)	-12.3 ^c	New Mexico	167.3 ^b	1.36	(48)	-11.9 ^c
Connecticut	184.3 ^b	0.98	(36)	-2.9	New York	179.9 ^b	0.42	(41)	-5.2 ^c
Delaware	204.3 ^b	2.19	(11)	7.6 ^c	North Carolina	196.1 ^b	0.69	(20)	3.3
D.C.	209.5 ^b	2.70	(06)	10.4 ^c	North Dakota	176.3 ^b	2.20	(44)	-7.1 ^c
Florida	180.1 ^b	0.41	(39)	-5.1 ^c	Ohio	204.1 ^b	0.58	(12)	7.5 ^c
Georgia	194.2 ^b	0.74	(24)	2.3	Oklahoma	200.1 ^b	1.04	(15)	5.4 ^c
Hawaii	152.2 ^b	1.50	(50)	-19.8 ^c	Oregon	192.0	1.02	(28)	1.2
Idaho	173.4 ^b	1.65	(45)	-8.6 ^c	Pennsylvania	198.5 ^b	0.52	(16)	4.6
Illinois	197.6 ^b	0.57	(17)	4.1	Rhode Island	194.7	1.80	(22)	2.6
Indiana	205.1 ^b	0.81	(10)	8.1 ^c	South Carolina	200.7 ^b	0.98	(14)	5.7 ^c
Iowa	185.9 ^b	1.05	(34)	-2.1	South Dakota	182.8	2.07	(38)	-3.7
Kansas	187.0	1.15	(30)	-1.5	Tennessee	211.2 ^b	0.84	(05)	11.3 ^c
Kentucky	222.7 ^b	1.03	(01)	17.3 ^c	Texas	184.9 ^b	0.45	(35)	-2.6
Louisiana	221.0 ^b	1.02	(02)	16.4 ^c	Utah	140.9 ^b	1.29	(51)	-25.8 ^c
Maine	206.8 ^b	1.65	(08)	9.0 ^c	Vermont	183.6	2.36	(37)	-3.3
Maryland	196.6 ^b	0.87	(19)	3.6	Virginia	194.7 ^b	0.75	(23)	2.6
Massachusetts	193.4 ^b	0.75	(26)	1.9	Washington	187.1	0.80	(29)	-1.4
Michigan	195.3 ^b	0.62	(21)	2.9	West Virginia	213.4 ^b	1.40	(03)	12.4 ^c
Minnesota	178.8 ^b	0.85	(43)	-5.8 ^c	Wisconsin	186.7 ^b	0.81	(31)	-1.6
Mississippi	211.9 ^b	1.23	(04)	11.6 ^c	Wyoming	180.0 ^b	2.70	(40)	-5.2 ^c
Missouri	201.9 ^b	0.81	(13)	6.4 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table II-14
ALL CANCER SITES (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	234.4	0.20		
High Five States				
Kentucky	286.9 ^b	1.86	(01)	22.4 ^c
Louisiana	285.9 ^b	1.84	(02)	22.0 ^c
Mississippi	282.6 ^b	2.25	(03)	20.6 ^c
D.C.	275.5 ^b	5.01	(04)	17.5 ^c
Tennessee	272.4 ^b	1.53	(05)	16.2 ^c
Low Five States				
New Mexico	203.4 ^b	2.29	(47)	-13.2 ^c
Arizona	201.3 ^b	1.27	(48)	-14.1 ^c
Colorado	199.3 ^b	1.63	(49)	-15.0 ^c
Hawaii	189.4 ^b	2.50	(50)	-19.2 ^c
Utah	172.5 ^b	2.18	(51)	-26.4 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	271.7 ^b	1.71	(06)	15.9 ^c	Montana	222.0 ^b	3.18	(38)	-5.3 ^c
Alaska	226.7	5.80	(34)	-3.3	Nebraska	220.6 ^b	2.38	(39)	-5.9 ^c
Arizona	201.3 ^b	1.27	(48)	-14.1 ^c	Nevada	229.4	2.30	(31)	-2.1
Arkansas	266.8 ^b	2.07	(07)	13.8 ^c	New Hampshire	237.6	3.02	(25)	1.4
California	206.4 ^b	0.56	(46)	-11.9 ^c	New Jersey	232.4	1.13	(26)	-0.9
Colorado	199.3 ^b	1.63	(49)	-15.0 ^c	New Mexico	203.4 ^b	2.29	(47)	-13.2 ^c
Connecticut	225.1 ^b	1.70	(36)	-4.0	New York	217.1 ^b	0.73	(42)	-7.4 ^c
Delaware	252.8 ^b	3.80	(13)	7.8 ^c	North Carolina	251.8 ^b	1.25	(16)	7.4 ^c
D.C.	275.5 ^b	5.01	(04)	17.5 ^c	North Dakota	216.3 ^b	3.69	(43)	-7.7 ^c
Florida	219.9 ^b	0.68	(40)	-6.2 ^c	Ohio	254.9 ^b	1.02	(12)	8.7 ^c
Georgia	250.9 ^b	1.37	(17)	7.0 ^c	Oklahoma	252.2 ^b	1.82	(14)	7.6 ^c
Hawaii	189.4 ^b	2.50	(50)	-19.2 ^c	Oregon	228.2 ^b	1.69	(32)	-2.6
Idaho	208.4 ^b	2.74	(45)	-11.1 ^c	Pennsylvania	246.4 ^b	0.90	(18)	5.1 ^c
Illinois	245.6 ^b	1.00	(19)	4.8	Rhode Island	243.3	3.17	(21)	3.8
Indiana	257.7 ^b	1.44	(10)	9.9 ^c	South Carolina	263.7 ^b	1.80	(09)	12.5 ^c
Iowa	231.3	1.80	(29)	-1.3	South Dakota	228.2	3.50	(33)	-2.6
Kansas	231.9	1.97	(28)	-1.1	Tennessee	272.4 ^b	1.53	(05)	16.2 ^c
Kentucky	286.9 ^b	1.86	(01)	22.4 ^c	Texas	232.3	0.80	(27)	-0.9
Louisiana	285.9 ^b	1.84	(02)	22.0 ^c	Utah	172.5 ^b	2.18	(51)	-26.4 ^c
Maine	256.0 ^b	2.87	(11)	9.2 ^c	Vermont	226.2	4.11	(35)	-3.5
Maryland	241.4 ^b	1.53	(22)	3.0	Virginia	244.8 ^b	1.35	(20)	4.4
Massachusetts	237.8	1.31	(24)	1.5	Washington	222.8 ^b	1.35	(37)	-4.9
Michigan	240.2 ^b	1.08	(23)	2.5	West Virginia	265.6 ^b	2.45	(08)	13.3 ^c
Minnesota	219.2 ^b	1.45	(41)	-6.5 ^c	Wisconsin	230.8	1.39	(30)	-1.5
Mississippi	282.6 ^b	2.25	(03)	20.6 ^c	Wyoming	211.2 ^b	4.47	(44)	-9.9 ^c
Missouri	252.1 ^b	1.42	(15)	7.6 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table II-15
ALL CANCER SITES (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	159.9	0.14		
High Five States				
Kentucky	180.5 ^b	1.23	(01)	12.9 ^c
West Virginia	178.2 ^b	1.71	(02)	11.4 ^c
Louisiana	178.0 ^b	1.21	(03)	11.3 ^c
Maine	173.8 ^b	2.02	(04)	8.7 ^c
Nevada	171.6 ^b	1.78	(05)	7.3 ^c
Low Five States				
Colorado	145.4 ^b	1.17	(47)	-9.1 ^c
Arizona	142.0 ^b	0.96	(48)	-11.2 ^c
New Mexico	140.9 ^b	1.67	(49)	-11.9 ^c
Hawaii	123.0 ^b	1.83	(50)	-23.1 ^c
Utah	117.9 ^b	1.58	(51)	-26.3 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	163.6	1.10	(24)	2.3	Montana	162.0	2.43	(27)	1.3
Alaska	155.9	4.10	(38)	-2.5	Nebraska	152.4 ^b	1.73	(41)	-4.7
Arizona	142.0 ^b	0.96	(48)	-11.2 ^c	Nevada	171.6 ^b	1.78	(05)	7.3 ^c
Arkansas	166.2 ^b	1.41	(17)	3.9	New Hampshire	164.2	2.13	(23)	2.7
California	150.5 ^b	0.41	(43)	-5.9 ^c	New Jersey	169.2 ^b	0.80	(11)	5.8 ^c
Colorado	145.4 ^b	1.17	(47)	-9.1 ^c	New Mexico	140.9 ^b	1.67	(49)	-11.9 ^c
Connecticut	158.6	1.21	(30)	-0.8	New York	156.8 ^b	0.52	(35)	-1.9
Delaware	171.0 ^b	2.66	(08)	6.9 ^c	North Carolina	160.1	0.82	(28)	0.1
D.C.	169.4	3.17	(10)	5.9 ^c	North Dakota	149.6 ^b	2.77	(45)	-6.4 ^c
Florida	149.9 ^b	0.51	(44)	-6.3 ^c	Ohio	171.2 ^b	0.70	(06)	7.1 ^c
Georgia	158.3	0.87	(31)	-1.0	Oklahoma	164.5 ^b	1.26	(21)	2.9
Hawaii	123.0 ^b	1.83	(50)	-23.1 ^c	Oregon	166.9 ^b	1.27	(16)	4.4
Idaho	147.6 ^b	2.06	(46)	-7.7 ^c	Pennsylvania	168.0 ^b	0.63	(13)	5.1 ^c
Illinois	167.3 ^b	0.69	(15)	4.6	Rhode Island	164.4	2.20	(22)	2.8
Indiana	171.1 ^b	0.98	(07)	7.0 ^c	South Carolina	159.0	1.15	(29)	-0.6
Iowa	156.1	1.29	(37)	-2.4	South Dakota	150.8 ^b	2.56	(42)	-5.7 ^c
Kansas	157.3	1.41	(34)	-1.6	Tennessee	170.9 ^b	1.00	(09)	6.9 ^c
Kentucky	180.5 ^b	1.23	(01)	12.9 ^c	Texas	152.6 ^b	0.54	(40)	-4.6
Louisiana	178.0 ^b	1.21	(03)	11.3 ^c	Utah	117.9 ^b	1.58	(51)	-26.3 ^c
Maine	173.8 ^b	2.02	(04)	8.7 ^c	Vermont	156.2	2.91	(36)	-2.3
Maryland	167.6 ^b	1.06	(14)	4.8	Virginia	162.5	0.90	(26)	1.6
Massachusetts	166.2 ^b	0.92	(18)	3.9	Washington	162.7	1.01	(25)	1.8
Michigan	165.7 ^b	0.76	(20)	3.6	West Virginia	178.2 ^b	1.71	(02)	11.4 ^c
Minnesota	153.0 ^b	1.05	(39)	-4.3	Wisconsin	157.6	0.99	(32)	-1.4
Mississippi	165.8 ^b	1.43	(19)	3.7	Wyoming	157.4	3.42	(33)	-1.6
Missouri	168.6 ^b	0.99	(12)	5.4 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table III-13
BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.4	0.02		
High Five States				
Washington	5.6 ^b	0.14	(01)	27.3 ^c
Tennessee	5.6 ^b	0.14	(02)	27.3 ^c
Oregon	5.5 ^b	0.17	(03)	25.0 ^c
Idaho	5.4 ^b	0.29	(04)	22.7 ^c
Kansas	5.4 ^b	0.20	(05)	22.7 ^c
Low Five States				
New Mexico	3.8	0.20	(47)	-13.6
New York	3.7 ^b	0.06	(48)	-15.9 ^c
New Jersey	3.7 ^b	0.09	(49)	-15.9 ^c
D.C.	3.2	0.34	(50)	-27.3 ^c
Hawaii	2.9 ^b	0.21	(51)	-34.1 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.4	0.14	(32)	0.0	Montana	4.8	0.31	(16)	9.1
Alaska	4.3	0.45	(36)	-2.3	Nebraska	4.9	0.23	(14)	11.4
Arizona	4.1	0.12	(39)	-6.8	Nevada	4.1	0.20	(42)	-6.8
Arkansas	5.0	0.18	(10)	13.6	New Hampshire	4.9	0.27	(12)	11.4
California	4.3	0.05	(34)	-2.3	New Jersey	3.7 ^b	0.09	(49)	-15.9 ^c
Colorado	4.4	0.15	(31)	0.0	New Mexico	3.8	0.20	(47)	-13.6
Connecticut	4.1	0.15	(40)	-6.8	New York	3.7 ^b	0.06	(48)	-15.9 ^c
Delaware	4.5	0.33	(26)	2.3	North Carolina	4.4	0.10	(33)	0.0
D.C.	3.2	0.34	(50)	-27.3 ^c	North Dakota	4.9	0.39	(11)	11.4
Florida	4.2	0.06	(38)	-4.5	Ohio	4.5	0.09	(28)	2.3
Georgia	3.9 ^b	0.10	(46)	-11.4	Oklahoma	4.9	0.16	(15)	11.4
Hawaii	2.9 ^b	0.21	(51)	-34.1 ^c	Oregon	5.5 ^b	0.17	(03)	25.0 ^c
Idaho	5.4 ^b	0.29	(04)	22.7 ^c	Pennsylvania	4.2	0.08	(37)	-4.5
Illinois	4.0 ^b	0.08	(44)	-9.1	Rhode Island	4.9	0.29	(13)	11.4
Indiana	4.7	0.12	(19)	6.8	South Carolina	4.4	0.15	(29)	0.0
Iowa	5.3 ^b	0.18	(07)	20.5 ^c	South Dakota	5.3	0.36	(06)	20.5 ^c
Kansas	5.4 ^b	0.20	(05)	22.7 ^c	Tennessee	5.6 ^b	0.14	(02)	27.3 ^c
Kentucky	4.5	0.15	(25)	2.3	Texas	4.5	0.07	(27)	2.3
Louisiana	4.6	0.15	(21)	4.5	Utah	4.4	0.22	(30)	0.0
Maine	5.1	0.26	(09)	15.9 ^c	Vermont	4.3	0.36	(35)	-2.3
Maryland	4.0	0.12	(43)	-9.1	Virginia	4.0 ^b	0.11	(45)	-9.1
Massachusetts	4.1	0.11	(41)	-6.8	Washington	5.6 ^b	0.14	(01)	27.3 ^c
Michigan	4.7 ^b	0.10	(17)	6.8	West Virginia	4.7	0.22	(18)	6.8
Minnesota	4.6	0.14	(22)	4.5	Wisconsin	4.7	0.13	(20)	6.8
Mississippi	5.2 ^b	0.19	(08)	18.2 ^c	Wyoming	4.5	0.43	(24)	2.3
Missouri	4.6	0.12	(23)	4.5					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table III-14
BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.4	0.03		
High Five States				
South Dakota	7.3	0.62	(01)	35.2 ^c
Oregon	6.7 ^b	0.28	(02)	24.1 ^c
Washington	6.7 ^b	0.22	(03)	24.1 ^c
Kansas	6.6 ^b	0.32	(04)	22.2 ^c
Idaho	6.5	0.46	(05)	20.4 ^c
Low Five States				
New York	4.6 ^b	0.10	(47)	-14.8
Nevada	4.6	0.30	(48)	-14.8
New Jersey	4.6 ^b	0.15	(49)	-14.8
D.C.	3.9	0.57	(50)	-27.8 ^c
Hawaii	3.8 ^b	0.35	(51)	-29.6 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.4	0.23	(32)	0.0	Montana	5.5	0.48	(23)	1.9
Alaska	5.2	0.73	(37)	-3.7	Nebraska	5.6	0.37	(22)	3.7
Arizona	4.9	0.19	(44)	-9.3	Nevada	4.6	0.30	(48)	-14.8
Arkansas	5.8	0.30	(17)	7.4	New Hampshire	6.5	0.47	(08)	20.4 ^c
California	5.3	0.09	(33)	-1.9	New Jersey	4.6 ^b	0.15	(49)	-14.8
Colorado	5.3	0.24	(34)	-1.9	New Mexico	4.9	0.34	(42)	-9.3
Connecticut	4.9	0.24	(43)	-9.3	New York	4.6 ^b	0.10	(47)	-14.8
Delaware	5.5	0.54	(27)	1.9	North Carolina	5.4	0.17	(30)	0.0
D.C.	3.9	0.57	(50)	-27.8 ^c	North Dakota	5.6	0.60	(19)	3.7
Florida	5.1	0.10	(41)	-5.6	Ohio	5.5	0.14	(29)	1.9
Georgia	4.7 ^b	0.17	(46)	-13.0	Oklahoma	6.1	0.27	(12)	13.0
Hawaii	3.8 ^b	0.35	(51)	-29.6 ^c	Oregon	6.7 ^b	0.28	(02)	24.1 ^c
Idaho	6.5	0.46	(05)	20.4 ^c	Pennsylvania	5.1	0.13	(39)	-5.6
Illinois	4.8 ^b	0.13	(45)	-11.1	Rhode Island	6.5	0.51	(06)	20.4 ^c
Indiana	5.5	0.20	(26)	1.9	South Carolina	5.2	0.24	(35)	-3.7
Iowa	6.2	0.29	(11)	14.8	South Dakota	7.3	0.62	(01)	35.2 ^c
Kansas	6.6 ^b	0.32	(04)	22.2 ^c	Tennessee	6.5 ^b	0.22	(07)	20.4 ^c
Kentucky	5.4	0.24	(31)	0.0	Texas	5.5	0.11	(24)	1.9
Louisiana	5.9	0.25	(14)	9.3	Utah	5.6	0.36	(20)	3.7
Maine	6.3	0.44	(09)	16.7 ^c	Vermont	5.8	0.62	(16)	7.4
Maryland	5.1	0.21	(40)	-5.6	Virginia	5.2	0.18	(36)	-3.7
Massachusetts	5.1	0.19	(38)	-5.6	Washington	6.7 ^b	0.22	(03)	24.1 ^c
Michigan	5.8	0.16	(15)	7.4	West Virginia	5.9	0.36	(13)	9.3
Minnesota	5.6	0.22	(21)	3.7	Wisconsin	5.7	0.21	(18)	5.6
Mississippi	6.3	0.32	(10)	16.7 ^c	Wyoming	5.5	0.69	(25)	1.9
Missouri	5.5	0.20	(28)	1.9					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table III-15
BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.6	0.02		
High Five States				
Tennessee	4.8 ^b	0.17	(01)	33.3 ^c
Washington	4.7 ^b	0.17	(02)	30.6 ^c
Oregon	4.4 ^b	0.21	(03)	22.2 ^c
Iowa	4.4 ^b	0.23	(04)	22.2 ^c
Kansas	4.4	0.24	(05)	22.2 ^c
Low Five States				
Virginia	3.0 ^b	0.12	(47)	-16.7 ^c
New Jersey	2.9 ^b	0.11	(48)	-19.4 ^c
New Mexico	2.9	0.24	(49)	-19.4 ^c
D.C.	2.5	0.41	(50)	-30.6 ^c
Hawaii	2.1 ^b	0.25	(51)	-41.7 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.5	0.16	(33)	-2.8	Montana	4.2	0.41	(11)	16.7 ^c
Alaska	3.5	0.58	(36)	-2.8	Nebraska	4.2	0.29	(09)	16.7 ^c
Arizona	3.5	0.15	(37)	-2.8	Nevada	3.5	0.25	(35)	-2.8
Arkansas	4.3	0.23	(08)	19.4 ^c	New Hampshire	3.6	0.32	(30)	0.0
California	3.5	0.06	(32)	-2.8	New Jersey	2.9 ^b	0.11	(48)	-19.4 ^c
Colorado	3.6	0.18	(26)	0.0	New Mexico	2.9	0.24	(49)	-19.4 ^c
Connecticut	3.5	0.19	(34)	-2.8	New York	3.0 ^b	0.07	(46)	-16.7 ^c
Delaware	3.7	0.40	(23)	2.8	North Carolina	3.5	0.12	(31)	-2.8
D.C.	2.5	0.41	(50)	-30.6 ^c	North Dakota	4.3	0.50	(06)	19.4 ^c
Florida	3.4	0.08	(39)	-5.6	Ohio	3.7	0.11	(25)	2.8
Georgia	3.2	0.12	(44)	-11.1	Oklahoma	3.9	0.20	(14)	8.3
Hawaii	2.1 ^b	0.25	(51)	-41.7 ^c	Oregon	4.4 ^b	0.21	(03)	22.2 ^c
Idaho	4.3	0.36	(07)	19.4 ^c	Pennsylvania	3.5	0.10	(38)	-2.8
Illinois	3.4	0.10	(41)	-5.6	Rhode Island	3.7	0.34	(22)	2.8
Indiana	4.0	0.15	(12)	11.1	South Carolina	3.9	0.18	(15)	8.3
Iowa	4.4 ^b	0.23	(04)	22.2 ^c	South Dakota	3.7	0.41	(24)	2.8
Kansas	4.4	0.24	(05)	22.2 ^c	Tennessee	4.8 ^b	0.17	(01)	33.3 ^c
Kentucky	3.8	0.18	(18)	5.6	Texas	3.6	0.08	(29)	0.0
Louisiana	3.7	0.18	(21)	2.8	Utah	3.4	0.27	(40)	-5.6
Maine	3.9	0.32	(13)	8.3	Vermont	3.1	0.43	(45)	-13.9
Maryland	3.2	0.15	(43)	-11.1	Virginia	3.0 ^b	0.12	(47)	-16.7 ^c
Massachusetts	3.3	0.13	(42)	-8.3	Washington	4.7 ^b	0.17	(02)	30.6 ^c
Michigan	3.8	0.12	(16)	5.6	West Virginia	3.6	0.26	(28)	0.0
Minnesota	3.7	0.17	(20)	2.8	Wisconsin	3.8	0.16	(19)	5.6
Mississippi	4.2	0.23	(10)	16.7 ^c	Wyoming	3.6	0.53	(27)	0.0
Missouri	3.8	0.15	(17)	5.6					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table IV-20
FEMALE BREAST CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	25.0	0.06		
High Five States				
D.C.	31.0 ^b	1.38	(01)	24.0 ^c
Louisiana	29.7 ^b	0.50	(02)	18.8 ^c
New Jersey	27.8 ^b	0.33	(03)	11.2 ^c
Ohio	27.6 ^b	0.28	(04)	10.4 ^c
Maryland	27.4 ^b	0.42	(05)	9.6
Low Five States				
Arizona	22.2 ^b	0.38	(47)	-11.2 ^c
New Mexico	22.1 ^b	0.67	(48)	-11.6 ^c
Idaho	22.1 ^b	0.80	(49)	-11.6 ^c
Alaska	20.5	1.42	(50)	-18.0 ^c
Hawaii	18.0 ^b	0.71	(51)	-28.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	26.0	0.44	(12)	4.0	Montana	23.8	0.94	(35)	-4.8
Alaska	20.5	1.42	(50)	-18.0 ^c	Nebraska	24.0	0.69	(33)	-4.0
Arizona	22.2 ^b	0.38	(47)	-11.2 ^c	Nevada	25.3	0.68	(17)	1.2
Arkansas	24.4	0.55	(27)	-2.4	New Hampshire	24.3	0.81	(29)	-2.8
California	23.7 ^b	0.16	(36)	-5.2	New Jersey	27.8 ^b	0.33	(03)	11.2 ^c
Colorado	22.6 ^b	0.46	(45)	-9.6	New Mexico	22.1 ^b	0.67	(48)	-11.6 ^c
Connecticut	25.1	0.49	(22)	0.4	New York	25.6	0.21	(13)	2.4
Delaware	25.1	1.03	(20)	0.4	North Carolina	25.4	0.33	(15)	1.6
D.C.	31.0 ^b	1.38	(01)	24.0 ^c	North Dakota	24.0	1.13	(32)	-4.0
Florida	23.0 ^b	0.20	(43)	-8.0	Ohio	27.6 ^b	0.28	(04)	10.4 ^c
Georgia	24.7	0.34	(25)	-1.2	Oklahoma	25.4	0.50	(16)	1.6
Hawaii	18.0 ^b	0.71	(51)	-28.0 ^c	Oregon	24.5	0.49	(26)	-2.0
Idaho	22.1 ^b	0.80	(49)	-11.6 ^c	Pennsylvania	27.0 ^b	0.26	(06)	8.0
Illinois	26.2 ^b	0.27	(11)	4.8	Rhode Island	23.5	0.84	(38)	-6.0
Indiana	25.0	0.38	(23)	0.0	South Carolina	25.3	0.46	(19)	1.2
Iowa	23.1 ^b	0.50	(41)	-7.6	South Dakota	22.9	1.02	(44)	-8.4
Kansas	24.3	0.56	(28)	-2.8	Tennessee	26.2	0.40	(10)	4.8
Kentucky	25.6	0.46	(14)	2.4	Texas	23.8 ^b	0.21	(34)	-4.8
Louisiana	29.7 ^b	0.50	(02)	18.8 ^c	Utah	23.1	0.70	(40)	-7.6
Maine	23.5	0.75	(39)	-6.0	Vermont	24.2	1.14	(30)	-3.2
Maryland	27.4 ^b	0.42	(05)	9.6	Virginia	26.5 ^b	0.36	(09)	6.0
Massachusetts	24.8	0.36	(24)	-0.8	Washington	23.6 ^b	0.38	(37)	-5.6
Michigan	25.3	0.30	(18)	1.2	West Virginia	25.1	0.65	(21)	0.4
Minnesota	23.0 ^b	0.41	(42)	-8.0	Wisconsin	24.0	0.39	(31)	-4.0
Mississippi	26.9	0.58	(07)	7.6	Wyoming	22.6	1.30	(46)	-9.6
Missouri	26.6 ^b	0.40	(08)	6.4					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 10% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table V-11
CERVIX UTERI CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.5	0.02		
High Five States				
D.C.	3.7	0.49	(01)	48.0 ^c
Mississippi	3.6 ^b	0.22	(02)	44.0 ^c
Arkansas	3.6 ^b	0.22	(03)	44.0 ^c
West Virginia	3.5 ^b	0.26	(04)	40.0 ^c
Texas	3.2 ^b	0.08	(05)	28.0 ^c
Low Five States				
Connecticut	1.7 ^b	0.13	(47)	-32.0 ^c
North Dakota	1.7	0.31	(48)	-32.0 ^c
Minnesota	1.6 ^b	0.11	(49)	-36.0 ^c
Massachusetts	1.5 ^b	0.09	(50)	-40.0 ^c
Utah	1.5 ^b	0.18	(51)	-40.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.1 ^b	0.16	(07)	24.0 ^c	Montana	1.9	0.27	(40)	-24.0 ^c
Alaska	1.9	0.39	(42)	-24.0 ^c	Nebraska	2.1	0.21	(36)	-16.0 ^c
Arizona	2.4	0.13	(26)	-4.0	Nevada	2.7	0.22	(17)	8.0
Arkansas	3.6 ^b	0.22	(03)	44.0 ^c	New Hampshire	2.3	0.25	(29)	-8.0
California	2.4	0.05	(25)	-4.0	New Jersey	2.7	0.10	(16)	8.0
Colorado	1.8 ^b	0.13	(44)	-28.0 ^c	New Mexico	2.6	0.23	(19)	4.0
Connecticut	1.7 ^b	0.13	(47)	-32.0 ^c	New York	2.6	0.07	(18)	4.0
Delaware	2.9	0.36	(09)	16.0 ^c	North Carolina	2.5	0.11	(21)	0.0
D.C.	3.7	0.49	(01)	48.0 ^c	North Dakota	1.7	0.31	(48)	-32.0 ^c
Florida	2.8 ^b	0.08	(14)	12.0	Ohio	2.4	0.09	(23)	-4.0
Georgia	2.8	0.11	(15)	12.0	Oklahoma	2.8	0.18	(12)	12.0
Hawaii	2.0	0.25	(38)	-20.0 ^c	Oregon	2.1	0.15	(35)	-16.0 ^c
Idaho	2.2	0.25	(34)	-12.0	Pennsylvania	2.3	0.08	(28)	-8.0
Illinois	2.8 ^b	0.09	(11)	12.0	Rhode Island	1.8	0.25	(43)	-28.0 ^c
Indiana	2.5	0.12	(22)	0.0	South Carolina	2.8	0.16	(13)	12.0
Iowa	2.2	0.17	(30)	-12.0	South Dakota	2.0	0.32	(37)	-20.0 ^c
Kansas	2.4	0.18	(24)	-4.0	Tennessee	3.2 ^b	0.14	(06)	28.0 ^c
Kentucky	2.9	0.16	(10)	16.0 ^c	Texas	3.2 ^b	0.08	(05)	28.0 ^c
Louisiana	3.0	0.16	(08)	20.0 ^c	Utah	1.5 ^b	0.18	(51)	-40.0 ^c
Maine	1.9	0.22	(39)	-24.0 ^c	Vermont	2.2	0.35	(32)	-12.0
Maryland	2.4	0.13	(27)	-4.0	Virginia	2.2	0.11	(31)	-12.0
Massachusetts	1.5 ^b	0.09	(50)	-40.0 ^c	Washington	1.8 ^b	0.11	(46)	-28.0 ^c
Michigan	2.2 ^b	0.09	(33)	-12.0	West Virginia	3.5 ^b	0.26	(04)	40.0 ^c
Minnesota	1.6 ^b	0.11	(49)	-36.0 ^c	Wisconsin	1.9 ^b	0.11	(41)	-24.0 ^c
Mississippi	3.6 ^b	0.22	(02)	44.0 ^c	Wyoming	1.8	0.37	(45)	-28.0 ^c
Missouri	2.6	0.13	(20)	4.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table VI-20
COLON AND RECTUM CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	18.8	0.04		
High Five States				
Louisiana	22.3 ^b	0.32	(01)	18.6 ^c
West Virginia	22.3 ^b	0.45	(02)	18.6 ^c
Kentucky	22.0 ^b	0.33	(03)	17.0 ^c
D.C.	21.4	0.86	(04)	13.8 ^c
Mississippi	21.0 ^b	0.39	(05)	11.7 ^c
Low Five States				
California	16.7 ^b	0.10	(47)	-11.2 ^c
Arizona	16.2 ^b	0.24	(48)	-13.8 ^c
Hawaii	16.1 ^b	0.49	(49)	-14.4 ^c
Idaho	15.6 ^b	0.49	(50)	-17.0 ^c
Utah	13.6 ^b	0.40	(51)	-27.7 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	18.7	0.28	(32)	-0.5	Montana	16.8	0.57	(46)	-10.6 ^c
Alaska	17.9	1.06	(37)	-4.8	Nebraska	19.7	0.46	(17)	4.8
Arizona	16.2 ^b	0.24	(48)	-13.8 ^c	Nevada	20.4	0.46	(12)	8.5
Arkansas	20.4 ^b	0.37	(11)	8.5	New Hampshire	18.9	0.54	(29)	0.5
California	16.7 ^b	0.10	(47)	-11.2 ^c	New Jersey	20.8 ^b	0.21	(06)	10.6 ^c
Colorado	17.2 ^b	0.31	(41)	-8.5	New Mexico	16.9 ^b	0.43	(44)	-10.1 ^c
Connecticut	17.7 ^b	0.30	(39)	-5.9	New York	19.2	0.14	(26)	2.1
Delaware	19.8	0.68	(16)	5.3	North Carolina	18.3	0.21	(35)	-2.7
D.C.	21.4	0.86	(04)	13.8 ^c	North Dakota	18.9	0.71	(30)	0.5
Florida	16.9 ^b	0.12	(43)	-10.1 ^c	Ohio	20.6 ^b	0.18	(09)	9.6
Georgia	18.5	0.23	(33)	-1.6	Oklahoma	19.5	0.33	(18)	3.7
Hawaii	16.1 ^b	0.49	(49)	-14.4 ^c	Oregon	17.6 ^b	0.31	(40)	-6.4
Idaho	15.6 ^b	0.49	(50)	-17.0 ^c	Pennsylvania	20.6 ^b	0.17	(10)	9.6
Illinois	20.7 ^b	0.18	(07)	10.1 ^c	Rhode Island	19.3	0.56	(21)	2.7
Indiana	20.6 ^b	0.26	(08)	9.6	South Carolina	19.2	0.31	(25)	2.1
Iowa	19.5	0.33	(20)	3.7	South Dakota	19.2	0.66	(24)	2.1
Kansas	18.9	0.36	(28)	0.5	Tennessee	20.0 ^b	0.26	(13)	6.4
Kentucky	22.0 ^b	0.33	(03)	17.0 ^c	Texas	17.9 ^b	0.14	(38)	-4.8
Louisiana	22.3 ^b	0.32	(01)	18.6 ^c	Utah	13.6 ^b	0.40	(51)	-27.7 ^c
Maine	19.5	0.51	(19)	3.7	Vermont	19.3	0.77	(22)	2.7
Maryland	20.0 ^b	0.28	(14)	6.4	Virginia	18.9	0.23	(27)	0.5
Massachusetts	19.3	0.23	(23)	2.7	Washington	16.9 ^b	0.24	(45)	-10.1 ^c
Michigan	18.7	0.19	(31)	-0.5	West Virginia	22.3 ^b	0.45	(02)	18.6 ^c
Minnesota	16.9 ^b	0.26	(42)	-10.1 ^c	Wisconsin	18.0	0.25	(36)	-4.3
Mississippi	21.0 ^b	0.39	(05)	11.7 ^c	Wyoming	18.4	0.86	(34)	-2.1
Missouri	19.9 ^b	0.25	(15)	5.9					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table VI-21
COLON AND RECTUM CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	22.7	0.06		
High Five States				
Louisiana	28.5 ^b	0.58	(01)	25.6 ^c
West Virginia	27.0 ^b	0.79	(02)	18.9 ^c
D.C.	27.0	1.58	(03)	18.9 ^c
Kentucky	26.6 ^b	0.58	(04)	17.2 ^c
Illinois	25.8 ^b	0.33	(05)	13.7 ^c
Low Five States				
Washington	19.7 ^b	0.40	(47)	-13.2 ^c
Wyoming	19.7	1.37	(48)	-13.2 ^c
Arizona	19.5 ^b	0.40	(49)	-14.1 ^c
Idaho	18.2 ^b	0.81	(50)	-19.8 ^c
Utah	15.9 ^b	0.67	(51)	-30.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	23.9	0.51	(17)	5.3	Montana	20.5	0.97	(42)	-9.7
Alaska	21.6	1.80	(37)	-4.8	Nebraska	23.5	0.78	(21)	3.5
Arizona	19.5 ^b	0.40	(49)	-14.1 ^c	Nevada	24.6	0.77	(13)	8.4
Arkansas	25.0 ^b	0.63	(10)	10.1 ^c	New Hampshire	23.0	0.95	(28)	1.3
California	19.8 ^b	0.17	(46)	-12.8 ^c	New Jersey	24.9 ^b	0.37	(12)	9.7
Colorado	20.0 ^b	0.52	(44)	-11.9 ^c	New Mexico	20.6	0.73	(41)	-9.3
Connecticut	20.7 ^b	0.52	(40)	-8.8	New York	23.1	0.24	(26)	1.8
Delaware	23.8	1.17	(18)	4.8	North Carolina	22.1	0.37	(33)	-2.6
D.C.	27.0	1.58	(03)	18.9 ^c	North Dakota	22.0	1.17	(34)	-3.1
Florida	20.3 ^b	0.21	(43)	-10.6 ^c	Ohio	24.9 ^b	0.32	(11)	9.7
Georgia	22.7	0.41	(30)	0.0	Oklahoma	24.3	0.57	(15)	7.0
Hawaii	21.1	0.83	(38)	-7.0	Oregon	21.0	0.51	(39)	-7.5
Idaho	18.2 ^b	0.81	(50)	-19.8 ^c	Pennsylvania	25.4 ^b	0.29	(08)	11.9 ^c
Illinois	25.8 ^b	0.33	(05)	13.7 ^c	Rhode Island	23.0	0.98	(27)	1.3
Indiana	25.7 ^b	0.46	(06)	13.2 ^c	South Carolina	23.7	0.54	(20)	4.4
Iowa	23.7	0.58	(19)	4.4	South Dakota	23.4	1.12	(22)	3.1
Kansas	22.4	0.61	(32)	-1.3	Tennessee	25.0 ^b	0.47	(09)	10.1 ^c
Kentucky	26.6 ^b	0.58	(04)	17.2 ^c	Texas	21.9	0.24	(36)	-3.5
Louisiana	28.5 ^b	0.58	(01)	25.6 ^c	Utah	15.9 ^b	0.67	(51)	-30.0 ^c
Maine	22.8	0.86	(29)	0.4	Vermont	23.1	1.32	(25)	1.8
Maryland	24.1	0.49	(16)	6.2	Virginia	23.4	0.42	(23)	3.1
Massachusetts	23.3	0.41	(24)	2.6	Washington	19.7 ^b	0.40	(47)	-13.2 ^c
Michigan	22.4	0.33	(31)	-1.3	West Virginia	27.0 ^b	0.79	(02)	18.9 ^c
Minnesota	19.8 ^b	0.44	(45)	-12.8 ^c	Wisconsin	22.0	0.43	(35)	-3.1
Mississippi	25.5 ^b	0.68	(07)	12.3 ^c	Wyoming	19.7	1.37	(48)	-13.2 ^c
Missouri	24.4 ^b	0.44	(14)	7.5					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table VI-22
COLON AND RECTUM CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	15.9	0.04		
High Five States				
Kentucky	18.9 ^b	0.39	(01)	18.9 ^c
West Virginia	18.8 ^b	0.55	(02)	18.2 ^c
D.C.	18.1	1.02	(03)	13.8 ^c
New Jersey	17.9 ^b	0.26	(04)	12.6 ^c
Louisiana	17.9 ^b	0.38	(05)	12.6 ^c
Low Five States				
Montana	13.8	0.70	(47)	-13.2 ^c
Arizona	13.5 ^b	0.29	(48)	-15.1 ^c
Idaho	13.4 ^b	0.61	(49)	-15.7 ^c
Hawaii	11.8 ^b	0.56	(50)	-25.8 ^c
Utah	11.8 ^b	0.50	(51)	-25.8 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	15.2	0.33	(36)	-4.4	Montana	13.8	0.70	(47)	-13.2 ^c
Alaska	14.8	1.30	(40)	-6.9	Nebraska	16.9	0.56	(14)	6.3
Arizona	13.5 ^b	0.29	(48)	-15.1 ^c	Nevada	16.8	0.56	(17)	5.7
Arkansas	16.8	0.44	(16)	5.7	New Hampshire	16.1	0.66	(27)	1.3
California	14.3 ^b	0.13	(44)	-10.1 ^c	New Jersey	17.9 ^b	0.26	(04)	12.6 ^c
Colorado	15.1	0.38	(38)	-5.0	New Mexico	13.8 ^b	0.52	(46)	-13.2 ^c
Connecticut	15.5	0.37	(35)	-2.5	New York	16.5 ^b	0.17	(23)	3.8
Delaware	16.7	0.83	(18)	5.0	North Carolina	15.5	0.25	(34)	-2.5
D.C.	18.1	1.02	(03)	13.8 ^c	North Dakota	16.5	0.88	(24)	3.8
Florida	14.2 ^b	0.15	(45)	-10.7 ^c	Ohio	17.5 ^b	0.22	(07)	10.1 ^c
Georgia	15.7	0.27	(33)	-1.3	Oklahoma	16.0	0.39	(28)	0.6
Hawaii	11.8 ^b	0.56	(50)	-25.8 ^c	Oregon	15.1	0.38	(37)	-5.0
Idaho	13.4 ^b	0.61	(49)	-15.7 ^c	Pennsylvania	17.3 ^b	0.20	(08)	8.8
Illinois	17.2 ^b	0.22	(09)	8.2	Rhode Island	17.1	0.69	(10)	7.5
Indiana	17.0 ^b	0.30	(12)	6.9	South Carolina	15.9	0.36	(30)	0.0
Iowa	16.5	0.40	(25)	3.8	South Dakota	15.8	0.79	(31)	-0.6
Kansas	16.2	0.44	(26)	1.9	Tennessee	16.5	0.31	(20)	3.8
Kentucky	18.9 ^b	0.39	(01)	18.9 ^c	Texas	14.7 ^b	0.17	(42)	-7.5
Louisiana	17.9 ^b	0.38	(05)	12.6 ^c	Utah	11.8 ^b	0.50	(51)	-25.8 ^c
Maine	17.0	0.62	(13)	6.9	Vermont	16.5	0.93	(21)	3.8
Maryland	16.9	0.33	(15)	6.3	Virginia	15.7	0.28	(32)	-1.3
Massachusetts	16.5	0.28	(22)	3.8	Washington	14.6 ^b	0.30	(43)	-8.2
Michigan	15.9	0.23	(29)	0.0	West Virginia	18.8 ^b	0.55	(02)	18.2 ^c
Minnesota	14.7 ^b	0.32	(41)	-7.5	Wisconsin	15.0	0.30	(39)	-5.7
Mississippi	17.9 ^b	0.47	(06)	12.6 ^c	Wyoming	17.0	1.12	(11)	6.9
Missouri	16.7	0.30	(19)	5.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table VII-11
CORPUS AND UTERUS, NOS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.1	0.02		
High Five States				
D.C.	5.9	0.59	(01)	43.9 ^c
Vermont	5.5	0.54	(02)	34.1 ^c
New Jersey	5.2 ^b	0.14	(03)	26.8 ^c
Nebraska	5.0	0.32	(04)	22.0 ^c
New York	4.9 ^b	0.09	(05)	19.5 ^c
Low Five States				
Kentucky	3.4 ^b	0.17	(47)	-17.1 ^c
Rhode Island	3.4	0.31	(48)	-17.1 ^c
Nevada	3.1 ^b	0.24	(49)	-24.4 ^c
Arizona	3.0 ^b	0.14	(50)	-26.8 ^c
Alaska	2.6	0.54	(51)	-36.6 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.6	0.16	(39)	-12.2	Montana	4.8	0.42	(09)	17.1 ^c
Alaska	2.6	0.54	(51)	-36.6 ^c	Nebraska	5.0	0.32	(04)	22.0 ^c
Arizona	3.0 ^b	0.14	(50)	-26.8 ^c	Nevada	3.1 ^b	0.24	(49)	-24.4 ^c
Arkansas	3.4 ^b	0.20	(46)	-17.1 ^c	New Hampshire	4.0	0.33	(27)	-2.4
California	3.9 ^b	0.07	(30)	-4.9	New Jersey	5.2 ^b	0.14	(03)	26.8 ^c
Colorado	3.5 ^b	0.18	(45)	-14.6	New Mexico	3.9	0.28	(31)	-4.9
Connecticut	4.3	0.20	(22)	4.9	New York	4.9 ^b	0.09	(05)	19.5 ^c
Delaware	4.8	0.45	(07)	17.1 ^c	North Carolina	4.0	0.13	(28)	-2.4
D.C.	5.9	0.59	(01)	43.9 ^c	North Dakota	4.6	0.48	(11)	12.2
Florida	3.6 ^b	0.08	(38)	-12.2	Ohio	4.4	0.11	(16)	7.3
Georgia	3.7 ^b	0.13	(37)	-9.8	Oklahoma	3.8	0.19	(32)	-7.3
Hawaii	3.7	0.32	(36)	-9.8	Oregon	3.8	0.19	(34)	-7.3
Idaho	3.5	0.32	(43)	-14.6	Pennsylvania	4.9 ^b	0.11	(06)	19.5 ^c
Illinois	4.5	0.11	(14)	9.8	Rhode Island	3.4	0.31	(48)	-17.1 ^c
Indiana	4.8 ^b	0.16	(08)	17.1 ^c	South Carolina	4.0	0.18	(26)	-2.4
Iowa	4.5	0.22	(13)	9.8	South Dakota	4.4	0.43	(18)	7.3
Kansas	3.8	0.22	(35)	-7.3	Tennessee	3.6 ^b	0.14	(40)	-12.2
Kentucky	3.4 ^b	0.17	(47)	-17.1 ^c	Texas	3.6 ^b	0.08	(41)	-12.2
Louisiana	3.6	0.17	(42)	-12.2	Utah	3.8	0.29	(33)	-7.3
Maine	4.5	0.33	(12)	9.8	Vermont	5.5	0.54	(02)	34.1 ^c
Maryland	4.3	0.17	(20)	4.9	Virginia	4.1	0.14	(23)	0.0
Massachusetts	4.4	0.15	(19)	7.3	Washington	4.1	0.16	(24)	0.0
Michigan	4.7 ^b	0.13	(10)	14.6	West Virginia	4.1	0.26	(25)	0.0
Minnesota	4.3	0.18	(21)	4.9	Wisconsin	4.4	0.17	(15)	7.3
Mississippi	3.9	0.22	(29)	-4.9	Wyoming	3.5	0.51	(44)	-14.6
Missouri	4.4	0.16	(17)	7.3					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table VIII-13
ESOPHAGUS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.4	0.02		
High Five States				
D.C.	6.1 ^b	0.46	(01)	38.6 ^c
Wyoming	6.0	0.49	(02)	36.4 ^c
Maine	6.0 ^b	0.28	(03)	36.4 ^c
New Hampshire	5.7 ^b	0.30	(04)	29.5 ^c
Massachusetts	5.3 ^b	0.12	(05)	20.5 ^c
Low Five States				
Texas	3.8 ^b	0.06	(47)	-13.6
California	3.7 ^b	0.05	(48)	-15.9 ^c
Arkansas	3.6 ^b	0.16	(49)	-18.2 ^c
Utah	3.3 ^b	0.19	(50)	-25.0 ^c
Hawaii	2.6 ^b	0.19	(51)	-40.9 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.0	0.13	(44)	-9.1	Montana	4.7	0.31	(16)	6.8
Alaska	4.9	0.54	(13)	11.4	Nebraska	4.4	0.22	(32)	0.0
Arizona	4.1	0.12	(41)	-6.8	Nevada	4.4	0.21	(29)	0.0
Arkansas	3.6 ^b	0.16	(49)	-18.2 ^c	New Hampshire	5.7 ^b	0.30	(04)	29.5 ^c
California	3.7 ^b	0.05	(48)	-15.9 ^c	New Jersey	4.3	0.10	(36)	-2.3
Colorado	4.0	0.15	(43)	-9.1	New Mexico	3.8	0.20	(45)	-13.6
Connecticut	4.7	0.16	(17)	6.8	New York	4.4	0.07	(28)	0.0
Delaware	5.2	0.35	(10)	18.2 ^c	North Carolina	4.2	0.10	(37)	-4.5
D.C.	6.1 ^b	0.46	(01)	38.6 ^c	North Dakota	4.2	0.34	(38)	-4.5
Florida	4.1 ^b	0.06	(39)	-6.8	Ohio	5.2 ^b	0.09	(09)	18.2 ^c
Georgia	4.4	0.11	(33)	0.0	Oklahoma	4.1	0.15	(42)	-6.8
Hawaii	2.6 ^b	0.19	(51)	-40.9 ^c	Oregon	5.2 ^b	0.17	(08)	18.2 ^c
Idaho	4.3	0.26	(34)	-2.3	Pennsylvania	5.0 ^b	0.08	(11)	13.6
Illinois	4.7	0.09	(20)	6.8	Rhode Island	5.2	0.30	(07)	18.2 ^c
Indiana	4.6	0.12	(22)	4.5	South Carolina	4.9	0.15	(14)	11.4
Iowa	4.7	0.17	(19)	6.8	South Dakota	4.4	0.32	(30)	0.0
Kansas	3.8 ^b	0.16	(46)	-13.6	Tennessee	4.1	0.12	(40)	-6.8
Kentucky	4.6	0.15	(21)	4.5	Texas	3.8 ^b	0.06	(47)	-13.6
Louisiana	4.4	0.14	(26)	0.0	Utah	3.3 ^b	0.19	(50)	-25.0 ^c
Maine	6.0 ^b	0.28	(03)	36.4 ^c	Vermont	4.8	0.38	(15)	9.1
Maryland	4.7	0.13	(18)	6.8	Virginia	4.4	0.11	(27)	0.0
Massachusetts	5.3 ^b	0.12	(05)	20.5 ^c	Washington	4.6	0.13	(23)	4.5
Michigan	5.0 ^b	0.10	(12)	13.6	West Virginia	4.5	0.20	(24)	2.3
Minnesota	4.4	0.13	(31)	0.0	Wisconsin	5.3 ^b	0.14	(06)	20.5 ^c
Mississippi	4.3	0.17	(35)	-2.3	Wyoming	6.0	0.49	(02)	36.4 ^c
Missouri	4.4	0.12	(25)	0.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table VIII-14
ESOPHAGUS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.8	0.04		
High Five States				
D.C.	11.5 ^b	0.99	(01)	47.4 ^c
Maine	10.8 ^b	0.57	(02)	38.5 ^c
Wyoming	10.6	0.98	(03)	35.9 ^c
New Hampshire	10.2 ^b	0.61	(04)	30.8 ^c
Massachusetts	9.9 ^b	0.26	(05)	26.9 ^c
Low Five States				
Texas	6.7 ^b	0.13	(47)	-14.1
California	6.5 ^b	0.10	(48)	-16.7 ^c
Arkansas	6.2 ^b	0.31	(49)	-20.5 ^c
Utah	6.1 ^b	0.40	(50)	-21.8 ^c
Hawaii	4.7 ^b	0.39	(51)	-39.7 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.2	0.27	(43)	-7.7	Montana	7.5	0.57	(35)	-3.8
Alaska	8.2	1.05	(20)	5.1	Nebraska	7.9	0.45	(27)	1.3
Arizona	7.2	0.24	(42)	-7.7	Nevada	7.3	0.39	(39)	-6.4
Arkansas	6.2 ^b	0.31	(49)	-20.5 ^c	New Hampshire	10.2 ^b	0.61	(04)	30.8 ^c
California	6.5 ^b	0.10	(48)	-16.7 ^c	New Jersey	7.5	0.20	(34)	-3.8
Colorado	7.3	0.30	(40)	-6.4	New Mexico	6.9	0.42	(45)	-11.5
Connecticut	8.4	0.32	(15)	7.7	New York	7.7	0.14	(31)	-1.3
Delaware	9.4	0.72	(07)	20.5 ^c	North Carolina	7.3	0.20	(41)	-6.4
D.C.	11.5 ^b	0.99	(01)	47.4 ^c	North Dakota	7.5	0.68	(38)	-3.8
Florida	7.2 ^b	0.12	(44)	-7.7	Ohio	9.4 ^b	0.19	(08)	20.5 ^c
Georgia	7.7	0.23	(30)	-1.3	Oklahoma	7.5	0.31	(36)	-3.8
Hawaii	4.7 ^b	0.39	(51)	-39.7 ^c	Oregon	9.0 ^b	0.33	(10)	15.4 ^c
Idaho	7.5	0.50	(33)	-3.8	Pennsylvania	9.1 ^b	0.17	(09)	16.7 ^c
Illinois	8.4	0.18	(17)	7.7	Rhode Island	8.7	0.59	(13)	11.5
Indiana	8.3	0.25	(18)	6.4	South Carolina	8.8	0.31	(12)	12.8
Iowa	8.6	0.34	(14)	10.3	South Dakota	8.0	0.65	(25)	2.6
Kansas	6.9	0.34	(46)	-11.5	Tennessee	7.5	0.24	(37)	-3.8
Kentucky	8.4	0.30	(16)	7.7	Texas	6.7 ^b	0.13	(47)	-14.1
Louisiana	8.1	0.30	(24)	3.8	Utah	6.1 ^b	0.40	(50)	-21.8 ^c
Maine	10.8 ^b	0.57	(02)	38.5 ^c	Vermont	8.0	0.75	(26)	2.6
Maryland	8.3	0.27	(19)	6.4	Virginia	7.7	0.23	(32)	-1.3
Massachusetts	9.9 ^b	0.26	(05)	26.9 ^c	Washington	8.1	0.25	(21)	3.8
Michigan	8.9 ^b	0.20	(11)	14.1	West Virginia	7.9	0.41	(28)	1.3
Minnesota	7.9	0.27	(29)	1.3	Wisconsin	9.4 ^b	0.28	(06)	20.5 ^c
Mississippi	8.1	0.36	(22)	3.8	Wyoming	10.6	0.98	(03)	35.9 ^c
Missouri	8.1	0.25	(23)	3.8					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table VIII-15
ESOPHAGUS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.7	0.01		
High Five States				
Rhode Island	2.4	0.26	(01)	41.2 ^c
Montana	2.4	0.29	(02)	41.2 ^c
Vermont	2.3	0.35	(03)	35.3 ^c
D.C.	2.2	0.37	(04)	29.4 ^c
Wyoming	2.2	0.41	(05)	29.4 ^c
Low Five States				
Oklahoma	1.3 ^b	0.11	(47)	-23.5 ^c
New Mexico	1.3	0.16	(48)	-23.5 ^c
Kansas	1.3 ^b	0.12	(49)	-23.5 ^c
Utah	0.8 ^b	0.13	(50)	-52.9 ^c
Hawaii	0.8 ^b	0.14	(51)	-52.9 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.5	0.10	(42)	-11.8	Montana	2.4	0.29	(02)	41.2 ^c
Alaska	2.1	0.49	(07)	23.5 ^c	Nebraska	1.6	0.17	(36)	-5.9
Arizona	1.4	0.10	(44)	-17.6 ^c	Nevada	1.7	0.18	(27)	0.0
Arkansas	1.6	0.14	(37)	-5.9	New Hampshire	2.2	0.24	(06)	29.4 ^c
California	1.6	0.04	(33)	-5.9	New Jersey	1.8	0.08	(22)	5.9
Colorado	1.4	0.12	(45)	-17.6 ^c	New Mexico	1.3	0.16	(48)	-23.5 ^c
Connecticut	1.9	0.13	(18)	11.8	New York	1.9	0.06	(16)	11.8
Delaware	1.9	0.28	(15)	11.8	North Carolina	1.8	0.09	(26)	5.9
D.C.	2.2	0.37	(04)	29.4 ^c	North Dakota	1.5	0.28	(38)	-11.8
Florida	1.6	0.05	(31)	-5.9	Ohio	2.0 ^b	0.08	(09)	17.6 ^c
Georgia	1.8	0.09	(24)	5.9	Oklahoma	1.3 ^b	0.11	(47)	-23.5 ^c
Hawaii	0.8 ^b	0.14	(51)	-52.9 ^c	Oregon	2.0	0.14	(08)	17.6 ^c
Idaho	1.5	0.21	(40)	-11.8	Pennsylvania	1.9	0.07	(17)	11.8
Illinois	1.9	0.07	(19)	11.8	Rhode Island	2.4	0.26	(01)	41.2 ^c
Indiana	1.7	0.10	(28)	0.0	South Carolina	1.8	0.12	(21)	5.9
Iowa	1.7	0.13	(29)	0.0	South Dakota	1.5	0.25	(41)	-11.8
Kansas	1.3 ^b	0.12	(49)	-23.5 ^c	Tennessee	1.5	0.09	(43)	-11.8
Kentucky	1.6	0.11	(35)	-5.9	Texas	1.5 ^b	0.05	(39)	-11.8
Louisiana	1.6	0.12	(30)	-5.9	Utah	0.8 ^b	0.13	(50)	-52.9 ^c
Maine	2.0	0.21	(11)	17.6 ^c	Vermont	2.3	0.35	(03)	35.3 ^c
Maryland	2.0	0.12	(10)	17.6 ^c	Virginia	1.8	0.10	(20)	5.9
Massachusetts	2.0	0.10	(13)	17.6 ^c	Washington	1.8	0.11	(25)	5.9
Michigan	1.9	0.08	(14)	11.8	West Virginia	1.8	0.17	(23)	5.9
Minnesota	1.6	0.11	(32)	-5.9	Wisconsin	2.0	0.11	(12)	17.6 ^c
Mississippi	1.4	0.13	(46)	-17.6 ^c	Wyoming	2.2	0.41	(05)	29.4 ^c
Missouri	1.6	0.09	(34)	-5.9					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table IX-13
HODGKIN LYMPHOMA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.4	0.01		
High Five States				
Arkansas	0.6	0.07	(01)	50.0 ^c
Mississippi	0.6	0.07	(02)	50.0 ^c
Kentucky	0.6	0.05	(03)	50.0 ^c
Alaska	0.6	0.17	(04)	50.0 ^c
Oklahoma	0.6	0.06	(05)	50.0 ^c
Low Five States				
Georgia	0.4	0.03	(47)	0.0
Nevada	0.4	0.06	(48)	0.0
North Dakota	0.4	0.11	(49)	0.0
Maine	0.3	0.07	(50)	-25.0 ^c
Utah	0.3	0.06	(51)	-25.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.5	0.05	(16)	25.0 ^c	Montana	0.5	0.10	(13)	25.0 ^c
Alaska	0.6	0.17	(04)	50.0 ^c	Nebraska	0.5	0.08	(06)	25.0 ^c
Arizona	0.4	0.04	(46)	0.0	Nevada	0.4	0.06	(48)	0.0
Arkansas	0.6	0.07	(01)	50.0 ^c	New Hampshire	0.4	0.08	(44)	0.0
California	0.5	0.02	(23)	25.0 ^c	New Jersey	0.5	0.03	(10)	25.0 ^c
Colorado	0.4	0.04	(42)	0.0	New Mexico	0.4	0.06	(43)	0.0
Connecticut	0.4	0.05	(39)	0.0	New York	0.5	0.02	(17)	25.0 ^c
Delaware	0.5	0.10	(20)	25.0 ^c	North Carolina	0.4	0.03	(37)	0.0
D.C.	0.4	0.12	(40)	0.0	North Dakota	0.4	0.11	(49)	0.0
Florida	0.4	0.02	(41)	0.0	Ohio	0.5	0.03	(11)	25.0 ^c
Georgia	0.4	0.03	(47)	0.0	Oklahoma	0.6	0.06	(05)	50.0 ^c
Hawaii	0.4	0.08	(30)	0.0	Oregon	0.4	0.05	(26)	0.0
Idaho	0.4	0.08	(25)	0.0	Pennsylvania	0.5	0.03	(07)	25.0 ^c
Illinois	0.4	0.03	(27)	0.0	Rhode Island	0.4	0.09	(31)	0.0
Indiana	0.4	0.04	(36)	0.0	South Carolina	0.4	0.04	(35)	0.0
Iowa	0.4	0.05	(38)	0.0	South Dakota	0.5	0.12	(09)	25.0 ^c
Kansas	0.5	0.06	(14)	25.0 ^c	Tennessee	0.4	0.04	(34)	0.0
Kentucky	0.6	0.05	(03)	50.0 ^c	Texas	0.5	0.02	(18)	25.0 ^c
Louisiana	0.5	0.05	(08)	25.0 ^c	Utah	0.3	0.06	(51)	-25.0 ^c
Maine	0.3	0.07	(50)	-25.0 ^c	Vermont	0.4	0.11	(45)	0.0
Maryland	0.4	0.04	(28)	0.0	Virginia	0.4	0.03	(33)	0.0
Massachusetts	0.5	0.04	(19)	25.0 ^c	Washington	0.4	0.04	(32)	0.0
Michigan	0.5	0.03	(15)	25.0 ^c	West Virginia	0.5	0.07	(21)	25.0 ^c
Minnesota	0.4	0.04	(24)	0.0	Wisconsin	0.4	0.04	(29)	0.0
Mississippi	0.6	0.07	(02)	50.0 ^c	Wyoming	0.5	0.14	(12)	25.0 ^c
Missouri	0.5	0.04	(22)	25.0 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table IX-14
HODGKIN LYMPHOMA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.5	0.01		
High Five States				
Kentucky	0.8	0.09	(01)	60.0 ^c
Arkansas	0.8	0.11	(02)	60.0 ^c
Mississippi	0.8	0.11	(03)	60.0 ^c
Pennsylvania	0.6	0.05	(04)	20.0 ^c
Ohio	0.6	0.05	(05)	20.0 ^c
Low Five States				
Georgia	0.4	0.05	(47)	-20.0 ^c
New Mexico	0.4	0.10	(48)	-20.0 ^c
Alaska	0.4	0.15	(49)	-20.0 ^c
Nevada	0.3	0.08	(50)	-40.0 ^c
North Dakota	0.3	0.13	(51)	-40.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.6	0.08	(12)	20.0 ^c	Montana	0.5	0.16	(23)	0.0
Alaska	0.4	0.15	(49)	-20.0 ^c	Nebraska	0.6	0.13	(06)	20.0 ^c
Arizona	0.5	0.06	(37)	0.0	Nevada	0.3	0.08	(50)	-40.0 ^c
Arkansas	0.8	0.11	(02)	60.0 ^c	New Hampshire	0.5	0.15	(22)	0.0
California	0.6	0.03	(19)	20.0 ^c	New Jersey	0.6	0.05	(13)	20.0 ^c
Colorado	0.4	0.07	(45)	-20.0 ^c	New Mexico	0.4	0.10	(48)	-20.0 ^c
Connecticut	0.6	0.09	(07)	20.0 ^c	New York	0.6	0.04	(18)	20.0 ^c
Delaware	0.5	0.17	(35)	0.0	North Carolina	0.5	0.05	(31)	0.0
D.C.	0.5	0.19	(34)	0.0	North Dakota	0.3	0.13	(51)	-40.0 ^c
Florida	0.5	0.03	(36)	0.0	Ohio	0.6	0.05	(05)	20.0 ^c
Georgia	0.4	0.05	(47)	-20.0 ^c	Oklahoma	0.6	0.09	(10)	20.0 ^c
Hawaii	0.5	0.13	(39)	0.0	Oregon	0.5	0.08	(24)	0.0
Idaho	0.6	0.14	(17)	20.0 ^c	Pennsylvania	0.6	0.05	(04)	20.0 ^c
Illinois	0.6	0.05	(15)	20.0 ^c	Rhode Island	0.5	0.14	(41)	0.0
Indiana	0.5	0.06	(30)	0.0	South Carolina	0.5	0.08	(25)	0.0
Iowa	0.5	0.08	(43)	0.0	South Dakota	0.6	0.18	(11)	20.0 ^c
Kansas	0.6	0.10	(09)	20.0 ^c	Tennessee	0.5	0.06	(38)	0.0
Kentucky	0.8	0.09	(01)	60.0 ^c	Texas	0.6	0.04	(20)	20.0 ^c
Louisiana	0.6	0.08	(08)	20.0 ^c	Utah	0.5	0.12	(29)	0.0
Maine	0.5	0.12	(40)	0.0	Vermont	0.4	0.18	(46)	-20.0 ^c
Maryland	0.5	0.07	(33)	0.0	Virginia	0.5	0.06	(32)	0.0
Massachusetts	0.6	0.06	(14)	20.0 ^c	Washington	0.5	0.06	(28)	0.0
Michigan	0.6	0.05	(16)	20.0 ^c	West Virginia	0.5	0.10	(42)	0.0
Minnesota	0.6	0.07	(21)	20.0 ^c	Wisconsin	0.5	0.07	(27)	0.0
Mississippi	0.8	0.11	(03)	60.0 ^c	Wyoming	0.5	0.19	(44)	0.0
Missouri	0.5	0.06	(26)	0.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table IX-15
HODGKIN LYMPHOMA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.4	0.01		
High Five States				
Alaska	0.7	0.28	(01)	75.0 ^c
Oklahoma	0.5	0.07	(02)	25.0 ^c
Mississippi	0.5	0.08	(03)	25.0 ^c
Arkansas	0.5	0.08	(04)	25.0 ^c
Wyoming	0.5	0.19	(05)	25.0 ^c
Low Five States				
D.C.	0.3	0.13	(47)	-25.0 ^c
Arizona	0.3	0.04	(48)	-25.0 ^c
Maine	0.2	0.08	(49)	-50.0 ^c
Connecticut	0.2	0.05	(50)	-50.0 ^c
Utah	0.2	0.06	(51)	-50.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.4	0.06	(19)	0.0	Montana	0.4	0.13	(13)	0.0
Alaska	0.7	0.28	(01)	75.0 ^c	Nebraska	0.5	0.10	(07)	25.0 ^c
Arizona	0.3	0.04	(48)	-25.0 ^c	Nevada	0.3	0.08	(38)	-25.0 ^c
Arkansas	0.5	0.08	(04)	25.0 ^c	New Hampshire	0.3	0.09	(46)	-25.0 ^c
California	0.4	0.02	(31)	0.0	New Jersey	0.5	0.04	(10)	25.0 ^c
Colorado	0.3	0.06	(40)	-25.0 ^c	New Mexico	0.4	0.09	(20)	0.0
Connecticut	0.2	0.05	(50)	-50.0 ^c	New York	0.4	0.03	(17)	0.0
Delaware	0.4	0.14	(11)	0.0	North Carolina	0.3	0.04	(42)	-25.0 ^c
D.C.	0.3	0.13	(47)	-25.0 ^c	North Dakota	0.5	0.18	(08)	25.0 ^c
Florida	0.3	0.02	(45)	-25.0 ^c	Ohio	0.4	0.04	(16)	0.0
Georgia	0.3	0.04	(43)	-25.0 ^c	Oklahoma	0.5	0.07	(02)	25.0 ^c
Hawaii	0.4	0.10	(21)	0.0	Oregon	0.4	0.06	(28)	0.0
Idaho	0.4	0.10	(36)	0.0	Pennsylvania	0.4	0.04	(12)	0.0
Illinois	0.3	0.03	(37)	-25.0 ^c	Rhode Island	0.4	0.11	(23)	0.0
Indiana	0.3	0.04	(41)	-25.0 ^c	South Carolina	0.3	0.05	(44)	-25.0 ^c
Iowa	0.4	0.07	(29)	0.0	South Dakota	0.5	0.16	(09)	25.0 ^c
Kansas	0.4	0.07	(26)	0.0	Tennessee	0.4	0.05	(30)	0.0
Kentucky	0.4	0.06	(18)	0.0	Texas	0.4	0.03	(22)	0.0
Louisiana	0.5	0.06	(06)	25.0 ^c	Utah	0.2	0.06	(51)	-50.0 ^c
Maine	0.2	0.08	(49)	-50.0 ^c	Vermont	0.3	0.14	(39)	-25.0 ^c
Maryland	0.4	0.05	(25)	0.0	Virginia	0.4	0.04	(32)	0.0
Massachusetts	0.4	0.04	(34)	0.0	Washington	0.4	0.05	(35)	0.0
Michigan	0.4	0.04	(15)	0.0	West Virginia	0.4	0.09	(14)	0.0
Minnesota	0.4	0.05	(27)	0.0	Wisconsin	0.4	0.05	(33)	0.0
Mississippi	0.5	0.08	(03)	25.0 ^c	Wyoming	0.5	0.19	(05)	25.0 ^c
Missouri	0.4	0.05	(24)	0.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.


Table XI-13
KIDNEY AND RENAL PELVIS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.2	0.02		
High Five States				
Louisiana	5.4 ^b	0.16	(01)	28.6 ^c
Oklahoma	5.3 ^b	0.17	(02)	26.2 ^c
North Dakota	5.1	0.37	(03)	21.4 ^c
Indiana	5.1 ^b	0.13	(04)	21.4 ^c
Iowa	4.9 ^b	0.17	(05)	16.7 ^c
Low Five States				
Florida	3.5 ^b	0.06	(47)	-16.7 ^c
New York	3.5 ^b	0.06	(48)	-16.7 ^c
Utah	3.2 ^b	0.19	(49)	-23.8 ^c
D.C.	3.1	0.33	(50)	-26.2 ^c
Hawaii	2.4 ^b	0.19	(51)	-42.9 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.9	0.13	(40)	-7.1	Montana	4.5	0.30	(18)	7.1
Alaska	3.9	0.48	(41)	-7.1	Nebraska	4.7	0.23	(10)	11.9
Arizona	4.1	0.12	(32)	-2.4	Nevada	3.9	0.20	(36)	-7.1
Arkansas	4.4	0.17	(21)	4.8	New Hampshire	4.0	0.25	(33)	-4.8
California	3.7 ^b	0.05	(45)	-11.9	New Jersey	4.0	0.09	(35)	-4.8
Colorado	3.9	0.14	(42)	-7.1	New Mexico	4.5	0.22	(17)	7.1
Connecticut	3.7 ^b	0.14	(46)	-11.9	New York	3.5 ^b	0.06	(48)	-16.7 ^c
Delaware	3.8	0.30	(43)	-9.5	North Carolina	4.3	0.10	(24)	2.4
D.C.	3.1	0.33	(50)	-26.2 ^c	North Dakota	5.1	0.37	(03)	21.4 ^c
Florida	3.5 ^b	0.06	(47)	-16.7 ^c	Ohio	4.7 ^b	0.09	(13)	11.9
Georgia	4.1	0.11	(30)	-2.4	Oklahoma	5.3 ^b	0.17	(02)	26.2 ^c
Hawaii	2.4 ^b	0.19	(51)	-42.9 ^c	Oregon	3.9	0.14	(37)	-7.1
Idaho	4.3	0.26	(25)	2.4	Pennsylvania	4.3	0.08	(28)	2.4
Illinois	4.4	0.08	(19)	4.8	Rhode Island	4.7	0.28	(11)	11.9
Indiana	5.1 ^b	0.13	(04)	21.4 ^c	South Carolina	4.2	0.14	(29)	0.0
Iowa	4.9 ^b	0.17	(05)	16.7 ^c	South Dakota	4.8	0.34	(09)	14.3
Kansas	4.6	0.18	(14)	9.5	Tennessee	4.4	0.12	(20)	4.8
Kentucky	4.8 ^b	0.15	(08)	14.3	Texas	4.8 ^b	0.07	(06)	14.3
Louisiana	5.4 ^b	0.16	(01)	28.6 ^c	Utah	3.2 ^b	0.19	(49)	-23.8 ^c
Maine	4.4	0.24	(22)	4.8	Vermont	3.9	0.34	(39)	-7.1
Maryland	3.9	0.12	(38)	-7.1	Virginia	4.0	0.11	(34)	-4.8
Massachusetts	3.8 ^b	0.10	(44)	-9.5	Washington	4.3	0.12	(26)	2.4
Michigan	4.4	0.09	(23)	4.8	West Virginia	4.5	0.20	(16)	7.1
Minnesota	4.3	0.13	(27)	2.4	Wisconsin	4.5	0.13	(15)	7.1
Mississippi	4.7	0.18	(12)	11.9	Wyoming	4.1	0.41	(31)	-2.4
Missouri	4.8 ^b	0.13	(07)	14.3					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XI-14
KIDNEY AND RENAL PELVIS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	6.0	0.03		
High Five States				
Louisiana	7.6 ^b	0.29	(01)	26.7 ^c
Oklahoma	7.4 ^b	0.30	(02)	23.3 ^c
Iowa	7.3 ^b	0.32	(03)	21.7 ^c
North Dakota	7.2	0.67	(04)	20.0 ^c
Indiana	7.2 ^b	0.24	(05)	20.0 ^c
Low Five States				
New York	5.1 ^b	0.11	(47)	-15.0 ^c
Vermont	4.9	0.59	(48)	-18.3 ^c
D.C.	4.5	0.63	(49)	-25.0 ^c
Utah	4.5 ^b	0.34	(50)	-25.0 ^c
Hawaii	3.3 ^b	0.33	(51)	-45.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.8	0.24	(36)	-3.3	Montana	6.7	0.55	(12)	11.7
Alaska	5.9	0.90	(33)	-1.7	Nebraska	6.4	0.40	(19)	6.7
Arizona	5.7	0.21	(39)	-5.0	Nevada	5.6	0.35	(42)	-6.7
Arkansas	6.1	0.31	(29)	1.7	New Hampshire	5.3	0.45	(45)	-11.7
California	5.3 ^b	0.09	(44)	-11.7	New Jersey	5.8	0.18	(34)	-3.3
Colorado	5.8	0.27	(35)	-3.3	New Mexico	6.4	0.40	(20)	6.7
Connecticut	5.5	0.26	(43)	-8.3	New York	5.1 ^b	0.11	(47)	-15.0 ^c
Delaware	5.7	0.57	(41)	-5.0	North Carolina	6.6	0.20	(15)	10.0
D.C.	4.5	0.63	(49)	-25.0 ^c	North Dakota	7.2	0.67	(04)	20.0 ^c
Florida	5.2 ^b	0.10	(46)	-13.3	Ohio	6.8 ^b	0.16	(11)	13.3
Georgia	6.0	0.21	(32)	0.0	Oklahoma	7.4 ^b	0.30	(02)	23.3 ^c
Hawaii	3.3 ^b	0.33	(51)	-45.0 ^c	Oregon	5.7	0.26	(40)	-5.0
Idaho	6.1	0.46	(31)	1.7	Pennsylvania	6.1	0.14	(30)	1.7
Illinois	6.5	0.16	(16)	8.3	Rhode Island	6.8	0.53	(10)	13.3
Indiana	7.2 ^b	0.24	(05)	20.0 ^c	South Carolina	6.2	0.27	(23)	3.3
Iowa	7.3 ^b	0.32	(03)	21.7 ^c	South Dakota	7.1	0.62	(06)	18.3 ^c
Kansas	6.7	0.33	(13)	11.7	Tennessee	6.5	0.23	(17)	8.3
Kentucky	7.0	0.29	(07)	16.7 ^c	Texas	6.9 ^b	0.13	(09)	15.0 ^c
Louisiana	7.6 ^b	0.29	(01)	26.7 ^c	Utah	4.5 ^b	0.34	(50)	-25.0 ^c
Maine	6.2	0.44	(25)	3.3	Vermont	4.9	0.59	(48)	-18.3 ^c
Maryland	5.7	0.23	(38)	-5.0	Virginia	6.1	0.21	(28)	1.7
Massachusetts	5.8	0.20	(37)	-3.3	Washington	6.2	0.22	(24)	3.3
Michigan	6.2	0.17	(22)	3.3	West Virginia	6.2	0.37	(21)	3.3
Minnesota	6.2	0.24	(26)	3.3	Wisconsin	6.5	0.23	(18)	8.3
Mississippi	6.7	0.34	(14)	11.7	Wyoming	6.1	0.74	(27)	1.7
Missouri	6.9 ^b	0.23	(08)	15.0 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XI-15
 KIDNEY AND RENAL PELVIS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.7	0.02		
High Five States				
Louisiana	3.6 ^b	0.17	(01)	33.3 ^c
North Dakota	3.6	0.43	(02)	33.3 ^c
Indiana	3.5 ^b	0.14	(03)	29.6 ^c
Oklahoma	3.5 ^b	0.18	(04)	29.6 ^c
Nebraska	3.4	0.26	(05)	25.9 ^c
Low Five States				
Florida	2.1 ^b	0.06	(47)	-22.2 ^c
Utah	2.1	0.21	(48)	-22.2 ^c
D.C.	2.1	0.35	(49)	-22.2 ^c
Alaska	2.0	0.47	(50)	-25.9 ^c
Hawaii	1.7 ^b	0.21	(51)	-37.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.5	0.13	(38)	-7.4	Montana	2.6	0.31	(34)	-3.7
Alaska	2.0	0.47	(50)	-25.9 ^c	Nebraska	3.4	0.26	(05)	25.9 ^c
Arizona	2.7	0.13	(31)	0.0	Nevada	2.5	0.22	(39)	-7.4
Arkansas	3.0	0.19	(19)	11.1	New Hampshire	3.1	0.29	(10)	14.8
California	2.5 ^b	0.05	(40)	-7.4	New Jersey	2.6	0.10	(33)	-3.7
Colorado	2.4	0.15	(44)	-11.1	New Mexico	2.9	0.24	(24)	7.4
Connecticut	2.4	0.15	(43)	-11.1	New York	2.4 ^b	0.06	(45)	-11.1
Delaware	2.6	0.33	(35)	-3.7	North Carolina	2.7	0.11	(32)	0.0
D.C.	2.1	0.35	(49)	-22.2 ^c	North Dakota	3.6	0.43	(02)	33.3 ^c
Florida	2.1 ^b	0.06	(47)	-22.2 ^c	Ohio	3.1 ^b	0.09	(11)	14.8
Georgia	2.8	0.12	(25)	3.7	Oklahoma	3.5 ^b	0.18	(04)	29.6 ^c
Hawaii	1.7 ^b	0.21	(51)	-37.0 ^c	Oregon	2.5	0.15	(37)	-7.4
Idaho	2.8	0.28	(28)	3.7	Pennsylvania	2.9	0.08	(22)	7.4
Illinois	2.9	0.09	(23)	7.4	Rhode Island	3.3	0.31	(06)	22.2 ^c
Indiana	3.5 ^b	0.14	(03)	29.6 ^c	South Carolina	2.8	0.15	(27)	3.7
Iowa	3.1	0.18	(13)	14.8	South Dakota	2.9	0.36	(20)	7.4
Kansas	3.1	0.20	(15)	14.8	Tennessee	2.8	0.13	(26)	3.7
Kentucky	3.1	0.16	(14)	14.8	Texas	3.2 ^b	0.08	(07)	18.5 ^c
Louisiana	3.6 ^b	0.17	(01)	33.3 ^c	Utah	2.1	0.21	(48)	-22.2 ^c
Maine	3.0	0.27	(17)	11.1	Vermont	3.1	0.41	(16)	14.8
Maryland	2.6	0.13	(36)	-3.7	Virginia	2.4	0.11	(41)	-11.1
Massachusetts	2.4	0.11	(42)	-11.1	Washington	2.7	0.13	(30)	0.0
Michigan	2.9	0.10	(21)	7.4	West Virginia	3.1	0.22	(12)	14.8
Minnesota	2.7	0.14	(29)	0.0	Wisconsin	3.0	0.14	(18)	11.1
Mississippi	3.2	0.20	(09)	18.5 ^c	Wyoming	2.3	0.40	(46)	-14.8
Missouri	3.2	0.14	(08)	18.5 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XII-13
LARYNX CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.3	0.01		
High Five States				
D.C.	2.9 ^b	0.32	(01)	123.1 ^c
West Virginia	1.9 ^b	0.13	(02)	46.2 ^c
Louisiana	1.8 ^b	0.09	(03)	38.5 ^c
Delaware	1.6	0.20	(04)	23.1 ^c
Kentucky	1.6 ^b	0.09	(05)	23.1 ^c
Low Five States				
Colorado	0.8 ^b	0.07	(47)	-38.5 ^c
Nebraska	0.8 ^b	0.09	(48)	-38.5 ^c
Hawaii	0.8 ^b	0.11	(49)	-38.5 ^c
Utah	0.5 ^b	0.08	(50)	-61.5 ^c
Alaska	0.4 ^b	0.15	(51)	-69.2 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.4	0.08	(13)	7.7	Montana	0.9	0.13	(44)	-30.8 ^c
Alaska	0.4 ^b	0.15	(51)	-69.2 ^c	Nebraska	0.8 ^b	0.09	(48)	-38.5 ^c
Arizona	1.0 ^b	0.06	(42)	-23.1 ^c	Nevada	1.0	0.10	(40)	-23.1 ^c
Arkansas	1.2	0.09	(30)	-7.7	New Hampshire	1.0	0.12	(41)	-23.1 ^c
California	1.0 ^b	0.03	(38)	-23.1 ^c	New Jersey	1.3	0.05	(18)	0.0
Colorado	0.8 ^b	0.07	(47)	-38.5 ^c	New Mexico	1.2	0.11	(27)	-7.7
Connecticut	1.2	0.08	(28)	-7.7	New York	1.3	0.04	(19)	0.0
Delaware	1.6	0.20	(04)	23.1 ^c	North Carolina	1.5	0.06	(10)	15.4 ^c
D.C.	2.9 ^b	0.32	(01)	123.1 ^c	North Dakota	1.1	0.18	(33)	-15.4 ^c
Florida	1.4 ^b	0.04	(11)	7.7	Ohio	1.4	0.05	(12)	7.7
Georgia	1.5	0.06	(09)	15.4 ^c	Oklahoma	1.3	0.08	(24)	0.0
Hawaii	0.8 ^b	0.11	(49)	-38.5 ^c	Oregon	1.0	0.07	(39)	-23.1 ^c
Idaho	0.9	0.12	(43)	-30.8 ^c	Pennsylvania	1.4	0.04	(17)	7.7
Illinois	1.3	0.05	(20)	0.0	Rhode Island	1.4	0.15	(16)	7.7
Indiana	1.2	0.06	(26)	-7.7	South Carolina	1.5	0.08	(07)	15.4 ^c
Iowa	1.1	0.08	(31)	-15.4 ^c	South Dakota	0.9	0.15	(45)	-30.8 ^c
Kansas	1.1	0.09	(32)	-15.4 ^c	Tennessee	1.6 ^b	0.07	(06)	23.1 ^c
Kentucky	1.6 ^b	0.09	(05)	23.1 ^c	Texas	1.3	0.04	(21)	0.0
Louisiana	1.8 ^b	0.09	(03)	38.5 ^c	Utah	0.5 ^b	0.08	(50)	-61.5 ^c
Maine	1.1	0.12	(36)	-15.4 ^c	Vermont	1.2	0.18	(29)	-7.7
Maryland	1.5	0.07	(08)	15.4 ^c	Virginia	1.2	0.06	(25)	-7.7
Massachusetts	1.4	0.06	(15)	7.7	Washington	1.1 ^b	0.06	(37)	-15.4 ^c
Michigan	1.4	0.05	(14)	7.7	West Virginia	1.9 ^b	0.13	(02)	46.2 ^c
Minnesota	0.9 ^b	0.06	(46)	-30.8 ^c	Wisconsin	1.1	0.06	(34)	-15.4 ^c
Mississippi	1.3	0.09	(23)	0.0	Wyoming	1.1	0.21	(35)	-15.4 ^c
Missouri	1.3	0.06	(22)	0.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XII-14
LARYNX CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	2.3	0.02		
High Five States				
D.C.	6.0 ^b	0.73	(01)	160.9 ^c
Louisiana	3.4 ^b	0.19	(02)	47.8 ^c
West Virginia	3.1	0.25	(03)	34.8 ^c
Kentucky	3.0 ^b	0.18	(04)	30.4 ^c
Delaware	2.9	0.39	(05)	26.1 ^c
Low Five States				
Hawaii	1.5 ^b	0.22	(47)	-34.8 ^c
Nebraska	1.5 ^b	0.19	(48)	-34.8 ^c
Montana	1.4 ^b	0.25	(49)	-39.1 ^c
Utah	0.8 ^b	0.15	(50)	-65.2 ^c
Alaska	0.8 ^b	0.32	(51)	-65.2 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.6	0.16	(12)	13.0	Montana	1.4 ^b	0.25	(49)	-39.1 ^c
Alaska	0.8 ^b	0.32	(51)	-65.2 ^c	Nebraska	1.5 ^b	0.19	(48)	-34.8 ^c
Arizona	1.8 ^b	0.12	(38)	-21.7 ^c	Nevada	1.7	0.19	(40)	-26.1 ^c
Arkansas	2.2	0.19	(26)	-4.3	New Hampshire	1.6	0.24	(42)	-30.4 ^c
California	1.9 ^b	0.05	(33)	-17.4 ^c	New Jersey	2.5	0.11	(20)	8.7
Colorado	1.5 ^b	0.14	(45)	-34.8 ^c	New Mexico	2.0	0.22	(30)	-13.0
Connecticut	2.2	0.17	(28)	-4.3	New York	2.5	0.08	(18)	8.7
Delaware	2.9	0.39	(05)	26.1 ^c	North Carolina	2.9 ^b	0.13	(08)	26.1 ^c
D.C.	6.0 ^b	0.73	(01)	160.9 ^c	North Dakota	1.5	0.31	(46)	-34.8 ^c
Florida	2.6	0.07	(14)	13.0	Ohio	2.6	0.10	(13)	13.0
Georgia	2.8	0.14	(09)	21.7 ^c	Oklahoma	2.2	0.17	(27)	-4.3
Hawaii	1.5 ^b	0.22	(47)	-34.8 ^c	Oregon	1.7 ^b	0.14	(41)	-26.1 ^c
Idaho	1.5	0.23	(44)	-34.8 ^c	Pennsylvania	2.5	0.09	(15)	8.7
Illinois	2.4	0.10	(21)	4.3	Rhode Island	2.4	0.31	(22)	4.3
Indiana	2.3	0.13	(24)	0.0	South Carolina	2.9	0.18	(07)	26.1 ^c
Iowa	2.1	0.17	(29)	-8.7	South Dakota	1.8	0.31	(37)	-21.7 ^c
Kansas	2.0	0.18	(31)	-13.0	Tennessee	2.9 ^b	0.15	(06)	26.1 ^c
Kentucky	3.0 ^b	0.18	(04)	30.4 ^c	Texas	2.5	0.08	(17)	8.7
Louisiana	3.4 ^b	0.19	(02)	47.8 ^c	Utah	0.8 ^b	0.15	(50)	-65.2 ^c
Maine	1.9	0.24	(34)	-17.4 ^c	Vermont	2.0	0.36	(32)	-13.0
Maryland	2.8	0.16	(10)	21.7 ^c	Virginia	2.3	0.13	(25)	0.0
Massachusetts	2.5	0.13	(16)	8.7	Washington	1.8 ^b	0.12	(36)	-21.7 ^c
Michigan	2.5	0.11	(19)	8.7	West Virginia	3.1	0.25	(03)	34.8 ^c
Minnesota	1.6 ^b	0.12	(43)	-30.4 ^c	Wisconsin	1.8 ^b	0.12	(35)	-21.7 ^c
Mississippi	2.7	0.21	(11)	17.4 ^c	Wyoming	1.7	0.38	(39)	-26.1 ^c
Missouri	2.4	0.13	(23)	4.3					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XII-15
LARYNX CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.5	0.01		
High Five States				
West Virginia	0.9	0.12	(01)	80.0 ^c
North Dakota	0.8	0.21	(02)	60.0 ^c
D.C.	0.8	0.23	(03)	60.0 ^c
Rhode Island	0.7	0.14	(04)	40.0 ^c
Louisiana	0.6	0.07	(05)	20.0 ^c
Low Five States				
Mississippi	0.3	0.06	(47)	-40.0 ^c
Hawaii	0.2	0.08	(48)	-60.0 ^c
Utah	0.2 ^b	0.06	(49)	-60.0 ^c
South Dakota	0.1 ^b	0.07	(50)	-80.0 ^c
Alaska	0.1 ^b	0.08	(51)	-80.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.5	0.06	(14)	0.0	Montana	0.6	0.15	(08)	20.0 ^c
Alaska	0.1 ^b	0.08	(51)	-80.0 ^c	Nebraska	0.3	0.08	(46)	-40.0 ^c
Arizona	0.3	0.05	(45)	-40.0 ^c	Nevada	0.4	0.09	(35)	-20.0 ^c
Arkansas	0.3	0.06	(43)	-40.0 ^c	New Hampshire	0.5	0.12	(25)	0.0
California	0.4 ^b	0.02	(41)	-20.0 ^c	New Jersey	0.5	0.05	(20)	0.0
Colorado	0.3	0.06	(44)	-40.0 ^c	New Mexico	0.5	0.10	(27)	0.0
Connecticut	0.4	0.06	(37)	-20.0 ^c	New York	0.5	0.03	(23)	0.0
Delaware	0.6	0.16	(06)	20.0 ^c	North Carolina	0.5	0.04	(31)	0.0
D.C.	0.8	0.23	(03)	60.0 ^c	North Dakota	0.8	0.21	(02)	60.0 ^c
Florida	0.5	0.03	(12)	0.0	Ohio	0.6	0.04	(10)	20.0 ^c
Georgia	0.5	0.05	(22)	0.0	Oklahoma	0.5	0.07	(19)	0.0
Hawaii	0.2	0.08	(48)	-60.0 ^c	Oregon	0.5	0.07	(21)	0.0
Idaho	0.5	0.12	(15)	0.0	Pennsylvania	0.5	0.04	(17)	0.0
Illinois	0.5	0.04	(24)	0.0	Rhode Island	0.7	0.14	(04)	40.0 ^c
Indiana	0.5	0.05	(29)	0.0	South Carolina	0.5	0.06	(32)	0.0
Iowa	0.4	0.07	(40)	-20.0 ^c	South Dakota	0.1 ^b	0.07	(50)	-80.0 ^c
Kansas	0.5	0.08	(28)	0.0	Tennessee	0.5	0.06	(13)	0.0
Kentucky	0.6	0.07	(11)	20.0 ^c	Texas	0.4	0.03	(38)	-20.0 ^c
Louisiana	0.6	0.07	(05)	20.0 ^c	Utah	0.2 ^b	0.06	(49)	-60.0 ^c
Maine	0.4	0.10	(36)	-20.0 ^c	Vermont	0.4	0.14	(39)	-20.0 ^c
Maryland	0.5	0.06	(18)	0.0	Virginia	0.5	0.05	(26)	0.0
Massachusetts	0.6	0.05	(07)	20.0 ^c	Washington	0.4	0.05	(34)	-20.0 ^c
Michigan	0.6	0.05	(09)	20.0 ^c	West Virginia	0.9	0.12	(01)	80.0 ^c
Minnesota	0.4	0.05	(42)	-20.0 ^c	Wisconsin	0.5	0.05	(33)	0.0
Mississippi	0.3	0.06	(47)	-40.0 ^c	Wyoming	0.5	0.20	(16)	0.0
Missouri	0.5	0.05	(30)	0.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XIII-21
LEUKEMIA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	7.4	0.02		
High Five States				
Wisconsin	8.4 ^b	0.17	(01)	13.5
Iowa	8.4 ^b	0.22	(02)	13.5
Idaho	8.3	0.36	(03)	12.2
Arkansas	8.1	0.23	(04)	9.5
Delaware	8.0	0.44	(05)	8.1
Low Five States				
Arizona	6.6 ^b	0.15	(47)	-10.8
New Mexico	6.5	0.27	(48)	-12.2
Alaska	6.4	0.63	(49)	-13.5
D.C.	6.4	0.47	(50)	-13.5
Hawaii	5.5 ^b	0.29	(51)	-25.7 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.6	0.18	(21)	2.7	Montana	7.5	0.39	(24)	1.4
Alaska	6.4	0.63	(49)	-13.5	Nebraska	7.5	0.28	(25)	1.4
Arizona	6.6 ^b	0.15	(47)	-10.8	Nevada	7.2	0.27	(33)	-2.7
Arkansas	8.1	0.23	(04)	9.5	New Hampshire	7.0	0.33	(40)	-5.4
California	6.9 ^b	0.07	(46)	-6.8	New Jersey	7.3	0.13	(30)	-1.4
Colorado	7.0	0.19	(42)	-5.4	New Mexico	6.5	0.27	(48)	-12.2
Connecticut	7.1	0.19	(37)	-4.1	New York	6.9 ^b	0.08	(44)	-6.8
Delaware	8.0	0.44	(05)	8.1	North Carolina	7.1	0.13	(38)	-4.1
D.C.	6.4	0.47	(50)	-13.5	North Dakota	7.0	0.43	(39)	-5.4
Florida	7.2	0.08	(36)	-2.7	Ohio	7.7	0.11	(18)	4.1
Georgia	7.2	0.14	(34)	-2.7	Oklahoma	7.9	0.21	(13)	6.8
Hawaii	5.5 ^b	0.29	(51)	-25.7 ^c	Oregon	7.4	0.20	(27)	0.0
Idaho	8.3	0.36	(03)	12.2	Pennsylvania	7.5	0.10	(23)	1.4
Illinois	7.9 ^b	0.11	(10)	6.8	Rhode Island	7.0	0.34	(41)	-5.4
Indiana	7.9	0.16	(11)	6.8	South Carolina	7.3	0.19	(31)	-1.4
Iowa	8.4 ^b	0.22	(02)	13.5	South Dakota	8.0	0.43	(07)	8.1
Kansas	7.9	0.23	(12)	6.8	Tennessee	7.7	0.16	(20)	4.1
Kentucky	7.8	0.19	(16)	5.4	Texas	7.4	0.09	(28)	0.0
Louisiana	7.5	0.19	(26)	1.4	Utah	7.3	0.29	(29)	-1.4
Maine	7.8	0.32	(15)	5.4	Vermont	8.0	0.50	(06)	8.1
Maryland	7.3	0.17	(32)	-1.4	Virginia	7.2	0.14	(35)	-2.7
Massachusetts	6.9	0.14	(45)	-6.8	Washington	7.7	0.16	(19)	4.1
Michigan	7.8 ^b	0.12	(14)	5.4	West Virginia	7.6	0.27	(22)	2.7
Minnesota	8.0	0.18	(09)	8.1	Wisconsin	8.4 ^b	0.17	(01)	13.5
Mississippi	7.8	0.24	(17)	5.4	Wyoming	7.0	0.53	(43)	-5.4
Missouri	8.0 ^b	0.16	(08)	8.1					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XIII-22
LEUKEMIA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	9.9	0.04		
High Five States				
Iowa	11.6 ^b	0.40	(01)	17.2 ^c
Maine	11.1	0.61	(02)	12.1
Wisconsin	11.1 ^b	0.30	(03)	12.1
Arkansas	11.0	0.43	(04)	11.1
South Dakota	11.0	0.77	(05)	11.1
Low Five States				
New Mexico	8.6	0.47	(47)	-13.1
Arizona	8.6 ^b	0.26	(48)	-13.1
Alaska	8.4	1.14	(49)	-15.2 ^c
D.C.	8.2	0.87	(50)	-17.2 ^c
Hawaii	7.1 ^b	0.48	(51)	-28.3 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	10.3	0.34	(20)	4.0	Montana	10.1	0.69	(24)	2.0
Alaska	8.4	1.14	(49)	-15.2 ^c	Nebraska	10.5	0.52	(13)	6.1
Arizona	8.6 ^b	0.26	(48)	-13.1	Nevada	9.5	0.47	(38)	-4.0
Arkansas	11.0	0.43	(04)	11.1	New Hampshire	10.1	0.63	(25)	2.0
California	9.1 ^b	0.12	(45)	-8.1	New Jersey	9.6	0.23	(34)	-3.0
Colorado	9.4	0.36	(40)	-5.1	New Mexico	8.6	0.47	(47)	-13.1
Connecticut	9.4	0.35	(41)	-5.1	New York	9.3 ^b	0.15	(43)	-6.1
Delaware	10.5	0.79	(15)	6.1	North Carolina	9.6	0.25	(36)	-3.0
D.C.	8.2	0.87	(50)	-17.2 ^c	North Dakota	10.6	0.82	(11)	7.1
Florida	9.6	0.14	(35)	-3.0	Ohio	10.4	0.21	(16)	5.1
Georgia	9.6	0.27	(37)	-3.0	Oklahoma	10.9	0.38	(08)	10.1
Hawaii	7.1 ^b	0.48	(51)	-28.3 ^c	Oregon	9.9	0.35	(32)	0.0
Idaho	10.0	0.60	(28)	1.0	Pennsylvania	10.3	0.19	(22)	4.0
Illinois	11.0 ^b	0.21	(06)	11.1	Rhode Island	8.8	0.60	(46)	-11.1
Indiana	10.8	0.30	(10)	9.1	South Carolina	10.2	0.36	(23)	3.0
Iowa	11.6 ^b	0.40	(01)	17.2 ^c	South Dakota	11.0	0.77	(05)	11.1
Kansas	10.4	0.42	(17)	5.1	Tennessee	10.5	0.30	(14)	6.1
Kentucky	10.5	0.36	(12)	6.1	Texas	10.0	0.17	(29)	1.0
Louisiana	10.1	0.35	(27)	2.0	Utah	9.9	0.52	(30)	0.0
Maine	11.1	0.61	(02)	12.1	Vermont	10.3	0.89	(21)	4.0
Maryland	9.9	0.31	(31)	0.0	Virginia	9.5	0.27	(39)	-4.0
Massachusetts	9.1	0.26	(44)	-8.1	Washington	10.4	0.29	(19)	5.1
Michigan	10.4	0.23	(18)	5.1	West Virginia	10.1	0.48	(26)	2.0
Minnesota	10.8	0.32	(09)	9.1	Wisconsin	11.1 ^b	0.30	(03)	12.1
Mississippi	9.8	0.42	(33)	-1.0	Wyoming	9.3	0.96	(42)	-6.1
Missouri	10.9	0.30	(07)	10.1					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XIII-23
LEUKEMIA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.6	0.03		
High Five States				
Idaho	6.9	0.44	(01)	23.2 ^c
Delaware	6.5	0.52	(02)	16.1 ^c
Vermont	6.4	0.59	(03)	14.3
Mississippi	6.4	0.28	(04)	14.3
Wisconsin	6.4 ^b	0.20	(05)	14.3
Low Five States				
New Mexico	4.9	0.31	(47)	-12.5
Alaska	4.9	0.73	(48)	-12.5
New Hampshire	4.9	0.37	(49)	-12.5
North Dakota	4.6	0.47	(50)	-17.9 ^c
Hawaii	4.1 ^b	0.34	(51)	-26.8 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.8	0.21	(18)	3.6	Montana	5.6	0.45	(29)	0.0
Alaska	4.9	0.73	(48)	-12.5	Nebraska	5.4	0.32	(36)	-3.6
Arizona	4.9 ^b	0.18	(46)	-12.5	Nevada	5.3	0.32	(40)	-5.4
Arkansas	6.0	0.27	(08)	7.1	New Hampshire	4.9	0.37	(49)	-12.5
California	5.3 ^b	0.08	(42)	-5.4	New Jersey	5.7	0.15	(24)	1.8
Colorado	5.3	0.22	(38)	-5.4	New Mexico	4.9	0.31	(47)	-12.5
Connecticut	5.6	0.23	(27)	0.0	New York	5.3	0.10	(39)	-5.4
Delaware	6.5	0.52	(02)	16.1 ^c	North Carolina	5.4	0.15	(35)	-3.6
D.C.	5.1	0.54	(45)	-8.9	North Dakota	4.6	0.47	(50)	-17.9 ^c
Florida	5.2 ^b	0.10	(43)	-7.1	Ohio	5.8	0.13	(16)	3.6
Georgia	5.7	0.17	(21)	1.8	Oklahoma	5.8	0.24	(19)	3.6
Hawaii	4.1 ^b	0.34	(51)	-26.8 ^c	Oregon	5.6	0.23	(28)	0.0
Idaho	6.9	0.44	(01)	23.2 ^c	Pennsylvania	5.7	0.12	(25)	1.8
Illinois	6.0	0.13	(13)	7.1	Rhode Island	5.7	0.41	(20)	1.8
Indiana	6.1	0.18	(07)	8.9	South Carolina	5.4	0.21	(37)	-3.6
Iowa	6.1	0.25	(06)	8.9	South Dakota	5.7	0.49	(23)	1.8
Kansas	6.0	0.27	(10)	7.1	Tennessee	5.8	0.19	(17)	3.6
Kentucky	5.9	0.22	(14)	5.4	Texas	5.6	0.10	(30)	0.0
Louisiana	5.7	0.21	(22)	1.8	Utah	5.3	0.33	(41)	-5.4
Maine	5.6	0.36	(32)	0.0	Vermont	6.4	0.59	(03)	14.3
Maryland	5.5	0.19	(33)	-1.8	Virginia	5.6	0.17	(31)	0.0
Massachusetts	5.5	0.17	(34)	-1.8	Washington	5.6	0.19	(26)	0.0
Michigan	6.0	0.14	(11)	7.1	West Virginia	5.9	0.31	(15)	5.4
Minnesota	6.0	0.21	(09)	7.1	Wisconsin	6.4 ^b	0.20	(05)	14.3
Mississippi	6.4	0.28	(04)	14.3	Wyoming	5.2	0.61	(44)	-7.1
Missouri	6.0	0.18	(12)	7.1					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XIV-13
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.0	0.02		
High Five States				
Hawaii	7.4 ^b	0.33	(01)	48.0 ^c
Louisiana	6.7 ^b	0.18	(02)	34.0 ^c
Texas	6.6 ^b	0.08	(03)	32.0 ^c
Alaska	6.4	0.61	(04)	28.0 ^c
California	6.2 ^b	0.06	(05)	24.0 ^c
Low Five States				
Montana	3.6 ^b	0.27	(47)	-28.0 ^c
Iowa	3.5 ^b	0.15	(48)	-30.0 ^c
Nebraska	3.5 ^b	0.20	(49)	-30.0 ^c
Vermont	3.5 ^b	0.32	(50)	-30.0 ^c
Utah	3.3 ^b	0.19	(51)	-34.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.2	0.15	(12)	4.0	Montana	3.6 ^b	0.27	(47)	-28.0 ^c
Alaska	6.4	0.61	(04)	28.0 ^c	Nebraska	3.5 ^b	0.20	(49)	-30.0 ^c
Arizona	4.8	0.13	(19)	-4.0	Nevada	5.4	0.23	(10)	8.0
Arkansas	5.5	0.19	(09)	10.0	New Hampshire	4.5	0.26	(30)	-10.0
California	6.2 ^b	0.06	(05)	24.0 ^c	New Jersey	4.9	0.10	(16)	-2.0
Colorado	4.3 ^b	0.15	(33)	-14.0	New Mexico	6.2 ^b	0.26	(06)	24.0 ^c
Connecticut	4.5	0.15	(28)	-10.0	New York	5.0	0.07	(13)	0.0
Delaware	4.5	0.33	(29)	-10.0	North Carolina	4.3 ^b	0.10	(32)	-14.0
D.C.	5.8	0.45	(08)	16.0 ^c	North Dakota	3.7 ^b	0.32	(45)	-26.0 ^c
Florida	4.8	0.07	(18)	-4.0	Ohio	4.3 ^b	0.08	(36)	-14.0
Georgia	4.3 ^b	0.11	(34)	-14.0	Oklahoma	4.3 ^b	0.15	(38)	-14.0
Hawaii	7.4 ^b	0.33	(01)	48.0 ^c	Oregon	4.2 ^b	0.15	(40)	-16.0 ^c
Idaho	3.7 ^b	0.24	(46)	-26.0 ^c	Pennsylvania	4.6 ^b	0.08	(27)	-8.0
Illinois	4.9	0.09	(15)	-2.0	Rhode Island	4.8	0.28	(21)	-4.0
Indiana	4.5 ^b	0.12	(31)	-10.0	South Carolina	4.7	0.15	(24)	-6.0
Iowa	3.5 ^b	0.15	(48)	-30.0 ^c	South Dakota	4.1	0.31	(41)	-18.0 ^c
Kansas	4.0 ^b	0.17	(42)	-20.0 ^c	Tennessee	4.9	0.13	(17)	-2.0
Kentucky	4.8	0.15	(20)	-4.0	Texas	6.6 ^b	0.08	(03)	32.0 ^c
Louisiana	6.7 ^b	0.18	(02)	34.0 ^c	Utah	3.3 ^b	0.19	(51)	-34.0 ^c
Maine	3.9 ^b	0.23	(44)	-22.0 ^c	Vermont	3.5 ^b	0.32	(50)	-30.0 ^c
Maryland	4.7	0.13	(25)	-6.0	Virginia	4.3 ^b	0.11	(37)	-14.0
Massachusetts	5.0	0.12	(14)	0.0	Washington	5.2	0.13	(11)	4.0
Michigan	4.8	0.10	(22)	-4.0	West Virginia	4.3 ^b	0.20	(35)	-14.0
Minnesota	4.3 ^b	0.13	(39)	-14.0	Wisconsin	4.7	0.13	(26)	-6.0
Mississippi	6.1 ^b	0.21	(07)	22.0 ^c	Wyoming	3.9	0.39	(43)	-22.0 ^c
Missouri	4.7	0.12	(23)	-6.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XIV-14
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.3	0.03		
High Five States				
Hawaii	10.6 ^b	0.59	(01)	45.2 ^c
Louisiana	10.0 ^b	0.34	(02)	37.0 ^c
Texas	9.8 ^b	0.16	(03)	34.2 ^c
Mississippi	9.0 ^b	0.39	(04)	23.3 ^c
California	8.9 ^b	0.11	(05)	21.9 ^c
Low Five States				
Montana	5.1 ^b	0.47	(47)	-30.1 ^c
Iowa	5.1 ^b	0.27	(48)	-30.1 ^c
North Dakota	4.8 ^b	0.55	(49)	-34.2 ^c
Wyoming	4.6 ^b	0.63	(50)	-37.0 ^c
Utah	4.4 ^b	0.34	(51)	-39.7 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.5	0.28	(11)	2.7	Montana	5.1 ^b	0.47	(47)	-30.1 ^c
Alaska	8.1	1.02	(08)	11.0	Nebraska	5.1 ^b	0.36	(46)	-30.1 ^c
Arizona	7.0	0.23	(24)	-4.1	Nevada	7.4	0.39	(13)	1.4
Arkansas	7.4	0.34	(12)	1.4	New Hampshire	6.8	0.50	(29)	-6.8
California	8.9 ^b	0.11	(05)	21.9 ^c	New Jersey	7.4	0.20	(14)	1.4
Colorado	5.8 ^b	0.27	(42)	-20.5 ^c	New Mexico	8.6	0.45	(07)	17.8 ^c
Connecticut	6.9	0.29	(26)	-5.5	New York	7.6	0.13	(10)	4.1
Delaware	7.3	0.62	(17)	0.0	North Carolina	6.5 ^b	0.19	(31)	-11.0
D.C.	8.7	0.85	(06)	19.2 ^c	North Dakota	4.8 ^b	0.55	(49)	-34.2 ^c
Florida	7.3	0.12	(16)	0.0	Ohio	6.3 ^b	0.16	(34)	-13.7
Georgia	6.3 ^b	0.21	(35)	-13.7	Oklahoma	6.1 ^b	0.28	(37)	-16.4 ^c
Hawaii	10.6 ^b	0.59	(01)	45.2 ^c	Oregon	5.9 ^b	0.26	(41)	-19.2 ^c
Idaho	5.2 ^b	0.42	(45)	-28.8 ^c	Pennsylvania	6.9	0.15	(27)	-5.5
Illinois	7.2	0.17	(19)	-1.4	Rhode Island	7.2	0.53	(20)	-1.4
Indiana	6.4 ^b	0.22	(33)	-12.3	South Carolina	7.2	0.29	(18)	-1.4
Iowa	5.1 ^b	0.27	(48)	-30.1 ^c	South Dakota	5.9	0.56	(39)	-19.2 ^c
Kansas	5.9 ^b	0.31	(40)	-19.2 ^c	Tennessee	7.0	0.24	(23)	-4.1
Kentucky	6.9	0.28	(28)	-5.5	Texas	9.8 ^b	0.16	(03)	34.2 ^c
Louisiana	10.0 ^b	0.34	(02)	37.0 ^c	Utah	4.4 ^b	0.34	(51)	-39.7 ^c
Maine	5.6 ^b	0.42	(43)	-23.3 ^c	Vermont	5.5	0.62	(44)	-24.7 ^c
Maryland	7.0	0.25	(25)	-4.1	Virginia	6.5 ^b	0.21	(32)	-11.0
Massachusetts	7.9	0.23	(09)	8.2	Washington	7.3	0.24	(15)	0.0
Michigan	6.8	0.18	(30)	-6.8	West Virginia	5.9 ^b	0.36	(38)	-19.2 ^c
Minnesota	6.2 ^b	0.24	(36)	-15.1 ^c	Wisconsin	7.0	0.24	(22)	-4.1
Mississippi	9.0 ^b	0.39	(04)	23.3 ^c	Wyoming	4.6 ^b	0.63	(50)	-37.0 ^c
Missouri	7.2	0.24	(21)	-1.4					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XIV-15
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.1	0.02		
High Five States				
Alaska	4.9	0.73	(01)	58.1 ^c
Hawaii	4.6 ^b	0.35	(02)	48.4 ^c
Louisiana	4.1 ^b	0.18	(03)	32.3 ^c
Texas	4.1 ^b	0.09	(04)	32.3 ^c
New Mexico	4.0	0.28	(05)	29.0 ^c
Low Five States				
Utah	2.3 ^b	0.22	(47)	-25.8 ^c
Iowa	2.3 ^b	0.16	(48)	-25.8 ^c
Delaware	2.2	0.30	(49)	-29.0 ^c
Nebraska	2.1 ^b	0.20	(50)	-32.3 ^c
Vermont	1.8 ^b	0.31	(51)	-41.9 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.4	0.16	(10)	9.7	Montana	2.4	0.29	(46)	-22.6 ^c
Alaska	4.9	0.73	(01)	58.1 ^c	Nebraska	2.1 ^b	0.20	(50)	-32.3 ^c
Arizona	2.9	0.14	(23)	-6.5	Nevada	3.4	0.25	(11)	9.7
Arkansas	3.8	0.21	(07)	22.6 ^c	New Hampshire	2.6	0.27	(40)	-16.1 ^c
California	4.0 ^b	0.07	(06)	29.0 ^c	New Jersey	3.0	0.11	(20)	-3.2
Colorado	3.0	0.17	(18)	-3.2	New Mexico	4.0	0.28	(05)	29.0 ^c
Connecticut	2.7	0.16	(33)	-12.9	New York	3.0	0.07	(19)	-3.2
Delaware	2.2	0.30	(49)	-29.0 ^c	North Carolina	2.6 ^b	0.10	(39)	-16.1 ^c
D.C.	3.4	0.45	(12)	9.7	North Dakota	2.9	0.38	(22)	-6.5
Florida	2.7 ^b	0.07	(32)	-12.9	Ohio	2.7 ^b	0.09	(36)	-12.9
Georgia	2.8	0.12	(29)	-9.7	Oklahoma	2.7	0.16	(35)	-12.9
Hawaii	4.6 ^b	0.35	(02)	48.4 ^c	Oregon	2.8	0.17	(30)	-9.7
Idaho	2.4	0.26	(44)	-22.6 ^c	Pennsylvania	2.8 ^b	0.08	(31)	-9.7
Illinois	3.2	0.09	(15)	3.2	Rhode Island	2.9	0.29	(26)	-6.5
Indiana	2.9	0.13	(24)	-6.5	South Carolina	2.7	0.15	(34)	-12.9
Iowa	2.3 ^b	0.16	(48)	-25.8 ^c	South Dakota	2.5	0.33	(43)	-19.4 ^c
Kansas	2.4 ^b	0.18	(45)	-22.6 ^c	Tennessee	3.1	0.13	(17)	0.0
Kentucky	3.3	0.16	(13)	6.5	Texas	4.1 ^b	0.09	(04)	32.3 ^c
Louisiana	4.1 ^b	0.18	(03)	32.3 ^c	Utah	2.3 ^b	0.22	(47)	-25.8 ^c
Maine	2.6	0.25	(41)	-16.1 ^c	Vermont	1.8 ^b	0.31	(51)	-41.9 ^c
Maryland	2.9	0.14	(25)	-6.5	Virginia	2.5 ^b	0.11	(42)	-19.4 ^c
Massachusetts	2.6 ^b	0.11	(38)	-16.1 ^c	Washington	3.4	0.15	(09)	9.7
Michigan	3.1	0.10	(16)	0.0	West Virginia	3.0	0.22	(21)	-3.2
Minnesota	2.7	0.14	(37)	-12.9	Wisconsin	2.8	0.13	(28)	-9.7
Mississippi	3.8	0.22	(08)	22.6 ^c	Wyoming	3.2	0.49	(14)	3.2
Missouri	2.8	0.13	(27)	-9.7					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XV-19
LUNG AND BRONCHUS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	54.1	0.06		
High Five States				
Kentucky	79.0 ^b	0.61	(01)	46.0 ^c
Arkansas	69.2 ^b	0.68	(02)	27.9 ^c
Tennessee	68.8 ^b	0.48	(03)	27.2 ^c
West Virginia	68.3 ^b	0.79	(04)	26.2 ^c
Mississippi	67.3 ^b	0.69	(05)	24.4 ^c
Low Five States				
Idaho	44.1 ^b	0.84	(47)	-18.5 ^c
Colorado	40.8 ^b	0.47	(48)	-24.6 ^c
New Mexico	38.0 ^b	0.65	(49)	-29.8 ^c
Hawaii	36.8 ^b	0.74	(50)	-32.0 ^c
Utah	24.4 ^b	0.54	(51)	-54.9 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	63.6 ^b	0.52	(09)	17.6 ^c	Montana	53.3	1.03	(31)	-1.5
Alaska	54.7	1.80	(25)	1.1	Nebraska	49.2 ^b	0.74	(38)	-9.1 ^c
Arizona	45.9 ^b	0.40	(44)	-15.2 ^c	Nevada	59.4 ^b	0.77	(16)	9.8 ^c
Arkansas	69.2 ^b	0.68	(02)	27.9 ^c	New Hampshire	54.2	0.92	(26)	0.2
California	44.1 ^b	0.17	(46)	-18.5 ^c	New Jersey	50.3 ^b	0.33	(35)	-7.0 ^c
Colorado	40.8 ^b	0.47	(48)	-24.6 ^c	New Mexico	38.0 ^b	0.65	(49)	-29.8 ^c
Connecticut	49.0 ^b	0.51	(40)	-9.4 ^c	New York	47.1 ^b	0.22	(42)	-12.9 ^c
Delaware	62.2 ^b	1.20	(11)	15.0 ^c	North Carolina	59.5 ^b	0.38	(15)	10.0 ^c
D.C.	52.1	1.35	(34)	-3.7	North Dakota	45.5 ^b	1.13	(45)	-15.9 ^c
Florida	54.2	0.22	(27)	0.2	Ohio	60.5 ^b	0.32	(14)	11.8 ^c
Georgia	58.8 ^b	0.41	(17)	8.7 ^c	Oklahoma	63.6 ^b	0.59	(08)	17.6 ^c
Hawaii	36.8 ^b	0.74	(50)	-32.0 ^c	Oregon	55.1	0.55	(22)	1.8
Idaho	44.1 ^b	0.84	(47)	-18.5 ^c	Pennsylvania	54.1	0.27	(28)	0.0
Illinois	55.3 ^b	0.30	(21)	2.2	Rhode Island	54.9	0.97	(24)	1.5
Indiana	63.9 ^b	0.45	(07)	18.1 ^c	South Carolina	60.6 ^b	0.54	(13)	12.0 ^c
Iowa	52.7	0.56	(33)	-2.6	South Dakota	49.5 ^b	1.08	(37)	-8.5 ^c
Kansas	55.0	0.63	(23)	1.7	Tennessee	68.8 ^b	0.48	(03)	27.2 ^c
Kentucky	79.0 ^b	0.61	(01)	46.0 ^c	Texas	53.0 ^b	0.24	(32)	-2.0
Louisiana	67.0 ^b	0.56	(06)	23.8 ^c	Utah	24.4 ^b	0.54	(51)	-54.9 ^c
Maine	61.4 ^b	0.90	(12)	13.5 ^c	Vermont	49.7 ^b	1.23	(36)	-8.1 ^c
Maryland	56.1 ^b	0.47	(20)	3.7	Virginia	56.8 ^b	0.41	(18)	5.0 ^c
Massachusetts	53.6	0.40	(30)	-0.9	Washington	53.6	0.43	(29)	-0.9
Michigan	56.7 ^b	0.34	(19)	4.8	West Virginia	68.3 ^b	0.79	(04)	26.2 ^c
Minnesota	46.6 ^b	0.44	(43)	-13.9 ^c	Wisconsin	49.1 ^b	0.42	(39)	-9.2 ^c
Mississippi	67.3 ^b	0.69	(05)	24.4 ^c	Wyoming	47.2 ^b	1.38	(41)	-12.8 ^c
Missouri	63.2 ^b	0.46	(10)	16.8 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XV-20
LUNG AND BRONCHUS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	72.0	0.11		
High Five States				
Kentucky	111.5 ^b	1.14	(01)	54.9 ^c
Mississippi	101.3 ^b	1.32	(02)	40.7 ^c
Tennessee	99.9 ^b	0.91	(03)	38.8 ^c
Arkansas	99.0 ^b	1.25	(04)	37.5 ^c
Louisiana	95.9 ^b	1.05	(05)	33.2 ^c
Low Five States				
California	54.8 ^b	0.29	(47)	-23.9 ^c
Colorado	51.0 ^b	0.82	(48)	-29.2 ^c
Hawaii	49.8 ^b	1.28	(49)	-30.8 ^c
New Mexico	48.8 ^b	1.11	(50)	-32.2 ^c
Utah	33.7 ^b	0.95	(51)	-53.2 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	94.8 ^b	0.99	(06)	31.7 ^c	Montana	64.9 ^b	1.70	(36)	-9.9 ^c
Alaska	67.1	3.03	(32)	-6.8 ^c	Nebraska	66.9 ^b	1.31	(33)	-7.1 ^c
Arizona	57.2 ^b	0.67	(45)	-20.6 ^c	Nevada	68.3	1.23	(28)	-5.1 ^c
Arkansas	99.0 ^b	1.25	(04)	37.5 ^c	New Hampshire	67.4	1.58	(30)	-6.4 ^c
California	54.8 ^b	0.29	(47)	-23.9 ^c	New Jersey	64.8 ^b	0.59	(37)	-10.0 ^c
Colorado	51.0 ^b	0.82	(48)	-29.2 ^c	New Mexico	48.8 ^b	1.11	(50)	-32.2 ^c
Connecticut	62.0 ^b	0.89	(40)	-13.9 ^c	New York	61.0 ^b	0.39	(41)	-15.3 ^c
Delaware	80.5 ^b	2.11	(15)	11.8 ^c	North Carolina	85.1 ^b	0.71	(13)	18.2 ^c
D.C.	76.3	2.61	(18)	6.0 ^c	North Dakota	60.5 ^b	1.95	(42)	-16.0 ^c
Florida	69.8 ^b	0.38	(27)	-3.1	Ohio	82.4 ^b	0.57	(14)	14.4 ^c
Georgia	86.0 ^b	0.78	(12)	19.4 ^c	Oklahoma	87.6 ^b	1.06	(09)	21.7 ^c
Hawaii	49.8 ^b	1.28	(49)	-30.8 ^c	Oregon	66.5 ^b	0.91	(34)	-7.6 ^c
Idaho	55.9 ^b	1.41	(46)	-22.4 ^c	Pennsylvania	73.7	0.49	(22)	2.4
Illinois	74.3 ^b	0.54	(20)	3.2	Rhode Island	72.8	1.72	(24)	1.1
Indiana	87.2 ^b	0.83	(10)	21.1 ^c	South Carolina	88.9 ^b	1.02	(08)	23.5 ^c
Iowa	72.6	1.00	(26)	0.8	South Dakota	67.6	1.90	(29)	-6.1 ^c
Kansas	73.7	1.11	(21)	2.4	Tennessee	99.9 ^b	0.91	(03)	38.8 ^c
Kentucky	111.5 ^b	1.14	(01)	54.9 ^c	Texas	72.7	0.44	(25)	1.0
Louisiana	95.9 ^b	1.05	(05)	33.2 ^c	Utah	33.7 ^b	0.95	(51)	-53.2 ^c
Maine	79.5 ^b	1.58	(16)	10.4 ^c	Vermont	64.6	2.16	(38)	-10.3 ^c
Maryland	73.1	0.83	(23)	1.5	Virginia	76.8 ^b	0.74	(17)	6.7 ^c
Massachusetts	67.3 ^b	0.69	(31)	-6.5 ^c	Washington	65.2 ^b	0.73	(35)	-9.4 ^c
Michigan	74.5 ^b	0.59	(19)	3.5	West Virginia	92.8 ^b	1.42	(07)	28.9 ^c
Minnesota	59.5 ^b	0.75	(43)	-17.4 ^c	Wisconsin	64.3 ^b	0.73	(39)	-10.7 ^c
Mississippi	101.3 ^b	1.32	(02)	40.7 ^c	Wyoming	58.8 ^b	2.33	(44)	-18.3 ^c
Missouri	87.2 ^b	0.83	(11)	21.1 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XV-21
LUNG AND BRONCHUS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	41.0	0.07		
High Five States				
Kentucky	55.9 ^b	0.69	(01)	36.3 ^c
Nevada	51.9 ^b	0.97	(02)	26.6 ^c
West Virginia	50.6 ^b	0.91	(03)	23.4 ^c
Delaware	48.7 ^b	1.42	(04)	18.8 ^c
Maine	48.4 ^b	1.07	(05)	18.0 ^c
Low Five States				
North Dakota	34.1 ^b	1.35	(47)	-16.8 ^c
Colorado	33.6 ^b	0.57	(48)	-18.0 ^c
New Mexico	29.7 ^b	0.77	(49)	-27.6 ^c
Hawaii	26.2 ^b	0.84	(50)	-36.1 ^c
Utah	16.9 ^b	0.61	(51)	-58.8 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	41.6	0.55	(26)	1.5	Montana	44.6	1.28	(16)	8.8 ^c
Alaska	44.2	2.20	(18)	7.8 ^c	Nebraska	36.2 ^b	0.85	(43)	-11.7 ^c
Arizona	36.9 ^b	0.49	(41)	-10.0 ^c	Nevada	51.9 ^b	0.97	(02)	26.6 ^c
Arkansas	47.1 ^b	0.75	(07)	14.9 ^c	New Hampshire	44.8	1.12	(15)	9.3 ^c
California	36.3 ^b	0.20	(42)	-11.5 ^c	New Jersey	40.4	0.40	(30)	-1.5
Colorado	33.6 ^b	0.57	(48)	-18.0 ^c	New Mexico	29.7 ^b	0.77	(49)	-27.6 ^c
Connecticut	40.2	0.61	(31)	-2.0	New York	37.6 ^b	0.26	(39)	-8.3 ^c
Delaware	48.7 ^b	1.42	(04)	18.8 ^c	North Carolina	41.3	0.42	(28)	0.7
D.C.	36.1	1.47	(44)	-12.0 ^c	North Dakota	34.1 ^b	1.35	(47)	-16.8 ^c
Florida	41.7	0.26	(25)	1.7	Ohio	45.2 ^b	0.36	(14)	10.2 ^c
Georgia	39.8	0.44	(33)	-2.9	Oklahoma	46.1 ^b	0.67	(11)	12.4 ^c
Hawaii	26.2 ^b	0.84	(50)	-36.1 ^c	Oregon	46.7 ^b	0.68	(08)	13.9 ^c
Idaho	34.9 ^b	1.01	(46)	-14.9 ^c	Pennsylvania	40.5	0.31	(29)	-1.2
Illinois	41.9	0.35	(24)	2.2	Rhode Island	42.7	1.14	(22)	4.1
Indiana	47.4 ^b	0.52	(06)	15.6 ^c	South Carolina	40.1	0.58	(32)	-2.2
Iowa	38.4 ^b	0.65	(37)	-6.3 ^c	South Dakota	35.9 ^b	1.26	(45)	-12.4 ^c
Kansas	41.5	0.74	(27)	1.2	Tennessee	46.7 ^b	0.52	(09)	13.9 ^c
Kentucky	55.9 ^b	0.69	(01)	36.3 ^c	Texas	38.5 ^b	0.27	(35)	-6.1 ^c
Louisiana	46.3 ^b	0.62	(10)	12.9 ^c	Utah	16.9 ^b	0.61	(51)	-58.8 ^c
Maine	48.4 ^b	1.07	(05)	18.0 ^c	Vermont	39.2	1.47	(34)	-4.4
Maryland	44.1 ^b	0.54	(19)	7.6 ^c	Virginia	42.5	0.46	(23)	3.7
Massachusetts	44.5 ^b	0.48	(17)	8.5 ^c	Washington	45.2 ^b	0.54	(13)	10.2 ^c
Michigan	44.1 ^b	0.40	(20)	7.6 ^c	West Virginia	50.6 ^b	0.91	(03)	23.4 ^c
Minnesota	37.3 ^b	0.53	(40)	-9.0 ^c	Wisconsin	38.4 ^b	0.50	(36)	-6.3 ^c
Mississippi	43.2	0.73	(21)	5.4 ^c	Wyoming	38.0	1.69	(38)	-7.3 ^c
Missouri	46.0 ^b	0.52	(12)	12.2 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 5% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XVI-13
MELANOMA OF THE SKIN (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.7	0.01		
High Five States				
Idaho	3.4	0.23	(01)	25.9 ^c
Oklahoma	3.4 ^b	0.14	(02)	25.9 ^c
Kentucky	3.4 ^b	0.13	(03)	25.9 ^c
Utah	3.4	0.20	(04)	25.9 ^c
Delaware	3.3	0.28	(05)	22.2 ^c
Low Five States				
Mississippi	2.2 ^b	0.12	(47)	-18.5 ^c
Alaska	1.9	0.30	(48)	-29.6 ^c
Hawaii	1.7 ^b	0.16	(49)	-37.0 ^c
North Dakota	1.6 ^b	0.21	(50)	-40.7 ^c
D.C.	1.4 ^b	0.22	(51)	-48.1 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.7	0.11	(29)	0.0	Montana	3.0	0.25	(12)	11.1
Alaska	1.9	0.30	(48)	-29.6 ^c	Nebraska	2.9	0.18	(19)	7.4
Arizona	2.8	0.10	(28)	3.7	Nevada	3.1	0.17	(10)	14.8
Arkansas	2.5	0.13	(39)	-7.4	New Hampshire	3.2	0.22	(07)	18.5 ^c
California	2.6	0.04	(36)	-3.7	New Jersey	2.7	0.08	(31)	0.0
Colorado	2.9	0.12	(23)	7.4	New Mexico	2.3	0.16	(42)	-14.8
Connecticut	2.9	0.12	(17)	7.4	New York	2.2 ^b	0.05	(46)	-18.5 ^c
Delaware	3.3	0.28	(05)	22.2 ^c	North Carolina	2.9	0.08	(22)	7.4
D.C.	1.4 ^b	0.22	(51)	-48.1 ^c	North Dakota	1.6 ^b	0.21	(50)	-40.7 ^c
Florida	2.8 ^b	0.05	(24)	3.7	Ohio	2.6	0.07	(32)	-3.7
Georgia	2.6	0.08	(34)	-3.7	Oklahoma	3.4 ^b	0.14	(02)	25.9 ^c
Hawaii	1.7 ^b	0.16	(49)	-37.0 ^c	Oregon	3.2 ^b	0.13	(06)	18.5 ^c
Idaho	3.4	0.23	(01)	25.9 ^c	Pennsylvania	2.9	0.06	(21)	7.4
Illinois	2.4 ^b	0.06	(40)	-11.1	Rhode Island	3.1	0.23	(08)	14.8
Indiana	2.8	0.09	(25)	3.7	South Carolina	2.5	0.11	(38)	-7.4
Iowa	2.4	0.12	(41)	-11.1	South Dakota	2.5	0.25	(37)	-7.4
Kansas	2.8	0.14	(27)	3.7	Tennessee	2.8	0.10	(26)	3.7
Kentucky	3.4 ^b	0.13	(03)	25.9 ^c	Texas	2.6	0.05	(35)	-3.7
Louisiana	2.2 ^b	0.10	(45)	-18.5 ^c	Utah	3.4	0.20	(04)	25.9 ^c
Maine	3.0	0.20	(14)	11.1	Vermont	3.0	0.30	(13)	11.1
Maryland	2.7	0.10	(30)	0.0	Virginia	2.9	0.09	(20)	7.4
Massachusetts	2.9	0.09	(15)	7.4	Washington	2.9	0.10	(16)	7.4
Michigan	2.3 ^b	0.07	(43)	-14.8	West Virginia	3.1	0.17	(09)	14.8
Minnesota	2.3 ^b	0.10	(44)	-14.8	Wisconsin	2.6	0.09	(33)	-3.7
Mississippi	2.2 ^b	0.12	(47)	-18.5 ^c	Wyoming	2.9	0.34	(18)	7.4
Missouri	3.1 ^b	0.10	(11)	14.8					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XVI-14
MELANOMA OF THE SKIN (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	3.9	0.03		
High Five States				
Delaware	5.2	0.54	(01)	33.3 ^c
Kentucky	5.0 ^b	0.24	(02)	28.2 ^c
Oklahoma	5.0 ^b	0.25	(03)	28.2 ^c
Idaho	5.0	0.41	(04)	28.2 ^c
Rhode Island	4.7	0.44	(05)	20.5 ^c
Low Five States				
New York	3.1 ^b	0.09	(47)	-20.5 ^c
Hawaii	2.4 ^b	0.28	(48)	-38.5 ^c
D.C.	2.4	0.48	(49)	-38.5 ^c
Alaska	2.3 ^b	0.45	(50)	-41.0 ^c
North Dakota	2.0 ^b	0.35	(51)	-48.7 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.9	0.20	(29)	0.0	Montana	3.7	0.40	(35)	-5.1
Alaska	2.3 ^b	0.45	(50)	-41.0 ^c	Nebraska	4.5	0.33	(11)	15.4 ^c
Arizona	4.3	0.18	(16)	10.3	Nevada	4.5	0.31	(10)	15.4 ^c
Arkansas	3.8	0.25	(32)	-2.6	New Hampshire	4.7	0.41	(06)	20.5 ^c
California	3.9	0.08	(30)	0.0	New Jersey	4.0	0.14	(27)	2.6
Colorado	4.1	0.23	(21)	5.1	New Mexico	3.2	0.28	(45)	-17.9 ^c
Connecticut	4.4	0.24	(13)	12.8	New York	3.1 ^b	0.09	(47)	-20.5 ^c
Delaware	5.2	0.54	(01)	33.3 ^c	North Carolina	4.1	0.16	(22)	5.1
D.C.	2.4	0.48	(49)	-38.5 ^c	North Dakota	2.0 ^b	0.35	(51)	-48.7 ^c
Florida	4.2	0.09	(18)	7.7	Ohio	3.7	0.12	(33)	-5.1
Georgia	3.7	0.16	(36)	-5.1	Oklahoma	5.0 ^b	0.25	(03)	28.2 ^c
Hawaii	2.4 ^b	0.28	(48)	-38.5 ^c	Oregon	4.7	0.24	(07)	20.5 ^c
Idaho	5.0	0.41	(04)	28.2 ^c	Pennsylvania	4.1	0.12	(25)	5.1
Illinois	3.5 ^b	0.12	(40)	-10.3	Rhode Island	4.7	0.44	(05)	20.5 ^c
Indiana	3.9	0.17	(28)	0.0	South Carolina	3.7	0.21	(34)	-5.1
Iowa	3.4	0.21	(42)	-12.8	South Dakota	3.2	0.41	(44)	-17.9 ^c
Kansas	4.1	0.26	(24)	5.1	Tennessee	4.2	0.19	(19)	7.7
Kentucky	5.0 ^b	0.24	(02)	28.2 ^c	Texas	3.8	0.10	(31)	-2.6
Louisiana	3.3	0.19	(43)	-15.4 ^c	Utah	4.7	0.35	(08)	20.5 ^c
Maine	4.4	0.37	(14)	12.8	Vermont	4.6	0.57	(09)	17.9 ^c
Maryland	4.1	0.20	(26)	5.1	Virginia	4.1	0.17	(23)	5.1
Massachusetts	4.3	0.17	(15)	10.3	Washington	4.2	0.18	(20)	7.7
Michigan	3.6	0.13	(39)	-7.7	West Virginia	4.4	0.31	(12)	12.8
Minnesota	3.2 ^b	0.17	(46)	-17.9 ^c	Wisconsin	3.7	0.17	(37)	-5.1
Mississippi	3.4	0.25	(41)	-12.8	Wyoming	3.7	0.56	(38)	-5.1
Missouri	4.3	0.18	(17)	10.3					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XVI-15
MELANOMA OF THE SKIN (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.7	0.01		
High Five States				
Montana	2.5	0.31	(01)	47.1 ^c
Utah	2.2	0.22	(02)	29.4 ^c
Wyoming	2.2	0.41	(03)	29.4 ^c
Kentucky	2.1	0.14	(04)	23.5 ^c
Missouri	2.1	0.11	(05)	23.5 ^c
Low Five States				
Mississippi	1.3	0.13	(47)	-23.5 ^c
Alaska	1.3	0.35	(48)	-23.5 ^c
North Dakota	1.2	0.26	(49)	-29.4 ^c
Hawaii	1.1	0.18	(50)	-35.3 ^c
D.C.	1.0	0.23	(51)	-41.2 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.9	0.12	(20)	11.8	Montana	2.5	0.31	(01)	47.1 ^c
Alaska	1.3	0.35	(48)	-23.5 ^c	Nebraska	1.7	0.19	(32)	0.0
Arizona	1.5	0.10	(44)	-11.8	Nevada	1.8	0.18	(28)	5.9
Arkansas	1.5	0.14	(43)	-11.8	New Hampshire	2.0	0.24	(14)	17.6 ^c
California	1.6 ^b	0.04	(42)	-5.9	New Jersey	1.8	0.08	(27)	5.9
Colorado	1.9	0.13	(19)	11.8	New Mexico	1.7	0.18	(35)	0.0
Connecticut	1.8	0.13	(24)	5.9	New York	1.7	0.05	(37)	0.0
Delaware	1.9	0.28	(21)	11.8	North Carolina	2.0	0.09	(15)	17.6 ^c
D.C.	1.0	0.23	(51)	-41.2 ^c	North Dakota	1.2	0.26	(49)	-29.4 ^c
Florida	1.8	0.06	(29)	5.9	Ohio	1.7	0.07	(31)	0.0
Georgia	1.7	0.09	(30)	0.0	Oklahoma	2.1	0.15	(06)	23.5 ^c
Hawaii	1.1	0.18	(50)	-35.3 ^c	Oregon	1.9	0.14	(17)	11.8
Idaho	2.0	0.24	(10)	17.6 ^c	Pennsylvania	2.0 ^b	0.07	(12)	17.6 ^c
Illinois	1.7	0.07	(36)	0.0	Rhode Island	2.0	0.25	(07)	17.6 ^c
Indiana	2.0	0.11	(16)	17.6 ^c	South Carolina	1.6	0.12	(39)	-5.9
Iowa	1.6	0.14	(41)	-5.9	South Dakota	2.0	0.31	(11)	17.6 ^c
Kansas	1.8	0.15	(26)	5.9	Tennessee	1.8	0.10	(25)	5.9
Kentucky	2.1	0.14	(04)	23.5 ^c	Texas	1.6	0.06	(40)	-5.9
Louisiana	1.4	0.11	(46)	-17.6 ^c	Utah	2.2	0.22	(02)	29.4 ^c
Maine	1.9	0.21	(22)	11.8	Vermont	1.8	0.32	(23)	5.9
Maryland	1.7	0.11	(33)	0.0	Virginia	2.0	0.10	(09)	17.6 ^c
Massachusetts	2.0	0.10	(08)	17.6 ^c	Washington	1.9	0.11	(18)	11.8
Michigan	1.4 ^b	0.07	(45)	-17.6 ^c	West Virginia	2.0	0.19	(13)	17.6 ^c
Minnesota	1.6	0.11	(38)	-5.9	Wisconsin	1.7	0.10	(34)	0.0
Mississippi	1.3	0.13	(47)	-23.5 ^c	Wyoming	2.2	0.41	(03)	29.4 ^c
Missouri	2.1	0.11	(05)	23.5 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XVIII-13
MYELOMA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.7	0.02		
High Five States				
D.C.	5.1	0.43	(01)	37.8 ^c
Alabama	4.5 ^b	0.14	(02)	21.6 ^c
South Carolina	4.4 ^b	0.15	(03)	18.9 ^c
North Carolina	4.2 ^b	0.10	(04)	13.5
Tennessee	4.2 ^b	0.12	(05)	13.5
Low Five States				
New York	3.3 ^b	0.06	(47)	-10.8
Vermont	3.2	0.32	(48)	-13.5
Arizona	3.2 ^b	0.11	(49)	-13.5
Florida	3.1 ^b	0.05	(50)	-16.2 ^c
Hawaii	2.3 ^b	0.18	(51)	-37.8 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.5 ^b	0.14	(02)	21.6 ^c	Montana	4.0	0.28	(11)	8.1
Alaska	3.6	0.47	(34)	-2.7	Nebraska	3.4	0.19	(42)	-8.1
Arizona	3.2 ^b	0.11	(49)	-13.5	Nevada	3.3	0.18	(46)	-10.8
Arkansas	3.8	0.16	(25)	2.7	New Hampshire	3.8	0.24	(24)	2.7
California	3.4 ^b	0.05	(43)	-8.1	New Jersey	3.7	0.09	(30)	0.0
Colorado	3.5	0.14	(41)	-5.4	New Mexico	3.5	0.20	(38)	-5.4
Connecticut	3.6	0.14	(36)	-2.7	New York	3.3 ^b	0.06	(47)	-10.8
Delaware	3.5	0.29	(39)	-5.4	North Carolina	4.2 ^b	0.10	(04)	13.5
D.C.	5.1	0.43	(01)	37.8 ^c	North Dakota	4.0	0.33	(15)	8.1
Florida	3.1 ^b	0.05	(50)	-16.2 ^c	Ohio	3.9	0.08	(19)	5.4
Georgia	4.1	0.11	(10)	10.8	Oklahoma	3.4	0.13	(45)	-8.1
Hawaii	2.3 ^b	0.18	(51)	-37.8 ^c	Oregon	3.9	0.15	(18)	5.4
Idaho	3.6	0.24	(35)	-2.7	Pennsylvania	3.6	0.07	(33)	-2.7
Illinois	3.8	0.08	(23)	2.7	Rhode Island	3.4	0.24	(44)	-8.1
Indiana	3.9	0.11	(22)	5.4	South Carolina	4.4 ^b	0.15	(03)	18.9 ^c
Iowa	3.7	0.15	(28)	0.0	South Dakota	4.2	0.31	(07)	13.5
Kansas	4.0	0.17	(12)	8.1	Tennessee	4.2 ^b	0.12	(05)	13.5
Kentucky	4.0	0.14	(14)	8.1	Texas	3.7	0.06	(31)	0.0
Louisiana	4.2 ^b	0.14	(06)	13.5	Utah	3.7	0.21	(27)	0.0
Maine	3.9	0.23	(20)	5.4	Vermont	3.2	0.32	(48)	-13.5
Maryland	4.1	0.13	(08)	10.8	Virginia	4.0	0.11	(13)	8.1
Massachusetts	3.6	0.10	(37)	-2.7	Washington	3.9	0.12	(17)	5.4
Michigan	4.1 ^b	0.09	(09)	10.8	West Virginia	3.6	0.18	(32)	-2.7
Minnesota	3.8	0.12	(26)	2.7	Wisconsin	3.5	0.11	(40)	-5.4
Mississippi	4.0	0.17	(16)	8.1	Wyoming	3.7	0.39	(29)	0.0
Missouri	3.9	0.11	(21)	5.4					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XVIII-14
MYELOMA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.6	0.03		
High Five States				
D.C.	6.0	0.75	(01)	30.4 ^c
Louisiana	5.5 ^b	0.26	(02)	19.6 ^c
Alabama	5.4	0.24	(03)	17.4 ^c
Tennessee	5.3	0.21	(04)	15.2 ^c
Michigan	5.3 ^b	0.16	(05)	15.2 ^c
Low Five States				
Arizona	4.1	0.18	(47)	-10.9
New York	4.1 ^b	0.10	(48)	-10.9
Florida	4.0 ^b	0.09	(49)	-13.0
Alaska	3.5	0.68	(50)	-23.9 ^c
Hawaii	2.8 ^b	0.30	(51)	-39.1 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.4	0.24	(03)	17.4 ^c	Montana	5.2	0.49	(08)	13.0
Alaska	3.5	0.68	(50)	-23.9 ^c	Nebraska	4.4	0.34	(40)	-4.3
Arizona	4.1	0.18	(47)	-10.9	Nevada	4.1	0.31	(46)	-10.9
Arkansas	4.8	0.28	(23)	4.3	New Hampshire	4.7	0.42	(26)	2.2
California	4.2 ^b	0.08	(44)	-8.7	New Jersey	4.5	0.16	(34)	-2.2
Colorado	4.3	0.24	(41)	-6.5	New Mexico	4.5	0.34	(30)	-2.2
Connecticut	4.2	0.23	(45)	-8.7	New York	4.1 ^b	0.10	(48)	-10.9
Delaware	4.4	0.50	(37)	-4.3	North Carolina	5.1	0.18	(11)	10.9
D.C.	6.0	0.75	(01)	30.4 ^c	North Dakota	5.0	0.56	(15)	8.7
Florida	4.0 ^b	0.09	(49)	-13.0	Ohio	4.6	0.14	(27)	0.0
Georgia	4.9	0.19	(20)	6.5	Oklahoma	4.3	0.24	(43)	-6.5
Hawaii	2.8 ^b	0.30	(51)	-39.1 ^c	Oregon	5.1	0.25	(12)	10.9
Idaho	4.6	0.41	(29)	0.0	Pennsylvania	4.4	0.12	(36)	-4.3
Illinois	4.5	0.14	(31)	-2.2	Rhode Island	4.3	0.42	(42)	-6.5
Indiana	5.1	0.20	(13)	10.9	South Carolina	5.3	0.26	(06)	15.2 ^c
Iowa	4.7	0.26	(25)	2.2	South Dakota	5.1	0.53	(09)	10.9
Kansas	5.0	0.29	(14)	8.7	Tennessee	5.3	0.21	(04)	15.2 ^c
Kentucky	5.3	0.25	(07)	15.2 ^c	Texas	4.5	0.11	(35)	-2.2
Louisiana	5.5 ^b	0.26	(02)	19.6 ^c	Utah	5.0	0.37	(19)	8.7
Maine	5.0	0.40	(17)	8.7	Vermont	4.5	0.59	(32)	-2.2
Maryland	5.0	0.22	(18)	8.7	Virginia	5.0	0.19	(16)	8.7
Massachusetts	4.4	0.18	(38)	-4.3	Washington	5.1	0.21	(10)	10.9
Michigan	5.3 ^b	0.16	(05)	15.2 ^c	West Virginia	4.9	0.34	(21)	6.5
Minnesota	4.9	0.22	(22)	6.5	Wisconsin	4.4	0.19	(39)	-4.3
Mississippi	4.6	0.29	(28)	0.0	Wyoming	4.5	0.66	(33)	-2.2
Missouri	4.8	0.20	(24)	4.3					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XVIII-15
MYELOMA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.0	0.02		
High Five States				
D.C.	4.6	0.52	(01)	53.3 ^c
Alabama	3.8 ^b	0.17	(02)	26.7 ^c
South Carolina	3.8 ^b	0.18	(03)	26.7 ^c
North Carolina	3.6 ^b	0.12	(04)	20.0 ^c
Alaska	3.6	0.63	(05)	20.0 ^c
Low Five States				
Nevada	2.6	0.22	(47)	-13.3
Vermont	2.5	0.36	(48)	-16.7 ^c
Arizona	2.4 ^b	0.12	(49)	-20.0 ^c
Florida	2.4 ^b	0.06	(50)	-20.0 ^c
Hawaii	2.0 ^b	0.23	(51)	-33.3 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.8 ^b	0.17	(02)	26.7 ^c	Montana	3.1	0.33	(23)	3.3
Alaska	3.6	0.63	(05)	20.0 ^c	Nebraska	2.7	0.23	(45)	-10.0
Arizona	2.4 ^b	0.12	(49)	-20.0 ^c	Nevada	2.6	0.22	(47)	-13.3
Arkansas	3.0	0.19	(31)	0.0	New Hampshire	3.1	0.29	(22)	3.3
California	2.8 ^b	0.06	(42)	-6.7	New Jersey	3.2	0.11	(19)	6.7
Colorado	2.9	0.17	(35)	-3.3	New Mexico	2.8	0.23	(43)	-6.7
Connecticut	3.2	0.17	(17)	6.7	New York	2.7 ^b	0.07	(46)	-10.0
Delaware	2.9	0.34	(38)	-3.3	North Carolina	3.6 ^b	0.12	(04)	20.0 ^c
D.C.	4.6	0.52	(01)	53.3 ^c	North Dakota	3.1	0.39	(25)	3.3
Florida	2.4 ^b	0.06	(50)	-20.0 ^c	Ohio	3.4 ^b	0.10	(10)	13.3
Georgia	3.5	0.13	(09)	16.7 ^c	Oklahoma	2.8	0.16	(44)	-6.7
Hawaii	2.0 ^b	0.23	(51)	-33.3 ^c	Oregon	3.0	0.17	(29)	0.0
Idaho	2.9	0.29	(37)	-3.3	Pennsylvania	3.1	0.08	(27)	3.3
Illinois	3.3	0.10	(14)	10.0	Rhode Island	2.8	0.28	(39)	-6.7
Indiana	3.0	0.13	(32)	0.0	South Carolina	3.8 ^b	0.18	(03)	26.7 ^c
Iowa	3.0	0.18	(30)	0.0	South Dakota	3.4	0.37	(12)	13.3
Kansas	3.4	0.20	(11)	13.3	Tennessee	3.5	0.14	(08)	16.7 ^c
Kentucky	3.1	0.16	(21)	3.3	Texas	3.1	0.08	(20)	3.3
Louisiana	3.3	0.16	(15)	10.0	Utah	2.8	0.25	(41)	-6.7
Maine	3.1	0.26	(26)	3.3	Vermont	2.5	0.36	(48)	-16.7 ^c
Maryland	3.5	0.15	(06)	16.7 ^c	Virginia	3.3	0.13	(13)	10.0
Massachusetts	3.0	0.12	(34)	0.0	Washington	3.1	0.14	(28)	3.3
Michigan	3.2	0.10	(18)	6.7	West Virginia	2.8	0.21	(40)	-6.7
Minnesota	3.0	0.15	(33)	0.0	Wisconsin	2.9	0.13	(36)	-3.3
Mississippi	3.5	0.21	(07)	16.7 ^c	Wyoming	3.1	0.49	(24)	3.3
Missouri	3.2	0.14	(16)	6.7					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XIX-17
NON-HODGKIN LYMPHOMA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	7.3	0.02		
High Five States				
Oregon	8.4 ^b	0.21	(01)	15.1 ^c
Michigan	8.3 ^b	0.13	(02)	13.7
Indiana	8.2 ^b	0.16	(03)	12.3
Kansas	8.1	0.24	(04)	11.0
West Virginia	8.0	0.27	(05)	9.6
Low Five States				
Nevada	6.4 ^b	0.26	(47)	-12.3
Alaska	6.4	0.62	(48)	-12.3
D.C.	6.4	0.47	(49)	-12.3
New Mexico	6.3 ^b	0.26	(50)	-13.7
Hawaii	5.8 ^b	0.29	(51)	-20.5 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	7.3	0.18	(28)	0.0	Montana	7.6	0.39	(23)	4.1
Alaska	6.4	0.62	(48)	-12.3	Nebraska	7.4	0.28	(27)	1.4
Arizona	6.9	0.16	(40)	-5.5	Nevada	6.4 ^b	0.26	(47)	-12.3
Arkansas	7.3	0.22	(32)	0.0	New Hampshire	7.7	0.35	(15)	5.5
California	6.9 ^b	0.07	(42)	-5.5	New Jersey	7.6	0.13	(20)	4.1
Colorado	6.9	0.19	(41)	-5.5	New Mexico	6.3 ^b	0.26	(50)	-13.7
Connecticut	7.3	0.20	(29)	0.0	New York	6.7 ^b	0.08	(44)	-8.2
Delaware	7.6	0.42	(24)	4.1	North Carolina	7.1	0.13	(34)	-2.7
D.C.	6.4	0.47	(49)	-12.3	North Dakota	7.2	0.44	(33)	-1.4
Florida	7.0 ^b	0.08	(38)	-4.1	Ohio	7.9 ^b	0.11	(11)	8.2
Georgia	6.6 ^b	0.14	(46)	-9.6	Oklahoma	7.6	0.20	(22)	4.1
Hawaii	5.8 ^b	0.29	(51)	-20.5 ^c	Oregon	8.4 ^b	0.21	(01)	15.1 ^c
Idaho	7.5	0.34	(26)	2.7	Pennsylvania	8.0 ^b	0.10	(06)	9.6
Illinois	7.6	0.11	(19)	4.1	Rhode Island	7.5	0.35	(25)	2.7
Indiana	8.2 ^b	0.16	(03)	12.3	South Carolina	6.7	0.18	(43)	-8.2
Iowa	8.0	0.22	(07)	9.6	South Dakota	7.3	0.41	(31)	0.0
Kansas	8.1	0.24	(04)	11.0	Tennessee	7.9	0.16	(10)	8.2
Kentucky	7.7	0.19	(14)	5.5	Texas	7.1	0.09	(35)	-2.7
Louisiana	7.8	0.19	(13)	6.8	Utah	7.0	0.29	(36)	-4.1
Maine	7.8	0.32	(12)	6.8	Vermont	7.3	0.47	(30)	0.0
Maryland	7.0	0.16	(37)	-4.1	Virginia	6.9	0.14	(39)	-5.5
Massachusetts	7.6	0.15	(18)	4.1	Washington	7.9	0.17	(09)	8.2
Michigan	8.3 ^b	0.13	(02)	13.7	West Virginia	8.0	0.27	(05)	9.6
Minnesota	7.9	0.18	(08)	8.2	Wisconsin	7.6	0.16	(21)	4.1
Mississippi	6.7	0.22	(45)	-8.2	Wyoming	7.7	0.56	(17)	5.5
Missouri	7.7	0.16	(16)	5.5					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XIX-18
NON-HODGKIN LYMPHOMA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	9.3	0.04		
HHigh Five States				
Michigan	10.5 ^b	0.23	(01)	12.9
West Virginia	10.4	0.48	(02)	11.8
Kansas	10.4	0.42	(03)	11.8
Oregon	10.3	0.36	(04)	10.8
Indiana	10.3 ^b	0.29	(05)	10.8
LLow Five States				
South Carolina	8.2	0.32	(47)	-11.8
Wyoming	7.9	0.88	(48)	-15.1 ^c
New Mexico	7.8	0.45	(49)	-16.1 ^c
Nevada	7.6 ^b	0.42	(50)	-18.3 ^c
Hawaii	7.4 ^b	0.49	(51)	-20.4 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.1	0.31	(33)	-2.2	Montana	9.8	0.67	(17)	5.4
Alaska	8.3	1.10	(46)	-10.8	Nebraska	9.1	0.49	(32)	-2.2
Arizona	8.5	0.26	(44)	-8.6	Nevada	7.6 ^b	0.42	(50)	-18.3 ^c
Arkansas	9.7	0.40	(19)	4.3	New Hampshire	9.6	0.61	(22)	3.2
California	8.8 ^b	0.12	(39)	-5.4	New Jersey	9.7	0.23	(20)	4.3
Colorado	8.7	0.34	(41)	-6.5	New Mexico	7.8	0.45	(49)	-16.1 ^c
Connecticut	9.4	0.35	(26)	1.1	New York	8.6 ^b	0.15	(42)	-7.5
Delaware	9.9	0.76	(12)	6.5	North Carolina	9.0	0.24	(34)	-3.2
D.C.	9.1	0.91	(30)	-2.2	North Dakota	9.1	0.76	(31)	-2.2
Florida	8.8	0.14	(36)	-5.4	Ohio	9.9	0.20	(11)	6.5
Georgia	8.3 ^b	0.25	(45)	-10.8	Oklahoma	9.8	0.36	(15)	5.4
Hawaii	7.4 ^b	0.49	(51)	-20.4 ^c	Oregon	10.3	0.36	(04)	10.8
Idaho	9.2	0.57	(28)	-1.1	Pennsylvania	10.2 ^b	0.18	(07)	9.7
Illinois	9.6	0.20	(23)	3.2	Rhode Island	9.3	0.62	(27)	0.0
Indiana	10.3 ^b	0.29	(05)	10.8	South Carolina	8.2	0.32	(47)	-11.8
Iowa	10.1	0.38	(08)	8.6	South Dakota	8.8	0.69	(38)	-5.4
Kansas	10.4	0.42	(03)	11.8	Tennessee	10.1	0.30	(10)	8.6
Kentucky	9.9	0.35	(13)	6.5	Texas	8.8	0.16	(37)	-5.4
Louisiana	9.8	0.34	(16)	5.4	Utah	9.2	0.50	(29)	-1.1
Maine	9.6	0.56	(24)	3.2	Vermont	9.9	0.87	(14)	6.5
Maryland	8.9	0.30	(35)	-4.3	Virginia	8.7	0.25	(40)	-6.5
Massachusetts	9.5	0.26	(25)	2.2	Washington	10.3	0.29	(06)	10.8
Michigan	10.5 ^b	0.23	(01)	12.9	West Virginia	10.4	0.48	(02)	11.8
Minnesota	10.1	0.31	(09)	8.6	Wisconsin	9.6	0.28	(21)	3.2
Mississippi	8.6	0.39	(43)	-7.5	Wyoming	7.9	0.88	(48)	-15.1 ^c
Missouri	9.8	0.28	(18)	5.4					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XIX-19
NON-HODGKIN LYMPHOMA

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	5.9	0.03		
High Five States				
Wyoming	7.7	0.76	(01)	30.5 ^c
Oregon	7.0 ^b	0.26	(02)	18.6 ^c
Indiana	6.7 ^b	0.19	(03)	13.6
Michigan	6.7 ^b	0.15	(04)	13.6
Maine	6.6	0.39	(05)	11.9
Low Five States				
Utah	5.2	0.33	(47)	-11.9
New Mexico	5.1	0.32	(48)	-13.6
Alaska	4.9	0.74	(49)	-16.9 ^c
D.C.	4.5	0.51	(50)	-23.7 ^c
Hawaii	4.4 ^b	0.34	(51)	-25.4 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.0	0.21	(25)	1.7	Montana	5.8	0.46	(28)	-1.7
Alaska	4.9	0.74	(49)	-16.9 ^c	Nebraska	6.2	0.34	(18)	5.1
Arizona	5.7	0.19	(34)	-3.4	Nevada	5.4	0.32	(45)	-8.5
Arkansas	5.5	0.25	(40)	-6.8	New Hampshire	6.4	0.42	(12)	8.5
California	5.4 ^b	0.08	(43)	-8.5	New Jersey	6.0	0.15	(24)	1.7
Colorado	5.6	0.23	(37)	-5.1	New Mexico	5.1	0.32	(48)	-13.6
Connecticut	5.9	0.23	(27)	0.0	New York	5.4 ^b	0.10	(44)	-8.5
Delaware	5.8	0.48	(32)	-1.7	North Carolina	5.8	0.15	(31)	-1.7
D.C.	4.5	0.51	(50)	-23.7 ^c	North Dakota	5.7	0.52	(35)	-3.4
Florida	5.4 ^b	0.10	(41)	-8.5	Ohio	6.4 ^b	0.13	(11)	8.5
Georgia	5.4	0.16	(42)	-8.5	Oklahoma	5.8	0.23	(30)	-1.7
Hawaii	4.4 ^b	0.34	(51)	-25.4 ^c	Oregon	7.0 ^b	0.26	(02)	18.6 ^c
Idaho	6.0	0.41	(26)	1.7	Pennsylvania	6.5 ^b	0.12	(07)	10.2
Illinois	6.2	0.13	(17)	5.1	Rhode Island	6.2	0.41	(19)	5.1
Indiana	6.7 ^b	0.19	(03)	13.6	South Carolina	5.7	0.22	(33)	-3.4
Iowa	6.6	0.26	(06)	11.9	South Dakota	6.1	0.49	(23)	3.4
Kansas	6.5	0.28	(08)	10.2	Tennessee	6.4	0.19	(13)	8.5
Kentucky	6.2	0.23	(20)	5.1	Texas	5.8	0.11	(29)	-1.7
Louisiana	6.4	0.23	(10)	8.5	Utah	5.2	0.33	(47)	-11.9
Maine	6.6	0.39	(05)	11.9	Vermont	5.5	0.54	(39)	-6.8
Maryland	5.6	0.19	(38)	-5.1	Virginia	5.6	0.17	(36)	-5.1
Massachusetts	6.4	0.18	(09)	8.5	Washington	6.1	0.19	(21)	3.4
Michigan	6.7 ^b	0.15	(04)	13.6	West Virginia	6.3	0.32	(16)	6.8
Minnesota	6.3	0.21	(14)	6.8	Wisconsin	6.1	0.19	(22)	3.4
Mississippi	5.3	0.25	(46)	-10.2	Wyoming	7.7	0.76	(01)	30.5 ^c
Missouri	6.3	0.19	(15)	6.8					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XX-13
 CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.6	0.01		
High Five States				
D.C.	3.8	0.36	(01)	46.2 ^c
Mississippi	3.6 ^b	0.16	(02)	38.5 ^c
Louisiana	3.5 ^b	0.13	(03)	34.6 ^c
South Carolina	3.3 ^b	0.12	(04)	26.9 ^c
Alaska	3.2	0.43	(05)	23.1 ^c
Low Five States				
Montana	2.1	0.20	(47)	-19.2 ^c
Vermont	2.1	0.25	(48)	-19.2 ^c
Nebraska	2.0 ^b	0.15	(49)	-23.1 ^c
New Mexico	1.9 ^b	0.14	(50)	-26.9 ^c
Utah	1.8 ^b	0.14	(51)	-30.8 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.9	0.11	(12)	11.5	Montana	2.1	0.20	(47)	-19.2 ^c
Alaska	3.2	0.43	(05)	23.1 ^c	Nebraska	2.0 ^b	0.15	(49)	-23.1 ^c
Arizona	2.2 ^b	0.09	(43)	-15.4 ^c	Nevada	3.0	0.17	(10)	15.4 ^c
Arkansas	3.1	0.14	(07)	19.2 ^c	New Hampshire	3.0	0.22	(08)	15.4 ^c
California	2.7	0.04	(22)	3.8	New Jersey	2.5	0.07	(32)	-3.8
Colorado	2.1 ^b	0.11	(46)	-19.2 ^c	New Mexico	1.9 ^b	0.14	(50)	-26.9 ^c
Connecticut	2.2 ^b	0.11	(45)	-15.4 ^c	New York	2.4 ^b	0.05	(38)	-7.7
Delaware	2.7	0.25	(21)	3.8	North Carolina	2.8	0.08	(18)	7.7
D.C.	3.8	0.36	(01)	46.2 ^c	North Dakota	2.9	0.29	(15)	11.5
Florida	2.9 ^b	0.05	(16)	11.5	Ohio	2.6	0.07	(26)	0.0
Georgia	2.9	0.09	(14)	11.5	Oklahoma	2.6	0.12	(27)	0.0
Hawaii	3.0	0.21	(09)	15.4 ^c	Oregon	2.7	0.12	(19)	3.8
Idaho	2.7	0.20	(20)	3.8	Pennsylvania	2.4 ^b	0.06	(36)	-7.7
Illinois	2.5	0.06	(31)	-3.8	Rhode Island	2.2	0.20	(42)	-15.4 ^c
Indiana	2.2 ^b	0.08	(44)	-15.4 ^c	South Carolina	3.3 ^b	0.12	(04)	26.9 ^c
Iowa	2.5	0.12	(33)	-3.8	South Dakota	2.3	0.23	(41)	-11.5
Kansas	2.3	0.13	(40)	-11.5	Tennessee	3.1 ^b	0.10	(06)	19.2 ^c
Kentucky	2.9	0.12	(13)	11.5	Texas	2.7	0.05	(24)	3.8
Louisiana	3.5 ^b	0.13	(03)	34.6 ^c	Utah	1.8 ^b	0.14	(51)	-30.8 ^c
Maine	2.8	0.19	(17)	7.7	Vermont	2.1	0.25	(48)	-19.2 ^c
Maryland	2.7	0.10	(25)	3.8	Virginia	2.4	0.08	(37)	-7.7
Massachusetts	2.6	0.09	(30)	0.0	Washington	3.0	0.10	(11)	15.4 ^c
Michigan	2.6	0.07	(28)	0.0	West Virginia	2.7	0.16	(23)	3.8
Minnesota	2.3	0.10	(39)	-11.5	Wisconsin	2.6	0.09	(29)	0.0
Mississippi	3.6 ^b	0.16	(02)	38.5 ^c	Wyoming	2.4	0.31	(35)	-7.7
Missouri	2.4	0.09	(34)	-7.7					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XX-14
 CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	4.0	0.03		
High Five States				
D.C.	6.5	0.73	(01)	62.5 ^c
Mississippi	5.8 ^b	0.31	(02)	45.0 ^c
Louisiana	5.8 ^b	0.25	(03)	45.0 ^c
South Carolina	5.5 ^b	0.24	(04)	37.5 ^c
Arkansas	5.3 ^b	0.29	(05)	32.5 ^c
Low Five States				
Colorado	3.0 ^b	0.19	(47)	-25.0 ^c
Montana	2.9	0.35	(48)	-27.5 ^c
Wyoming	2.8	0.49	(49)	-30.0 ^c
New Mexico	2.8 ^b	0.26	(50)	-30.0 ^c
Utah	2.5 ^b	0.26	(51)	-37.5 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.7	0.22	(07)	17.5 ^c	Montana	2.9	0.35	(48)	-27.5 ^c
Alaska	3.9	0.69	(29)	-2.5	Nebraska	3.1	0.28	(46)	-22.5 ^c
Arizona	3.5	0.16	(40)	-12.5	Nevada	4.2	0.30	(17)	5.0
Arkansas	5.3 ^b	0.29	(05)	32.5 ^c	New Hampshire	4.2	0.39	(15)	5.0
California	4.0	0.08	(24)	0.0	New Jersey	3.8	0.14	(32)	-5.0
Colorado	3.0 ^b	0.19	(47)	-25.0 ^c	New Mexico	2.8 ^b	0.26	(50)	-30.0 ^c
Connecticut	3.5	0.21	(38)	-12.5	New York	3.6 ^b	0.09	(35)	-10.0
Delaware	4.2	0.47	(18)	5.0	North Carolina	4.1	0.15	(20)	2.5
D.C.	6.5	0.73	(01)	62.5 ^c	North Dakota	3.9	0.50	(27)	-2.5
Florida	4.3	0.10	(12)	7.5	Ohio	4.1	0.13	(21)	2.5
Georgia	4.3	0.17	(14)	7.5	Oklahoma	3.9	0.22	(28)	-2.5
Hawaii	4.4	0.38	(10)	10.0	Oregon	3.7	0.21	(34)	-7.5
Idaho	3.9	0.37	(26)	-2.5	Pennsylvania	3.8	0.11	(31)	-5.0
Illinois	4.0	0.12	(23)	0.0	Rhode Island	3.4	0.37	(43)	-15.0 ^c
Indiana	3.5	0.16	(41)	-12.5	South Carolina	5.5 ^b	0.24	(04)	37.5 ^c
Iowa	3.6	0.22	(36)	-10.0	South Dakota	3.1	0.41	(45)	-22.5 ^c
Kansas	3.5	0.24	(39)	-12.5	Tennessee	5.0 ^b	0.20	(06)	25.0 ^c
Kentucky	4.6	0.23	(08)	15.0 ^c	Texas	4.2	0.10	(16)	5.0
Louisiana	5.8 ^b	0.25	(03)	45.0 ^c	Utah	2.5 ^b	0.26	(51)	-37.5 ^c
Maine	4.5	0.37	(09)	12.5	Vermont	3.2	0.47	(44)	-20.0 ^c
Maryland	4.3	0.20	(11)	7.5	Virginia	3.7	0.16	(33)	-7.5
Massachusetts	4.0	0.17	(22)	0.0	Washington	4.3	0.18	(13)	7.5
Michigan	4.0	0.13	(25)	0.0	West Virginia	4.1	0.29	(19)	2.5
Minnesota	3.4	0.18	(42)	-15.0 ^c	Wisconsin	3.8	0.17	(30)	-5.0
Mississippi	5.8 ^b	0.31	(02)	45.0 ^c	Wyoming	2.8	0.49	(49)	-30.0 ^c
Missouri	3.6	0.16	(37)	-10.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XX-15
 CANCER OF THE ORAL CAVITY AND PHARYNX (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	1.5	0.01		
High Five States				
Alaska	2.7	0.54	(01)	80.0 ^c
North Dakota	2.1	0.33	(02)	40.0 ^c
New Hampshire	2.1	0.24	(03)	40.0 ^c
Wyoming	1.9	0.37	(04)	26.7 ^c
Oregon	1.9	0.13	(05)	26.7 ^c
Low Five States				
Vermont	1.2	0.25	(47)	-20.0 ^c
New Mexico	1.1	0.15	(48)	-26.7 ^c
Nebraska	1.1	0.15	(49)	-26.7 ^c
Connecticut	1.1 ^b	0.10	(50)	-26.7 ^c
Utah	1.1	0.15	(51)	-26.7 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.6	0.11	(17)	6.7	Montana	1.4	0.23	(34)	-6.7
Alaska	2.7	0.54	(01)	80.0 ^c	Nebraska	1.1	0.15	(49)	-26.7 ^c
Arizona	1.2 ^b	0.09	(46)	-20.0 ^c	Nevada	1.9	0.19	(06)	26.7 ^c
Arkansas	1.4	0.13	(37)	-6.7	New Hampshire	2.1	0.24	(03)	40.0 ^c
California	1.7 ^b	0.04	(13)	13.3	New Jersey	1.4	0.07	(29)	-6.7
Colorado	1.4	0.11	(35)	-6.7	New Mexico	1.1	0.15	(48)	-26.7 ^c
Connecticut	1.1 ^b	0.10	(50)	-26.7 ^c	New York	1.4	0.05	(38)	-6.7
Delaware	1.5	0.25	(23)	0.0	North Carolina	1.7	0.08	(11)	13.3
D.C.	1.8	0.33	(09)	20.0 ^c	North Dakota	2.1	0.33	(02)	40.0 ^c
Florida	1.6	0.05	(18)	6.7	Ohio	1.4	0.06	(33)	-6.7
Georgia	1.8	0.09	(08)	20.0 ^c	Oklahoma	1.6	0.12	(19)	6.7
Hawaii	1.7	0.21	(12)	13.3	Oregon	1.9	0.13	(05)	26.7 ^c
Idaho	1.6	0.21	(16)	6.7	Pennsylvania	1.3 ^b	0.06	(44)	-13.3
Illinois	1.3	0.06	(42)	-13.3	Rhode Island	1.4	0.21	(39)	-6.7
Indiana	1.2 ^b	0.08	(45)	-20.0 ^c	South Carolina	1.5	0.11	(24)	0.0
Iowa	1.5	0.13	(20)	0.0	South Dakota	1.5	0.24	(28)	0.0
Kansas	1.3	0.13	(43)	-13.3	Tennessee	1.6	0.10	(15)	6.7
Kentucky	1.5	0.11	(22)	0.0	Texas	1.4	0.05	(30)	-6.7
Louisiana	1.7	0.12	(14)	13.3	Utah	1.1	0.15	(51)	-26.7 ^c
Maine	1.5	0.18	(25)	0.0	Vermont	1.2	0.25	(47)	-20.0 ^c
Maryland	1.4	0.09	(41)	-6.7	Virginia	1.4	0.08	(40)	-6.7
Massachusetts	1.5	0.09	(26)	0.0	Washington	1.8	0.11	(07)	20.0 ^c
Michigan	1.5	0.07	(27)	0.0	West Virginia	1.4	0.15	(36)	-6.7
Minnesota	1.4	0.10	(32)	-6.7	Wisconsin	1.5	0.10	(21)	0.0
Mississippi	1.7	0.14	(10)	13.3	Wyoming	1.9	0.37	(04)	26.7 ^c
Missouri	1.4	0.09	(31)	-6.7					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XXI-11
OVARY CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	8.8	0.03		
High Five States				
Wyoming	10.4	0.88	(01)	18.2 ^c
Washington	10.3 ^b	0.25	(02)	17.0 ^c
Oregon	10.0 ^b	0.31	(03)	13.6
West Virginia	9.6	0.40	(04)	9.1
Alabama	9.6	0.27	(05)	9.1
Low Five States				
Mississippi	7.9	0.31	(47)	-10.2
South Carolina	7.9 ^b	0.26	(48)	-10.2
North Dakota	7.9	0.64	(49)	-10.2
Hawaii	7.0 ^b	0.44	(50)	-20.5 ^c
Alaska	6.0 ^b	0.78	(51)	-31.8 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.6	0.27	(05)	9.1	Montana	9.5	0.59	(06)	8.0
Alaska	6.0 ^b	0.78	(51)	-31.8 ^c	Nebraska	8.8	0.42	(31)	0.0
Arizona	8.9	0.24	(28)	1.1	Nevada	9.1	0.41	(17)	3.4
Arkansas	8.3	0.32	(41)	-5.7	New Hampshire	8.7	0.49	(34)	-1.1
California	8.9	0.10	(24)	1.1	New Jersey	9.4	0.19	(11)	6.8
Colorado	9.5	0.30	(08)	8.0	New Mexico	8.2	0.40	(43)	-6.8
Connecticut	8.4	0.28	(38)	-4.5	New York	8.9	0.12	(27)	1.1
Delaware	8.8	0.60	(30)	0.0	North Carolina	8.7	0.19	(35)	-1.1
D.C.	8.0	0.69	(45)	-9.1	North Dakota	7.9	0.64	(49)	-10.2
Florida	8.2 ^b	0.12	(44)	-6.8	Ohio	9.1	0.16	(18)	3.4
Georgia	9.0	0.21	(22)	2.3	Oklahoma	8.4	0.29	(39)	-4.5
Hawaii	7.0 ^b	0.44	(50)	-20.5 ^c	Oregon	10.0 ^b	0.31	(03)	13.6
Idaho	8.9	0.51	(23)	1.1	Pennsylvania	9.5 ^b	0.15	(07)	8.0
Illinois	8.8	0.16	(29)	0.0	Rhode Island	9.4	0.54	(10)	6.8
Indiana	9.4	0.23	(12)	6.8	South Carolina	7.9 ^b	0.26	(48)	-10.2
Iowa	9.4	0.32	(09)	6.8	South Dakota	9.3	0.64	(13)	5.7
Kansas	9.2	0.34	(15)	4.5	Tennessee	9.2	0.23	(14)	4.5
Kentucky	8.7	0.27	(33)	-1.1	Texas	8.0 ^b	0.12	(46)	-9.1
Louisiana	8.3	0.26	(40)	-5.7	Utah	8.7	0.43	(32)	-1.1
Maine	9.1	0.47	(16)	3.4	Vermont	8.3	0.67	(42)	-5.7
Maryland	8.6	0.24	(36)	-2.3	Virginia	8.9	0.21	(26)	1.1
Massachusetts	9.0	0.22	(21)	2.3	Washington	10.3 ^b	0.25	(02)	17.0 ^c
Michigan	9.0	0.18	(20)	2.3	West Virginia	9.6	0.40	(04)	9.1
Minnesota	8.9	0.26	(25)	1.1	Wisconsin	9.1	0.24	(19)	3.4
Mississippi	7.9	0.31	(47)	-10.2	Wyoming	10.4	0.88	(01)	18.2 ^c
Missouri	8.6	0.22	(37)	-2.3					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXII-13
PANCREAS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	10.6	0.03		
High Five States				
D.C.	12.2	0.65	(01)	15.1 ^c
Louisiana	12.1 ^b	0.24	(02)	14.2
Maine	11.5	0.39	(03)	8.5
Massachusetts	11.4 ^b	0.18	(04)	7.5
Connecticut	11.4	0.24	(05)	7.5
Low Five States				
Arizona	9.3 ^b	0.18	(47)	-12.3
Nebraska	9.3 ^b	0.32	(48)	-12.3
West Virginia	9.1 ^b	0.29	(49)	-14.2
Vermont	9.1	0.52	(50)	-14.2
Utah	8.9 ^b	0.32	(51)	-16.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	10.6	0.21	(30)	0.0	Montana	9.9	0.44	(45)	-6.6
Alaska	10.4	0.79	(37)	-1.9	Nebraska	9.3 ^b	0.32	(48)	-12.3
Arizona	9.3 ^b	0.18	(47)	-12.3	Nevada	10.6	0.32	(28)	0.0
Arkansas	10.4	0.26	(34)	-1.9	New Hampshire	10.9	0.41	(16)	2.8
California	10.3 ^b	0.08	(38)	-2.8	New Jersey	11.1	0.16	(08)	4.7
Colorado	9.9	0.23	(46)	-6.6	New Mexico	10.0	0.33	(42)	-5.7
Connecticut	11.4	0.24	(05)	7.5	New York	11.0 ^b	0.10	(11)	3.8
Delaware	10.5	0.50	(31)	-0.9	North Carolina	10.8	0.16	(20)	1.9
D.C.	12.2	0.65	(01)	15.1 ^c	North Dakota	10.1	0.53	(40)	-4.7
Florida	9.9 ^b	0.10	(43)	-6.6	Ohio	10.6	0.13	(27)	0.0
Georgia	10.5	0.17	(32)	-0.9	Oklahoma	9.9	0.23	(44)	-6.6
Hawaii	10.7	0.40	(25)	0.9	Oregon	10.9	0.24	(17)	2.8
Idaho	10.7	0.41	(23)	0.9	Pennsylvania	11.0	0.12	(15)	3.8
Illinois	11.3 ^b	0.14	(07)	6.6	Rhode Island	10.6	0.42	(29)	0.0
Indiana	11.0	0.19	(12)	3.8	South Carolina	10.8	0.23	(21)	1.9
Iowa	10.4	0.25	(36)	-1.9	South Dakota	10.9	0.51	(18)	2.8
Kansas	10.4	0.27	(35)	-1.9	Tennessee	11.0	0.19	(14)	3.8
Kentucky	10.4	0.22	(33)	-1.9	Texas	10.1 ^b	0.11	(41)	-4.7
Louisiana	12.1 ^b	0.24	(02)	14.2	Utah	8.9 ^b	0.32	(51)	-16.0 ^c
Maine	11.5	0.39	(03)	8.5	Vermont	9.1	0.52	(50)	-14.2
Maryland	11.4 ^b	0.21	(06)	7.5	Virginia	10.7	0.18	(22)	0.9
Massachusetts	11.4 ^b	0.18	(04)	7.5	Washington	11.0	0.19	(13)	3.8
Michigan	10.8	0.15	(19)	1.9	West Virginia	9.1 ^b	0.29	(49)	-14.2
Minnesota	10.2	0.20	(39)	-3.8	Wisconsin	11.1	0.20	(10)	4.7
Mississippi	11.1	0.28	(09)	4.7	Wyoming	10.7	0.66	(24)	0.9
Missouri	10.6	0.19	(26)	0.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXII-14
PANCREAS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	12.2	0.04		
High Five States				
D.C.	14.4	1.14	(01)	18.0 ^c
Louisiana	13.8 ^b	0.40	(02)	13.1 ^c
Maine	13.7	0.66	(03)	12.3 ^c
Connecticut	13.3	0.41	(04)	9.0
Massachusetts	13.3 ^b	0.31	(05)	9.0
Low Five States				
Colorado	11.0	0.37	(47)	-9.8
West Virginia	10.9	0.49	(48)	-10.7 ^c
Utah	10.9	0.54	(49)	-10.7 ^c
Arizona	10.8 ^b	0.29	(50)	-11.5 ^c
Vermont	10.5	0.87	(51)	-13.9 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	12.4	0.36	(19)	1.6	Montana	11.1	0.70	(46)	-9.0
Alaska	12.2	1.33	(27)	0.0	Nebraska	11.2	0.54	(44)	-8.2
Arizona	10.8 ^b	0.29	(50)	-11.5 ^c	Nevada	11.7	0.51	(34)	-4.1
Arkansas	12.3	0.44	(23)	0.8	New Hampshire	11.2	0.65	(43)	-8.2
California	11.6 ^b	0.13	(38)	-4.9	New Jersey	12.5	0.26	(16)	2.5
Colorado	11.0	0.37	(47)	-9.8	New Mexico	11.2	0.53	(45)	-8.2
Connecticut	13.3	0.41	(04)	9.0	New York	12.4	0.17	(20)	1.6
Delaware	11.5	0.79	(40)	-5.7	North Carolina	12.8	0.28	(10)	4.9
D.C.	14.4	1.14	(01)	18.0 ^c	North Dakota	11.7	0.86	(35)	-4.1
Florida	11.6 ^b	0.16	(36)	-4.9	Ohio	12.3	0.22	(24)	0.8
Georgia	12.5	0.30	(18)	2.5	Oklahoma	12.0	0.39	(30)	-1.6
Hawaii	11.9	0.63	(31)	-2.5	Oregon	12.3	0.39	(25)	0.8
Idaho	11.4	0.63	(42)	-6.6	Pennsylvania	12.8	0.20	(11)	4.9
Illinois	13.1 ^b	0.23	(06)	7.4	Rhode Island	11.6	0.69	(37)	-4.9
Indiana	12.9	0.32	(08)	5.7	South Carolina	12.7	0.39	(13)	4.1
Iowa	11.5	0.40	(39)	-5.7	South Dakota	11.4	0.78	(41)	-6.6
Kansas	12.2	0.45	(26)	0.0	Tennessee	12.5	0.32	(17)	2.5
Kentucky	12.0	0.37	(29)	-1.6	Texas	11.8	0.18	(32)	-3.3
Louisiana	13.8 ^b	0.40	(02)	13.1 ^c	Utah	10.9	0.54	(49)	-10.7 ^c
Maine	13.7	0.66	(03)	12.3 ^c	Vermont	10.5	0.87	(51)	-13.9 ^c
Maryland	12.9	0.35	(09)	5.7	Virginia	12.4	0.30	(21)	1.6
Massachusetts	13.3 ^b	0.31	(05)	9.0	Washington	12.4	0.31	(22)	1.6
Michigan	12.7	0.24	(12)	4.1	West Virginia	10.9	0.49	(48)	-10.7 ^c
Minnesota	11.8	0.33	(33)	-3.3	Wisconsin	12.6	0.32	(14)	3.3
Mississippi	13.1	0.48	(07)	7.4	Wyoming	12.1	1.06	(28)	-0.8
Missouri	12.6	0.31	(15)	3.3					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXII-15
PANCREAS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	9.3	0.03		
High Five States				
Louisiana	10.7 ^b	0.29	(01)	15.1 ^c
New Hampshire	10.7	0.54	(02)	15.1 ^c
D.C.	10.6	0.78	(03)	14.0
South Dakota	10.4	0.67	(04)	11.8
Maryland	10.2 ^b	0.26	(05)	9.7
Low Five States				
Vermont	8.0	0.66	(47)	-14.0
Arizona	8.0 ^b	0.23	(48)	-14.0
Nebraska	7.9	0.39	(49)	-15.1 ^c
West Virginia	7.7 ^b	0.35	(50)	-17.2 ^c
Utah	7.3 ^b	0.40	(51)	-21.5 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	9.1	0.26	(33)	-2.2	Montana	8.8	0.56	(42)	-5.4
Alaska	9.0	0.99	(35)	-3.2	Nebraska	7.9	0.39	(49)	-15.1 ^c
Arizona	8.0 ^b	0.23	(48)	-14.0	Nevada	9.5	0.42	(21)	2.2
Arkansas	9.0	0.32	(36)	-3.2	New Hampshire	10.7	0.54	(02)	15.1 ^c
California	9.2	0.10	(30)	-1.1	New Jersey	10.0 ^b	0.19	(06)	7.5
Colorado	8.9	0.29	(39)	-4.3	New Mexico	8.9	0.42	(41)	-4.3
Connecticut	9.9	0.30	(09)	6.5	New York	9.9 ^b	0.13	(10)	6.5
Delaware	9.5	0.62	(23)	2.2	North Carolina	9.2	0.20	(29)	-1.1
D.C.	10.6	0.78	(03)	14.0	North Dakota	8.7	0.66	(43)	-6.5
Florida	8.5 ^b	0.12	(45)	-8.6	Ohio	9.2	0.16	(31)	-1.1
Georgia	9.0	0.21	(34)	-3.2	Oklahoma	8.3 ^b	0.28	(46)	-10.8
Hawaii	9.6	0.50	(20)	3.2	Oregon	9.7	0.31	(16)	4.3
Idaho	9.9	0.53	(08)	6.5	Pennsylvania	9.5	0.15	(22)	2.2
Illinois	9.9 ^b	0.17	(11)	6.5	Rhode Island	9.8	0.53	(13)	5.4
Indiana	9.5	0.23	(24)	2.2	South Carolina	9.4	0.28	(25)	1.1
Iowa	9.4	0.32	(26)	1.1	South Dakota	10.4	0.67	(04)	11.8
Kansas	8.9	0.33	(40)	-4.3	Tennessee	9.6	0.24	(18)	3.2
Kentucky	9.1	0.28	(32)	-2.2	Texas	8.7 ^b	0.13	(44)	-6.5
Louisiana	10.7 ^b	0.29	(01)	15.1 ^c	Utah	7.3 ^b	0.40	(51)	-21.5 ^c
Maine	9.6	0.47	(17)	3.2	Vermont	8.0	0.66	(47)	-14.0
Maryland	10.2 ^b	0.26	(05)	9.7	Virginia	9.3	0.22	(27)	0.0
Massachusetts	10.0	0.22	(07)	7.5	Washington	9.8	0.25	(12)	5.4
Michigan	9.3	0.18	(28)	0.0	West Virginia	7.7 ^b	0.35	(50)	-17.2 ^c
Minnesota	9.0	0.25	(38)	-3.2	Wisconsin	9.8	0.24	(14)	5.4
Mississippi	9.7	0.34	(15)	4.3	Wyoming	9.6	0.85	(19)	3.2
Missouri	9.0	0.22	(37)	-3.2					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXIII-11
PROSTATE CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	26.7	0.07		
High Five States				
D.C.	43.8 ^b	2.09	(01)	64.0 ^c
Mississippi	36.2 ^b	0.86	(02)	35.6 ^c
Alabama	33.0 ^b	0.64	(03)	23.6 ^c
South Carolina	32.4 ^b	0.69	(04)	21.3 ^c
Louisiana	32.2 ^b	0.66	(05)	20.6 ^c
Low Five States				
Oklahoma	24.9	0.60	(47)	-6.7
Missouri	24.3 ^b	0.46	(48)	-9.0
Arizona	23.2 ^b	0.45	(49)	-13.1 ^c
Florida	22.3 ^b	0.21	(50)	-16.5 ^c
Hawaii	18.8 ^b	0.80	(51)	-29.6 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	33.0 ^b	0.64	(03)	23.6 ^c	Montana	29.2	1.20	(12)	9.4
Alaska	26.1	2.27	(39)	-2.2	Nebraska	25.1	0.82	(43)	-6.0
Arizona	23.2 ^b	0.45	(49)	-13.1 ^c	Nevada	26.6	0.86	(33)	-0.4
Arkansas	28.9	0.71	(14)	8.2	New Hampshire	28.3	1.11	(19)	6.0
California	25.0 ^b	0.21	(46)	-6.4	New Jersey	26.2	0.40	(38)	-1.9
Colorado	26.5	0.64	(36)	-0.7	New Mexico	26.9	0.88	(30)	0.7
Connecticut	27.0	0.60	(27)	1.1	New York	26.0	0.26	(40)	-2.6
Delaware	29.4	1.37	(09)	10.1 ^c	North Carolina	29.9 ^b	0.47	(08)	12.0 ^c
D.C.	43.8 ^b	2.09	(01)	64.0 ^c	North Dakota	28.1	1.33	(22)	5.2
Florida	22.3 ^b	0.21	(50)	-16.5 ^c	Ohio	27.9	0.36	(23)	4.5
Georgia	30.6 ^b	0.52	(06)	14.6 ^c	Oklahoma	24.9	0.60	(47)	-6.7
Hawaii	18.8 ^b	0.80	(51)	-29.6 ^c	Oregon	27.7	0.61	(25)	3.7
Idaho	29.0	1.07	(13)	8.6	Pennsylvania	26.9	0.31	(29)	0.7
Illinois	28.1 ^b	0.35	(20)	5.2	Rhode Island	27.0	1.08	(28)	1.1
Indiana	27.8	0.50	(24)	4.1	South Carolina	32.4 ^b	0.69	(04)	21.3 ^c
Iowa	27.5	0.63	(26)	3.0	South Dakota	28.8	1.26	(16)	7.9
Kansas	25.0	0.66	(44)	-6.4	Tennessee	29.3 ^b	0.54	(11)	9.7
Kentucky	26.9	0.62	(31)	0.7	Texas	25.5 ^b	0.28	(42)	-4.5
Louisiana	32.2 ^b	0.66	(05)	20.6 ^c	Utah	26.5	0.90	(34)	-0.7
Maine	28.1	1.00	(21)	5.2	Vermont	28.3	1.54	(18)	6.0
Maryland	29.3 ^b	0.57	(10)	9.7	Virginia	30.1 ^b	0.51	(07)	12.7 ^c
Massachusetts	26.4	0.45	(37)	-1.1	Washington	26.8	0.49	(32)	0.4
Michigan	26.5	0.38	(35)	-0.7	West Virginia	25.0	0.79	(45)	-6.4
Minnesota	28.6	0.54	(17)	7.1	Wisconsin	28.8 ^b	0.51	(15)	7.9
Mississippi	36.2 ^b	0.86	(02)	35.6 ^c	Wyoming	25.8	1.66	(41)	-3.4
Missouri	24.3 ^b	0.46	(48)	-9.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 10% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXIV-13
STOMACH CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.1	0.02		
High Five States				
Hawaii	7.3 ^b	0.33	(01)	78.0 ^c
D.C.	6.2 ^b	0.46	(02)	51.2 ^c
Rhode Island	5.6 ^b	0.30	(03)	36.6 ^c
Alaska	5.3	0.57	(04)	29.3 ^c
Louisiana	5.0 ^b	0.15	(05)	22.0 ^c
Low Five States				
Wyoming	3.0	0.35	(47)	-26.8 ^c
Iowa	2.9 ^b	0.13	(48)	-29.3 ^c
Utah	2.8 ^b	0.18	(49)	-31.7 ^c
Idaho	2.7 ^b	0.20	(50)	-34.1 ^c
Nebraska	2.6 ^b	0.17	(51)	-36.6 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	4.0	0.13	(18)	-2.4	Montana	3.3	0.26	(37)	-19.5 ^c
Alaska	5.3	0.57	(04)	29.3 ^c	Nebraska	2.6 ^b	0.17	(51)	-36.6 ^c
Arizona	3.4 ^b	0.11	(34)	-17.1 ^c	Nevada	3.7	0.19	(27)	-9.8
Arkansas	3.8	0.16	(25)	-7.3	New Hampshire	3.1 ^b	0.22	(45)	-24.4 ^c
California	4.9 ^b	0.06	(08)	19.5 ^c	New Jersey	4.9 ^b	0.10	(07)	19.5 ^c
Colorado	3.3 ^b	0.13	(39)	-19.5 ^c	New Mexico	4.7	0.23	(09)	14.6
Connecticut	4.7 ^b	0.16	(10)	14.6	New York	5.0 ^b	0.07	(06)	22.0 ^c
Delaware	3.7	0.29	(26)	-9.8	North Carolina	3.8	0.10	(22)	-7.3
D.C.	6.2 ^b	0.46	(02)	51.2 ^c	North Dakota	3.8	0.33	(23)	-7.3
Florida	3.8 ^b	0.06	(24)	-7.3	Ohio	3.5 ^b	0.08	(31)	-14.6
Georgia	3.9	0.10	(20)	-4.9	Oklahoma	3.2 ^b	0.13	(40)	-22.0 ^c
Hawaii	7.3 ^b	0.33	(01)	78.0 ^c	Oregon	3.1 ^b	0.13	(43)	-24.4 ^c
Idaho	2.7 ^b	0.20	(50)	-34.1 ^c	Pennsylvania	3.9	0.07	(19)	-4.9
Illinois	4.5 ^b	0.09	(11)	9.8	Rhode Island	5.6 ^b	0.30	(03)	36.6 ^c
Indiana	3.2 ^b	0.10	(41)	-22.0 ^c	South Carolina	4.5	0.15	(12)	9.8
Iowa	2.9 ^b	0.13	(48)	-29.3 ^c	South Dakota	3.1	0.27	(42)	-24.4 ^c
Kansas	3.1 ^b	0.15	(44)	-24.4 ^c	Tennessee	3.5 ^b	0.11	(32)	-14.6
Kentucky	3.5 ^b	0.13	(33)	-14.6	Texas	4.5 ^b	0.07	(14)	9.8
Louisiana	5.0 ^b	0.15	(05)	22.0 ^c	Utah	2.8 ^b	0.18	(49)	-31.7 ^c
Maine	3.6	0.22	(28)	-12.2	Vermont	3.0 ^b	0.30	(46)	-26.8 ^c
Maryland	4.1	0.13	(16)	0.0	Virginia	4.0	0.11	(17)	-2.4
Massachusetts	4.4	0.11	(15)	7.3	Washington	3.6 ^b	0.11	(30)	-12.2
Michigan	3.8	0.09	(21)	-7.3	West Virginia	3.6	0.18	(29)	-12.2
Minnesota	3.4 ^b	0.12	(35)	-17.1 ^c	Wisconsin	3.3 ^b	0.11	(36)	-19.5 ^c
Mississippi	4.5	0.18	(13)	9.8	Wyoming	3.0	0.35	(47)	-26.8 ^c
Missouri	3.3 ^b	0.10	(38)	-19.5 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXIV-14
STOMACH CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	5.7	0.03		
High Five States				
Hawaii	10.0 ^b	0.58	(01)	75.4 ^c
D.C.	8.9 ^b	0.89	(02)	56.1 ^c
Rhode Island	8.2 ^b	0.58	(03)	43.9 ^c
Alaska	7.6	1.07	(04)	33.3 ^c
Louisiana	7.4 ^b	0.30	(05)	29.8 ^c
Low Five States				
Wyoming	4.2	0.63	(47)	-26.3 ^c
Oregon	4.1 ^b	0.23	(48)	-28.1 ^c
Nebraska	3.7 ^b	0.31	(49)	-35.1 ^c
Utah	3.5 ^b	0.31	(50)	-38.6 ^c
Idaho	3.5 ^b	0.35	(51)	-38.6 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	5.6	0.24	(17)	-1.8	Montana	4.7	0.47	(34)	-17.5 ^c
Alaska	7.6	1.07	(04)	33.3 ^c	Nebraska	3.7 ^b	0.31	(49)	-35.1 ^c
Arizona	4.7 ^b	0.19	(38)	-17.5 ^c	Nevada	4.9	0.34	(31)	-14.0
Arkansas	5.1	0.29	(28)	-10.5	New Hampshire	4.7	0.42	(36)	-17.5 ^c
California	6.7 ^b	0.10	(09)	17.5 ^c	New Jersey	7.1 ^b	0.20	(06)	24.6 ^c
Colorado	4.5 ^b	0.24	(43)	-21.1 ^c	New Mexico	6.7	0.42	(08)	17.5 ^c
Connecticut	6.6	0.29	(10)	15.8 ^c	New York	7.0 ^b	0.13	(07)	22.8 ^c
Delaware	5.4	0.55	(19)	-5.3	North Carolina	5.3	0.18	(23)	-7.0
D.C.	8.9 ^b	0.89	(02)	56.1 ^c	North Dakota	5.2	0.58	(25)	-8.8
Florida	5.3 ^b	0.11	(22)	-7.0	Ohio	5.0 ^b	0.14	(29)	-12.3
Georgia	5.4	0.20	(20)	-5.3	Oklahoma	4.5 ^b	0.24	(42)	-21.1 ^c
Hawaii	10.0 ^b	0.58	(01)	75.4 ^c	Oregon	4.1 ^b	0.23	(48)	-28.1 ^c
Idaho	3.5 ^b	0.35	(51)	-38.6 ^c	Pennsylvania	5.6	0.14	(18)	-1.8
Illinois	6.4 ^b	0.16	(12)	12.3	Rhode Island	8.2 ^b	0.58	(03)	43.9 ^c
Indiana	4.6 ^b	0.19	(41)	-19.3 ^c	South Carolina	6.3	0.27	(13)	10.5
Iowa	4.2 ^b	0.24	(46)	-26.3 ^c	South Dakota	4.5	0.49	(44)	-21.1 ^c
Kansas	4.6 ^b	0.27	(40)	-19.3 ^c	Tennessee	4.7 ^b	0.20	(33)	-17.5 ^c
Kentucky	5.0	0.24	(30)	-12.3	Texas	6.2 ^b	0.13	(14)	8.8
Louisiana	7.4 ^b	0.30	(05)	29.8 ^c	Utah	3.5 ^b	0.31	(50)	-38.6 ^c
Maine	5.1	0.41	(27)	-10.5	Vermont	4.2	0.56	(45)	-26.3 ^c
Maryland	5.3	0.22	(24)	-7.0	Virginia	5.9	0.21	(16)	3.5
Massachusetts	6.2	0.21	(15)	8.8	Washington	4.8 ^b	0.20	(32)	-15.8 ^c
Michigan	5.3	0.16	(21)	-7.0	West Virginia	5.2	0.35	(26)	-8.8
Minnesota	4.7 ^b	0.21	(37)	-17.5 ^c	Wisconsin	4.7 ^b	0.20	(35)	-17.5 ^c
Mississippi	6.5	0.34	(11)	14.0	Wyoming	4.2	0.63	(47)	-26.3 ^c
Missouri	4.6 ^b	0.19	(39)	-19.3 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXIV-15
STOMACH CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.9	0.02		
High Five States				
Hawaii	5.1 ^b	0.37	(01)	75.9 ^c
D.C.	4.2	0.49	(02)	44.8 ^c
Rhode Island	3.8	0.32	(03)	31.0 ^c
New York	3.7 ^b	0.08	(04)	27.6 ^c
California	3.6 ^b	0.06	(05)	24.1 ^c
Low Five States				
Kansas	2.0 ^b	0.16	(47)	-31.0 ^c
Wyoming	1.9	0.37	(48)	-34.5 ^c
Iowa	1.9 ^b	0.14	(49)	-34.5 ^c
Nebraska	1.9 ^b	0.19	(50)	-34.5 ^c
New Hampshire	1.8 ^b	0.22	(51)	-37.9 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	2.8	0.14	(20)	-3.4	Montana	2.3	0.28	(40)	-20.7 ^c
Alaska	3.4	0.61	(07)	17.2 ^c	Nebraska	1.9 ^b	0.19	(50)	-34.5 ^c
Arizona	2.4 ^b	0.13	(32)	-17.2 ^c	Nevada	2.7	0.22	(26)	-6.9
Arkansas	2.8	0.18	(19)	-3.4	New Hampshire	1.8 ^b	0.22	(51)	-37.9 ^c
California	3.6 ^b	0.06	(05)	24.1 ^c	New Jersey	3.5 ^b	0.11	(06)	20.7 ^c
Colorado	2.4 ^b	0.15	(35)	-17.2 ^c	New Mexico	3.2	0.25	(10)	10.3
Connecticut	3.3	0.17	(09)	13.8	New York	3.7 ^b	0.08	(04)	27.6 ^c
Delaware	2.4	0.31	(36)	-17.2 ^c	North Carolina	2.8	0.11	(18)	-3.4
D.C.	4.2	0.49	(02)	44.8 ^c	North Dakota	2.7	0.36	(27)	-6.9
Florida	2.6 ^b	0.07	(28)	-10.3	Ohio	2.5 ^b	0.08	(31)	-13.8
Georgia	2.8	0.12	(17)	-3.4	Oklahoma	2.2 ^b	0.15	(41)	-24.1 ^c
Hawaii	5.1 ^b	0.37	(01)	75.9 ^c	Oregon	2.3 ^b	0.15	(37)	-20.7 ^c
Idaho	2.0 ^b	0.24	(46)	-31.0 ^c	Pennsylvania	2.8	0.08	(21)	-3.4
Illinois	3.2	0.09	(11)	10.3	Rhode Island	3.8	0.32	(03)	31.0 ^c
Indiana	2.2 ^b	0.11	(43)	-24.1 ^c	South Carolina	3.1	0.16	(15)	6.9
Iowa	1.9 ^b	0.14	(49)	-34.5 ^c	South Dakota	2.1	0.31	(44)	-27.6 ^c
Kansas	2.0 ^b	0.16	(47)	-31.0 ^c	Tennessee	2.7	0.12	(25)	-6.9
Kentucky	2.4 ^b	0.14	(34)	-17.2 ^c	Texas	3.1	0.08	(13)	6.9
Louisiana	3.4	0.16	(08)	17.2 ^c	Utah	2.2	0.22	(42)	-24.1 ^c
Maine	2.6	0.25	(29)	-10.3	Vermont	2.1	0.33	(45)	-27.6 ^c
Maryland	3.2	0.14	(12)	10.3	Virginia	2.7	0.12	(24)	-6.9
Massachusetts	3.1	0.12	(14)	6.9	Washington	2.7	0.13	(23)	-6.9
Michigan	2.7	0.10	(22)	-6.9	West Virginia	2.6	0.20	(30)	-10.3
Minnesota	2.4 ^b	0.13	(33)	-17.2 ^c	Wisconsin	2.3 ^b	0.12	(38)	-20.7 ^c
Mississippi	3.0	0.19	(16)	3.4	Wyoming	1.9	0.37	(48)	-34.5 ^c
Missouri	2.3 ^b	0.12	(39)	-20.7 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXV-11
TESTIS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.3	0.01		
High Five States				
New Hampshire	0.4	0.12	(01)	33.3 ^c
Idaho	0.4	0.12	(02)	33.3 ^c
Nevada	0.4	0.09	(03)	33.3 ^c
North Dakota	0.4	0.16	(04)	33.3 ^c
Oregon	0.4	0.07	(05)	33.3 ^c
Low Five States				
Georgia	0.2 ^b	0.03	(47)	-33.3 ^c
Delaware	0.1	0.09	(48)	-66.7 ^c
Kentucky	0.1 ^b	0.04	(49)	-66.7 ^c
Wyoming	0.1	0.07	(50)	-66.7 ^c
Alaska	0.0 ^b	0.05	(51)	-100.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.2	0.04	(40)	-33.3 ^c	Montana	0.2	0.09	(43)	-33.3 ^c
Alaska	0.0 ^b	0.05	(51)	-100.0 ^c	Nebraska	0.3	0.09	(08)	0.0
Arizona	0.3	0.05	(14)	0.0	Nevada	0.4	0.09	(03)	33.3 ^c
Arkansas	0.2	0.06	(41)	-33.3 ^c	New Hampshire	0.4	0.12	(01)	33.3 ^c
California	0.3	0.02	(07)	0.0	New Jersey	0.2	0.03	(38)	-33.3 ^c
Colorado	0.2	0.05	(34)	-33.3 ^c	New Mexico	0.3	0.08	(16)	0.0
Connecticut	0.3	0.06	(17)	0.0	New York	0.2	0.02	(28)	-33.3 ^c
Delaware	0.1	0.09	(48)	-66.7 ^c	North Carolina	0.3	0.04	(21)	0.0
D.C.	0.2	0.12	(39)	-33.3 ^c	North Dakota	0.4	0.16	(04)	33.3 ^c
Florida	0.3	0.02	(22)	0.0	Ohio	0.3	0.03	(11)	0.0
Georgia	0.2 ^b	0.03	(47)	-33.3 ^c	Oklahoma	0.2	0.05	(31)	-33.3 ^c
Hawaii	0.2	0.09	(30)	-33.3 ^c	Oregon	0.4	0.07	(05)	33.3 ^c
Idaho	0.4	0.12	(02)	33.3 ^c	Pennsylvania	0.3	0.03	(19)	0.0
Illinois	0.2	0.03	(35)	-33.3 ^c	Rhode Island	0.3	0.11	(15)	0.0
Indiana	0.2	0.04	(27)	-33.3 ^c	South Carolina	0.2	0.05	(36)	-33.3 ^c
Iowa	0.3	0.07	(10)	0.0	South Dakota	0.2	0.09	(46)	-33.3 ^c
Kansas	0.2	0.06	(26)	-33.3 ^c	Tennessee	0.3	0.04	(20)	0.0
Kentucky	0.1 ^b	0.04	(49)	-66.7 ^c	Texas	0.3	0.02	(13)	0.0
Louisiana	0.3	0.05	(12)	0.0	Utah	0.2	0.07	(29)	-33.3 ^c
Maine	0.3	0.10	(09)	0.0	Vermont	0.2	0.10	(42)	-33.3 ^c
Maryland	0.3	0.05	(18)	0.0	Virginia	0.2	0.03	(45)	-33.3 ^c
Massachusetts	0.2	0.04	(33)	-33.3 ^c	Washington	0.2	0.04	(24)	-33.3 ^c
Michigan	0.2	0.03	(37)	-33.3 ^c	West Virginia	0.2	0.08	(25)	-33.3 ^c
Minnesota	0.2	0.04	(32)	-33.3 ^c	Wisconsin	0.2	0.04	(44)	-33.3 ^c
Mississippi	0.3	0.08	(06)	0.0	Wyoming	0.1	0.07	(50)	-66.7 ^c
Missouri	0.3	0.04	(23)	0.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXVI-13
 THYROID CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.5	0.01		
High Five States				
New Mexico	0.6	0.08	(01)	20.0 ^c
Idaho	0.6	0.09	(02)	20.0 ^c
Kansas	0.6	0.06	(03)	20.0 ^c
Louisiana	0.6	0.05	(04)	20.0 ^c
Kentucky	0.5	0.05	(05)	0.0
Low Five States				
Nevada	0.4	0.06	(47)	-20.0 ^c
Rhode Island	0.4	0.08	(48)	-20.0 ^c
D.C.	0.3	0.11	(49)	-40.0 ^c
Vermont	0.3	0.09	(50)	-40.0 ^c
Alaska	0.3	0.14	(51)	-40.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.4	0.04	(46)	-20.0 ^c	Montana	0.4	0.09	(43)	-20.0 ^c
Alaska	0.3	0.14	(51)	-40.0 ^c	Nebraska	0.4	0.07	(35)	-20.0 ^c
Arizona	0.5	0.04	(07)	0.0	Nevada	0.4	0.06	(47)	-20.0 ^c
Arkansas	0.5	0.06	(12)	0.0	New Hampshire	0.5	0.09	(13)	0.0
California	0.5	0.02	(10)	0.0	New Jersey	0.5	0.03	(06)	0.0
Colorado	0.5	0.05	(27)	0.0	New Mexico	0.6	0.08	(01)	20.0 ^c
Connecticut	0.5	0.05	(09)	0.0	New York	0.5	0.02	(17)	0.0
Delaware	0.5	0.10	(24)	0.0	North Carolina	0.4	0.03	(44)	-20.0 ^c
D.C.	0.3	0.11	(49)	-40.0 ^c	North Dakota	0.5	0.11	(25)	0.0
Florida	0.4	0.02	(41)	-20.0 ^c	Ohio	0.5	0.03	(21)	0.0
Georgia	0.4	0.03	(45)	-20.0 ^c	Oklahoma	0.4	0.05	(32)	-20.0 ^c
Hawaii	0.5	0.08	(14)	0.0	Oregon	0.4	0.05	(39)	-20.0 ^c
Idaho	0.6	0.09	(02)	20.0 ^c	Pennsylvania	0.5	0.03	(22)	0.0
Illinois	0.5	0.03	(16)	0.0	Rhode Island	0.4	0.08	(48)	-20.0 ^c
Indiana	0.5	0.04	(28)	0.0	South Carolina	0.4	0.05	(38)	-20.0 ^c
Iowa	0.4	0.05	(33)	-20.0 ^c	South Dakota	0.5	0.10	(23)	0.0
Kansas	0.6	0.06	(03)	20.0 ^c	Tennessee	0.5	0.04	(20)	0.0
Kentucky	0.5	0.05	(05)	0.0	Texas	0.5	0.02	(08)	0.0
Louisiana	0.6	0.05	(04)	20.0 ^c	Utah	0.4	0.07	(34)	-20.0 ^c
Maine	0.4	0.07	(40)	-20.0 ^c	Vermont	0.3	0.09	(50)	-40.0 ^c
Maryland	0.5	0.04	(19)	0.0	Virginia	0.4	0.03	(42)	-20.0 ^c
Massachusetts	0.4	0.04	(36)	-20.0 ^c	Washington	0.5	0.04	(18)	0.0
Michigan	0.5	0.03	(11)	0.0	West Virginia	0.5	0.06	(26)	0.0
Minnesota	0.5	0.04	(30)	0.0	Wisconsin	0.5	0.04	(29)	0.0
Mississippi	0.4	0.06	(37)	-20.0 ^c	Wyoming	0.5	0.13	(31)	0.0
Missouri	0.5	0.04	(15)	0.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XXVI-14
 THYROID CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	0.5	0.01		
High Five States				
New Mexico	0.6	0.12	(01)	20.0 ^c
Arkansas	0.6	0.09	(02)	20.0 ^c
Maine	0.6	0.13	(03)	20.0 ^c
Kansas	0.6	0.10	(04)	20.0 ^c
Arizona	0.5	0.07	(05)	0.0
Low Five States				
Alaska	0.3	0.28	(47)	-40.0 ^c
Washington	0.3	0.05	(48)	-40.0 ^c
Vermont	0.3	0.15	(49)	-40.0 ^c
Minnesota	0.3	0.06	(50)	-40.0 ^c
Montana	0.2	0.10	(51)	-60.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.5	0.07	(21)	0.0	Montana	0.2	0.10	(51)	-60.0 ^c
Alaska	0.3	0.28	(47)	-40.0 ^c	Nebraska	0.3	0.09	(46)	-40.0 ^c
Arizona	0.5	0.07	(05)	0.0	Nevada	0.4	0.09	(40)	-20.0 ^c
Arkansas	0.6	0.09	(02)	20.0 ^c	New Hampshire	0.5	0.13	(25)	0.0
California	0.5	0.03	(23)	0.0	New Jersey	0.5	0.05	(13)	0.0
Colorado	0.4	0.07	(30)	-20.0 ^c	New Mexico	0.6	0.12	(01)	20.0 ^c
Connecticut	0.5	0.08	(19)	0.0	New York	0.4	0.03	(28)	-20.0 ^c
Delaware	0.5	0.16	(15)	0.0	North Carolina	0.5	0.06	(12)	0.0
D.C.	0.4	0.19	(42)	-20.0 ^c	North Dakota	0.4	0.16	(34)	-20.0 ^c
Florida	0.4	0.03	(36)	-20.0 ^c	Ohio	0.5	0.05	(14)	0.0
Georgia	0.4	0.05	(45)	-20.0 ^c	Oklahoma	0.5	0.08	(16)	0.0
Hawaii	0.5	0.12	(26)	0.0	Oregon	0.4	0.07	(39)	-20.0 ^c
Idaho	0.5	0.13	(18)	0.0	Pennsylvania	0.5	0.04	(27)	0.0
Illinois	0.4	0.04	(38)	-20.0 ^c	Rhode Island	0.4	0.13	(37)	-20.0 ^c
Indiana	0.5	0.07	(10)	0.0	South Carolina	0.5	0.08	(20)	0.0
Iowa	0.5	0.08	(22)	0.0	South Dakota	0.5	0.17	(09)	0.0
Kansas	0.6	0.10	(04)	20.0 ^c	Tennessee	0.4	0.06	(35)	-20.0 ^c
Kentucky	0.5	0.08	(07)	0.0	Texas	0.5	0.04	(06)	0.0
Louisiana	0.4	0.07	(29)	-20.0 ^c	Utah	0.5	0.12	(08)	0.0
Maine	0.6	0.13	(03)	20.0 ^c	Vermont	0.3	0.15	(49)	-40.0 ^c
Maryland	0.4	0.06	(32)	-20.0 ^c	Virginia	0.4	0.05	(43)	-20.0 ^c
Massachusetts	0.4	0.06	(31)	-20.0 ^c	Washington	0.3	0.05	(48)	-40.0 ^c
Michigan	0.5	0.05	(11)	0.0	West Virginia	0.4	0.10	(33)	-20.0 ^c
Minnesota	0.3	0.06	(50)	-40.0 ^c	Wisconsin	0.4	0.06	(41)	-20.0 ^c
Mississippi	0.4	0.08	(44)	-20.0 ^c	Wyoming	0.5	0.21	(24)	0.0
Missouri	0.5	0.06	(17)	0.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XXVI-15
 THYROID CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	0.5	0.01		
High Five States				
New Mexico	0.7	0.11	(01)	40.0 ^c
Louisiana	0.6	0.07	(02)	20.0 ^c
Idaho	0.6	0.13	(03)	20.0 ^c
Kansas	0.6	0.09	(04)	20.0 ^c
Washington	0.6	0.06	(05)	20.0 ^c
Low Five States				
North Carolina	0.3 ^b	0.04	(47)	-40.0 ^c
Maine	0.3	0.08	(48)	-40.0 ^c
D.C.	0.3	0.13	(49)	-40.0 ^c
Alaska	0.2	0.17	(50)	-60.0 ^c
Vermont	0.2	0.11	(51)	-60.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	0.3	0.05	(46)	-40.0 ^c	Montana	0.5	0.13	(11)	0.0
Alaska	0.2	0.17	(50)	-60.0 ^c	Nebraska	0.5	0.10	(23)	0.0
Arizona	0.5	0.06	(15)	0.0	Nevada	0.4	0.08	(43)	-20.0 ^c
Arkansas	0.4	0.07	(31)	-20.0 ^c	New Hampshire	0.5	0.12	(14)	0.0
California	0.5	0.02	(12)	0.0	New Jersey	0.5	0.05	(07)	0.0
Colorado	0.5	0.07	(28)	0.0	New Mexico	0.7	0.11	(01)	40.0 ^c
Connecticut	0.5	0.07	(08)	0.0	New York	0.5	0.03	(18)	0.0
Delaware	0.4	0.12	(40)	-20.0 ^c	North Carolina	0.3 ^b	0.04	(47)	-40.0 ^c
D.C.	0.3	0.13	(49)	-40.0 ^c	North Dakota	0.4	0.14	(30)	-20.0 ^c
Florida	0.4	0.03	(37)	-20.0 ^c	Ohio	0.4	0.03	(33)	-20.0 ^c
Georgia	0.4	0.04	(38)	-20.0 ^c	Oklahoma	0.4	0.06	(41)	-20.0 ^c
Hawaii	0.5	0.12	(09)	0.0	Oregon	0.5	0.07	(29)	0.0
Idaho	0.6	0.13	(03)	20.0 ^c	Pennsylvania	0.5	0.03	(27)	0.0
Illinois	0.5	0.04	(06)	0.0	Rhode Island	0.3	0.09	(45)	-40.0 ^c
Indiana	0.4	0.05	(36)	-20.0 ^c	South Carolina	0.4	0.06	(39)	-20.0 ^c
Iowa	0.4	0.06	(35)	-20.0 ^c	South Dakota	0.4	0.13	(42)	-20.0 ^c
Kansas	0.6	0.09	(04)	20.0 ^c	Tennessee	0.5	0.05	(21)	0.0
Kentucky	0.5	0.07	(10)	0.0	Texas	0.5	0.03	(16)	0.0
Louisiana	0.6	0.07	(02)	20.0 ^c	Utah	0.3	0.08	(44)	-40.0 ^c
Maine	0.3	0.08	(48)	-40.0 ^c	Vermont	0.2	0.11	(51)	-60.0 ^c
Maryland	0.5	0.06	(19)	0.0	Virginia	0.4	0.05	(34)	-20.0 ^c
Massachusetts	0.4	0.05	(32)	-20.0 ^c	Washington	0.6	0.06	(05)	20.0 ^c
Michigan	0.5	0.04	(24)	0.0	West Virginia	0.5	0.09	(25)	0.0
Minnesota	0.5	0.06	(13)	0.0	Wisconsin	0.5	0.06	(17)	0.0
Mississippi	0.5	0.08	(26)	0.0	Wyoming	0.5	0.19	(20)	0.0
Missouri	0.5	0.05	(22)	0.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
 SE Standard error of the rate.
 PD Percent difference between state rate and total U.S. rate.


Table XXVII-13
URINARY BLADDER CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males and Females

State	Rate	SE	Rank	PD
TOTAL U.S.	4.3	0.02		
High Five States				
Maine	5.7 ^b	0.27	(01)	32.6 ^c
Delaware	5.6	0.36	(02)	30.2 ^c
Vermont	5.5	0.41	(03)	27.9 ^c
Nevada	5.5 ^b	0.24	(04)	27.9 ^c
Rhode Island	5.3	0.29	(05)	23.3 ^c
Low Five States				
Texas	3.7 ^b	0.07	(47)	-14.0
Alabama	3.6 ^b	0.12	(48)	-16.3 ^c
Mississippi	3.5 ^b	0.16	(49)	-18.6 ^c
Utah	3.4 ^b	0.20	(50)	-20.9 ^c
Hawaii	2.6 ^b	0.20	(51)	-39.5 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.6 ^b	0.12	(48)	-16.3 ^c	Montana	4.7	0.30	(14)	9.3
Alaska	4.9	0.59	(10)	14.0	Nebraska	3.7	0.20	(46)	-14.0
Arizona	4.1	0.12	(35)	-4.7	Nevada	5.5 ^b	0.24	(04)	27.9 ^c
Arkansas	3.9	0.16	(42)	-9.3	New Hampshire	5.1	0.28	(07)	18.6 ^c
California	4.1 ^b	0.05	(33)	-4.7	New Jersey	4.8 ^b	0.10	(12)	11.6
Colorado	4.0	0.15	(39)	-7.0	New Mexico	3.9	0.21	(45)	-9.3
Connecticut	4.7	0.15	(16)	9.3	New York	4.4	0.07	(23)	2.3
Delaware	5.6	0.36	(02)	30.2 ^c	North Carolina	4.0	0.10	(36)	-7.0
D.C.	4.8	0.40	(13)	11.6	North Dakota	4.0	0.32	(41)	-7.0
Florida	4.3	0.06	(27)	0.0	Ohio	5.0 ^b	0.09	(09)	16.3 ^c
Georgia	3.9 ^b	0.11	(43)	-9.3	Oklahoma	4.2	0.15	(31)	-2.3
Hawaii	2.6 ^b	0.20	(51)	-39.5 ^c	Oregon	5.1 ^b	0.16	(08)	18.6 ^c
Idaho	4.4	0.26	(24)	2.3	Pennsylvania	4.6 ^b	0.08	(17)	7.0
Illinois	4.4	0.08	(25)	2.3	Rhode Island	5.3	0.29	(05)	23.3 ^c
Indiana	4.4	0.12	(22)	2.3	South Carolina	4.0	0.14	(38)	-7.0
Iowa	4.0	0.15	(37)	-7.0	South Dakota	4.3	0.31	(26)	0.0
Kansas	4.1	0.17	(32)	-4.7	Tennessee	4.3	0.12	(29)	0.0
Kentucky	4.5	0.15	(21)	4.7	Texas	3.7 ^b	0.07	(47)	-14.0
Louisiana	3.9	0.14	(44)	-9.3	Utah	3.4 ^b	0.20	(50)	-20.9 ^c
Maine	5.7 ^b	0.27	(01)	32.6 ^c	Vermont	5.5	0.41	(03)	27.9 ^c
Maryland	4.6	0.13	(19)	7.0	Virginia	4.3	0.11	(28)	0.0
Massachusetts	5.1 ^b	0.12	(06)	18.6 ^c	Washington	4.6	0.13	(20)	7.0
Michigan	4.6	0.10	(18)	7.0	West Virginia	4.7	0.21	(15)	9.3
Minnesota	4.2	0.13	(30)	-2.3	Wisconsin	4.8 ^b	0.13	(11)	11.6
Mississippi	3.5 ^b	0.16	(49)	-18.6 ^c	Wyoming	4.0	0.41	(40)	-7.0
Missouri	4.1	0.12	(34)	-4.7					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXVII-14
URINARY BLADDER CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Males

State	Rate	SE	Rank	PD
TOTAL U.S.	7.5	0.04		
High Five States				
Vermont	9.9	0.88	(01)	32.0 ^c
Delaware	9.8	0.77	(02)	30.7 ^c
New Hampshire	9.7	0.63	(03)	29.3 ^c
Rhode Island	9.6	0.64	(04)	28.0 ^c
Maine	9.5 ^b	0.56	(05)	26.7 ^c
Low Five States				
Utah	6.4	0.43	(47)	-14.7
Mississippi	6.4	0.35	(48)	-14.7
Texas	6.3 ^b	0.14	(49)	-16.0 ^c
New Mexico	6.1 ^b	0.41	(50)	-18.7 ^c
Hawaii	4.2 ^b	0.38	(51)	-44.0 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.6	0.28	(45)	-12.0	Montana	8.0	0.61	(17)	6.7
Alaska	9.0	1.26	(08)	20.0 ^c	Nebraska	6.6	0.42	(44)	-12.0
Arizona	6.8	0.24	(39)	-9.3	Nevada	9.1	0.49	(06)	21.3 ^c
Arkansas	6.7	0.34	(41)	-10.7	New Hampshire	9.7	0.63	(03)	29.3 ^c
California	7.1 ^b	0.11	(34)	-5.3	New Jersey	8.2	0.22	(14)	9.3
Colorado	6.9	0.32	(38)	-8.0	New Mexico	6.1 ^b	0.41	(50)	-18.7 ^c
Connecticut	8.4	0.33	(11)	12.0	New York	7.8	0.14	(20)	4.0
Delaware	9.8	0.77	(02)	30.7 ^c	North Carolina	7.0	0.22	(37)	-6.7
D.C.	6.8	0.80	(40)	-9.3	North Dakota	7.6	0.69	(23)	1.3
Florida	7.3	0.12	(32)	-2.7	Ohio	9.1 ^b	0.20	(07)	21.3 ^c
Georgia	6.6 ^b	0.24	(43)	-12.0	Oklahoma	7.0	0.31	(36)	-6.7
Hawaii	4.2 ^b	0.38	(51)	-44.0 ^c	Oregon	8.8 ^b	0.34	(10)	17.3 ^c
Idaho	7.4	0.53	(27)	-1.3	Pennsylvania	8.2 ^b	0.17	(16)	9.3
Illinois	7.5	0.18	(25)	0.0	Rhode Island	9.6	0.64	(04)	28.0 ^c
Indiana	7.7	0.26	(21)	2.7	South Carolina	7.2	0.31	(33)	-4.0
Iowa	7.1	0.32	(35)	-5.3	South Dakota	8.2	0.67	(15)	9.3
Kansas	7.6	0.36	(24)	1.3	Tennessee	7.4	0.26	(30)	-1.3
Kentucky	7.6	0.31	(22)	1.3	Texas	6.3 ^b	0.14	(49)	-16.0 ^c
Louisiana	6.7	0.29	(42)	-10.7	Utah	6.4	0.43	(47)	-14.7
Maine	9.5 ^b	0.56	(05)	26.7 ^c	Vermont	9.9	0.88	(01)	32.0 ^c
Maryland	7.8	0.29	(19)	4.0	Virginia	7.4	0.25	(29)	-1.3
Massachusetts	8.9 ^b	0.26	(09)	18.7 ^c	Washington	7.8	0.26	(18)	4.0
Michigan	8.3 ^b	0.21	(13)	10.7	West Virginia	7.5	0.42	(26)	0.0
Minnesota	7.4	0.27	(28)	-1.3	Wisconsin	8.3	0.27	(12)	10.7
Mississippi	6.4	0.35	(48)	-14.7	Wyoming	6.5	0.81	(46)	-13.3
Missouri	7.3	0.25	(31)	-2.7					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.


Table XXVII-15
URINARY BLADDER CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2001-2005

Females

State	Rate	SE	Rank	PD
TOTAL U.S.	2.3	0.02		
High Five States				
D.C.	3.4	0.43	(01)	47.8 ^c
Maine	2.9	0.26	(02)	26.1 ^c
West Virginia	2.9	0.21	(03)	26.1 ^c
Rhode Island	2.8	0.28	(04)	21.7 ^c
Massachusetts	2.8 ^b	0.11	(05)	21.7 ^c
Low Five States				
Mississippi	1.7 ^b	0.14	(47)	-26.1 ^c
Alabama	1.7 ^b	0.11	(48)	-26.1 ^c
North Dakota	1.5	0.26	(49)	-34.8 ^c
Hawaii	1.5 ^b	0.19	(50)	-34.8 ^c
Utah	1.1 ^b	0.16	(51)	-52.2 ^c

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	1.7 ^b	0.11	(48)	-26.1 ^c	Montana	2.3	0.28	(19)	0.0
Alaska	1.7	0.47	(45)	-26.1 ^c	Nebraska	1.9	0.18	(44)	-17.4 ^c
Arizona	2.0	0.11	(39)	-13.0	Nevada	2.7	0.22	(08)	17.4 ^c
Arkansas	2.1	0.15	(38)	-8.7	New Hampshire	2.1	0.24	(35)	-8.7
California	2.1	0.05	(33)	-8.7	New Jersey	2.7 ^b	0.10	(07)	17.4 ^c
Colorado	2.1	0.14	(37)	-8.7	New Mexico	2.2	0.21	(28)	-4.3
Connecticut	2.3	0.14	(21)	0.0	New York	2.4	0.06	(17)	4.3
Delaware	2.7	0.33	(06)	17.4 ^c	North Carolina	2.2	0.09	(27)	-4.3
D.C.	3.4	0.43	(01)	47.8 ^c	North Dakota	1.5	0.26	(49)	-34.8 ^c
Florida	2.1	0.06	(32)	-8.7	Ohio	2.5	0.08	(11)	8.7
Georgia	2.3	0.10	(23)	0.0	Oklahoma	2.2	0.14	(26)	-4.3
Hawaii	1.5 ^b	0.19	(50)	-34.8 ^c	Oregon	2.5	0.15	(12)	8.7
Idaho	2.1	0.24	(34)	-8.7	Pennsylvania	2.4	0.07	(15)	4.3
Illinois	2.4	0.08	(18)	4.3	Rhode Island	2.8	0.28	(04)	21.7 ^c
Indiana	2.4	0.11	(16)	4.3	South Carolina	2.0	0.13	(42)	-13.0
Iowa	2.1	0.14	(31)	-8.7	South Dakota	1.7	0.26	(46)	-26.1 ^c
Kansas	1.9	0.15	(43)	-17.4 ^c	Tennessee	2.3	0.11	(22)	0.0
Kentucky	2.5	0.14	(14)	8.7	Texas	2.0 ^b	0.06	(41)	-13.0
Louisiana	2.2	0.13	(30)	-4.3	Utah	1.1 ^b	0.16	(51)	-52.2 ^c
Maine	2.9	0.26	(02)	26.1 ^c	Vermont	2.5	0.36	(13)	8.7
Maryland	2.6	0.13	(09)	13.0	Virginia	2.3	0.11	(20)	0.0
Massachusetts	2.8 ^b	0.11	(05)	21.7 ^c	Washington	2.3	0.12	(24)	0.0
Michigan	2.3	0.09	(25)	0.0	West Virginia	2.9	0.21	(03)	26.1 ^c
Minnesota	2.1	0.12	(36)	-8.7	Wisconsin	2.5	0.12	(10)	8.7
Mississippi	1.7 ^b	0.14	(47)	-26.1 ^c	Wyoming	2.2	0.41	(29)	-4.3
Missouri	2.0	0.11	(40)	-13.0					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.