

Table I-9
SEER INCIDENCE AND U.S. MORTALITY TRENDS BY PRIMARY CANCER SITE AND SEX
Blacks, 1996-2005

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.1*	-1.9*	-0.4*	-1.9*	-2.5*	-1.2*
Oral Cavity & Pharynx:	-3.0*	-3.9*	-1.1	-3.4*	-3.0*	-4.5*
Lip	-	-	-	-	-	-
Tongue	-1.8	-2.8*	0.0	-4.5*	-3.6*	-6.8*
Salivary gland	0.8	-2.8	4.1	-4.1*	-5.1*	-3.5
Floor of mouth	-8.6*	-9.4*	-	-11.0*	-9.1*	-
Gum & other oral cavity	-3.7*	-4.1	-2.6	-5.7*	-5.5*	-6.4*
Nasopharynx	-0.9	-	-	0.0	1.0	-1.5
Tonsil	-3.0	-3.3	-	-4.3*	-4.2*	-4.2
Oropharynx	-0.1	-0.9	-	-1.7*	-1.5	-1.1
Hypopharynx	-4.1	-4.4	-	-5.2*	-4.6*	-
Other oral cavity & pharynx	-	-	-	-1.5	-1.4	-2.0
Digestive System:	-0.8*	-1.0*	-0.8*	-1.6*	-1.7*	-1.7*
Esophagus	-5.3*	-5.3*	-5.0*	-4.4*	-4.6*	-3.9*
Stomach	-3.4*	-3.6*	-3.1*	-3.5*	-3.8*	-3.3*
Small intestine	0.9	0.5	0.8	-1.2	-1.5	-1.0
Colon & Rectum:	-0.9	-0.8	-0.9*	-1.7*	-1.6*	-1.9*
Colon	-0.9	-0.7	-1.0*	-	-	-
Rectum	-0.7	-1.1	-0.4	-	-	-
Anus, anal canal & anorectum	2.3	0.5	4.0*	0.2	0.1	0.8
Liver & Intrahep:	3.9*	4.3*	2.3	1.9*	2.4*	0.6
Liver	4.4*	5.0*	2.3	1.4*	2.1*	-0.5
Intrahep bile duct	-2.3	-	-	4.9*	5.2*	4.8*
Gallbladder	-1.1	-	-0.7	-1.2	-0.1	-1.6
Other biliary	5.1*	6.1	3.4*	-2.4*	-3.7	-1.5
Pancreas	-0.7	-1.3	-0.2	-0.8*	-0.9*	-0.8*
Retroperitoneum	-	-	-	-4.5	-	-
Peritoneum, omentum & mesentery	-	-	-	2.9*	-	3.7*
Other digestive system	-	-	-	1.3	1.1	1.7
Respiratory System:	-1.9*	-3.2*	0.1	-1.8*	-2.8*	0.0
Nose, nasal cavity & middle ear	-5.4*	-	-	-4.7	-2.2	-8.3*
Larynx	-3.7*	-3.4*	-4.9*	-3.3*	-3.3*	-3.0*
Lung & bronchus	-1.7*	-3.2*	0.4	-1.7*	-2.8*	0.1
Pleura	-	-	-	-9.0*	-	-
Trachea & other respiratory organs	-	-	-	-8.9*	-	-
Bones & joints	0.1	-4.2	5.1*	-1.8	-2.7*	-1.3
Soft tissue (incl heart)	-1.2	-1.1	-1.6	-4.0*	-3.1*	-4.4*
Skin (ex basal & squam):	-2.8	-6.3*	-0.2	-2.7*	-2.8*	-2.6
Melanoma of the skin	-5.3*	-9.1*	-	-1.7	-0.4	-2.8
Other non-epithelial skin	-1.0	-3.5	1.1	-3.5*	-4.0*	-2.4
Breast	-0.8*	-	-0.7*	-1.5*	-3.4*	-1.5*
Breast (<i>in situ</i>)	2.0*	-	2.0*	-	-	-

^a The APC is the Annual Percent Change over the time interval.
SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table I-9 - continued
 SEER INCIDENCE AND U.S. MORTALITY TRENDS BY PRIMARY CANCER SITE AND SEX
 Blacks, 1996-2005

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	-0.6	-	-0.4	-1.3*	-	-1.2*
Cervix uteri	-4.9*	-	-4.8*	-4.4*	-	-4.3*
Corpus uteri	1.3	-	1.4	-0.9	-	-0.7
Uterus, NOS	-	-	-	0.8	-	1.1*
Ovary ^c	0.3	-	0.5	-0.2	-	-0.1
Vagina	-0.2	-	0.4	-2.7	-	-2.7
Vulva	-0.7	-	-0.5	-0.3	-	0.0
Other female genital system	-0.2	-	-0.3	-3.9	-	-3.8
Male Genital System:	-1.9*	-2.2*	-	-3.8*	-4.0*	-
Prostate	-1.9*	-2.2*	-	-3.8*	-4.0*	-
Testis	-2.4	-2.8	-	2.4	2.5	-
Penis	-	-	-	-1.4	-1.6	-
Other male genital system	-	-	-	-	-	-
Urinary System:	1.1*	1.4*	0.3	-0.9*	-0.7	-1.1*
Urinary bladder	0.6	0.5	0.4	-1.3*	-1.0	-1.7*
Kidney & renal pelvis	1.9*	2.8*	0.4	-0.5	-0.5	-0.6
Ureter	-	-	-	-	-	-
Other urinary system	-	-	-	0.4	-	-0.1
Eye & Orbit	-	-	-	-	-	-
Brain & Nervous System: ^d	0.5	-0.4	0.9	-1.0*	-0.4	-1.8*
Brain	0.4	-1.0	1.5	-	-	-
Cranial nerves & other nervous system	-	-	-	-	-	-
Endocrine System:	5.7*	5.1*	6.0*	0.1	-0.9	0.6
Thyroid	6.4*	6.5*	6.5*	0.2	-0.2	0.5
Other endocrine & thymus	0.5	-	-	-0.1	-1.4	0.8
Lymphoma:	0.6	0.0	1.4*	-3.0*	-3.7*	-2.3*
Hodgkin lymphoma	1.9	1.1	3.1	-2.7*	-2.9*	-2.3
Non-Hodgkin lymphoma	0.4	-0.1	1.0	-3.1*	-3.7*	-2.3*
Myeloma	-0.5	0.2	-1.2	-1.8*	-2.0*	-1.7*
Leukemia:	-0.4	-1.2	0.1	-1.5*	-1.8*	-1.3*
Lymphocytic:	-0.2	-0.8	0.5	-2.3*	-2.2*	-2.9*
Acute lymphocytic	1.0	-2.1	3.9	-1.8	-1.1	-3.3
Chronic lymphocytic	-0.9	-1.2	-0.5	-2.4*	-2.2*	-3.0*
Other lymphocytic	-	-	-	-3.2	-5.3	-
Myeloid & Monocytic:	-0.8	-1.9	-0.1	-1.2*	-1.6*	-0.7
Acute myeloid	0.8	-0.7	1.6	1.1	0.8	1.3
Chronic myeloid	-4.2*	-4.5*	-4.8	-9.0*	-9.1*	-8.8*
Acute monocytic	-	-	-	-	-	-
Other myeloid & monocytic	-	-	-	7.4*	-	-
Other:	0.6	-	-	-1.0*	-1.4	-0.9
Other acute	-	-	-	-3.7*	-2.7*	-4.8*
Aleukemic, subleuk & NOS	-	-	-	0.6	-0.6	1.5*
Kaposi Sarcoma ^e	-9.6*	-9.5*	-	-	-	-
Mesothelioma ^e	-5.3	-	-	-	-	-
Ill-defined & unspecified	-4.3*	-4.3*	-4.5*	-1.2	-0.9	-1.5

^a The APC is the Annual Percent Change over the time interval.
 SEER 13 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.

* The APC is significantly different from zero (p<.05).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.