

Certificate

Standard Reference Material 4224

Beta-Ray Standard

n-Hexadecane-1-Carbon-14

This standard consists of approximately 3 grams of 99+ percent hexadecane (olefin free) containing carbon-14 labeled *n*-hexadecane in a flame-sealed glass ampoule.

The activity in disintegrations per second per gram of solution as of June 1967, was

$$*3.69_2 \times 10^2 \pm 3.0_6 \% *$$

Aliquots of the solution from which this standard was prepared, were converted to sodium carbonate-¹⁴C and compared, by liquid-scintillation counting, with the National Bureau of Standards sodium carbonate-¹⁴C standard.

The value reported above is based on 37 liquid-scintillation comparisons, and the uncertainty, $\pm 3.0_6$ percent, is the sum of the computed standard error, 0.5_5 percent, at the 99.73-percent confidence level (3.23σ), the estimated limits of systematic error, $\pm 1.0_1$ percent, and the overall uncertainty, $\pm 1.5_0$ percent, assigned to the National Bureau of Standards sodium carbonate-¹⁴C standard.

The beta-ray and gamma-ray spectra were examined with anthracene-crystal and NaI(Tl)-crystal spectrometers and no radioactive impurities were observed.

A half life of 5730 ± 40 years was adopted as the best value at the Fifth Radiocarbon Dating Conference, Cambridge, England, 1962 [Nature 195, 984 (1962)].

This standard was prepared and calibrated in the Institute for Basic Standards, Radiation Physics Division, by members of the Radioactivity Section, W. B. Mann, Chief.

WASHINGTON, D.C. 20234
August 1967

W. Wayne Meinke, Chief
Office of Standard Reference Materials