

9 Factors that Influence Prescribers' Response to Alerts at the Point-of-Care: Implications for VA Order Check Design

Alissa L. Russ, PhD
alissa.russ@va.gov

Research Scientist
Roudebush VA HSR&D Center of Excellence on
Implementing Evidence-Based Practice (CIEBP)
Regenstrief Institute, Inc.

Adjunct Assistant Professor
Purdue University College of Pharmacy

My Background

- Human Factors Engineering:
 - **design** of technology, processes, and work systems so they are compatible with **human** cognitive and physical **capabilities and limitations**

Audience Question #1

What is your primary role in the VA?

- a. informatics
- b. patient care
- c. research
- d. other
- e. work outside the VA

Outline

1. Overview of VA alerts
2. Results from field observations and interviews with VA prescribers
3. Framework for prescriber–alert interaction
4. Describe potential implications for VA alert design and medication safety

Computerized Medication Alerts

Background:

- 1.5 million preventable adverse drug events annually in U.S. (IOM, 2006)

Literature:

- Alert fatigue (Grizzle et al, 2007)
- Database analyses, surveys (Van der Sijs et al, 2006)

Study Aim:

Identify factors that influence how prescribers perceive and respond to alerts at the point-of-care

Medication Alerts (i.e. order checks)



Medication Alerts (i.e. order checks)

Order Checking

Previous adverse reaction to: PENICILLIN (LOCAL)

Accept Order Cancel Order

Medication Alerts (i.e. order checks)

Order Checks

PENICILLIN VK TAB 250MG
TAKE ONE TABLET ORALLY EVERY 6 HOURS
Quantity: 120 Refills: 0 *UNSIGNED*


[Previous adverse reaction to: PENICILLIN \(LOCAL\)](#)

Cancel Selected Order(s)

Enter justification for overriding critical order checks -

Continue

Ordering Process (simplified)



Audience Question #2:

- According to human factors science, what is the primary way that providers learn about how health IT is designed and intended to be used for patient care?
 - a. operations manual
 - b. from their peers
 - c. training sessions
 - d. software interface design
 - e. talk to the developers

Answer for Question #2:

- According to human factors science, what is the primary way that providers learn about how health IT is designed and intended to be used for patient care?
 - a. operations manual
 - b. from their peers
 - c. training sessions
 - d. software interface design
 - e. talk to the developers



Methods

- Field observations and interviews
 - Major Midwestern VA Medical Center
 - ½ day per prescriber
 - NPs, MDs, Clinical Pharmacists
- Qualitative analysis
 - Inductive, emergent themes (Campbell 2006; Patterson 2002)
 - Team consensus
 - MAXQDA software

Data Analysis

Observations	
Total hrs	102.8 hrs
Typed Notes	351 pgs
Patients	146
Alerts	320
Qualitative Analysis	
Meetings	30 (27.5 hrs)



Jason
Saleem PhD



Alan Zillich
PharmD



Sue
McManus
PhD, NP


Recruiting Results

Prescribers	N=30
Primary Care (PC)	20 prescribers: 5 PC teams 4 per team
Specialty clinics	10 prescribers 8 clinics
Expertise	18 physicians 7 nurse practitioners 5 pharmacists
Age	42 (27-63) yrs
Years in VA	10 (<1-24) yrs
Gender	14M/16F

Results: prescriber-alert interaction

- 44 themes
- 9 overarching factors
- framework

Underlying Framework



From Norman 1990 and adapted by Russ et al 2012 for prescribing

Prescriber-Alert Interaction



Norman 1990
Russ et al, *IJMI* 2012

1. Alert System Logic

- External Crosschecks:
 - Compares VA meds across VAMCs
 - Compares VA meds to entered non-VA meds
 - Perceived as strength of VA alert system
- Detection (over/under detection)
 - Identified gaps; prescribers wanted more alerts for:
 - lithium
 - thyroid and liver function
 - non-steroidal anti-inflammatory drugs (NSAIDs)
 - medications that can affect psychiatry patients

2. Alert System Redundancy

Repetition within an encounter

- Example:
 - 1: NP orders niacin; alert for niacin/pravastatin
 - 2: NP signs order, alert appears again
 - 3: NP orders pravastatin; alert for pravastatin/niacin
 - 4: NP signs order, alert appears again

– Observer noted: *We have now seen the same alert 4 times in last 10 min or less.*

Prescriber-Alert Interaction



Norman 1990
Russ et al, *IJMI* 2012

3. Alert Display

- 5 themes included:
 - Format
 - Saliency
 - Timing
 - Level of intrusiveness
 - Retrievability
- “Redesigning Medication Alerts to Support Prescriber Workflow”
 - HSR&D grant PPO #09-298
 - Coming soon to a journal near you...

4. Alert Content

Specification: *explanation of why* an alert was triggered

Order check appears:

Duplicate drug class, non- opioids [antitussives]/expectorants.
Non-VA medication guaifenesin.

MD: “I don’t even know what that means. It says expectorants, but it doesn’t say what the other [medication] is. It says non-VA medication guaifenesin, but you’re [pt 4] getting that here, right?”

Audience Question #3

Out of the 30 prescribers in the study, data from _____ prescribers indicated that there was confusion about why alerts were triggered.

- a. less than 5
- b. 5-9
- c. 10-19
- d. more than 20

Audience Question #3

Out of the 30 prescribers in the study, data from

21 prescribers indicated that there was confusion about why alerts were triggered.

- a. less than 5
- b. 5-9
- c. 10-19
- d. more than 20

Implication: a lack of specification poses a substantial barrier to resolving alerts; additional attention is needed on the clinical content presented by alerts

Prescriber-Alert Interaction



Norman 1988
Russ et al, *IJMI* 2012

5. Cognitive factors

Audience Question #4:

In this study, alerts sometimes supported awareness by providing *new* information for prescribers. Prescribers found alerts particularly helpful for:


- a. new patients
- b. new medications
- c. medications they rarely prescribed
- d. allergies documented by someone else
- e. all of the above

Answer for #4:

In this study, alerts sometimes supported awareness by providing *new* information for prescribers. Prescribers found alerts particularly helpful for:

- a. new patients
- b. new medications
- c. medications they rarely prescribed
- d. allergies documented by someone else
- e. all of the above

Prescriber-Alert Interaction



Norman 1990
Russ et al, *IJMI* 2012

6. Pharmaceutical Knowledge:

"If I'm not in the room, the doctors don't know what to do with the alert."



"I talk to the clinical pharmacist to resolve order checks. I like having the pharmacist in the room here with me."



"I'll ask the pharmacist in the clinic, 'Is this an important interaction?' ."



6. Pharmaceutical Knowledge:


- Pharmacist consultation and proximity: pharmacists consulted in real-time to help resolve alerts

“Physicians are not trained like pharmacists. We have to learn what a significant, clinically-relevant interaction is.”

- VA physician

- **Implications:**
 - Alert language, information currently inadequate
 - Universal design needed for alerts to help support prescribers with different training and levels of experience

Prescriber-Alert Interaction



Norman 1990
Russ et al, *IJMI* 2012

7. Medication Management

- Computerized Provider Order Entry (CPOE) design
 - CPOE design sometimes hindered alert resolution
 - MD orders mycophenolate
 - Duplicate drug class alert:
mycophenolate/azathioprine
 - MD overrides alert
 - MD explains: “I couldn’t discontinue the [azathioprine] because it is coming from [another VAMC]....I can only tell the patient to stop it.”
 - **Implications:** need more advanced CPOE and alert for EHR interoperability

8. Workflow

Audience Question #5:

In this study, after computer delays reached _____, prescribers began expressing frustration.

- a. 10-15 sec
- b. 16-30 sec
- c. 31-60 sec
- d. 61-90 sec




Answer for #5:

In this study, after computer delays reached _____, prescribers began expressing frustration.

- a. 10-15 sec
- b. 16-30 sec
- c. 31-60 sec
- d. 61-90 sec



Prescriber-Alert Interaction



Norman 1990
Russ et al, *IJMI* 2012

9. Alert System Reliability

- Common care practices:
 - alerts inappropriately warn against practices that apply to **broad** patient populations

Alert Trigger

- NPH* insulin/regular insulin/metformin
- Mometasone with albuterol [inhalers].
- Duplicate drug class, antiretrovirals

Prescriber Response

- “There are tons of [diabetic] patients on this combination and that is safe.”
- “We have a lot of patients on multiple inhalers.”
- “Each patient is on at least 3 antiretrovirals. The cocktail is how we treat patients.”

*Neutral Protamine Hagedorn


9. Alert System Reliability

- ▶ Common care practices:
 - alerts inappropriately warn against practices that apply to **broad** patient populations

Implications:

- Reduce alerts that conflict with evidence

Prescriber-Alert Interaction



Norman 1990
Russ et al, *IJMI* 2012

Summary: alert design implications

- ▶ Alert interface is often a barrier for prescribers, and should be addressed along with alert fatigue
- ▶ Additional work is needed to understand how to present clinical content on alerts, so that alerts aid more prescriber types
- ▶ Need to prepare alert/CPOE systems for increased EHR interoperability

Conclusions

- One of first studies to examine alerts real-time at point-of-care
- Novel framework for prescriber-alert interaction
- Findings may inform alert redesigns to enhance patient safety

“Some are critical interactions....For example, nitrates [and] phosphodiesterase inhibitors...

It has happened before where I didn't catch this interaction, but the computer did.”

- *Physician*



ELSEVIER

journal homepage: www.ijmijournal.com



Prescribers' interactions with medication alerts at the point of prescribing: A multi-method, in situ investigation of the human–computer interaction

Alissa L. Russ^{a,b,c,d,*}, Alan J. Zillich^{a,c,d}, M. Sue McManus^e, Bradley N. Doebbeling^{a,b,d,f}, Jason J. Saleem^{a,b,d,g}

^a Department of Veterans Affairs, Health Services Research and Development Center of Excellence on Implementing Evidence-Based Practice, Richard L. Roudebush Veterans Affairs Medical Center, Indianapolis, IN, United States

^b Regenstrief Institute, Inc., Indianapolis, IN, United States

^c Department of Pharmacy Practice, Purdue University College of Pharmacy, West Lafayette, IN, United States

^d Indiana University (IU) Center for Health Services & Outcomes Research, Indianapolis, IN, United States

^e Department of Nephrology, Roudebush VAMC, Indianapolis, IN, United States

^f Department of Medicine, IU School of Medicine, Indianapolis, IN, United States

^g Department of Electrical & Computer Engineering, Indiana University-Purdue University at Indianapolis (IUPUI), Indianapolis, IN, United States

Funding Support

- U.S. Department of Veterans Affairs, Veterans Health Administration
 - Health Services Research & Development (HSR&D)
- HSR&D Center of Excellence on Implementing Evidence-Based Practice (CIEBP)
 - Center grant #HFP 04-148
- Dr. Russ was supported by a VA HSR&D Associated Health Postdoctoral Fellowship
- Drs. Saleem and Zillich were supported by VA HSR&D Career Development Awards

Acknowledgements

- Study Participants
- Project Team
 - Jason Saleem, PhD
 - Alan Zillich, PharmD
 - Sue McManus, NP, PhD
 - Brad Doebbeling, MD, MSc
- Additional thanks:
 - Rich Frankel, PhD
 - Marshall Jones, PharmD
 - Kim Zipper, IT specialist
 - David Bickel, IT specialist
 - Bruce Molitoris, MD

Bibliography

- Campbell, E. M., Sittig, D. F., Ash, J. S., Guappone, K. P., & Dykstra, R. H. (2006). Types of unintended consequences related to computerized provider order entry. *J Am Med Inform Assoc, 13(5), 547-556.*
- Grizzle, A. J., Mahmood, M. H., Ko, Y., Murphy, J. E., Armstrong, E. P., Skrepnek, G. H., et al. (2007). Reasons provided by prescribers when overriding drug-drug interaction alerts. *Am J Manag Care, 13(10), 573-578.*
- Institute of Medicine of the national academies. (July 2006), Preventing medication errors. Report brief.
- Norman, D. A. (1990). *The Design of Everyday Things* (1st ed.). New York, NY: Doubleday.
- Patterson, E. S., Cook, R. I., & Render, M. L. (2002). Improving patient safety by identifying side effects from introducing bar coding in medication administration. *J Am Med Inform Assoc, 9(5), 540-553.*
- Russ AL, Zillich AJ, McManus MS, Doebbeling BD, Saleem JJ. (2012) “Prescribers’ Interactions with Medication Alerts at the Point of Prescribing: A Multi-Method, In Situ Investigation of the Human-Computer Interaction”, *International Journal of Medical Informatics, 81(4):232-243.*
- Van der Sijs, H., Aarts, J., Vulto, A., & Berg, M. (2006). Overriding of drug safety alerts in computerized physician order entry. *J Am Med Inform Assoc, 13(2), 138-147.*

Thank you!

alissa.russ@va.gov

