

Treatment Adherence

What is treatment adherence?

Treatment adherence means following your treatment regimen closely every day—taking the correct dose of each anti-HIV medication at the correct time and exactly as prescribed. Adherence is very important for successful HIV treatment.

Why is adherence important?

Adherence affects HIV treatment in two ways:

- Close adherence to an HIV treatment regimen allows anti-HIV medications to work effectively to reduce the amount of HIV in the body. Skipping medications, even occasionally, gives HIV the chance to multiply rapidly. Preventing the virus from multiplying is the best way to protect your health.
- Close adherence to an HIV treatment regimen also helps prevent **drug resistance**. Drug resistance develops when the virus mutates (changes form), becoming “resistant” to certain anti-HIV medications. One or more anti-HIV medications in a treatment regimen can become ineffective as a result of drug resistance.

Skipping medications makes it easier for drug resistance to develop. HIV can become resistant to the anti-HIV medications in a person’s current regimen or to other, similar anti-HIV medications not yet taken, limiting options for successful HIV treatment. And drug-resistant strains of HIV can be transmitted to others, too.

There are many different anti-HIV medications and treatment regimens, but studies show that **a person’s first regimen offers the best chance for long-term treatment success**. And good adherence from the start is key to successful treatment.

Why is treatment adherence sometimes difficult?

Adhering to an HIV treatment regimen can be difficult for several reasons. Some treatment regimens involve taking several pills every day—with or without food, or before or after other medications. Other factors that can make treatment adherence difficult include:

- Difficulty taking medications (such as trouble swallowing pills)
- Side effects from medications (for example, nausea or diarrhea)

- A busy schedule, shift work, or travel away from home that makes it easy to forget to take pills
- Being sick or depressed
- Alcohol or drug abuse

What can I do to adhere to my HIV treatment regimen?

Before you start treatment, be certain you’re committed to taking anti-HIV medications every day as directed. Talk to your health care provider about any issues that can make adherence difficult, including:

- Possible side effects from the anti-HIV medications in your regimen
- How other medications you take may interact with your anti-HIV medications
- Your schedule at home and at work
- Any personal issues such as depression or alcohol or drug abuse
- Lack of health insurance to pay for anti-HIV medications

Understanding issues that can make adherence difficult will help you and your health care provider select the best regimen for you. Some people find that adhering to an HIV treatment regimen becomes more difficult over time. So, every time you see your health care provider, make it a point to talk about adherence. (See the [Following an HIV Treatment Regimen](#) fact sheet for tips on adherence.)

For more information:

Contact an *AIDSinfo* health information specialist at 1-800-448-0440 or visit <http://aidsinfo.nih.gov>. See your health care provider for medical advice.

Terms Used in This Fact Sheet:

Drug resistance: When HIV mutates (changes form), causing one or more anti-HIV medications to be ineffective.

Regimen: A combination of three or more anti-HIV medications from at least two different drug classes.

Treatment adherence: Closely following an HIV treatment regimen—taking the correct dose of each anti-HIV medication at the correct time and exactly as prescribed.