

List of Tables and Figures

Chapter 1

Introduction, Summary, and Conclusions

- Table 1.1 Conclusions from previous Surgeon General's reports on the health effects of secondhand smoke exposure 4
- Table 1.2 Major conclusions of the 1986 Surgeon General's report, *The Health Consequences of Involuntary Smoking* 6
- Table 1.3 Selected major reports, other than those of the U.S. Surgeon General, addressing adverse effects from exposure to tobacco smoke 7
- Table 1.4 Four-level hierarchy for classifying the strength of causal inferences based on available evidence 10
- Figure 1.1 The determinants of exposure, dose, and biologically effective dose that underlie the development of health effects from smoking 18
- Figure 1.2 Model for socioeconomic status (SES) and secondhand smoke (SHS) exposure 22

Chapter 2

Toxicology of Secondhand Smoke

- Table 2.1 Levels of carcinogens in sidestream and secondhand cigarette smoke 31
- Table 2.2 Inhalation studies of secondhand smoke (89% sidestream smoke and 11% mainstream smoke) in A/J mice 33
- Table 2.3 Representative biomarker studies of carcinogens in persons exposed to secondhand smoke 35
- Table 2.4 Relationship of specific biomarkers of carcinogen uptake to secondhand smoke exposure 40
- Table 2.5 4-(Methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) and NNAL-glucuronide (NNAL-Gluc) in the urine of nonsmokers exposed to secondhand smoke 41

- Table 2.6 Studies of experimental atherosclerosis in animals exposed to secondhand smoke 60

- Figure 2.1 The correlation between levels of cotinine plus cotinine 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL) plus NNAL-glucuronide (NNAL-Gluc) conjugates in the urine of 74 school-age children exposed to secondhand smoke 42

- Figure 2.2 Scheme showing the steps linking secondhand smoke exposure and cancer via tobacco smoke carcinogens 43

- Figure 2.3 Metabolic pathways of benzo[a]pyrene (B[a]P) and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK) 44

- Figure 2.4 Effect of active and involuntary smoking on platelet aggregation in smokers and nonsmokers 53

- Figure 2.5 Flow-mediated (endothelium-dependent) and nitroglycerin-induced (endothelium-independent) vasodilation in human brachial arteries 56

- Figure 2.6 Coronary flow velocity changes before and after secondhand smoke exposure 57

- Figure 2.7 Secondhand smoke exposure and lipid deposits in rabbits 62

- Figure 2.8 Secondhand smoke exposure and infarct size in rats 63

Chapter 3

Assessment of Exposure to Secondhand Smoke

- Table 3.1 Outdoor air requirements for ventilation 90

- Table 3.2 Correlations between various secondhand smoke constituents as selective markers of exposures 95

- Table 3.3 Expired air carbon monoxide (CO), carboxyhemoglobin, and thiocyanate levels following exposure to secondhand smoke 102

Table 3.4	Half-life of cotinine in smokers and nonsmokers from several studies 104
Table 3.5	Analytical methods for measuring cotinine in nonsmokers 106
Table 3.6	Calculation of nicotine dosage from plasma cotinine concentrations 111
Figure 3.1	Schematic of a typical air handling unit 87
Figure 3.2	Anticipated changes in concentrations of airborne materials for various air exchange rates 88
Figure 3.3	Sensitivity of markers for secondhand smoke exposure 96
Figure 3.4	Estimates of time-varying respirable suspended particle (RSP) concentrations associated with secondhand smoke 98

**Chapter 4
Prevalence of Exposure to Secondhand Smoke**

Table 4.1	Trends in serum cotinine levels (nanograms per milliliter) of nonsmokers stratified by age, gender, race, and ethnicity, United States, 1988–2002 134
Table 4.2	Serum cotinine levels among nonsmokers aged 3 years and older, NHANES 1999–2002 136
Table 4.3	Concentrations of nicotine in homes of U.S. smokers 142
Table 4.4	Occupational exposures to nicotine among nonsmoking office workers stratified by the smoking policy in effect at the time of the measurements 146
Table 4.5	Occupational exposures to nicotine in nonoffice workplace settings among nonsmokers only, stratified by the smoking policy in effect at the time of the measurements 150
Table 4.6	Concentrations of nicotine in restaurants 152
Table 4.7	Concentrations of nicotine in bars, lounges, and other public venues 156
Figure 4.1	Trends in exposure of nonsmokers to secondhand smoke in the U.S. population, NHANES 1988–2002 133

Figure 4.2	Concentrations of nicotine in homes of U.S. smokers 141
Figure 4.3	Concentrations of nicotine away from work in 12 locations 144
Figure 4.4	Occupational exposures to nicotine among groups of nonsmoking office workers 148
Figure 4.5	Mean concentrations of nicotine in nonoffice workplace settings with different smoking policies 149
Figure 4.6	Average concentrations of nicotine in homes, offices, other workplaces, and restaurants where smoking is permitted 155

**Chapter 5
Reproductive and Developmental Effects
from Exposure to Secondhand Smoke**

Table 5.1	Findings on secondhand smoke exposure and reproductive and perinatal effects 168
Table 5.2	Potentially relevant exposure periods for reproductive and perinatal outcomes 171
Table 5.3	Studies of secondhand smoke exposure and fertility 174
Table 5.4	Studies of secondhand smoke exposure and pregnancy loss 176
Table 5.5	Studies of secondhand smoke exposure and sudden infant death syndrome (SIDS) 182
Table 5.6	Studies of secondhand smoke exposure and preterm delivery 196
Table 5.7	Summary of published literature on secondhand smoke and low birth weight (LBW) 200
Table 5.8	Studies of secondhand smoke exposure and congenital malformations 206
Table 5.9	Studies of secondhand smoke exposure and cognitive development 212
Table 5.10	Studies of secondhand smoke exposure and behavioral problems among children 218
Table 5.11	Studies of secondhand smoke exposure and children's growth 222

Table 5.12 Case-control studies of childhood cancer by cancer type 226

Chapter 6
Respiratory Effects in Children from Exposure to Secondhand Smoke

Table 6.1 Design, sample size, and recruitment criteria for studies of illness associated with parental smoking included in meta-analyses 269

Table 6.2 Unadjusted relative risks (odds ratios) of illness associated with parental smoking 274

Table 6.3 Pooled odds ratios (ORs), 95% confidence intervals (CIs), and heterogeneity tests from meta-analyses of lower respiratory illnesses associated with parental smoking 282

Table 6.4 Effects of adjusting for potential confounders of illness associated with parental smoking 284

Table 6.5 Design, sample size, and recruitment criteria of studies of illness associated with parental smoking excluded from meta-analyses 294

Table 6.6 Design, sample size, and recruitment criteria of studies of illness associated with parental smoking included in meta-analyses 297

Table 6.7 Unadjusted relative risks for updated meta-analysis of illness associated with parental smoking 301

Table 6.8 Effects of adjusting for potential confounders in each study of illness associated with parental smoking 304

Table 6.9 List of secondhand smoke exposure analyses included in the meta-analysis 312

Table 6.10 Studies of asthma prevalence associated with parental smoking 320

Table 6.11 Studies of wheeze prevalence associated with parental smoking 330

Table 6.12 Studies of cough prevalence associated with parental smoking 342

Table 6.13 Studies of phlegm and breathlessness associated with parental smoking 350

Table 6.14 Summary of pooled random effects (odds ratios) of respiratory symptoms associated with parental smoking 353

Table 6.15 Summary of pooled random effects (odds ratios) associated with parental smoking restricted to studies of children aged ≤ 11 years 354

Table 6.16 Design, sample size, and recruitment criteria for studies of asthma incidence and prognosis associated with parental smoking included in this overview 357

Table 6.17 Incidence and prognosis of asthma or wheeze in relation to parental smoking 361

Table 6.18 Unadjusted relative risks associated with parental smoking for asthma (meta-analysis of case-control studies) 365

Table 6.19 Design, sample size, and severity index for studies of asthma severity associated with parental smoking included in this overview 369

Table 6.20 Summary of studies on asthma severity associated with parental smoking 371

Table 6.21 Atopy studies of markers for exposure to secondhand smoke 377

Table 6.22 Studies relating parental smoking to skin-prick positivity in children 381

Table 6.23 Summary of pooled odds ratios (95% confidence intervals) in skin-prick positivity comparing unexposed children with children exposed to secondhand smoke at various time points 382

Table 6.24 Cross-sectional and cohort studies that used lung function level as the primary indicator of adverse effects of exposure to secondhand smoke 386

Table 6.25 Summary of pooled percentage differences in cross-sectional studies of lung function in children exposed to secondhand smoke compared with unexposed children 397

Table 6.26 Pooled percentage differences in lung function according to secondhand smoke exposure category (random effects results) 398

Table 6.27 Pooled percentage differences in lung function according to confounders adjusted for (random effects results) 399

Figure 6.1 Odds ratios for the effect of smoking by either parent on lower respiratory illnesses during infancy 279

Figure 6.2 Odds ratios for the effect of maternal smoking on lower respiratory illnesses during infancy 280

Figure 6.3 Odds ratios for the effect of paternal smoking on lower respiratory illnesses during infancy 281

Figure 6.4 Odds ratios for the effect of smoking by either parent on middle ear disease in children 307

Figure 6.5 Odds ratios for the effect of smoking by either parent on asthma prevalence 328

Figure 6.6 Odds ratios for the effect of smoking by either parent on wheeze prevalence 340

Figure 6.7 Odds ratios for the effect of smoking by either parent on cough prevalence 341

Figure 6.8 Odds ratios for the effect of smoking by either parent on phlegm and breathlessness 352

Figure 6.9 Odds ratios for the effect of maternal smoking on asthma or wheeze incidence throughout childhood (cohort studies) 363

Figure 6.10 Odds ratios for the effect of smoking by either parent on childhood asthma or wheeze prevalence (case-control studies) 367

Figure 6.11 Odds ratios for the association between parental smoking during pregnancy and infancy and skin-prick positivity 383

Figure 6.12 Odds ratios for the association between parental smoking during childhood and skin-prick positivity 383

Figure 6.13 Odds ratios for the association between parental smoking and skin-prick positivity 384

Figure 6.14 Percentage difference in the forced vital capacity (FVC) between children of smokers and children of nonsmokers in studies included in the meta-analysis 394

Figure 6.15 Percentage difference in the forced expiratory volume in 1 second (FEV₁) between children of smokers and children of nonsmokers in studies included in the meta-analysis 395

Figure 6.16 Percentage difference in the mid-expiratory flow rate (MEFR) between children of smokers and children of nonsmokers in studies included in the meta-analysis 396

Figure 6.17 Percentage difference in the end-expiratory flow rate (EEFR) between children of smokers and children of nonsmokers in studies included in the meta-analysis 397

Chapter 7
Cancer Among Adults from Exposure to Secondhand Smoke

Table 7.1 Conclusions of selected authoritative scientific bodies on the role of secondhand smoke and the risk of lung cancer among lifetime nonsmokers 424

Table 7.2 Cohort studies of the associations between adult exposure to secondhand smoke and the relative risks for lung cancer incidence and mortality among women who had never smoked 426

Table 7.3 Case-control studies by geographic area of exposure to secondhand smoke and the relative risks for lung cancer among lifetime nonsmokers 428

Table 7.4 Quantitative estimate of lung cancer risk with differing sources of exposure to secondhand smoke 436

Table 7.5 Relative risks for lung cancer associated with any workplace exposure to secondhand smoke among lifetime nonsmokers 437

Table 7.6 Dose-response relationships between workplace secondhand smoke exposure and lung cancer risk among lifetime nonsmokers 440

Table 7.7 Relative risks for lung cancer associated with exposure to secondhand smoke during childhood among lifetime nonsmokers 441

Table 7.8 Cohort studies of associations between exposures to secondhand smoke and the relative risks for breast cancer incidence and mortality among women who had never smoked 450

Table 7.9 Case-control studies of the association between exposures to secondhand smoke and relative risks for breast cancer incidence and mortality among women who had never smoked 456

Table 7.10 Pooled risk estimates and 95% confidence intervals (CI) for breast cancer meta-analysis 471

Figure 7.1 Relative risks (with 95% confidence intervals) of breast cancer associated with all sources of adult exposure to secondhand smoke 472

Figure 7.2 Relative risks (with 95% confidence intervals) of breast cancer associated with adult exposure to secondhand smoke from spouses' smoking 473

Figure 7.3 Relative risks (with 95% confidence intervals) of breast cancer associated with all sources of adult exposure to secondhand smoke among premenopausal women 474

Figure 7.4 Relative risks (with 95% confidence intervals) of breast cancer associated with all sources of adult exposure to secondhand smoke among postmenopausal women 475

Figure 7.5 Begg's funnel plot with pseudo 95% confidence limits for 21 studies of breast cancer and secondhand smoke exposure 476

Chapter 8
Cardiovascular Diseases from Exposure to Secondhand Smoke

Table 8.1 Cohort studies of secondhand smoke exposure and the risk of coronary heart disease (CHD) among nonsmokers 510

Table 8.2 Case-control studies of exposure to secondhand smoke and the risk of coronary heart disease (CHD) among nonsmokers 514

Table 8.3 Meta-analyses of secondhand smoke exposure and coronary heart disease 522

Table 8.4 Studies included in the dose-response meta-analysis and pooled results 526

Table 8.5 Epidemiologic studies of secondhand smoke exposure and stroke 528

Figure 8.1 Relative risks of coronary heart disease associated with secondhand smoke exposure among nonsmokers 524

Figure 8.2 Pooled relative risks of coronary heart disease (CHD) associated with secondhand smoke exposure among nonsmokers in various subgroups 525

Figure 8.3 Pooled relative risks of coronary heart disease associated with various levels of exposure to secondhand smoke among nonsmokers 526

Chapter 9
Respiratory Effects in Adults from Exposure to Secondhand Smoke

Table 9.1 Major conclusions from reports on adverse respiratory effects of secondhand smoke exposure in adults 540

Table 9.2 Chamber studies of exposure to secondhand smoke and acute respiratory symptoms 548

Table 9.3 Observational studies of exposure to secondhand smoke and chronic respiratory symptoms 550

Chapter 10
Control of Secondhand Smoke Exposure

Table 10.1 Summary of milestones in establishing clean indoor air policies in the United States 572

Table 10.2 Percentage of respondents aged 18 years or older who believe that secondhand smoke is harmful, by selected characteristics, United States, 1992 and 2000 589

Table 10.3 Percentage of respondents aged 18 years or older who support smoke-free restaurants, by selected characteristics, United States, 1992–2002 590

Table 10.4 Percentage of respondents aged 18 years or older who support smoke-free bars, by selected characteristics, United States, 1992–2002 591

Table 10.5 Percentage of respondents aged 18 years or older who believe smoking should be allowed in some areas of bars, by selected characteristics, United States, 1992–2002 593

Table 10.6	Percentage of respondents aged 18 years or older who support smoke-free sports arenas, by selected characteristics, United States, 1992–2002 595	nonsmoking areas in United Kingdom public houses 643
Table 10.7	Percentage of respondents aged 18 years or older who support smoke-free malls, by selected characteristics, United States, 1992–2002 596	Figure 10.1 Number of municipalities with local laws covering smoking in workplaces, restaurants, and enclosed public places, generally, 1978–2006 579
Table 10.8	Percentage of respondents aged 18 years or older who support smoke-free indoor workplaces, by selected characteristics, United States, 1992–2002 597	Figure 10.2 Cumulative number of state laws and amendments enacted for clean indoor air, 1963–2005 583
Table 10.9	Percentage of indoor workers aged 18 years or older who reported smoke-free workplace smoking policies by state, United States, 1992–2002 600	Figure 10.3 Percentage of indoor workers aged 18 years or older who reported smoke-free workplace policies, by region, United States, 1992–2002 601
Table 10.10	Percentage of indoor workers aged 18 years or older who reported smoke-free workplace policies, by geographic region and gender, United States, 1992–2002 602	Figure 10.4 Percentage of indoor workers aged 18 years or older who reported smoke-free workplace policies, by gender, United States, 1992–2002 603
Table 10.11	Percentage of indoor workers aged 18 years or older who reported smoke-free workplace policies, by age and gender, United States, 1992–2002 605	Figure 10.5 Percentage of indoor workers aged 18 years or older who reported smoke-free workplace policies, by occupational status, United States, 1992–2002 604
Table 10.12	Percentage of indoor workers aged 18 years or older who reported smoke-free workplace policies, by smoking status, United States, 1992–2002 606	Figure 10.6 Percentage of indoor workers aged 18 years or older who reported smoke-free workplace policies, by race and ethnicity, United States, 1992–2002 606
Table 10.13	Percentage of indoor workers aged 18 years or older who reported compliance with workplace smoking bans, by geographic region and occupational status, United States, 1995–2002 608	Figure 10.7 Percentage of indoor workers aged 18 years or older who reported smoke-free workplace policies, by level of education, United States, 1992–2002 607
Table 10.14	Prevalence of smoke-free households, by state, United States, 1992–2002 618	Figure 10.8 Nicotine levels measured in 10 Vancouver, British Columbia, pubs for the Heart and Stroke Foundation of British Columbia and Yukon, 1995 644
Table 10.15	Prevalence of smoke-free households, by geographic region, socioeconomic status, and household smoking status, United States, 1992–2002 619	Figure 10.9 Cumulative frequency distributions of weekly average nicotine concentrations at the desks of nonsmoking workers in open offices 646
Table 10.16	Studies assessing the effectiveness of interventions to reduce home-based secondhand smoke exposure of children 622	Figure 10.10 Cumulative frequency distributions of weekly average nicotine concentrations in nonsmokers' work areas in shops and other nonoffice settings 647
Table 10.17	Comparison of air-cleaning systems 640	Figure 10.11 Cumulative distribution of indoor PM _{2.5} concentrations in 100 United States office buildings by smoking policy 648
Table 10.18	Untransformed secondhand smoke marker concentrations ($\mu\text{g}/\text{m}^3$) for smoking and	Figure 10.12 Cumulative distribution of ventilation rates in 100 United States office buildings by smoking policy 648