

**Table 1. Evaluation of under reporting in sampled communities during the 2011 Greenland salmon fishery. The total number of Atlantic salmon (*Salmo salar*) documented by the sampling teams (salmon that have been sampled, seen but not sampled and seen and checked for an adipose fin clip only) is converted to a total whole weight and compared to the reported landings for each community.**

<b>Community (NAFO Division)</b>	<b>Number sampled</b>	<b>Number seen</b>	<b>Average sampled guttled wt (kg)</b>	<b>Average converted whole wt (kg)</b>
Ilulissat (1A)	55	20	3.94	4.37
Sisimiut (1B)	272	190	2.90	3.22
Nuuk (1D)	387	371	2.85	3.16
Qaqortoq (1F)	256	40	3.12	3.46
<b>Total</b>	<b>970</b>	<b>621</b>	<b>3.00</b>	<b>3.33</b>

<b>Community</b>	<b>Estimated whole wt sampled/seen (kg)</b>	<b>Reported landings (kg)</b>	<b>Adjusted Landings (kg)</b>	<b>Difference (kg)</b>	<b>Difference</b>
Ilulissat (1A)	328	1018	<b>1018.0</b>	0.0	0%
Sisimiut (1B)	1487	3649	<b>3649.0</b>	0.0	0%
Nuuk (1D)	2398	7977	<b>7977.0</b>	0.0	0%
Qaqortoq (1F)	1025	1039	<b>1039.0</b>	0.0	0%
<b>Total</b>	<b>5238</b>	<b>13683</b>	<b>13683.0</b>	<b>0.0</b>	<b>0%</b>