

Table 22. Estimated summer flounder recreational landings (catch types A + B1), live discard (catch type B2), and total catch (catch types A + B1 + B2) in numbers (000s), Proportional Standard Error (PSE) of the total catch estimate, and live discard (catch type B2) as a proportion of total catch. Catch type B2 uses estimates for North Carolina (NC) from NCDMF (C.Batsavage, pers. comm.)

Year	A+B1	B2	A+B1+B2	PSE (%)	B2 / (A+B1+B2)
1982	15,473	8,084	23,557	59	0.343
1983	20,996	11,026	32,022	16	0.344
1984	17,475	12,307	29,782	11	0.413
1985	11,066	2,460	13,526	15	0.182
1986	11,621	13,655	25,276	8	0.540
1987	7,865	13,472	21,337	6	0.631
1988	9,960	7,201	17,161	6	0.420
1989	1,717	908	2,625	10	0.346
1990	3,794	5,283	9,077	5	0.582
1991	6,068	9,870	15,938	5	0.619
1992	5,002	7,540	12,542	5	0.601
1993	6,494	17,741	24,235	5	0.732
1994	6,703	12,332	19,035	5	0.648
1995	3,326	13,568	16,894	5	0.803
1996	6,997	12,987	19,984	4	0.650
1997	7,167	13,854	21,021	4	0.659
1998	6,979	16,960	23,939	4	0.708
1999	4,107	17,833	21,940	5	0.813
2000	7,801	18,643	26,444	4	0.705
2001	5,294	24,049	29,343	3	0.820
2002	3,262	13,386	16,648	3	0.804
2003	4,559	15,776	20,335	4	0.776
2004	4,557	17,009	21,566	4	0.789
2005	4,036	23,135	27,171	5	0.851
2006	4,138	17,516	21,654	5	0.809
2007	3,394	20,428	23,822	5	0.858
2008	2,295	22,204	24,499	5	0.906
2009	1,910	23,749	25,659	5	0.926
2010	1,484	22,432	23,916	5	0.938
2011	1,782	20,361	22,143	7	0.920