



Northeast Fisheries Science Center Reference Document 11-15

Frequency of Whale and Vessel Collisions on the US Eastern Seaboard: Ten Years Prior and Two Years Post Ship Strike Rule

by Richard M. Pace

August 2011

Frequency of Whale and Vessel Collisions on the US Eastern Seaboard: Ten Years Prior and Two Years Post Ship Strike Rule

by Richard M. Pace

NOAA National Marine Fisheries Service
Northeast Fisheries Science Center, 166 Water St., Woods Hole, MA 02543

US DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Science Center
Woods Hole, Massachusetts

August 2011

Northeast Fisheries Science Center Reference Documents

This series is a secondary scientific series designed to assure the long-term documentation and to enable the timely transmission of research results by Center and/or non-Center researchers, where such results bear upon the research mission of the Center (see the outside back cover for the mission statement). These documents receive internal scientific review, and most receive copy editing. The National Marine Fisheries Service does not endorse any proprietary material, process, or product mentioned in these documents.

All documents issued in this series since April 2001, and several documents issued prior to that date, have been copublished in both paper and electronic versions. To access the electronic version of a document in this series, go to <http://www.nefsc.noaa.gov/nefsc/publications/>. The electronic version is available in PDF format to permit printing of a paper copy directly from the Internet. If you do not have Internet access, or if a desired document is one of the pre-April 2001 documents available only in the paper version, you can obtain a paper copy by contacting the senior Center author of the desired document. Refer to the title page of the document for the senior Center author's name and mailing address. If there is no Center author, or if there is corporate (*i.e.*, non-individualized) authorship, then contact the Center's Woods Hole Laboratory Library (166 Water St., Woods Hole, MA 02543-1026).

Editorial Treatment: To distribute this report quickly, it has not undergone the normal technical and copy editing by the Northeast Fisheries Science Center's (NEFSC's) Editorial Office as have most other issues in the NOAA Technical Memorandum NMFS-NE series. Other than the four covers and first two preliminary pages, all writing and editing have been performed by the authors listed within.

Information Quality Act Compliance: In accordance with section 515 of Public Law 106-554, the Northeast Regional Office completed both technical and policy reviews for this report. These predissemination reviews are on file at the Northeast Regional Office.

This document may be cited as:

Pace RM. 2011. Frequency of whale and vessel collisions on the US Eastern seaboard: Ten years prior and two years post ship strike rule. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 11-15; 12 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://www.nefsc.noaa.gov/nefsc/publications/>

TABLE OF CONTENTS

Introduction.....	1
Methods.....	1
Results.....	2
Conclusions.....	3
Acknowledgements.....	3
References Cited.....	4
Appendix A. Classical Statistical comparisons	9
Appendix B. Classical Change Point Analysis.....	11
Appendix C. Bayesian Change Point Analysis.....	12

INTRODUCTION

Protected species science programs are frequently asked to provide management advice based on imperfect data associated with occurrence rates of rare events such as strandings, road kills or other rarely detected mortalities. Ship strikes of large whales and right whales are such settings, and they are of particular interest because economically significant management actions have been enacted to hopefully reduce their occurrence. These measures are of unknown effectiveness while possibly causing annual industry costs ranging from tens of thousands to exceeding \$100 million (shipping regulations). Following implementation of what have been termed “the Ship Strike Rules” (Federal Register 2006) which became effective 9 December 2008, the National Marine Fisheries Service (NMFS) will likely be challenged to demonstrate the recovery benefits of these expensive conservation measures in terms of effectiveness measures (e.g. whales saved). An added question on the minds of managers, industry representatives and conservation organizations is, “How long need actions be in place before we know if they are effective?” For the analyst, this entails evaluating a Poisson process of relatively rare events for significant decreases in rates of occurrence. Data on ship strikes include a highly scrutinized time series of dates when whale mortalities that resulted from whale-ship collisions were detected during 2000-2010. With only 8 years of data prior to implementation of the Ship Strike Rule, uncertainty about the *status quo* rate will still be large. Further, whale-ship collisions that produce whale deaths will likely not be eliminated by management actions. Therefore, it is the amount by which they may have been reduced concomitant with adherence to regulations that must be investigated. Herein, I examine the timing of detected ship strikes of large whales to see whether there has been any reduction in their rate of occurrence detected ship-strike related mortalities. I also provide some advice on increasing the length of the time series after rule enactment to detect different effect sizes.

METHODS

Serious injury and mortality data for large whale stocks in the US Atlantic were evaluated for evidence of collisions with ships from necropsy and gross observations reported to the Northeast Fisheries Science Center (NEFSC) (see Glass et al. 2009 for a description). From these data, I included all reports judged to be mortalities or serious injuries (hereafter mortalities) to fin (*Balaenoptera physalus*), sei (*B. borealis*), right (*Eubalaena glacialis*) and humpback (*Megaptera novaeangliae*) whales during the period 1 January 1999-31 December 2010. Strikes of each species should resemble a Poisson process, each with its own inherent rate, and because Poisson processes are summable, events pooled across species should also resemble a Poisson process. Using the discovery date associated with each strike, I calculated the time elapsed since the previous event, which I refer to as “waiting time,” and I refer to the times since the Ship Strike Rule went into effect as event times (events occurring prior to the rule were coded as negative event times). I first examined the waiting time data relative to fits of models of exponential waiting times. Competing models included, in descending order of complexity:

1. Variable rates among years (i.e., 12 rates, 1 per year 1999-2010),
2. 2 rates, one prior to the rule and one after, and
3. A single rate.

Preliminary evaluations of similar data suggested that a more powerful approach at detecting changes may be to develop regressions of event times against order of occurrence, and to compare models with and without change points. I fit both classical linear models and their Bayesian counterparts to examine the evidence for a change in the rate of ship strikes since the implementation date of the Ship Strike Rule. Competing models included:

4. a single slope (a constant ship strike rate)
5. a fixed change point having 2 slopes on either side of the implementation date
6. 2 distinct regression models for before and after the rule, and
7. a free-floating, single change point analysis with 2 slopes on either side of an arbitrary change date, where that date was also allowed to vary and achieve the best fit.

The latter 2 models were only evaluated in the Bayesian framework. All Bayesian models were evaluated using WinBugs (ver. 1.4.3) (Lunn, et al. 2000) and were structured with broad flat priors on all parameters (Carlin and Louis 2000). Model Selection was based on DIC, an information criterion similar to AIC for likelihood models (Spiegelhalter 2002).

In addition to examining the available data on detected ship strike mortalities, I examined the potential to detect a change in rates of ship strikes using a set of simulation trials. Specifically, I estimated the mean of the exponential distribution that best fit the pre-Rule waiting times. I simulated sets (1000 each) of waiting times that would occur, if the estimated rate of occurrence of ship strikes were 66, 50 and 33% of the pre-Rule rate for 2, 5 and 7 years post implementation. I then tested the hypothesis that a change point model with rates differing before and after implementation of the rule (model 5 above) fit these simulated data better than a constant regression model (model 4 above). The percent rejections ($\alpha=0.05$) were taken as measure of power to detect a true change for the 9 combinations of 3 study durations and 3 effect sizes.

RESULTS

A total of 58 ship strikes of large whales that were deemed to be serious injuries or mortalities were included in NEFSC data during 1 Jan 1999 – 31 December 2011. These included 17 humpback, 16 fin, 21 northern right, and 4 sei whales. The most consistent evaluation of these data occurred beginning in 2000 (TVN Cole, Pers. Comm.), so I limited analysis to event times starting with the first strike in 2000 ($n=55$). A simple plot of the data gives an appearance of heterogeneity among years with 2005 appearing as a particularly nasty one (Figure 1). However, there was no statistical support for heterogeneity in event waiting times among years (Appendix A). As with most biological data, waiting times between detected ship strikes appear somewhat more variable than those associated with a simple Poisson process (ship strikes per year).

Comparing change point models for these data offered a meager amount of evidence for an increase in the time between events after rule implementation, which equates to fewer ship strike mortalities detected per annum. Based on AICc, the classical regression model with a fixed change after the rule (model 5 above) received weight of 0.75 vs. the single rate regression (model 4 above) weight of 0.25, with an estimated effect size of only 3 days longer between

strikes after rule implementation (Appendix B). Similarly, only weak distinctions were possible among Bayesian change point models with DIC values of 64.7, 63.0, 63.2, and 53.3 (Appendix C) for single slope (model 4 above), two slopes on either side of the implementation date (model 5 above), 2 distinct regression models (model 6 above), and a free-floating, single change point analysis (model 7 above), respectively (smaller values are better). The one exception was the free-floating change point model, which rather convincingly suggested that, if one change occurred in these data, it was a significant decrease in time between strikes starting in early 2004 (Appendix C; Figure 2). Using the Bayesian framework to evaluate the before and after rule model (fixed change point referred to as model 5 above), the estimated times between ship strikes were 62 days before the rule and 88 days after the rule (Figure 3). Although this effect size differed considerably from the classical framework estimates, the posterior distribution for the rate of mortalities after the ship strike rule was enacted included a relatively large amount of variance (Figure 3).

Clearly there would be more power to detect change the large that change is and the longer the period of evaluation after the rule is enacted. In my simulations, correct detections of significant changes in times between ship strikes ranged from 1% when a 33% reduction in the rate of ship strikes occurred and post-rule monitoring existed for only 2 years to a 99.7 correct detection rate when a 66% reduction in ship strikes occurred and monitoring included 7 years of data after the ship strike rule was enacted (Table 1).

CONCLUSIONS

Based on the analysis of change points, there was only weak evidence to support an increase in the time between detected ship strike mortalities of large whales on the eastern U.S. seaboard after enactment of the Ship Strike Rule. Rates of detected serious injuries and mortalities of large whales resulting from ship-whale collisions appeared to show somewhat greater variability during the 11 years evaluated than what might be expected by chance alone. The estimated size of the effect, if one existed, depended heavily on the frame work (classical regression or Bayesian MCMC) in which the time series of ship strike dates were evaluated. Due to the lack of a clear outcome from the evaluation of ship strike event times when coupled with the results of the simulation study, I suggest at least 5 years of data be evaluated prior to passing judgment on the biological effectiveness of the Ship Strike Rule.

ACKNOWLEDGEMENTS

Data used in this paper come from numerous sources. A. G. Henry and T.V. N. Cole are largely responsible for collating and often evaluating the level of evidence from a report to determine if it warrants a serious injury and were it not for their diligence and consistent treatment of reports, my evaluation would have little meaning. Determinations of causes of mortality are due in large part to a few highly skilled biologists that form a part of the stranding network and were essential in developing these data.

REFERENCES CITED

- Carlin, B.P. and T.A Louis. 2000. Bayes and empirical Bayes methods for data analysis. Second Edition. Chapman and Hall, New York, New York, USA.
- Federal Register. 2006. Endangered Fish and Wildlife; Proposed Rule to Implement Speed Restrictions to Reduce the Threat of Ship Collisions with North Atlantic Right Whales. 71:36299-36313.
- Glass, A.H, T.V.N. Cole, M. Garron. 2009. Mortality and serious injury determinations for baleen whale stocks along the United States eastern seaboard and adjacent Canadian Maritimes, 2003-2007 (2nd Edition). U.S. Department of Commerce, Northeast Fish Science Center Reference Document 09-04, 19 pp.
- Lunn, DJ, Thomas A, Best N, and Spiegelhalter D. 2000. WinBUGS -- a Bayesian modelling framework: concepts, structure, and extensibility. *Statistics and Computing* 10:325--337.
- Spiegelhalter, D. J., N. G. Best, B. P. Carlin, A. van der Linde. 2002. Bayesian measures of model complexity and fit (with discussion). [Journal of the Royal Statistical Society, Series B \(Statistical Methodology\)](#) 64: 583–639.

Table 1. Detection rates (%) of false null hypotheses for simulated times between ship strikes assuming that the rates estimated for serious injuries and mortalities detected between 1 January 2000 and 8 December 2008 were reduced as indicated.

REDUCTION IN RATE	YEARS OF POST RULE MONITORING		
	2	5	7
33%	1	50.8	65.9
50%	2.5	80.5	92.8
66%	6.1	94.6	99.7




Figure 1. Whale and ship collisions resulting in serious injuries or mortalities detected along the US Eastern seaboard 2000-2010. Graphs represent timing of events in chronological order (A) and the cumulative distribution (B) resulting from the best generalized linear model fit to time between events (model 4 above).




Figure 2. Whale and ship collisions resulting in serious injuries or mortalities detected along the US Eastern seaboard 2000-2010. Graphs depict fit resulting from a Bayesian framework used to estimate a free floating change point for the timing of events in chronological order (A) and the posterior distributions of estimated of rates (1/years between events) for change point model (model 7 above).




Figure 3. Whale and ship collisions resulting in serious injuries or mortalities detected along the US Eastern seaboard 2000-2010. Graphs depict Bayesian model fit of rate of events that included a change point fixed at Ship Strike Rule enactment date estimated for the timing of events in chronological order (A) and the posterior distributions of estimated of rates (1/years between events) for change point model (model 5 above).

APPENDIX A. CLASSICAL STATISTICAL COMPARISONS

summary(model_1, dispersion=1)

Call:

```
glm(formula = TimeBetween ~ as.factor(Year), family = Gamma,  
     data = LW_00)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-2.4383	-0.6774	-0.2990	0.4531	1.5846

Coefficients:

	<u>Estimate</u>	<u>Std. Error</u>	<u>z value</u>	<u>Pr(> z)</u>
(Intercept)	0.007576	0.004374	1.732	0.0833 .
as.factor(Year)2001	0.016562	0.010118	1.637	0.1016
as.factor(Year)2002	0.001770	0.006946	0.255	0.7989
as.factor(Year)2003	-0.002576	0.005624	-0.458	0.6470
as.factor(Year)2004	0.006154	0.007110	0.866	0.3867
as.factor(Year)2005	0.022959	0.011648	1.971	0.0487 *
as.factor(Year)2006	0.011614	0.007749	1.499	0.1339
as.factor(Year)2007	0.026907	0.014741	1.825	0.0680 .
as.factor(Year)2008	-0.001453	0.005624	-0.258	0.7961
as.factor(Year)2009	0.002424	0.006643	0.365	0.7152
as.factor(Year)2010	0.012626	0.011007	1.147	0.2513

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for Gamma family taken to be 1)

Null deviance: 59.892 on 54 degrees of freedom
Residual deviance: 41.073 on 44 degrees of freedom
AIC: 577.83

Number of Fisher Scoring iterations: 6

summary(model_2, dispersion=1)

Call:

```
glm(formula = TimeBetween ~ as.factor(Rule), family = Gamma,  
     data = LW_00)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-2.5103	-0.9778	-0.2913	0.3310	2.0157

Coefficients:

	<u>Estimate</u>	<u>Std. Error</u>	<u>z value</u>	<u>Pr(> z)</u>
(Intercept)	0.016006	0.002413	6.633	3.28e-11 ***
as.factor(Rule)1	-0.005896	0.003888	-1.516	0.129

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for Gamma family taken to be 1)

Null deviance: 59.892 on 54 degrees of freedom
Residual deviance: 57.864 on 53 degrees of freedom
AIC: 581.3

Number of Fisher Scoring iterations: 6

summary(model_3, dispersion=1)

Call:

glm(formula = TimeBetween ~ 1, family = Gamma, data = LW_00)

Deviance Residuals:

Min	1Q	Median	3Q	Max
-2.5532	-0.9853	-0.3895	0.2659	2.2856

Coefficients:

	<u>Estimate</u>	<u>Std. Error</u>	<u>z value</u>	<u>Pr(> z)</u>	
(Intercept)	0.014334	0.001933	7.416	1.20e-13	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for Gamma family taken to be 1)

Null deviance: 59.892 on 54 degrees of freedom
Residual deviance: 59.892 on 54 degrees of freedom
AIC: 581.51
Number of Fisher Scoring iterations: 6

Confidence set for the best model

Method: raw sum of model probabilities

95% confidence set:

	<u>Model</u>	<u>K</u>	<u>AICc</u>	<u>Delta_AICc</u>	<u>AICcWt</u>
intercept only	3	2	581.74	0.00	0.46
Before and After Rule	2	3	581.77	0.04	0.46
All Years	1	12	585.26	3.52	0.08

Model probabilities sum to 1

Conclusion --- Note that the AICc for intercept only model and 2 rate model are the same even though 1 parameter was added: only one rate is supported

APPENDIX B. CLASSICAL CHANGE POINT ANALYSIS

summary(model.oneslope) (Model 4)

Call:

```
glm(formula = DaysSince2 ~ count, data = LW_00)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-328.93	-112.88	-15.83	98.70	299.95

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-3041.774	41.798	-72.77	<2e-16 ***
count	64.675	1.299	49.80	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Null deviance: 59213198 on 54 degrees of freedom
Residual deviance: 1238754 on 53 degrees of freedom
AIC: 713.31

summary(model.change) (Model 5)

Call:

```
glm(formula = DaysSince2 ~ 1 + count:as.factor(Rule), data = LW_00)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-258.13	-88.46	-29.35	72.82	306.03

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-2998.680	45.336	-66.14	<2e-16 ***
count:as.factor(Rule)0	62.087	1.756	35.36	<2e-16 ***
count:as.factor(Rule)1	65.025	1.269	51.24	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Null deviance: 59213198 on 54 degrees of freedom
Residual deviance: 1140740 on 52 degrees of freedom
AIC: 710.78

AICc Comparison --- Confidence set for the best model

Method: raw sum of model probabilities

95% confidence set:

	K	AICc	Delta_AICc	AICcWt
Change After Rule	4	711.58	0.0	0.75
One Slope	3	713.78	2.2	0.25

Conclusion - with an evidence ratio of 3:1, the change point is somewhat preferred, but the estimated difference in rates before and after the Rule (62 vs 65 days between) was small.

APPENDIX C. BAYESIAN CHANGE POINT ANALYSIS

(smaller DIC indicate BETTER fit)

<u>Model</u>	<u>DIC</u>
Free Change point	53.287
Fixed Change point	63.012
2 Regressions	63.170
1 Slope	64.706

Conclusion - Fixed change point is slightly preferred over a constant rate. Free change point is much preferred over the rest which indicates some unidentified heterogeneity is dominant over any rate change that might have occurred post-Rule.

Procedures for Issuing Manuscripts in the *Northeast Fisheries Science Center Reference Document (CRD) Series*

Clearance

All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If any author is not a federal employee, he/she will be required to sign an "NEFSC Release-of-Copyright Form." If your manuscript includes material from another work which has been copyrighted, then you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

For more information, NEFSC authors should see the NEFSC's online publication policy manual, "Manuscript/abstract/webpage preparation, review, and dissemination: NEFSC author's guide to policy, process, and procedure," located in the Publications/Manuscript Review section of the NEFSC intranet page.

Organization

Manuscripts must have an abstract and table of contents, and (if applicable) lists of figures and tables. As much as possible, use traditional scientific manuscript organization for sections: "Introduction," "Study Area" and/or "Experimental Apparatus," "Methods," "Results," "Discussion," "Conclusions," "Acknowledgments," and "Literature/References Cited."

Style

The CRD series is obligated to conform with the style contained in the current edition of the United States Government Printing Office Style Manual. That style manual is silent on many aspects of scientific manuscripts. The CRD series relies more on the CSE Style Manual. Manuscripts should be prepared to conform with these style manuals.

The CRD series uses the American Fisheries Society's guides to names of fishes, mollusks, and decapod

crustaceans, the Society for Marine Mammalogy's guide to names of marine mammals, the Biosciences Information Service's guide to serial title abbreviations, and the ISO's (International Standardization Organization) guide to statistical terms.

For in-text citation, use the name-date system. A special effort should be made to ensure that all necessary bibliographic information is included in the list of cited works. Personal communications must include date, full name, and full mailing address of the contact.

Preparation

Once your document has cleared the review process, the Editorial Office will contact you with publication needs – for example, revised text (if necessary) and separate digital figures and tables if they are embedded in the document. Materials may be submitted to the Editorial Office as files on zip disks or CDs, email attachments, or intranet downloads. Text files should be in Microsoft Word, tables may be in Word or Excel, and graphics files may be in a variety of formats (JPG, GIF, Excel, PowerPoint, etc.).

Production and Distribution

The Editorial Office will perform a copy-edit of the document and may request further revisions. The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages of the document.

Once both the PDF (print) and Web versions of the CRD are ready, the Editorial Office will contact you to review both versions and submit corrections or changes before the document is posted online.

A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the document online.

Research Communications Branch
Northeast Fisheries Science Center
National Marine Fisheries Service, NOAA
166 Water St.
Woods Hole, MA 02543-1026

**MEDIA
MAIL**

Publications and Reports of the Northeast Fisheries Science Center

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of living marine resources for the benefit of the nation through their science-based conservation and management and promotion of the health of their environment." As the research arm of the NMFS's Northeast Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (*e.g.*, anonymously-peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own media. Currently, there are three such media:

NOAA Technical Memorandum NMFS-NE -- This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review and most issues receive technical and copy editing.

Northeast Fisheries Science Center Reference Document -- This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review and most issues receive copy editing.

Resource Survey Report (formerly *Fishermen's Report*) -- This information report is a regularly-issued, quick-turnaround report on the distribution and relative abundance of selected living marine resources as derived from each of the NEFSC's periodic research vessel surveys of the Northeast's continental shelf. This report undergoes internal review, but receives no technical or copy editing.

TO OBTAIN A COPY of a *NOAA Technical Memorandum NMFS-NE* or a *Northeast Fisheries Science Center Reference Document*, either contact the NEFSC Editorial Office (166 Water St., Woods Hole, MA 02543-1026; 508-495-2350) or consult the NEFSC webpage on "Reports and Publications" (<http://www.nefsc.noaa.gov/nefsc/publications/>). To access *Resource Survey Report*, consult the Ecosystem Surveys Branch webpage (<http://www.nefsc.noaa.gov/femad/ecosurvey/mainpage/>).

ANY USE OF TRADE OR BRAND NAMES IN ANY NEFSC PUBLICATION OR REPORT DOES NOT IMPLY ENDORSEMENT.