






SPECIES: SURFCLAM - OCEAN QUAHOG



























































































Note: Selected fleets include Rows 2, 4-8, 13, 14, 16, 19-24, and 26-36; these represent fleets where discards were estimated in the majority of years.



FLEET: Otter Trawl OPEN all NE sm (Row 7)

Note: Selected fleets include Rows 2, 4-8, 13, 14, 16, 19-24, and 26-36; these represent fleets where discards were estimated in the majority of years.




FLEET: Otter Trawl OPEN all NE lg (Row 8)

Note: Selected fleets include Rows 2, 4-8, 13, 14, 16, 19-24, and 26-36; these represent fleets where discards were estimated in the majority of years.

FLEET: Otter Trawl, Ruhle OPEN all NE lg (Row 13)

SBRM 2009

SBRM 2010





Note: Selected fleets include Rows 2, 4-8, 13, 14, 16, 19-24, and 26-36; these represent fleets where discards were estimated in the majority of years.

FLEET: Otter Trawl, Haddock Separator OPEN all NE Ig (Row 14)


SBRM 2009

SBRM 2010







Note: Selected fleets include Rows 2, 4-8, 13, 14, 16, 19-24, and 26-36; these represent fleets where discards were estimated in the majority of years.




FLEET: Gillnet OPEN all MA sm (Row 19)







Note: Selected fleets include Rows 2, 4-8, 13, 14, 16, 19-24, and 26-36; these represent fleets where discards were estimated in the majority of years.




SBRM 2011


Total








Note: Selected fleets include Rows 2, 4-8, 13, 14, 16, 19-24, and 26-36; these represent fleets where discards were estimated in the majority of years.



FLEET: Scallop Dredge AA GEN MA all (Row 27)


Note: Selected fleets include Rows 2, 4-8, 13, 14, 16, 19-24, and 26-36; these represent fleets where discards were estimated in the majority of years.




SBRM 2011


Total

Discards
























Note: Selected fleets include Rows 2, 4-8, 13, 14, 16, 19-24, and 26-36; these represent fleets where discards were estimated in the majority of years.



FLEET: Scallop Dredge OPEN LIM NE all (Row 34)

Note: Selected fleets include Rows 2, 4-8, 13, 14, 16, 19-24, and 26-36; these represent fleets where discards were estimated in the majority of years.

FLEET: Mid-water Trawl OPEN all MA all (Row 35)






Figure 8A. Precision (coefficient of variation, CV) of estimated discards for each of the 14 SBRM species groups by selected fleet for all SBRM years (2009 through 2011). The 24 selected fleets are: Rows 2, 4-8, 16, 19-24, and 26-36. Each point represents a separate species group and SBRM year. Red circles denote New England fleets and black stars denote Mid-Atlantic fleets. Dash line represents 30 % CV. See Table 14 for fleet abbreviations.




Figure 8B. Precision (coefficient of variation, CV) of estimated discards for each of the SBRM individual species by selected fleet for all SBRM years (2009 through 2011). The 24 selected fleets are: Rows 2, 4-8, 16, 19-24, and 26-36. Each point represents a separate species and SBRM year. Red circles denote New England fleets and black stars denote Mid-Atlantic fleets. Dash line represents 30 % CV. See Table 14 for fleet abbreviations.




Figure 9A. Precision (coefficient of variation, CV) of estimated discards for 9 of the 14 SBRM species groups by selected fleet for all SBRM years (2009 through 2011). The 24 selected fleets are: Rows 2, 4-8, 16, 19-24, and 26-36. Each point represents a separate fleet and SBRM year. Red circles denote New England fleets and black stars denote Mid-Atlantic fleets. Dash line represents 30 % CV.




Figure 9B. Precision (coefficient of variation, CV) of estimated discards for each of the individual species that comprise the 14 SBRM species groups by selected fleet for all SBRM years (2009 through 2011). The 24 selected fleets are: Rows 2, 4-8, 16, 19-24, and 26-36. Each point represents a separate fleet and SBRM year. Red circles denote New England fleets and black stars denote Mid-Atlantic fleets. Dash line represents 30 % CV.

