

Publications

- Bacheler NM, Bailey KM, Ciannelli L, Bartolino V, Chan K-S (2009) Density-dependent, landscape, and climate effects on spawning distribution of walleye pollock *Theragra chalcogramma* Mar. Ecol. Prog. Ser. 391: 1-12.
- Bacheler NM, Ciannelli L, Bailey KM, Duffy-Anderson JT (2010) Spatial and temporal patterns of walleye pollock (*Theragra chalcogramma*) spawning in the eastern Bering Sea inferred from egg and larval distributions. Fish. Oceanogr. 19: 107-120.
- Boldt JL, Rooper CN (2009) Juvenile Pacific ocean perch abundance, condition, and diet in the Aleutian Islands. Fish. Bull. 170:278-285.
- Chan K-S, Zhang T, Bailey KM (2010) Otolith biochronology reveals factors underlying dynamics in marine fish larvae. Mar. Ecol. Prog. Ser. 412: 1-10.
- Collins WT, McConnaughey RA (1998) Acoustic classification of the sea floor to address essential fish habitat and marine protected area requirements. Pages 369-377 in Proceedings of the 1998 Canadian Hydrographic Conference, Victoria, B.C.
- Harris PM, Johnson SW, Holland LG, Neff AD, Thedinga JF, Rice SD (2005) Hydrocarbons and fisheries habitat in Berners Bay, Alaska: baseline monitoring associated with the Kensington Gold Mine. U.S. Dep. Commer., NOAA Proc. Rep., AFSC 2005-06. 44 p.
- Harris PM, Neff SD, Johnson SW, Thedinga JF (2008) Eelgrass habitat and faunal assemblages in the City and Borough of Juneau, Alaska. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-182. 46 p.
- Hoff GR (2008) A nursery site of the Alaska skate (*Bathyraja parmifera*) in the eastern Bering Sea. Fish. Bull. 106: 233-244.
- Hoff GR (2009a) Embryo developmental events and the egg case of the Aleutian skate *Bathyraja aleutica* (Gilbert) and the Alaska skate *Bathyraja parmifera* (Bean). J. Fish. Biol. 74: 483-501
- Hoff GR (2009b) Skate *Bathyraja* spp. egg predation in the eastern Bering Sea. J. Fish. Biol. 74: 250-269.
- Hurst TP, Ryer CH, Ramsey JA, Haines SA (2007) Divergent foraging strategies of three co-occurring north Pacific flatfishes. Mar. Biol. 151:1087-1098.

- Hurst TP, Abookire AA, Knoth B (2010) Quantifying thermal effects on contemporary growth variability to predict responses to climate change in northern rock sole (*Lepidopsetta polyxystra*) Can. J. Fish. Aquat. Sci. 67: 97-107.
- Johnson SW, Murphy ML, Csepp DJ (2003a) Distribution, habitat, and behavior of rockfishes, *Sebastes* spp., in nearshore waters of southeastern Alaska: observations from a remotely operated vehicle. *Envir. Biol. Fish.* 66:259-270.
- Johnson SW, Murphy ML, Csepp DJ, Harris PM, Thedinga JF (2003b) A survey of fish assemblages in eelgrass and kelp habitats of southeastern Alaska. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-139. 39 p.
- Johnson SW, Neff AD, Thedinga JF (2005a) An atlas on the distribution and habitat of common fishes in shallow nearshore waters of southeastern Alaska. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-157. 89 p.
- Johnson SW, Thedinga JF (2005b) Fish use and size of eelgrass meadows in southeastern Alaska: a baseline for long-term assessment of biotic change. *Northwest Sci.* 79:141-155.
- Johnson SW, Thedinga JF, Munk KM (2008) Distribution and use of shallow-water habitats by Pacific sand lances in southeastern Alaska. *Trans. Am. Fish. Soc.* 137:1455-1463.
- Johnson SW, Thedinga JT, Neff AD, Hoffman CA (2010) Fish fauna in nearshore waters of a barrier island in the western Beaufort Sea, Alaska. NOAA Tech. Memo. NMFS-AFSC-210.
- Johnson SW, Thedinga JT, Neff AD, Harris PM, Lindeberg MR, Maselko JM, Rice SD (2010) Fish assemblages in nearshore habitats of Prince William Sound, Alaska. *Northwest Science* 84: 266-280.
- Johnson SW, Thedinga JF, Neff AD (in press) Invasion by saffron cod *Eleginus gracilis* into nearshore habitats of Prince William Sound, Alaska. *Mar. Ecol. Prog. Ser.*
- Johnson SW (2010) Nearshore fish assemblages in upper Cook Inlet, Alaska. Trip report for August 18, 2009. NOAA Alaska Fisheries Science Center, Juneau, Alaska.
- Johnson SW (2010) Nearshore fish assemblages in Resurrection Bay, Alaska. Trip report for August 20, 2009. NOAA Alaska Fisheries Science Center, Juneau, Alaska.
- Johnson SW (2010) Nearshore fish assemblages in Passage Canal, Alaska. Trip report for August 21, 2009. NOAA Alaska Fisheries Science Center, Juneau, Alaska.
- Johnson SW (2010) Nearshore fish assemblages in Upper Cook Inlet, Alaska. Trip report for July 14, 2010. NOAA Alaska Fisheries Science Center, Juneau, Alaska.

- Johnson SW (2010) Fish fauna in nearshore waters of Port Valdez, Alaska. Trip report for 12-13 July, 2010. NOAA Alaska Fisheries Science Center, Juneau, Alaska.
- Laurel BJ, Stoner AW, Hurst TP (2007a) Density-dependent habitat selection in marine flatfish: the dynamic role of ontogeny and temperature. *Mar. Ecol. Prog. Ser.* 338:183-192.
- Laurel BJ, Stoner AW, Ryer CH, Hurst TP, Abookire AA (2007b) Comparative habitat associations in juvenile Pacific cod and other gadids using seines, baited cameras and laboratory techniques. *J. Exp. Mar. Biol. Ecol.* 351: 42-55.
- Laurel BJ, Ryer CH, Knoth B, Stoner AW (2009) Temporal and ontogenetic shifts in habitat use of juvenile Pacific cod (*Gadus macrocephalus*). *J. Exp. Mar. Biol. Ecol.* 377: 28-35.
- Laurel BJ, Ryer CH, Spencer M, Iseri P, Knoth B, Stoner A. 2012. Effects of natural and anthropogenic disturbance on polychaete worm tubes and age-0 flatfish distribution. *Mar. Ecol. Prog. Ser.* 46: 193-203.
- Marlow MS, Stevenson AJ, Chezar H, McConnaughey RA (1999) Tidally generated sea-floor lineations in Bristol Bay, Alaska. *Geo-Marine Letters* 19: 219-226.
- McConnaughey RA, Mier K, Dew CB (2000) An examination of chronic trawling effects on soft-bottom benthos of the eastern Bering Sea. *ICES J. Mar. Sci.* 57: 1377-1388.
- McConnaughey RA, Syrjala SE, Dew CB (2005) Effects of chronic bottom trawling on the size structure of soft-bottom benthic invertebrates. Pages 425-437 in P. W. Barnes and J. P. Thomas, editors. *Benthic habitats and the effects of fishing*. American Fisheries Society, Symposium 41, Bethesda, Maryland.
- McConnaughey RA, Syrjala SE (2009) Statistical relationships between the distributions of groundfish and crabs in the eastern Bering Sea and processed returns from a single-beam echosounder. *ICES J. Mar. Sci.* 66: 1425-1432.
- McConnaughey, RA, Olson JV, Sigler MF (2009) Alaska Fisheries Science Center Essential Fish Habitat Data Inventory. AFSC Processed Rep. 2009-01, 40 p.
- Murphy ML, Johnson SW, Csepp DJ (2000) A comparison of fish assemblages in eelgrass and adjacent non-eelgrass habitats near Craig, Alaska. *Alaska Fish. Res. Bull.* 7:11-21.
- Murphy ML, Johnson SW (1999) Distribution of salmon in relation to eelgrass during seaward migration. Proc. of the 19th NorthEast Pacific Pink and Chum Salmon Workshop. March 3-5, 1999, Juneau, AK. pp.-58-63.

- Preston JM, Christney AC, Collins WT, McConnaughey RA, Syrjala SE (2004) Considerations in large-scale acoustic seabed characterization for mapping benthic habitats. ICES CM 2004/T:13, 8 p.
- Reynolds BF, Powers SP, Bishop MA (2010) Application of Acoustic Telemetry to Assess Residency and Movements of Rockfish and Lingcod at Created and Natural Habitats in Prince William Sound. PlosOne 5(8):e12130
- Rooper, CN, Boldt JL, Zimmermann M (2007) An assessment of juvenile Pacific ocean perch (*Sebastes alutus*) habitat use in a deepwater nursery. Est. Coast. Shelf Sci. 75:371-380.
- Rooper CN, Zimmermann M (2007) A bottom-up methodology for integrating underwater video and acoustic mapping for seafloor substrate classification. Cont. Shelf Res. 27:947-957.
- Rooper CN, Hoff GR, De Robertis A (2010) Assessing habitat utilization and rockfish (*Sebastes* spp.) biomass on an isolated rocky ridge using acoustics and stereo image analysis. Can. J. Fish. Aquat. Sci. 67(10): 1658–1670.
- Rooper CN, Wilkins ME, Rose CS, Coon C. 2011. Modeling the impacts of bottom trawling and the subsequent recovery rates of sponges and corals in the Aleutian Islands, Alaska. Cont. Shelf Res. 31: 1827-1834.
- Rooper CN, Boldt JL, Batten S, Gburski C. 2012. Growth and production of Pacific ocean perch (*Sebastes alutus*) in nursery habitats of the Gulf of Alaska. Fish. Oceanogr. 21: 415-429.
- Rose C (2006) Development and evaluation of trawl groundgear modifications to reduce damage to living structure in soft bottom areas. Available NOAA Alaska Fisheries Science Center 7600 Sand Point Way NE, Seattle WA 98115.
- Ryer CH, Hurst TP (2008) Indirect predator effects upon age-0 northern rock sole *Lepidopsetta polyxystra*: growth suppression and temporal reallocation of feeding. Mar. Ecol. Prog. Ser. 357:207-212.
- Ryer CH, Stoner AW, Spencer ML, Abookire AA (2007) Presence of larger flatfish modifies habitat preference by age-0 northern rock sole *Lepidopsetta polyxystra*. Mar. Ecol. Prog. Ser. 342:227-238.
- Ryer CH, Laurel BJ, Stoner AW (In press) Testing the shallow water refuge hypothesis in flatfish nurseries. Mar. Ecol. Prog. Ser. Doi: 10.3354/meps08732

- Stoner AW, Spencer ML, Ryer CH (2007) Flatfish-habitat associations in Alaska nursery grounds: use of continuous video records for multi-scale spatial analysis. *J. Sea Res.* 57:137-150.
- Stoner AW, Laurel BJ, Hurst TP (2008) Using a baited camera to assess relative abundance of juvenile Pacific cod: Field and laboratory trials. *J. Exp. Mar. Biol. Ecol.* 354: 202-211.
- Thedinga, J. F., Johnson SW, Mortensen DG (2006) Habitat, age, and diet of a forage fish in southeastern Alaska: Pacific sandfish (*Trichodon trichodon*). *Fish. Bull.* 104:631-637.
- Thedinga JF, Johnson SW, Neff AD (2011) Diel differences in fish assemblages in nearshore eelgrass and kelp habitats in Prince William Sound, Alaska. *Environ. Biol. Fish.* 90: 61-70.
- von Szalay PG (1998) The feasibility of using single beam seabed classification systems to identify and quantify slope rockfish habitat in the Gulf of Alaska. MS Thesis. University of Washington, Seattle, WA, 158 pp.
- von Szalay PG, McConnaughey RA (2002) The effect of slope and vessel speed on the performance of a single beam acoustic seabed classification system. *Fish. Res. (Amst.)* 56: 99-112.
- von Szalay PG, McConnaughey RA (2002) Erratum to: "The effect of slope and vessel speed on the performance of a single beam acoustic seabed classification system". *Fish. Res. (Amst.)* 56: 97.
- Williams K, Rooper CN, Towler R (2010) Use of stereo camera systems for assessment of rockfish abundance in untrawlable areas and for recording pollock behavior during midwater trawls. *Fish. Bull.* 108:352– 362.
- Yeung C, McConnaughey RA (2008) Using acoustic backscatter from a sidescan sonar to explain fish and invertebrate distributions: a case study in Bristol Bay, Alaska. *ICES J. Mar. Sci.* 65: 242–254.
- Yeung C, Yang M-S, McConnaughey RA (2010) Polychaete community in southeastern Bering Sea in relation to groundfish distribution and diet. *J. Mar. Biol. Assoc. U.K.* 90: 903-917.
- Zhang T, Bailey KM, Chan K-S (2010) Recruitment forecast models of walleye pollock (*Theragra chalcogramma*) fine-tuned from juvenile survey data, predator abundance and environmental phase shifts. *Mar. Ecol. Prog. Ser.* 417: 237-248.

Web sites

Fish Atlas: <http://www.fakr.noaa.gov/habitat/fishatlas/>

ShoreZone: <http://www.fakr.noaa.gov/maps/szintro.htm>

Pacific Ocean Perch:

http://www.afsc.noaa.gov/race/groundfish/habitat/POP/pop_intro.htm