

Figure 1c. Comparison of **bluefish discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1d. Comparison of **bluefish discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1e. Comparison of **bluefish** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1f. Comparison of **bluefish** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1g. Comparison of **herring discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1h. Comparison of **herring discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1i. Comparison of **herring discards** (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1j. Comparison of **herring discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1k. Comparison of **red crab discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1l. Comparison of **red crab discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1m. Comparison of **red crab** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1n. Comparison of **red crab** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1o. Comparison of **scallop discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1p. Comparison of **scallop discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1q. Comparison of **scallop** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1r. Comparison of **scallop** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1s. Comparison of **squid-butterfish-mackerel discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1t. Comparison of **squid-butterfish-mackerel discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1u. Comparison of **squid-butterfish-mackerel** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1v. Comparison of **squid-butterfish-mackerel** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1w. Comparison of **monkfish discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1x. Comparison of **monkfish discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1y. Comparison of **monkfish** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1z. Comparison of **monkfish** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1aa. Comparison of **NE multispecies (large mesh) discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1bb. Comparison of **NE multispecies (large mesh) discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1cc. Comparison of **NE multispecies (large mesh)** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1dd. Comparison of **NE multispecies (large mesh)** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1ee. Comparison of **NE multispecies (small mesh) discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1ff. Comparison of **NE multispecies (small mesh) discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1gg. Comparison of **NE multispecies (small mesh)** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1hh. Comparison of **NE multispecies (small mesh)** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1ii. Comparison of **skates discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1jj. Comparison of **skates discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1kk. Comparison of **skates** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1ll. Comparison of **skates** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1mm. Comparison of **spiny dogfish discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1nn. Comparison of **spiny dogfish discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 100. Comparison of **spiny dogfish** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1pp. Comparison of **spiny dogfish** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1qq. Comparison of **fluke-scup-black sea bass discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (; sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1rr. Comparison of **fluke-scup-black sea bass discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1ss. Comparison of **fluke-scup-black sea bass** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1tt. Comparison of **fluke-scup-black sea bass** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1uu. Comparison of **surf clams/quahogs discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1vv. Comparison of **surf clams/quahogs discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1ww. Comparison of **tilefish discards** (pounds) and **trip duration** (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1xx. Comparison of **tilefish discards** (pounds) and **kept weight** of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure 1yy. Comparison of **tilefish** discards (pounds) and **trip duration** (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure 1zz. Comparison of **tilefish** discards (pounds) and **kept weight** of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.