

ABC overages from 2006 to 2008

	Catch	TAC	ABC	ABC Overage	% Over ABC	OFL (bold - specific OFL) 1, 2
2006 Western AI Pacific Ocean Perch	5,506	5,084	5,372	134	2%	17,600
2006 Central GOA Pacific Ocean Perch	8,285	7,418	7,418	867	12%	8,806
2006 West Yakutat Pacific Ocean Perch	1,259	1,101	1,101	158	14%	3,190
2006 Central GOA Northern Rockfish	3,985	3,608	3,608	377	10%	7,673
2007 Western AI Pacific Ocean Perch	7,824	7,720	7,720	104	1%	26,100
2007 Eastern AI Pacific Ocean Perch	5,101	4,970	4,970	131	3%	26,100
2007 Western GOA Pacific Ocean Perch	4,430	4,244	4,244	186	4%	4,976
2007 West Yakutat Pacific Ocean Perch	1,242	1,140	1,140	102	9%	8,922
2007 Western GOA Shortraker Rockfish	194	153	153	41	27%	1,124
2008 Bering Sea Greenland Turbot	1,916	1,750	1,750	166	9%	15,600
2008 AI Greenland Turbot	824	790	790	34	4%	15,600
2008 BSAI Rougheye Rockfish	212	202	202	10	5%	269
2008 630 Pollock	14,629	13,640	13,640	989	7%	95,940
2008 Western GOA Shortraker Rockfish	133	120	120	13	11%	1,197
2008 Western GOA Thornyhead Rockfish	274	267	267	7	3%	2,540

1/Most OFLs are management area wide except POP in the Western and Central GOA and BSAI Rougheye Rockfish

2/ bold OFL where OFL = ABC =TAC. These (area-wide) OFLs have more than one (sub-area) TAC associated with them. POP in the Central and Western GOA each have their own OFLs. Rougheye rockfish specifications are set for the entire BSAI.

Source: NMFS In-Season Management