




North Pacific Fishery
Management Council

Groundfish Species Profiles

Biology • Management • Catch History • Economics • Assessment • Fishery



For over 30 years, the North Pacific Fishery Management Council has provided responsible stewardship of the groundfish resources under its jurisdiction, resulting in sustainable and profitable fisheries off Alaska. The foundation for this success is the scientifically based annual catch limits that are established for each target groundfish stock, species, or species complex. The NMFS Alaska Fisheries Science Center provides the necessary scientific information, ranging from basic research data on life history parameters to fishery independent surveys and rigorous stock assessments. These stock assessments are peer reviewed by the BSAI and GOA Groundfish Plan Teams and the Scientific and Statistical Committee. Using this information, the Council establishes total allowable catch levels that do not exceed biologically sustainable catch limits set by the scientists. All catch accrues towards the total allowable catch levels, and catches are closely monitored by the NMFS Alaska Regional Office during the season based on data from mandatory electronic reporting by vessels and processing plants, and a comprehensive observer program.

This publication was developed to provide the public with readily available and accessible information about the groundfish species managed by the Council. For more information on the Council's management program, I invite you to visit the website at www.alaskafisheries.noaa.gov/npfmc

Dave Witherell
Deputy Director, NPFMC

This 2011 publication was prepared by David Witherell (NPFMC) and Megan Peterson (NPFMC intern, University of Alaska Fairbanks) based on the annual Stock Assessment and Fishery Evaluation (SAFE) reports, which are assembled by the groundfish plan teams and include contributions from numerous assessment authors. Front cover image of shortraker rockfish coming up on a longline courtesy of Julianne Curry, Petersburg Vessel Owners Association. Back cover image of yellowfin sole and rock sole on deck courtesy of Carwyn Hammond, AFSC. Special thanks to those who provided editorial revisions and suggestions to improve the report: Jane DiCosimo, Phil Rigby, Jon Heifetz, Dana Hanselman, Kalei Shotwell, Chris Lunsford, Cindy Tribuzio, Sandra Lowe, Tom Wilderbuer, Buck Stockhausen, Teresa D'mar, Dave Clausen, and Grant Thompson.

Species Profiles

Bering Sea/Aleutian Islands

Pollock.....	1
Pacific cod.....	3
Sablefish.....	5
Yellowfin sole.....	7
Arrowtooth flounder and Kamchatka flounder.....	9
Rock sole.....	11
Greenland turbot.....	13
Alaska plaice.....	14
Flathead sole.....	15
Other flatfish.....	16
Pacific ocean perch.....	17
Northern rockfish.....	19
Rougheye/Blackspotted rockfish.....	20
Shortraker rockfish.....	21
Other rockfish.....	22
Atka mackerel.....	23
Squid.....	24
Sculpins.....	25
Skates.....	26
Sharks.....	27
Octopus.....	28

Diana Evans, NPFMC



Megan Peterson, UAF



Gulf of Alaska

Pollock.....	29
Pacific cod.....	31
Sablefish.....	33
Shallow water flatfish.....	35
Deepwater flatfish.....	37
Rex sole.....	38
Arrowtooth flounder.....	39
Flathead sole.....	40
Pacific ocean perch.....	41
Northern rockfish.....	43
Shortraker rockfish.....	44
Other slope rockfish.....	45
Pelagic shelf rockfish.....	46
Rougheye/Blackspotted rockfish.....	47
Demersal shelf rockfish.....	48
Thornyhead rockfish.....	49
Atka mackerel.....	50
Longnose skate, Big skate, and Other Skates.....	51
Sharks.....	53
Squid.....	55
Octopus.....	56
Sculpins.....	57

Carwyn Hammond, AFSC





AFSC, NOAA Fisheries



Common Acronyms


ABC	Acceptable Biological Catch
ACL	Annual Catch Limit
AFA	American Fisheries Act
AI	Aleutian Islands
AP	Advisory Panel
ADF&G	Alaska Department of Fish and Game
AFSC	Alaska Fisheries Science Center
BSAI	Bering Sea and Aleutian Islands
CDQ	Community Development Quota
CP	Catcher Processor
CV	Catcher Vessel
EBS	Eastern Bering Sea
ESA	Endangered Species Act
F/V	Fishing Vessel
FMP	Fishery Management Plan
GOA	Gulf of Alaska
IFQ	Individual Fishing Quotas
LLP	License Limitation Program
MSA	Magnuson-Stevens Fishery Conservation and Management Act
MSST	Minimum Stock Size Threshold
MSY	Maximum Sustainable Yield
mt	Metric Ton
NMFS	National Marine Fisheries Service
NPFMC	North Pacific Fishery Management Council
OFL	Overfishing Level
POP	Pacific ocean perch
PSC	Prohibited species catch
QS	Quota Share
SAFE	Stock Assessment and Fishery Evaluation
SSC	Scientific and Statistical Committee
TAC	Total allowable catch

Regulatory Areas



Forward

Strict annual catch limits for every target fishery have proven an effective management tool for achieving sustainable fisheries. In the North Pacific, a rigorous process in place for over 30 years ensures that annual quotas are set at conservative, sustainable levels for each of our managed groundfish stocks. Below is a brief summary of the process for setting annual catch limits for Gulf of Alaska groundfish (comprised of 134 species) Bering Sea and Aleutian Islands groundfish (comprised of 131 species).



Three reference points are used for management of groundfish fisheries in the North Pacific. The overfishing level (OFL) is the catch limit which should never be exceeded. It is based on the fishing mortality rate associated with producing the maximum sustainable yield on a continuing basis. The acceptable biological catch (ABC) is the annual sustainable catch limit, and is set lower than the OFL. The buffer between these reference points allows for scientific uncertainty in single species stock assessments, ecosystem considerations, and operational management of the fishery. The total allowable catch (TAC) is the annual catch target that incorporates economic considerations and management uncertainty. The fishery management plans prescribe that TAC may equal but never exceed ABC, such that $TAC \leq ABC < OFL$. The sum of TACs for all groundfish stocks must also remain within the optimum yield range defined in the FMP. In the BSAI, the upper limit is 2 million mt, which can be constraining. TAC may be set lower than ABC for a variety of reasons, such as to remain under the 2 million mt optimum yield limit; to increase a rebuilding rate or address other conservation issues; to limit incidental bycatch; or to account for state water removals. Fisheries are managed in-season to achieve the TACs without exceeding the ABC or OFL. All catch taken in directed fisheries or caught incidentally in other fisheries, whether retained or discarded, accrues towards the TAC.

The optimum yield range defined in the FMP. In the BSAI, the upper limit is 2 million mt, which can be constraining. TAC may be set lower than ABC for a variety of reasons, such as to remain under the 2 million mt optimum yield limit; to increase a rebuilding rate or address other conservation issues; to limit incidental bycatch; or to account for state water removals. Fisheries are managed in-season to achieve the TACs without exceeding the ABC or OFL. All catch taken in directed fisheries or caught incidentally in other fisheries, whether retained or discarded, accrues towards the TAC.

Catch Limit Control Rules for North Pacific Groundfish.

Tier 1: Reliable point estimates of B and B_{MSY} and pdf of F_{MSY} .

- 1a) Stock status: $B/B_{MSY} > 1$
 $F_{OFL} = mA$, the arithmetic mean of the pdf
 $F_{ABC} \leq mH$, the harmonic mean of the pdf
- 1b) Stock status: $\alpha < B/B_{MSY} \leq 1$
 $F_{OFL} = mA \times (B/B_{MSY} - \alpha)/(1 - \alpha)$
 $F_{ABC} \leq mH \times (B/B_{MSY} - \alpha)/(1 - \alpha)$
- 1c) Stock status: $B/B_{MSY} \leq \alpha$
 $F_{OFL} = 0$; $F_{ABC} = 0$

Tier 2: Reliable point estimates of B , B_{MSY} , F_{MSY} , $F_{35\%}$, and $F_{40\%}$.

- 2a) Stock status: $B/B_{MSY} > 1$
 $F_{OFL} = F_{MSY}$
 $F_{ABC} \leq F_{MSY} \times (F_{40\%}/F_{35\%})$
- 2b) Stock status: $\alpha < B/B_{MSY} \leq 1$
 $F_{OFL} = F_{MSY} \times (B/B_{MSY} - \alpha)/(1 - \alpha)$
 $F_{ABC} \leq F_{MSY} \times (F_{40\%}/F_{35\%}) \times (B/B_{MSY} - \alpha)/(1 - \alpha)$
- 2c) Stock status: $B/B_{MSY} \leq \alpha$
 $F_{OFL} = 0$; $F_{ABC} = 0$

Tier 3: Reliable point estimates of B , $B_{40\%}$, $F_{35\%}$, and $F_{40\%}$.

- 3a) Stock status: $B/B_{40\%} > 1$
 $F_{OFL} = F_{35\%}$; $F_{ABC} \leq F_{40\%}$
- 3b) Stock status: $\alpha < B/B_{40\%} \leq 1$
 $F_{OFL} = F_{35\%} \times (B/B_{40\%} - \alpha)/(1 - \alpha)$
 $F_{ABC} \leq F_{40\%} \times (B/B_{40\%} - \alpha)/(1 - \alpha)$
- 3c) Stock status: $B/B_{40\%} \leq \alpha$
 $F_{OFL} = 0$; $F_{ABC} = 0$

Tier 4: Reliable point estimates of B , $F_{35\%}$, and $F_{40\%}$.

$$F_{OFL} = F_{35\%}; F_{ABC} \leq F_{40\%}$$

Tier 5: Reliable point estimates of B and natural mortality rate M .

$$F_{OFL} = M; F_{ABC} \leq 0.75 \times M$$


Tier 6: Reliable catch history from 1978 through 1995.

$$OFL = \text{the average catch, unless an alternative value is established by the SSC.}$$


$$ABC \leq 0.75 \times OFL$$

The catch limits are specified annually through an established public process. The annual process of determining OFL and ABC specifications begins with the assignment of each stock to one of six “tiers” based on the availability of information about that stock. Stocks in Tier 1 have the most information, and those in Tier 6, the least. Application of a control rule for each tier prescribes the resulting OFL and maximum ABC for each stock. For many groundfish stocks, the estimate of $F_{40\%}$ is used as a surrogate for F_{ABC} . $F_{40\%}$ is the fishing mortality rate at which the spawning biomass per recruit is reduced to 40% of its value in the equivalent unfished stock. The control rules for Tiers 1-3 also provide for automatic rebuilding, because if a stock falls below target biomass levels, ABC and OFL are drastically reduced.


Scientists prepare an assessment of the status of each stock (or stock complex), and include alternate model simulations and tier assignments to arrive at recommendations for OFLs and ABCs. The Groundfish Plan Teams compile these assessments into Stock Assessment and Fishery Evaluation (SAFE) reports, develop their own OFL and ABC recommendations (which may or may not agree with the stock assessment author), and present this information to the Council and its Scientific and Statistical Committee (SSC) and Advisory Panel (AP). The SSC is responsible for setting the Council's OFL and ABC limits, using the SAFE reports and Plan Team recommendations. The SSC retains the flexibility to adjust ABC and OFL values from the control rule, based on factors such as multispecies interactions, ecosystem considerations, and additional scientific uncertainty. The Council then sets the TAC levels at or below the ABC levels, incorporating recommendations from the Advisory Panel and industry stakeholders.




Gulf of Alaska Species Complex History




Bering Sea Aleutian Islands Species Complex History







Groundfish stock groupings for establishing catch limits have evolved over time as new scientific information has become available and new markets have developed for certain species. The original fishery management plans set catch limits for the few major target species (e.g., pollock, cod, sablefish), with the remaining species managed in a few complex groups (e.g., flounders, rockfish, other species). Over time, with new information and new fisheries developing, species were separated out from the complexes and assigned their own catch limits. Currently, there are nearly 50 separate single species groundfish stocks or species complexes that are assigned annual catch limits. For many of these stocks, catch limits are further subdivided into each regulatory area as a precautionary measure to prevent disproportionate exploitation rates in small areas, in case the stock consists of multiple populations.



Bering Sea and Aleutian Islands




Gulf of Alaska



The Council's conservative catch limit policies, combined with favorable environmental conditions, have resulted in abundant fish stocks and sustainable fisheries. No groundfish stock is overfished or undergoing overfishing. Further, most stocks are well above target biomass levels that produces maximum sustainable yield (B_{msy}).


The total catch and species composition of the catch has remained relatively stable since 1976 with the formation of the Council and development of the fishery management program. Prior to 1976, fisheries were only minimally regulated by bilateral agreements between the U.S. and foreign nations with fishing fleets off Alaska (Japan, USSR, South Korea, and Taiwan). Very high catches of yellowfin sole, rockfish, and pollock were taken during this time. Catches and targets began to stabilize with the development of the U.S. fishing fleet through joint ventures in the 1980s. By the time the U.S. fishery was fully developed in 1991, the catch composition was more dependent on the TAC limits than on certain species being targeted. The variability in total groundfish catch in the Bering Sea and Gulf of Alaska is now due mainly to changes in pollock biomass and resulting changes in annual catch limits.

Bering Sea & Aleutian Islands Groundfish Catch Limits 1981-2010



Catch Specifications

At each December meeting, the Council specifies catch limits for a two year period, which when implemented (in early March) supersede the limits that were set the prior year to start the fishery (which opens January 1). For example, the adjacent specification tables adopted by the Council in December 2010 will be implemented for 2011 and 2012 fisheries, effectively replacing the catch limits that were recommended in December 2009. The catch limits specified for the start of the 2012 season will be superseded by those limits set by the Council in December 2011. The 2-year cycle allows for the use of the most recent biological information in the stock assessment while eliminating any potential delay or gap in setting the second year's limits.



Bering Sea & Aleutian Islands								
Catch Specifications for 2011-2012 BSAI Groundfish								
Species	Area	2011			2012			
		OFL	ABC	TAC	OFL	ABC	TAC	
Pollock	EBS	2,450,000	1,270,000	1,252,000	3,170,000	1,600,000	1,253,658	
	AI	44,500	36,700	19,000	50,400	41,600	19,000	
	Bogoslof	22,000	156	150	22,000	156	150	
Pacific cod	BSAI	272,000	235,000	227,950	329,000	281,000	229,608	
Sablefish	BS	3,360	2,850	2,850	3,080	2,610	2,610	
	AI	2,250	1,900	1,900	2,060	1,740	1,740	
Yellowfin sole	BSAI	262,000	239,000	196,000	266,000	242,000	197,660	
Greenland turbot	Total	7,220	6,140	5,050	6,760	5,750	4,950	
	BS	n/a	4,590	3,500	n/a	4,300	3,500	
	AI	n/a	1,550	1,550	n/a	1,450	1,450	
Arrowtooth flounder	BSAI	186,000	153,000	25,900	191,000	157,000	25,900	
Kamchatka flounder	BSAI	23,600	17,700	17,700	23,600	17,700	17,700	
Northern rock sole	BSAI	248,000	224,000	85,000	243,000	219,000	85,000	
Flathead sole	BSAI	83,300	69,300	41,548	82,100	68,300	41,548	
Alaska plaice	BSAI	79,100	65,100	16,000	83,800	69,100	16,000	
Other flatfish	BSAI	19,500	14,500	3,000	19,500	14,500	3,000	
Pacific Ocean perch	BSAI	36,300	24,700	24,700	34,300	24,700	24,700	
	BS	n/a	5,710	5,710	n/a	5,710	5,710	
	EAI	n/a	5,660	5,660	n/a	5,660	5,660	
	CAI	n/a	4,960	4,960	n/a	4,960	4,960	
	WAI	n/a	8,370	8,370	n/a	8,370	8,370	
Northern rockfish	BSAI	10,600	8,670	4,000	10,400	8,330	4,000	
Blackspotted/Rougheye Rockfish	BSAI	549	454	454	563	465	465	
Shortraker rockfish	EBS/EAI	n/a	234	234	n/a	240	240	
	CAI/WAI	n/a	220	220	n/a	225	225	
	BSAI	524	393	393	524	393	393	
Other rockfish	BSAI	1,700	1,280	1,000	1,700	1,280	1,000	
	BS	n/a	710	500	n/a	710	500	
	AI	n/a	570	500	n/a	570	500	
Atka mackerel	Total	101,000	85,300	53,080	92,200	77,900	48,593	
	EAI/BS	n/a	40,300	40,300	n/a	36,800	36,800	
	CAI	n/a	24,000	11,280	n/a	21,900	10,293	
	WAI	n/a	21,000	1,500	n/a	19,200	1,500	
	BSAI	2,620	1,970	425	2,620	1,970	425	
Squid	BSAI	n/a	n/a	n/a	n/a	n/a	n/a	
Other species	BSAI	n/a	n/a	n/a	n/a	n/a	n/a	
Skate	BSAI	37,800	31,500	16,500	37,200	31,000	16,500	
Shark	BSAI	1,360	1,020	50	1,360	1,020	50	
Octopus	BSAI	528	396	150	528	396	150	
Sculpin	BSAI	58,300	43,700	5,200	58,300	43,700	5,200	
Total	BSAI	3,954,111	2,534,729	2,000,000	4,731,995	2,911,610	2,000,000	

Gulf of Alaska							
Catch Specifications for 2011-2012 GOA Groundfish							
Stock/Assemblage	Area	2011			2012		
		OFL	ABC	TAC	OFL	ABC	TAC
Pollock	W (61)	27,031	27,031		34,932	34,932	
	C (62)	37,365	37,365		48,293	48,293	
	C (63)	20,235	20,235		26,155	26,155	
	WYAK	2,339	2,339		3,024	3,024	
	Subtotal	118,030	86,970	86,970	151,030	112,404	112,404
	EYAK/SEO	12,326	9,245	9,245	12,326	9,245	9,245
	Total	130,356	96,215	96,215	163,356	121,649	121,649
Pacific Cod	W	30,380	22,785		27,370	20,528	
	C	53,816	40,362		48,484	36,363	
	E	2,604	1,953		2,346	1,759	
	Total	102,600	86,800	65,100	92,300	78,200	58,650
Sablefish	W	1,620	1,620		1,484	1,484	
	C	4,740	4,740		4,343	4,343	
	WYAK	1,990	1,990		1,818	1,818	
	SEO	2,940	2,940		2,700	2,700	
Total	13,340	11,290	11,290	12,232	10,345	10,345	
Shallow-water Flatfish	W	23,681	4,500		23,681	4,500	
	C	29,999	13,000		29,999	13,000	
	WYAK	1,228	1,228		1,228	1,228	
	EYAK/SEO	1,334	1,334		1,334	1,334	
Total	67,768	56,242	20,062	67,768	56,242	20,062	
Deep-water Flatfish	W	529	529		541	541	
	C	2,919	2,919		3,004	3,004	
	WYAK	2,083	2,083		2,144	2,144	
	EYAK/SEO	774	774		797	797	
Total	7,823	6,305	6,305	8,046	6,486	6,486	
Rex sole	W	1,517	1,517		1,490	1,490	
	C	6,294	6,294		6,184	6,184	
	WYAK	868	868		853	853	
	EYAK/SEO	886	886		869	869	
Total	12,499	9,565	9,565	12,279	9,396	9,396	
Arrowtooth Flounder	W	34,317	8,000		33,975	8,000	
	C	144,559	30,000		143,119	30,000	
	WYAK	22,551	2,500		22,327	2,500	
	EYAK/SEO	11,723	2,500		11,606	2,500	
	Total	251,068	213,150	43,000	248,576	211,027	43,000
Flathead Sole	W	17,442	2,000		17,960	2,000	
	C	28,104	5,000		28,938	5,000	
	WYAK	2,064	2,064		2,125	2,125	
	EYAK/SEO	1,523	1,523		1,568	1,568	
Total	61,412	49,133	10,587	63,202	50,591	10,693	

Stock/Assemblage	Area	2011			2012		
		OFL	ABC	TAC	OFL	ABC	TAC
Pacific Ocean Perch	W	3,221	2,798		3,068	2,665	2,665
	C	11,948	10,379		11,379	9,884	9,884
	WYAK		1,937			1,845	1,845
	SED		1,883			1,793	1,793
	Total	4,397	3,820	3,820	4,188	3,638	3,638
Northern Rockfish	W		2,573	2,573		2,446	2,446
	C		2,281	2,281		2,168	2,168
	Total	5,784	4,854	4,854	5,498	4,614	4,614
Shortraker	W		134	134		134	134
	C		325	325		325	325
	E		455	455		455	455
	Total	1,219	914	914	1,219	914	914
Other Slope Rockfish	W		212	212		212	212
	C		507	507		507	507
	WYAK		276	276		275	275
	EYAK/SEO		2,757	200		2,757	200
Total	4,881	3,752	1,195	4,881	3,751	1,194	
Pelagic Shelf Rockfish	W		611			570	570
	C		3,052	3,052		2,850	2,850
	WYAK		407	407		380	380
	EYAK/SEO		684	684		638	638
Total	5,570	4,754	4,754	5,387	4,438	4,438	
Rougheye and Blackspotted Rockfish	W		81	81		81	81
	C		868	868		868	868
	E		363	363		363	363
	Total	1,579	1,312	1,312	1,579	1,312	1,312
Demersal Rockfish	Total	479	300	300	479	300	300
Thornhead Rockfish	W		425	425		425	425
	C		637	637		637	637
	E		708	708		708	708
	Total	2,360	1,770	1,770	2,360	1,770	1,770
Atka mackerel	Total	6,200	4,700	2,000	6,200	4,700	2,000
Big Skate	W		598	598		598	598
	C		2,049	2,049		2,049	2,049
	E		681	681		681	681
	Total	4,438	3,328	3,328	4,438	3,328	3,328
Longnose Skate	W		81	81		81	81
	C		2,009	2,009		2,009	2,009
	E		762	762		762	762
	Total	3,803	2,852	2,852	3,803	2,852	2,852
Other skates	Total	2,791	2,093	2,093	2,791	2,093	2,093
Squid	GOA-wide	1,530	1,148	1,148	1,530	1,148	1,148
Sharks	GOA-wide	8,262	6,197	6,197	8,262	6,197	6,197
Octopus	GOA-wide	1,272	954	954	1,272	954	954
Sculpins	GOA-wide	7,328	5,496	5,496	7,328	5,496	5,496
Total		723,928	590,121	318,288	743,421	603,990	335,078



Megan Peterson, UAF

Walleye Pollock

Biology: Walleye pollock *Theragra chalcogramma* is the most abundant fish species in the Bering Sea. In the Eastern Bering Sea (EBS), pollock are found throughout the water column and adults are concentrated along the outer continental shelf. Seasonal migrations occur from overwintering areas along the outer shelf to shallower waters to spawn. Pollock feed on copepods, euphausiids (krill) and fish (primarily juvenile pollock) and are prey for other fish, marine mammals and seabirds.



Diana Stram, NPFMC




For More Information:

J. Ianelli, S. Barbeaux, T. Honkalehto, S. Kotwicki, K. Aydin, and N. Williamson. 2010. Assessment of Walleye Pollock in the Eastern Bering Sea. http://www.afsc.noaa.gov/re_fm/stocks/assessments.htm

Pollock is a relatively fast growing and short lived species. They begin to recruit to the fishery at age 4 and longevity extends to 12 years or more. Annual natural mortality is estimated at 25% (M=0.30). Most fish reach maturity between ages 3 and 5. Females produce 60,000 to 400,000 pelagic eggs. Peak spawning occurs in the in the southeastern BS and eastern AI along the outer continental shelf in late February. Smaller spawning aggregations also occur in the northern Bering Sea in mid-late April.

Fishery Management: The U.S. manages pollock as 3 separate stocks; the Eastern Bering Sea stock (Unimak Pass to the US-Russia Convention line), the Aleutian Islands stock (the Aleutian Islands shelf region from 170°W to the US-Russia Convention line), and the Central Bering Sea - Bogoslof Island stock.

The American Fisheries Act (1998) established eligibility to participate in the BSAI pollock fishery and settled the contentious inshore/offshore allocation issue by establishing permanent allocations of pollock quota among sectors. CDQ groups are allocated 10% of EBS pollock TAC. The remaining TAC is divided up as follows; catcher vessels delivering inshore (50%), catcher processors offshore (40%) and catcher vessels delivering to motherships (10%). The 2004 Appropriations Act established that the non-CDQ pollock fishery in the AI is fully allocated to the Aleut Corporation, for the purpose of economic development in Adak, with a percentage allocated to vessels 60 feet or less in length overall..



The EBS pollock fishery has been redistributed spatially and seasonally to reduce the potential competition for prey with the endangered western stock of Steller sea lions, with fishery exclusion zones around sea lion rookeries and reductions in TAC taken from critical habitat zones. TACs have also been divided into separate seasons since 2000; the “A-season” (Jan-Apr) and the “B-season” (Jun-Oct).

Catch History: Fisheries for Bering Sea pollock developed in 1964, and catches increased rapidly in the early 1970s and peaked in 1972 at 1.9 million mt. Early 1980s joint ventures were phased out by the domestic fleet by 1991. The international zone or “Donut Hole” also supported significant harvests of pollock through 1987, followed by a

sharp decline and a fishing moratorium for the international zone beginning in 1993.

Stock Assessment: The EBS pollock assessment is based on a statistical age-structured model that incorporates fishery data and fishery independent data from annual bottom trawl surveys and biennial acoustic trawl surveys. EBS pollock fall under Tier 1b of the ABC/OFL control rules. The 2011 age 3+ biomass in the EBS is estimated at 9,620,000 mt and B_{msy} is equal to 1,948,000 mt. Catch specifications for 2011 are as follows; OFL=2,447,000, mt, ABC=1,267,000 mt, TAC=1,252,000 mt. The AI pollock ABC =36,700 mt and the Bogoslof ABC = 156 mt.

Biomass of EBS pollock declined steadily from 2004-2009 due to poor recruitment from the 2000-2005 year classes. The biomass is now increasing with recruitment of above average 2006 and 2008 year-classes.


Total catches, pre-season catch specifications, and exploitable biomass of age 3+ Walleye Pollock in the EBS 1976-2011 (in mt).

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1976	1,177,822	-	-	-	-
1977	978,370	950,000	-	-	-
1978	979,431	950,000	950,000	-	-
1979	935,714	950,000	1,100,000	-	-
1980	958,280	1,000,000	1,300,000	-	-
1981	973,502	1,000,000	1,300,000	-	-
1982	955,964	1,000,000	1,300,000	-	-
1983	981,450	1,000,000	1,300,000	-	-
1984	1,092,055	1,200,000	1,300,000	-	-
1985	1,139,676	1,200,000	1,300,000	-	-
1986	1,141,993	1,200,000	1,300,000	-	-
1987	859,416	1,200,000	1,300,000	-	-
1988	1,228,721	1,300,000	1,500,000	-	6,500,000
1989	1,229,600	1,340,000	1,340,000	-	5,300,000
1990	1,455,193	1,280,000	1,450,000	-	5,843,800
1991	1,195,646	1,300,000	1,676,000	-	6,667,146
1992	1,390,331	1,300,000	1,490,000	1,770,000	6,190,000
1993	1,326,601	1,300,000	1,340,000	1,340,000	5,900,000
1994	1,329,350	1,330,000	1,330,000	1,590,000	8,020,000
1995	1,264,245	1,250,000	1,250,000	1,500,000	8,080,000
1996	1,192,778	1,190,000	1,190,000	1,460,000	7,360,000
1997	1,124,430	1,130,000	1,130,000	1,980,000	6,120,000
1998	1,101,165	1,110,000	1,110,000	2,060,000	5,820,000
1999	989,816	992,000	992,000	1,720,000	7,040,000
2000	1,132,707	1,139,000	1,139,000	1,680,000	7,700,000
2001	1,387,194	1,400,000	1,842,000	3,536,000	10,060,000
2002	1,480,195	1,485,000	2,110,000	3,530,000	9,800,000
2003	1,490,899	1,491,760	2,330,000	3,530,000	11,100,000
2004	1,480,543	1,492,000	2,560,000	2,740,000	11,000,000
2005	1,483,286	1,478,500	1,960,000	2,100,000	8,410,000
2006	1,486,435	1,485,000	1,930,000	2,090,000	8,050,000
2007	1,354,097	1,394,000	1,394,000	1,640,000	6,360,000
2008	990,566	1,000,000	1,000,000	1,440,000	4,357,000
2009	810,731	815,000	815,000	977,000	6,240,000
2010	809,238	813,000	813,000	918,000	4,620,000
2011	-	1,252,000	1,267,000	2,447,000	9,620,000

¹Catch data current through November 2010.

²TAC, ABC and OFL data from Federal Register Harvest Specifications.

³Biomass from annual SAFE report projections issued the previous year.



Fishery: The BSAI pollock fishery is prosecuted by relatively large vessels using pelagic trawls. A total of 89 catcher vessels and 17 catcher processors participated in the 2008 fishery. The A-season fishery is focused in the southeast portion of the EBS and targets pre-spawning pollock. Roe, fillets and surimi are the main product forms of the A-season fishery, and approximately 40% of the TAC is caught during the A-season. The B-season fishery takes the remaining 60% of the quota and is distributed over the outer shelf edge of the Bering Sea extending to the Russian border.

Economics: Pollock fishery products include whole fish, head and gut (H&G), roe, deep-skin fillets, other fillets, surimi, minced fish, and fish meal. In 2009, ex-vessel catch was \$293 million for BSAI pollock. Production was 357,390 mt for all pollock products in Alaska, with a gross value of \$1 billion. Surimi products comprised approximately 25% of the gross value of pollock products, roe comprised around 17%, and fillets about 44% of the gross value.

Ecosystem Components: Pollock are an important prey for fish, seabirds, and marine mammals (including Steller sea lions) in the BSAI.



Diana Evans, NPFMC

Pacific Cod

Biology: Pacific cod *Gadus macrocephalus* is a demersal species found in the eastern BS, the AI and GOA south to California. Pacific cod are distributed over the continental shelf at depths from shoreline to 500 m. Mature fish tend to concentrate on the outer continental shelf and prefer muddy or sandy soft sediment substrate. Juvenile Pacific feed primarily on small invertebrates and euphausiids, whereas adults feed on fish such as juvenile pollock, and invertebrates such as polychaetes, amphipods and crangonid shrimp. Predators of Pacific cod include adult Pacific cod, Pacific halibut, salmon sharks and Steller sea lions.

Pacific cod are a relatively fast growing and short lived fish. Longevity can extend to 19 years. The size at 50% maturity is 58 cm (about 5 years). Females are highly fecund and can produce more than 1 million eggs. Adults form spawning aggregations from January to May in the BS. Natural mortality is estimated at $M=0.34$. Pacific cod begin to recruit to the fisheries at age 3 and are 50% recruited by ages 4-5.



For More Information:

G. Thompson, J. Ianelli, and R. Lauth. 2010. Assessment of Pacific Cod Stock in the Eastern Bering Sea and Aleutian Islands Area.

<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Catch History: Pacific cod were taken by Japanese longline and trawl fisheries beginning in the early 1960s. Vessels from the USSR entered the fishery in 1971. Japanese and Russian fisheries harvested around 50,000 mt annually in the 1970s. Joint ventures became more prevalent in the early 1980s until they were entirely phased out by the domestic fleet a few years later. Catches have remained fairly stable since 1988, averaging 193,000 mt annually.




Jackie Patt, UAF

Fishery Management: Like most other groundfish, 10.7% of the TAC is allocated to CDQ fisheries.

Since 2007 with implementation of Amendment 85, the remaining TAC is allocated among sectors as follows: 1.4% to jig gear; 2% to hook and line/pot catcher vessels < 60', 0.2% to hook and line/pot catcher vessels ≥ 60' LOA; 48.7% to hook and line catcher processors; 8.4% to pot catcher vessels > 60'; 1.5% to pot catcher processors; 2.3% to AFA trawl catcher processors; 13.4% to non-AFA trawl catcher processors; and 22.1% to trawl catcher vessels.

Stock Assessment: The Pacific cod assessment is based on a Stock Synthesis model that uses



both length-structured and age-structured data. This model incorporates fishery data and fishery independent data from the NMFS EBS trawl surveys. Pacific cod fall under Tier 3b of the ABC/OFL control rules. The 2011 age 3+ biomass is estimated at 1,560,000 mt for the BSAI. MSY reference points are not available for Pacific cod. Catch specifications for 2011 are as follows: OFL=272,000 mt, $F_{OFL}=0.29$, ABC=235,000 mt, TAC=227,950 mt.

Estimated biomass of Pacific cod has fluctuated over the last 35 years. The stock increased rapidly and peaked in 1985, then declined slightly and stabilized. An increase in biomass is projected in the short term due to

relatively good year classes produced in 2006 and 2008.

Fishery: Pacific cod are taken with trawl, longline, pot and jig gear. In 2009, a total of 55 vessels using fixed gear (17 catcher vessels, 38 catcher processors), 47 pot gear vessels (43 catcher vessels, 4 catcher processors), and 70 vessels using trawl gear (54 catcher vessels, 16 catcher processors) caught Pacific cod in the BSAI.

Economics: In 2009, ex-vessel value of Pacific cod catch in the BSAI was \$88 million, and production for all Pacific cod products in Alaska was 100,340 mt. worth \$281 million. Products included whole fish, headed and gutted fish, and fillets. Exvessel price averaged \$0.19/lb for trawl gear and \$0.26/lb for fixed gear.

Total catches, pre-season catch specifications, and exploitable biomass of Pacific Cod in the BSAI, 1976-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1976	54,671	-	-	-	-
1977	36,597	58,000	-	-	-
1978	45,838	70,500	-	-	-
1979	39,354	70,500	-	-	-
1980	51,649	70,700	148,000	-	-
1981	63,941	78,700	160,000	-	-
1982	69,501	78,700	168,000	-	-
1983	103,231	120,000	298,200	-	-
1984	133,084	210,000	291,300	-	-
1985	150,384	220,000	347,400	-	-
1986	142,511	229,000	249,300	-	-
1987	163,110	280,000	400,000	-	-
1988	208,236	200,000	385,300	-	1,481,000
1989	182,865	230,681	370,600	-	1,190,000
1990	179,608	227,000	417,000	-	1,389,500
1991	172,158	229,000	229,000	-	1,030,000
1992	206,129	182,000	182,000	188,000	910,000
1993	167,390	164,500	164,500	192,000	655,000
1994	196,572	191,000	191,000	228,000	925,000
1995	245,030	250,000	328,000	390,000	1,620,000
1996	240,590	270,000	305,000	420,000	1,640,000
1997	234,641	270,000	306,000	418,000	1,590,000
1998	195,645	210,000	210,000	336,000	1,340,000
1999	162,361	177,000	177,000	264,000	1,210,000
2000	191,056	193,000	193,000	240,000	1,300,000
2001	176,659	188,000	188,000	248,000	1,320,000
2002	197,353	200,000	223,000	294,000	1,540,000
2003	211,059	207,500	223,000	324,000	1,680,000
2004	212,161	215,500	223,000	350,000	1,660,000
2005	205,635	206,000	206,000	265,000	1,290,000
2006	193,017	194,000	194,000	230,000	922,000
2007	174,124	170,720	176,000	207,000	960,000
2008	170,661	170,720	176,000	207,000	1,080,000
2009	175,746	176,540	182,000	212,000	1,260,000
2010	144,924	168,780	174,000	205,000	1,140,000
2011	-	227,950	235,000	272,000	1,560,000

¹Catch data current through November 2010.

²TAC, ABC and OFL data from annual SAFE report.


³Biomass from annual SAFE report projections issued the previous year.



SeaAlliance/AGDB

Ecosystem Components: Bottom-up controls and/or climactic shifts may be affecting Pacific cod distribution and recruitment in the BSAI. Trends in Pacific cod estimated biomass reflect potential groundfish regime shifts associated with the Pacific Decadal Oscillation (PDO). In 1978, the PDO switched from a negative or cooler phase to a warmer, positive phase. Model estimates show increased Pacific cod biomass throughout the early 1980s following the PDO switch to a positive phase (Hunt et al. 2002). Increased Pacific cod abundance associated with warmer temperatures may have caused a shift towards top-down ecological forcing in the EBS, where Pacific cod predation impacts EBS juvenile pollock survivorship.

Pacific cod are an important prey item for SSLs, especially in winter months. The draft Steller Sea Lion Biological Opinion (Biop) released in August 2010 recommended Pacific cod fishery closure in area 543 and additional restrictions in 541 and 542.



AFSC, NOAA Fisheries




For More Information:

D. Hanselman, C. Lunsford, and C. Rodgveller. 2010. Assessment of the Sablefish in Alaska. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: Sablefish *Anoplopoma fimbria* distribution extends from the northern Mexico through the Gulf of Alaska, along the Aleutian Islands and into the Bering Sea. Adult sablefish are generally found at depths greater than 200 m along the continental slope, shelf gullies and deep fjords. Juveniles (less than 40 cm) spend the first 2-3 years farther inshore along the continental shelf and begin to move out to the continental slope around age 4-5. Young-of-the-year feed primarily on euphausiids and copepods while adults are more opportunistic feeders, relying more heavily on fish such as pollock, Pacific herring and Pacific cod. Squid and jellyfish are important invertebrates in the adult sablefish diet. Coho and Chinook salmon are the main predators of young-of-the-year.

Sablefish are relatively long lived. They begin to recruit to the fishery at age 4 or 5 and longevity often reaches 40 years (oldest recorded sablefish in Alaska was 94 years old). Female sablefish size at 50% maturity is approximately 65 cm (age 6). Females are slightly larger than males, and fish in the BSAI generally tend to be smaller than in the GOA. Natural mortality is estimated at $M=0.10$. Alaskan sablefish spawn near the edges of the continental slope between 300-500 m in late March.


Catch History: US fishermen have harvested sablefish since the end of the 19th century as a byproduct of halibut fisheries. Harvests were relatively small, averaging 1,666 mt from 1930-1957. Japanese longlining began in the EBS around 1958 and expanded into the AI and GOA through the 1970s. Japanese fleet catches increased throughout the 1960s, and catch peaked 36,776 mt in 1972. High fishing pressure in the early 1970s may have resulted in a population decline of sablefish in the mid-1970s. By 1988, US fishermen took the majority of the sablefish harvested in the GOA and BSAI. The fishery was a derby-style fishery in the late 1980s and early 1990s until Individual Fishing Quotas were implemented for the hook and line fishery in 1995.



Fishery Management: BSAI and GOA sablefish are managed as one population in federal waters due to their highly migratory behavior during certain life history stages. The

sablefish fishery is regulated under the BSAI groundfish FMP through permits, limited entry, catch quotas (TACs), seasons, in-season adjustments, gear restrictions, closed waters, bycatch limits and rates, allocations, regulatory areas, record keeping, reporting requirements and observer monitoring.

In 1990, Amendment 13 to the BSAI FMP similarly allocated sablefish quota by gear type; 50% to fixed gear and 50% to trawl gear in the BS; 75% to fixed gear and 25% to trawl gear in the AI. Amendment 20 to the GOA FMP and 15 to BSAI FMP established IFQ management for



the sablefish fishery and allocated 20% of the fixed gear quota to a CDQ reserve for the BSAI, effective 1990.

Stock Assessment: The sablefish assessment is based on a statistical sex-specific age-structured model. This model incorporates fishery data and fishery independent data from domestic and Japan-US cooperative longline surveys and the NMFS GOA trawl survey. Sablefish fall under Tier 3b of the ABC/OFL control rules. The 2010 age 4+ biomass is estimated at 205,000 mt for the GOA and BSAI combined. The catch specifications for 2011 Bering Sea sablefish are as follows; OFL=3,360 mt, ABC=2,850 mt, TAC=2,850 mt. For the Aleutian Islands, OFL=2,250 mt, ABC=1,900 mt, TAC=1,900 mt.

Total catches, pre-season catch specifications, and exploitable biomass of Sablefish age 4+ in the BSAI, 1976-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1976	4,582	-	-	-	-
1977	4,615	-	-	-	-
1978	2,013	-	-	-	-
1979	2,158	-	-	-	-
1980	2,480	5,000	-	-	-
1981	3,137	5,000	-	-	-
1982	4,139	5,000	-	-	-
1983	3,368	5,000	-	-	-
1984	3,328	5,340	6,185	-	-
1985	3,796	4,500	6,080	-	-
1986	6,546	6,450	7,200	-	-
1987	8,012	7,700	7,700	-	-
1988	6,608	8,400	9,200	-	152,800
1989	4,500	5,270	6,200	-	93,300
1990	4,445	7,200	7,200	-	114,500
1991	3,199	6,300	6,300	-	53,100
1992	2,104	4,400	4,400	5,870	37,400
1993	2,747	4,100	4,100	4,500	37,000
1994	2,470	3,340	3,340	4,160	28,500
1995	2,048	3,800	3,800	4,900	30,400
1996	1,349	2,300	2,500	3,300	26,100
1997	1,326	2,300	2,675	5,610	36,500
1998	1,181	2,680	2,680	4,390	39,200
1999	1,211	3,200	3,200	4,980	43,000
2000	1,790	3,900	3,900	4,840	51,000
2001	1,937	4,060	4,060	4,980	58,000
2002	2,261	4,480	4,480	6,750	67,000
2003	2,048	6,000	6,000	8,880	70,000
2004	1,993	6,000	6,450	8,640	71,000
2005	2,539	5,060	5,060	6,120	66,000
2006	2,166	5,820	6,160	7,420	66,000
2007	2,322	5,790	5,790	9,840	66,000
2008	2,018	5,300	5,300	6,270	75,000
2009	1,939	4,920	4,920	5,810	67,000
2010	1,770	4,860	4,860	5,760	65,000
2011	-	4,750	4,750	5,610	62,000

¹Catch data current through November 2010.

²TAC, ABC and OFL from annual Federal Register.

³Biomass from annual SAFE report projections.

Biomass of sablefish has fluctuated over time. There were two high points in biomass in the early 1970s and mid-1980s and two decreases in the late 1970s and the mid-1990s. Relative abundance is near an all-time low for the domestic longline survey.

Fishery: Sablefish are taken with trawl, longline and pot gear. Most sablefish are taken with longline gear in the Aleutian Islands and pot gear in the Bering Sea. The sablefish season is open 7 months beginning in April, concurrent with the halibut fishing season. Primary species taken incidentally in the sablefish fishery include shortaker, rougheye and thornyhead rockfish.

Sperm whale and killer whale depredation is problematic for sablefish fisheries in the GOA and BSAI. Depredation occurs when whales remove sablefish from longline gear, damage the fish and/or fishing gear. Killer whale depredation predominates in the BSAI and sperm whale depredation is more common the GOA. Depredation can lead to significant economic losses in the form of reduced catch, extended travel distances, and damaged gear. Depredation may also reduce the accuracy of sablefish stock assessment models. Additionally, depredating whales may be at greater risk of mortality or injury through vessel strikes or entanglement in gear.

Economics: In 2009, the ex-vessel value of sablefish catch from the BSAI was \$11.6.5 million. Exvessel prices for BSAI sablefish in 2009 averaged \$2.80/lb for fish caught on longline gear and \$1.43/lb for fish taken with trawl gear. For both gear types, the primary product is frozen, head and gutted fish.



AFSC, NOAA Fisheries



Diana Evans, NPFMC

Biology: Yellowfin sole *Limanda aspera* are distributed from the Sea of Japan to British Columbia, with the highest abundance in the Bering Sea. Yellowfin sole are the target of the largest flatfish fishery in the US and are one of the most abundant flatfish species in the EBS. Adult yellowfin sole occupy the benthos and have separate winter spawning and summertime feeding grounds on the EBS shelf. Adults over-winter near the shelf margins and then migrate to inner shelf areas in April/May each year for spawning and feeding. Yellowfin sole predate on bivalves, polychaetes, amphipods, mollusks and fish. They are prey for Pacific cod, Pacific halibut and skates.



For More Information:

T. Wilderbuer, D. Nichol, and J. Ianelli. 2010. Assessment of Yellowfin Sole in the Bering Sea and Aleutian Islands Area. <http://www.afsc.noaa.gov/refm/stocks/assessments.htm>

Yellowfin sole are relatively slow-growing and long-lived. They begin to recruit to the fishery at age 6, are fully selected by age 13 and longevity extends to 30+ years. Females reach 50% maturity at 30 cm (10.5 years old) and are highly fecund, producing 1.3-3.3 million eggs depending on size. Annual natural mortality of adults is estimated at 0.12. Spawning occurs in June/July in shallow waters from Bristol Bay to Nunivak Island.




Diana Evans, NPFMC

Fishery Management:

BSAI Flatfish are regulated under the BSAI groundfish FMP through permits, limited entry, catch quotas (TACs), seasons, in-season adjustments, gear restrictions, closed waters, bycatch limits and rates (for halibut and crab), allocations, regulatory areas, record keeping, reporting requirements and observer monitoring.

In 1985, the Flounder (Flatfish) category was broken into four management groups (Greenland turbot, Arrowtooth flounder, Yellowfin sole, Other Flatfish) due to significant differences in stock robustness and product values. Northern rock sole was separated from the Other Flatfish complex in 1987. Flathead sole was separated from the Other Flatfish complex in 1995, and Alaska plaice was separated in 2002.

In 2008, BSAI FMP Amendment 80 established catch shares for the bottom trawl catcher-processor fleet. Flatfish resources were allocated among BSAI trawl harvesters according to their historic harvest patterns, groundfish retention standards were extended to the catcher/processor fleet,



Total catches, pre-season catch specifications, and exploitable biomass of Yellowfin Sole in the BSAI, 1976-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1976	56,221	-	-	-	-
1977	58,373	106,000	-	-	-
1978	138,433	126,000	-	-	-
1979	99,017	126,000	-	-	-
1980	87,391	117,000	169,000	-	-
1981	97,301	117,000	214,500	-	-
1982	95,712	117,000	214,500	-	-
1983	108,385	117,000	214,500	-	-
1984	159,526	230,000	310,000	-	-
1985	227,107	226,900	310,000	-	-
1986	208,597	209,500	230,000	-	-
1987	181,429	187,000	187,000	-	-
1988	223,156	254,000	254,000	-	1,408,000
1989	153,165	182,675	241,000	-	1,530,000
1990	80,584	207,650	278,900	-	1,640,000
1991	96,135	135,000	250,600	-	1,790,000
1992	146,946	235,000	372,000	452,000	2,660,000
1993	105,809	220,000	238,000	275,000	2,500,000
1994	144,544	150,325	230,000	269,000	1,925,000
1995	124,752	190,000	277,000	319,000	2,770,000
1996	130,163	200,000	278,000	342,000	2,850,000
1997	166,915	230,000	233,000	339,000	2,530,000
1998	101,315	220,000	220,000	314,000	3,010,000
1999	67,320	207,980	212,000	308,000	3,180,000
2000	84,070	123,262	191,000	222,600	2,820,000
2001	63,578	113,000	176,000	209,000	2,380,000
2002	74,985	86,000	115,000	136,000	1,597,000
2003	81,050	83,750	114,000	136,000	1,550,000
2004	75,510	86,075	114,000	135,000	1,560,000
2005	94,384	90,686	124,000	148,000	1,560,000
2006	99,138	95,701	121,000	144,000	1,680,000
2007	120,968	136,000	225,000	240,000	2,000,000
2008	148,894	225,000	248,000	265,000	2,200,000
2009	103,805	210,000	210,000	224,000	1,870,000
2010	114,600	219,000	219,000	234,000	1,960,000
2011	-	196,000	239,000	262,000	1,958,600

¹Catch data current through November 2010.


²1988-2010 TAC, ABC and OFL data from annual Federal Register Harvest Specifications. Pre-1988 TAC and ABC data from annual SAFE reports.

³Biomass from annual SAFE report projections.

monitoring requirements were increased, and fishermen were given the ability to form cooperatives. Up to 93% of the yellowfin sole TAC is allocated to the Amendment 80 fleet, depending on the TAC. Like other groundfish stocks except pollock, 10.7% of TAC is first allocated to CDQ groups.

Catch History: Yellowfin sole have been harvested annually since the inception of the BS bottom trawl fishery in 1954. Overharvesting by foreign vessels occurred from 1959-1962, and catches averaged 404,000 mt annually during that period. Catches declined during the late 1960s and early 1970s as a result of reduced abundance. Domestic and joint venture fisheries for yellowfin sole emerged in the 1980s, and only domestic harvesting has occurred since 1990.

Stock Assessment: The yellowfin sole assessment is a separable catch-age, sex-specific analysis. This model incorporates fishery data and fishery independent data from annual trawl surveys. Yellowfin sole fall under Tier 1a of the ABC/OFL control rules. The 2011 projected age 2+ biomass is 2,173,700 mt. Catch specifications for 2011 are as follows; OFL=262,000 mt, ABC= 239,000 mt, TAC= 196,000 mt.



Yellowfin sole biomass peaked in the early-1990s. The population has been in a slow decline as the strongest year classes have passed through the fishery, however, the population remains at fairly high/stable levels.

Fishery: Yellowfin sole are primarily caught with trawl gear. One catcher vessel and 29 catcher processors participated in 2009 flatfish fisheries in the BSAI. Fishing effort is focused on the mid and inner BS shelf during ice-free conditions. A small area in Bristol Bay is open to bottom trawling from April 1 – June 15 to allow the fishery to target this species when they are aggregated and can be taken with low incidental catches of other species. Yellowfin sole are usually headed and gutted or frozen whole for further processing. Retention by the Amendment 80 sector was 88% in 2009.

Economics: In 2009, production was 126,540 mt for all flatfish products for a total gross value of \$157 million. Ex-vessel value of all flatfish caught in the BSAI in 2009 was \$61.8 million.



Arrowtooth flounder
AFSC, NOAA Fisheries

Arrowtooth Flounder & Kamchatka Flounder

Biology: From 1986 until 2011, two species were managed together under the “Arrowtooth Flounder” complex (arrowtooth flounder *Atheresthes stomias* is and Kamchatka flounder *A. evermanni*). In 2011, separate catch specifications were established for these species.


Arrowtooth flounder are distributed from the Kamchatka Peninsula to the BSAI south to central California. Juveniles occupy continental shelf waters until age 4, at which point their range expands into the continental slope. Adults migrate seasonally from shelf margins in the winter to the outer shelf in April/May with the onset of warmer waters temperatures. In the BSAI, arrowtooth flounder predate on juvenile pollock (47%), adult pollock (19%) and euphausiids (9%). Predators of include Pacific cod, pollock, skates and Pacific sleeper sharks.




For More Information:

T. Wilderbuer, D. Nichol, and K. Aydin. 2010. Assessment of Arrowtooth Flounder in the Eastern Bering Sea and Aleutian Islands Area. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Arrowtooth flounder recruitment to the fishery begins at about 5 years, and females are fully recruited by age 9. The estimated length at 50% maturity is 28 cm for males (4 years) and 37 cm for females (5 years). Female natural mortality is estimated at $M=0.2$; for males, the best estimate is $M=0.35$. Adult males range in size from 30-50 cm, and females range in size from 30-70 cm. The spawning period for arrowtooth flounder is protracted and variable, ranging from September through March. Compared to Arrowtooth flounder, relatively little is known about the life history of Kamchatka flounder.



Catch History: USSR and Japan targeted Greenland turbot and arrowtooth flounder during the 1960s. Catches peaked from 1974-1976 at 19,000-25,000 mt. Arrowtooth flounder and Greenland turbot were managed as a complex until 1985 due to their similar life history characteristics and distribution. Catches decreased following implementation of the Magnuson-Stevens Act in 1977.




Stock Assessment: The arrowtooth flounder assessment uses a length-based approach to calculating estimates of population numbers-at-age. This model incorporates fishery data and fishery independent data from EBS and AI trawl surveys. Arrowtooth flounder fall under Tier 3a of the ABC/OFL control rules. The 2011 projected biomass is 1,124,200 mt. Catch specifications for 2011 are as follows; OFL=186,000 mt, ABC= 153,000 mt, TAC= 25,900 mt. Biomass of Kamchatka flounder is projected to be 128,800 mt in

2011. Catch specifications for Kamchatka flounder for 2011 are as follows; OFL=23,600 mt, ABC= 17,700 mt, TAC= 17,700 mt.

Fishery: Arrowtooth flounder remains a non-target fishery but catches and retention have increased in response to developing markets. Arrowtooth flounder are also an incidental catch in the more valuable target flatfish fisheries. One catcher vessels and 29

catcher processors participated in 2009 flatfish fisheries in the BSAI. From 2005-2007, at least 50% of arrowtooth flounder caught was discarded. Retention improved with Amendment 80 to the BSAI FMP in 2008. Retention rate for arrowtooth flounder by the Amendment 80 sector was 93% in 2009.



Total catches, pre-season catch specifications, and exploitable biomass of Arrowtooth Flounder in the BSAI, 1986-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1986	6,903	20,000	20,000	-	-
1987	4,539	9,795	30,900	-	-
1988	5,883	5,531	109,500	-	377,700
1989	3,222	6,000	163,700	-	528,200
1990	4,232	10,000	242,500	-	519,000
1991	13,686	20,000	116,400	-	590,400
1992	11,980	10,000	82,300	114,000	457,000
1993	9,298	10,000	72,000	96,000	480,000
1994	14,377	10,000	93,400	130,000	519,000
1995	9,283	10,227	113,000	138,000	625,000
1996	14,610	9,000	129,000	162,000	578,000
1997	9,651	20,760	108,000	167,000	587,000
1998	15,679	16,000	147,000	23,000	869,000
1999	10,573	134,354	140,000	219,000	819,000
2000	13,228	131,000	131,000	160,000	785,000
2001	14,056	22,011	117,000	141,500	701,000
2002	11,853	16,000	113,000	137,000	671,000
2003	14,580	12,000	112,000	139,000	597,000
2004	18,139	12,000	115,000	142,000	696,000
2005	14,237	12,000	108,000	132,000	684,000
2006	13,361	13,000	136,000	160,000	964,000
2007	11,917	20,000	158,000	193,000	1,280,000
2008	21,884	75,000	244,000	297,000	1,780,000
2009	28,914	75,000	156,000	190,000	1,140,000
2010	38,098	75,000	156,000	191,000	1,120,000
2011	-	25,900	153,000	186,000	1,124,200


¹Catch data current through November 2010.

²1988-2010 TAC, ABC and OFL data from annual Federal Register Harvest Specifications. Pre-1988 TAC and ABC data from annual SAFE reports.

³Biomass from annual SAFE report projections.

Ecosystem Components: Arrowtooth flounder biomass has been steadily increasing in the BS since the early 1980s. There is the potential that arrowtooth flounder could exert predation pressure on their main prey resource, juvenile pollock. However, simulation models suggest that the role of pollock as a predator of arrowtooth flounder (limiting population growth) is more important than the role of pollock as prey in the BSAI.

BS Arrowtooth diet





Diana Evans, NPFMC

Biology: Two species of rock sole, Northern rock sole *Lepidopsetta polyxstra* and Southern rock sole *L. bilineatus*, occur in the North Pacific Ocean and are managed together as one complex. Northern rock sole are the most commonly found species of rock sole in the BSAI. Adults are bottom dwellers and occupy separate winter and summer feeding ground along the continental shelf. As early juveniles, rock sole consume plankton and zooplankton, switching to bivalves, polychaetes, amphipods, mullosks and crustaceans as they age and become late juveniles and adults. Small rock sole are prey for Pacific cod, pollock, yellowfin sole, skates and Pacific halibut.



For More Information:


T. Wilderbuer and D. Nichol. 2010. Assessment of Northern Rock Sole in the Eastern Bering Sea and Aleutian Islands Area. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Recruitment to the fishery begins at age 4 and they are fully selected by age 11. Estimated length at 50% maturity is 35 cm (9 years). Natural mortality is estimated at $M=0.15$. Rock sole spawn from December to March in two separate concentrations in the BS along the continental shelf/slope break.

Catch History: Rock sole were harvested by Japanese and Soviet vessels beginning in 1963. Catches averaged 7,000 mt annually from 1963-1969 and increased during the early 1970s. Peak catch occurred in 1972 (61,000 mt). Catches declined until joint venture operations began in 1980. Catches again increased during the 1980s and peaked in 1988 (86,000 mt). The fishery was fully domesticated by 1990, and catches have remained fairly stable since 1990 (average 46,000 mt annually).

Fishery Management: Rock sole is regulated under the BSAI groundfish FMP through permits, limited entry, catch quotas (TACs), seasons, in-season adjustments, gear restrictions, closed waters, bycatch limits and rates, allocations, regulatory areas, record keeping, reporting requirements and observer monitoring. In 2008, BSAI FMP Amendment 80 modified rock sole fishery management, such that 100% of the directed fishery rock sole TAC is allocated among non-AFA trawl catcher processors according to their historic harvest patterns, groundfish retention standards were extended to catcher/processor fleet and fishermen were given the ability to form cooperatives. Like other groundfish, 10.7% of rock sole TAC is allocated to CDQ groups.

**Northern Rock Sole
Female Spawning Biomass**




Stock Assessment: The rock sole assessment uses AD Model builder software to conduct a separable catch-age analysis that estimates abundance, mortality and recruitment. This model incorporates fishery data and fishery independent data from EBS and AI trawl surveys. Rock sole fall under Tier 1a of the ABC/OFL control rules. The 2011 projected biomass is 1,868,400 mt and $B_{msy}=259,000$. Catch specifications for 2011 are as follows; OFL=248,000 mt, ABC= 224,000 mt, TAC= 85,000 mt.

Strong recruitment and low fishing effort enabled rock sole biomass to increase significantly from 1985-1995. Estimated biomass peaked in the late 1990s and then declined by about 20% through 2005. The decline during the early 2000s was attributed to below average recruitment to the adult population during the 1990s. Estimated biomass began increasing again in 2005 as a result of a series of above average year-classes.

Fishery: Rock sole are caught by trawl catcher-processors targeting roe-bearing females. The primary product for the rock sole fishery is the high value roe. The fishery

occurs from January-March and is focused in outer Bristol Bay and north of Unimak Island. A total of 3 catcher vessels and 34 catcher processors participated in the 2008 flatfish fisheries in the BSAI. From 1987-2000, over 50% of rock sole catch was discarded. Retention rate for rock sole by the Amendment 80 sector increased to 87% in 2009.



Ecosystem Components: Northern rock sole recruitment has been linked to decadal scale climate variability, especially ocean forcing by onshelf/offshelf winds in the BS. After spawning in March, northern rock sole larvae are subject to advection from wind, current and tidal forcing during the spring. Using an ocean surface current model, northern rock sole larvae advection towards favorable nursery areas and resultant above-average recruitment occurred during years with onshelf (easterly) winds during the 1980s and again in 2001-2003. Conversely, periods of offshelf (westerly) winds during the 1990s corresponded with average or poor recruitment.



Diana Evans, NPFMC

Total catches, pre-season catch specifications, and exploitable biomass of Rock Sole* in the BSAI, 1989-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1989	68,912	90,762	171,000	-	1,277,900
1990	35,253	60,000	216,300	-	1,193,900
1991	46,681	90,000	246,500	-	1,363,700
1992	51,956	40,000	260,800	260,800	1,481,000
1993	64,260	75,000	185,000	270,000	1,550,000
1994	60,584	75,000	313,000	363,000	1,790,000
1995	55,028	60,000	347,000	388,000	2,330,000
1996	47,146	70,000	361,000	420,000	2,360,000
1997	67,520	97,185	296,000	427,000	2,390,000
1998	33,667	100,000	312,000	449,000	2,360,000
1999	40,511	120,000	309,000	444,000	2,320,000
2000	49,666	137,760	230,000	273,000	2,070,000
2001	29,475	75,000	228,000	271,000	1,940,000
2002	41,865	54,000	225,000	268,000	1,850,000
2003	37,339	44,000	110,000	132,000	877,000
2004	48,680	41,000	139,000	166,000	1,160,000
2005	37,361	41,500	132,000	157,000	1,380,000
2006	36,456	41,500	126,000	150,000	1,490,000
2007	37,127	55,000	198,000	200,000	1,670,000
2008	51,277	75,000	301,000	304,000	1,880,000
2009	48,593	90,000	296,000	301,000	1,630,000
2010	53,111	90,000	240,000	243,000	1,770,000
2011	-	85,000	224,000	248,000	1,868,400

*Rock Sole included in Other Flatfish category before 1989.

¹Catch data current through November 2010.

²1989-2010 TAC, ABC and OFL data from annual Federal Register Harvest Specifications.

³Biomass from annual SAFE report projections.



NOAA Fisheries

Greenland Turbot

Biology: Greenland turbot *Reinhardtius hippoglossoides* has a circumpolar distribution, occurring in both the North Pacific and North Atlantic Oceans. Juveniles inhabit shallow continental shelf waters (<200 m) for the first 3-4 years and move out to the deeper waters of the continental slope (200-1,000 m). Greenland turbot predate on euphausiids, polychaetes and small fish (e.g. pollock) as they mature. In the North Pacific, juveniles are prey for Pacific cod and Pacific halibut.

Greenland turbot size at 50% maturity is around 60 cm (age 5-10). Greenland turbot begin to recruit to longline fisheries at about 60 cm and are fully recruited at 90 cm. Natural mortality is estimated at $M=0.112$. Peak spawning period is from November – February in the EBS. Female fecundity is fairly low; females less than 83 cm release 25,000-150,000 eggs.




For More Information:

J. Ianelli, T. Wilderbuer, and D. Nichol. 2010. Assessment of Greenland Turbot in the Eastern Bering Sea and Aleutian Islands Area. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Catch History: Catches averaged 30,000 mt annually during that during the 1960s when the USSR and Japan first targeted the Greenland turbot fishery. Catches peaked in the mid-1970s, and declined after 1986 due to poor recruitment.


Stock Assessment: The Greenland turbot assessment is based on a stock synthesis model that incorporates fishery data and fishery independent data from EBS slope and shelf bottom-trawl surveys and the NMFS longline survey. Greenland turbot fall under Tier 3a of the ABC/OFL control rules. The 2011 projected biomass is 74,000 mt. Catch specifications for 2011 are as follows; OFL=7,220 mt, ABC= 6,140 mt, TAC= 5,050 mt. Catch limits are further apportioned into BS (3,500 mt) and AI (1,550 mt) components.



Biomass has declined since the early 1970s, however it may have been abnormally high during the 1970s. Good year-classes have been produced in recent years, so biomass is

expected to increase in the near future.

Fishery: The Greenland turbot season opens in May and fishing effort is concentrated on the continental slope throughout the EBS and on both sides of the AI.



Current Issues: Killer whale depredation is problematic for Greenland turbot longline fisheries in the EBS.

Total catches, pre-season catch specifications, and exploitable biomass of Greenland Turbot in the BSAI, 2000-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2000	6,974	9,300	9,300	42,000	233,000
2001	5,312	8,400	8,400	31,000	210,000
2002	3,635	8,000	8,100	36,500	208,000
2003	3,530	4,000	5,800	17,800	112,000
2004	2,239	3,500	4,740	19,300	132,000
2005	2,579	3,500	3,930	19,200	98,300
2006	1,977	2,740	2,740	14,200	74,200
2007	2,003	2,440	2,440	15,600	119,000
2008	2,751	2,540	2,540	15,600	104,000
2009	4,283	7,380	7,380	14,800	105,000
2010	-	6,120	6,120	7,460	61,100
2011	-	5,050	6,140	7,220	73,981

¹Catch data current through November 2010.

²TAC, ABC and OFL from Federal Register Harvest Specifications.

³Biomass from annual SAFE report projections.



Diana Evans, NPFMC

Alaska Plaice

Biology: Alaska plaice *Pleuronectes quadrituberculatus* distribution extends through the Sea of Japan, Chukchi Sea, BSAI and GOA. Alaska plaice are generally found along the EBS continental shelf, with relatively few found in the AI region. Summer distribution of adults is generally confined to depths less than 110 m, with larger fish in deeper waters and smaller juveniles in shallower coastal waters. Alaska plaice predate on polychaetes and amphipods and are prey for Pacific cod, Pacific halibut and yellowfin sole.

Alaska plaice recruit to trawl fisheries at age 4, are full recruited by age 13. Females mature between ages 7 and 12. Natural mortality is estimated at $M=0.13$. Spawning usually occurs in March and April on hard sandy substrate in the EBS.

Catch History: Alaska plaice were harvested by Japanese and Soviet vessels beginning in 1963. Catches increased from 1,000 mt in 1971 to a peak of 62,000 mt in 1988. Joint ventures began in 1988, and the fishery was fully harvested by domestic vessels in 1991. Although a small directed target fishery exists, it is mainly a secondary catch in the yellowfin sole fishery.





For More Information:

T. Wilderbuer, D. Nichol, and P. Spencer. 2010. Assessment of Alaska Plaice in the Eastern Bering Sea and Aleutian Islands Area. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Stock Assessment:

The assessment uses a sex-specific, age-structured model. This model incorporates fishery data and fishery independent data from trawl surveys. Alaska plaice fall under Tier 3a of the ABC/OFL control rules. The 2011 projected biomass is 780,300 mt. Catch specifications for 2011 are as follows; OFL= 79,100 mt, ABC= 65,100 mt, TAC= 16,000 mt.



Recent strong year-classes should contribute to increasing total biomass and female spawning biomass in the near future.

Fishery: Alaska plaice are caught primarily by trawl catcher processors targeting higher-value flatfish species such as yellowfin sole. With the implementation of Amendment 80 in 2008, retention rates of Alaska plaice increased from about 5% (2003-2005 average) to 64% in 2008.

Recent strong year-classes should contribute to increasing total biomass and female spawning biomass in the near future.

Total catches, pre-season catch specifications, and exploitable biomass of Alaska Plaice* in the BSAI, 2002-2011.


Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2002	-	-	-	-	1,100,000
2003	10,118	10,000	137,000	165,000	1,080,000
2004	7,888	8,000	203,000	258,000	1,050,000
2005	11,194	8,000	189,000	237,000	913,000
2006	17,318	8,000	188,000	237,000	1,008,000
2007	19,522	25,000	190,000	241,000	1,340,000
2008	17,376	50,000	194,000	248,000	1,850,000
2009	13,699	50,000	232,000	298,000	1,500,000
2010	15,771	50,000	224,000	278,000	2,260,000
2011	-	16,000	65,100	79,100	780,300

*Alaska plaice removed from Other flatfish complex 2002.

¹Catch data current through November 2010.

²TAC, ABC and OFL data from Federal Register.

³Biomass from annual SAFE report projections.



Flathead Sole

AFSC, NOAA Fisheries



For More Information:

W. Stockhausen, D. Nichol, R. Lauth, and M. Wilkins. 2010. Assessment of Flathead Sole in the Eastern Bering Sea and Aleutian Islands Area. <http://www.afsc.noaa.gov/refm/stocks/assessments.htm>

Biology: Flathead Sole is managed as a two-species complex including flathead sole *Hippoglossoides elassodon* and Bering flounder *Hippoglossoides robustus*. Individuals of both species are morphologically similar; flathead sole are faster growing and achieve larger size. Flathead sole are distributed in the Kuril Islands, BS, GOA and down to northern California. In the northern part of its range, its distribution overlaps with Bering flounder. Bering flounder distribution extends from the Chukchi Sea into the western BS. Bering flounder generally represents less than 3% of the estimated survey biomass of the two species. Adult flathead sole overwinter near the shelf margins before migrating to the mid and outer continental shelf in April or May each year for feeding. Flathead sole predate on pollock, polychaetes, brittle stars and crustaceans. They are prey for adult pollock and Pacific cod.

Flathead sole recruitment to the fishery begins at age 4, and longevity extends to 32 years. Estimated length at 50% maturity is 32 cm. Natural mortality is estimated at $M=0.20$. Flathead sole spawn in March and April, primarily in deeper waters near the margins of the continental shelf. Females release from 70,000-600,000 eggs depending on size.

Catch History: Flathead sole were harvested by Japanese and Soviet vessels beginning in 1963. Flathead sole catches peaked in 1971 (51,000 mt). Catches declined to 15,000 mt in 1975 and remained under 10,000 mt until 1990. Catch levels have increased since the 1980s due to higher incidental catch rates and emerging markets for flathead sole, averaging 18,377 mt from 1995-2009.


Stock Assessment: The assessment uses a split-sex, age-based model with length-based formulations for fishery and survey selectivities. This model incorporates fishery data and fishery independent data from trawl surveys. Flathead sole fall under Tier 3a of the ABC/OFL control rules. The 2011 projected biomass is 791,000 mt. Catch specifications for 2011 are as follows; OFL=83,300 mt, ABC= 69,300 mt, TAC= 41,548 mt.

Fishery: In 2008, BSAI FMP Amendment 80 allocated 100% of the directed fishery flathead sole TAC is allocated among non-AFA trawl catcher processors according to their historic harvest patterns. The fishery mainly occurs from January-June. Primary products are H&G with roe-in and kirimi.

Total catches, pre-season catch specifications, and exploitable biomass of Flathead Sole* in the BSAI, 1995-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1995	14,713	30,000	138,000	167,000	677,000
1996	17,344	30,000	116,000	140,000	593,000
1997	20,681	43,500	101,000	145,000	632,000
1998	24,597	100,000	132,000	190,000	824,000
1999	18,555	77,300	77,300	118,000	710,000
2000	20,439	52,652	73,500	90,000	660,000
2001	17,809	40,000	84,000	102,000	618,000
2002	15,547	25,000	82,600	101,000	695,000
2003	13,792	20,000	66,000	81,000	550,000
2004	16,850	19,000	61,900	75,200	505,000
2005	16,151	19,500	58,500	70,200	560,000
2006	17,947	19,500	59,800	71,800	636,000
2007	18,744	30,000	79,200	95,300	875,000
2008	24,539	50,000	71,700	86,000	820,000
2009	19,549	60,000	71,400	83,500	834,000
2010	19,863	60,000	69,200	83,100	785,000
2011	-	41,548	69,300	83,300	791,000

*Flathead Sole removed from Other Flatfish category 1995. Flathead Sole category includes Bering Flounder and Flathead Sole.
¹Catch data from BSAI SAFE.
²TAC, ABC and OFL from annual Specifications.
³Biomass corresponds to the annual SAFE report projections.





Dover sole
AFSC, NOAA Fisheries

Biology: The Other Flatfish complex consists of 15 species. Starry flounder, rex sole, longhead dab, Dover sole, and butter sole comprise the majority of harvested “Other Flatfish.”

Data are limited for many of the species in this complex. Rex sole and Dover sole are distributed from Baja California, through the BSAI and widely throughout the GOA. Adult rex sole and Dover sole are bottom dwellers and are generally found in water deeper than 300 m.

Maturity studies from Oregon show that rex sole estimated length at 50% maturity is 16 cm (males) and 24 cm (females). Maturity studies from Oregon indicate that Dover sole estimated length at 50% maturity is 33 cm (females). Available natural mortalities are as follows; rex sole $M=0.17$, Dover sole $M=0.085$, remaining Other Flatfish $M=0.20$.

Catch History: Other Flatfish are non-target species but have been captured in target flatfish fisheries since Japanese and Soviet fleets began fishing in the Bering Sea in 1963. Prior to its removal from the “Other Flatfish” complex in 2002, Alaska plaice comprised the majority of harvested “Other Flatfish.” Catch of Alaska plaice and “Other Flatfish” peaked in 1988 at 137,418 mt. Since the removal of Alaska plaice from the complex, annual catches have averaged about 3,500 mt from 2003-2010.

Stock Assessment: The Other Flatfish assessment uses survey biomass estimates as principal data sources to estimate biomass. “Other Flatfish” are managed under Tier 5 of the ABC/OFL control rules. The 2011 projected biomass is 127,300 mt. Catch specifications for 2011 are as follows; OFL=19,500 mt, ABC=14,500 mt, TAC= 3,000 mt.

Common Name	Scientific Name
Arctic flounder	<i>Liopsetta glacialis</i>
Butter sole	<i>Isopsetta isolepis</i>
Curlfin sole	<i>Pleuronectes decurrens</i>
Deepsea sole	<i>Embassichthys bathybius</i>
Dover sole	<i>Microstomus pacificus</i>
English sole	<i>Parophrys vetulus</i>
Longhead dab	<i>Limanda proboscidea</i>
Pacific sanddab	<i>Citharichthys sordidus</i>
Petrale sole	<i>Eopsetta jordani</i>
Rex sole	<i>Glyptocephalus zachirus</i>
Roughscale sole	<i>Clidodoerma asperrimum</i>
Sand sole	<i>Psettichthys melanostictus</i>
Slender sole	<i>Lyopsetta exilis</i>
Starry flounder	<i>Platichthys stellatus</i>
Sakhalin sole	<i>Pleuronectes sakhalinensis</i>



For More Information:

T. Wilderbuer, D. Nichol, and P. Spencer. 2010. Assessment of Other Flatfish in the Eastern Bering Sea and Aleutian Islands Area.
<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Total catches, pre-season catch specifications, and exploitable of Other Flatfish* in the BSAI, 2002-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2002	2,631	3,000	18,100	21,800	78,300
2003	2,749	3,000	16,000	21,400	107,000
2004	4,669	3,000	13,500	18,100	90,300
2005	4,599	3,500	21,400	28,500	143,000
2006	3,233	3,500	18,100	24,200	121,000
2007	5,840	10,000	21,400	28,500	149,000
2008	3,623	21,600	21,600	28,800	150,000
2009	2,163	17,400	17,440	23,100	121,000
2010	2,179	17,300	17,300	23,000	121,000
2011	-	3,000	14,500	19,500	127,329

*Alaska plaice removed from Other flatfish complex 2002. Flathead sole removed from Other Flatfish complex 1995.

¹Catch data current through November 2010.

²1988-2010 TAC, ABC and OFL data from FR Specifications.

³Biomass from annual SAFE report projections.

Estimated biomass of “Other Flatfish” was relatively stable from 1983-1995 (averaging about 55,000 mt), then increased and has remained at a high level from 2004-2010, averaging about 130,000 mt.

Fishery: Other Flatfish are caught primarily by trawl catcher processors targeting higher value flatfish species. Nevertheless, 80% of the Other flatfish were retained by the Amendment 80 Sector in 2009. Other Flatfish are also caught in Pacific cod, bottom pollock, Pacific ocean perch and other flatfish fisheries.



Pacific Ocean Perch

AFSC, NOAA Fisheries

Biology: Pacific Ocean Perch (POP) *Sebastes alutus* distribution extends from Japan around the Pacific Rim south to California. POP are most abundant in AI, GOA and British Columbia and are found primarily offshore along the continental slope in depths from 180-420 m. POP are a demersal species found over cobble substrate. Seasonal changes in depth distribution occur, and adults migrate farther offshore to deeper waters during winter.

During late spring and summer, POP migrate to shallower waters inshore for summer feeding. Populations often occur in patchy aggregations. Juveniles feed on calanoid copepods, whereas adults prey on euphausiids, shrimp and squids. POP are prey for Pacific halibut, sablefish, Pacific cod and arrowtooth flounder.



AFSC, NOAA Fisheries




For More Information:

P. Spencer and J. Ianelli. 2010. Assessment of Pacific Ocean Perch in the Bering Sea and Aleutian Islands Area. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

POP is a slow-growing, long lived species. Recruitment to trawl fisheries begins at age 5, and they are fully recruited to the fishery around age 8. Females reach 50% maturity at 29 cm (10.5 years in the GOA) and longevity extends to 90 years (oldest recorded 98 years). Natural mortality is estimated to be $M=0.06$. Females are viviparous, retaining their fertilized eggs within the ovary until larval extrusion. Mating takes place in late fall, and larval extrusion occurs in early spring. Females release from 10,000-300,000 eggs each year, depending on size.

Catch History: Soviet and Japanese trawl fisheries targeted POP throughout the 1960s. Catches in the EBS peaked at 47,000 mt in 1961 and in the AI in 1965 at 109,100 mt. Intense harvesting pressure reduced the stock biomass during that time, and catches declined through the mid 1980s. Foreign fisheries were replaced by joint ventures in the late 1980s, and the fishery was fully domesticated by 1990, with catches reaching 20,289 mt. Catches averaged 14,311 mt annually from 2004-2009.



Stock Assessment: The assessment uses an age-structured population dynamics model using AD Model Builder software. This model incorporates fishery data and fishery independent data from biennial trawl surveys. POP are managed under Tier 3a of the ABC/OFL control rules. The 2011 projected biomass is 600,600 mt. Catch specifications for 2011 are as follows; OFL=36,300 mt, ABC= 24,700 mt, TAC= 24,700 mt. Catch limits are further apportioned by AI subarea.

Estimated biomass declined significantly from 890,000 mt in 1960 to 107,000 mt in 1980. Biomass recovered during the late 1980s due to above-average year classes in the AI and reduced

exploitation rates. Estimated biomass averaged 408,000 mt annually from 2004-2009.

Fishery: POP are caught primarily in bottom trawl fisheries. The AI fishery opens in July and is concentrated during the summer months. Since 1996, the majority of the catch (by weight) occurred in the western Aleutians. In 2009, the discard rate for POP was 5%.

Fishery Management: In 1991, the POP and Other Red Rockfish complexes were separated from the POP/Other Rockfish complex. In 2001, the POP complex was separated into three management units; POP, Shortaker/Rougheye, and Sharpchin/Northern rockfish. In 2002, sharpchin rockfish were dropped from the complex due to sparse catches, leaving Northern rockfish as a single species management unit. In 2004, Shortraker and Rougheye rockfish were split into single species management units. In 2008, the Rougheye rockfish category was reclassified as a two species complex, Blackspotted and Rougheye rockfishes.

Rockfish are regulated under the BSAI groundfish FMP through permits, limited entry, catch quotas (TACs), seasons, in-season adjustments, gear restrictions, closed waters, bycatch limits and rates, allocations, regulatory areas, record keeping, reporting requirements and observer monitoring. In 2008, BSAI FMP Amendment 80 allocated 90-98% (depending on sub-area) of the AI Pacific ocean perch TAC, along with flatfish and Atka mackerel as catch shares among non-AFA trawl catcher processors according to their historic harvest patterns. Like other groundfish, 10.7% of rockfish is first allocated to CDQ groups.



Mark Fina, NPFMC

Total catches, pre-season catch specifications, and exploitable biomass of Pacific Ocean Perch in the BSAI, 2004-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2004	11,896	12,580	13,300	15,800	375,000
2005	10,426	12,600	14,600	17,300	382,000
2006	12,859	12,600	14,800	17,600	385,000
2007	18,468	19,900	21,900	26,100	457,000
2008	17,436	21,700	21,700	25,700	453,000
2009	14,779	18,800	18,880	22,300	402,000
2010	16,567	18,860	18,860	22,400	403,000
2011	-	24,700	24,700	36,300	600,600

*POP removed from POP Complex 2004.

¹Catch data current through November 2010.

²TAC, ABC and OFL data from annual Federal Register Harvest Specifications.

³Biomass from annual SAFE report projections.

Economics: In 2009, ex-vessel value of catch was \$7.2 million for all BSAI Rockfish. Production was 18,910 mt for all rockfish products in Alaska, with a gross value \$37.2 million. Four catcher vessels and 12 catcher processor vessels participated in rockfish fisheries in the BSAI in 2009. Primary products are H&G and whole fish. Rockfish product price averaged \$0.87/lb for at-sea processors and \$0.97/lb for shoreside processors.

Ecosystem Components: POP habitat use shifts with ontogeny. Juveniles are thought to remain in more rugged, rocky benthic environments, whereas adults move into deeper, less rough habitats. POP were also found to be associated with epibenthic sea pens and sea whips along the BS slope.



Northern Rockfish



AFSC, NOAA Fisheries

Biology: Northern rockfish *Sebastes polypsinus* distribution extends from the Kamchatka Peninsula, through the BSAI, GOA and British Columbia. This species is most abundant in the central GOA to the western end of the AI. Northern rockfish are demersal and are generally found in discrete aggregations with patchy distributions along the outer continental shelf from 75-150 m. They prey on calanoid copepods, euphausiids and chaetognaths. Based on stomach content data for POP, Pacific halibut and sablefish likely prey on Northern rockfish.

Northern rockfish is a relatively slow-growing, long lived species. Age at 50% maturity is 12.8 years in the GOA, and longevity extends to 50 years (oldest recorded 57 years). Natural mortality is estimated to be $M=0.043$. Females are viviparous, retaining their fertilized eggs within the ovary until larval extrusion.

Catch History: Foreign trawl fisheries were replaced by joint ventures in the 1980s, and the fishery was fully domesticated by 1990. Catches of Northern rockfish peaked in 1995 at 6,724 mt and ranged from 859-6,724 mt from 1990-2009. Catches from 2004-2009 averaged 3,800 mt annually.

Stock Assessment: The Northern rockfish assessment uses an age-structured population dynamics model using AD Model Builder software. This model incorporates fishery data and fishery independent data from biennial trawl surveys. Northern Rockfish are managed under Tier 3a of the ABC/OFL control rules. The 2011 projected biomass is 201,000 mt.



specifications for 2011 are as follows; OFL=10,600 mt, ABC=8,670 mt, TAC=4,000 mt.

Fishery: Northern rockfish are generally caught in bottom trawl fisheries targeting other species. Catches in the BSAI primarily occur within the Atka mackerel fishery, and historically, most (>80%) were discarded. Discard rates of Northern Rockfish have been decreasing over time, with discard rates of 16% in the Aleutian Islands and 37% in the Bering Sea. Important fishing grounds include Petral Bank, Surdevant Rock, south to Amchikta Island and Seguam Pass.



For More Information:

P. Spencer and J. Ianelli. 2010. Assessment of Northern Rockfish in the Bering Sea and Aleutian Islands Area.

<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Total catches, pre-season catch specifications, and exploitable biomass of Northern Rockfish in the BSAI, 2001-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2001	-	-	-	-	150,000
2002	-	6,760	6,760	9,020	150,000
2003	-	6,000	7,101	9,468	156,000
2004	4,684	5,000	6,880	8,140	142,000
2005	3,964	5,000	8,260	9,810	200,000
2006	3,824	4,500	8,530	10,100	204,000
2007	4,021	8,190	8,190	9,750	212,000
2008	3,287	8,180	8,180	9,740	212,000
2009	3,088	7,160	7,160	8,540	200,000
2010	4,039	7,240	7,240	8,640	203,000
2011	-	4,000	8,670	10,600	201,000

*Northern Rockfish removed from Other Rockfish catery 2001.

¹Catch data current through November 2010.

²TAC, ABC and OFL from annual Federal Register.

³Biomass data corresponds to the annual SAFE report projections.



Rougheye and Blackspotted Rockfish

Blackspotted rockfish
AFSC, NOAA Fisheries


Biology: The rougheye and blackspotted rockfish complex consists of 2 species: rougheye rockfish *Sebastes aleutianus* and blackspotted rockfish *S. melanostictus*. Blackspotted and rougheye rockfish are distributed from Japan, through the BSAI and GOA to southern California. Adults inhabit a narrow band along the upper continental slope at depths from 300-500 m. Data from recent bottom trawl surveys suggests that although the two species distributions overlap, blackspotted rockfish are predominant in the AI, while rougheye rockfish are more common in the GOA and southeastern BS.



Because both species are data limited, the stock assessments refer to the “Blackspotted and Rougheye” complex as one unit - rougheye rockfish. Rougheye rockfish length at 50% maturity is 44 cm., and longevity may extend to 200 years. Natural mortality is estimated at $M=0.034$. Rougheye rockfish prey primarily on shrimps, squids and myctophids.

Catch History: Rougheye rockfish catches were relatively high during the late 1970s and peaked in 1979 at 3,553 mt. Catches then declined in the 1980s as the foreign fishery was reduced. Catches increased again during the 1990s with the domestication of the fishery, averaging 800 mt from 1990-1999. Catches have decreased since 2001, averaging 179 mt from 2004-2009.

Stock Assessment: The rougheye rockfish assessment uses an age-structured population dynamics model using AD Model Builder software. This model incorporates fishery data and fishery independent data from biennial trawl surveys. Rougheye rockfish are managed under Tier 3, and blackspotted rockfish are managed under Tier 5 of the ABC/OFL control rules. The 2011



projected biomass is 21,200

mt. Catch specifications for 2011 are as follows; $OFL=549$ mt, $F_{OFL}=0.047$, $ABC=454$ mt, $TAC=454$ mt. The catch limits are further split into two units: Bering Sea/Eastern AI combined (2011 $ABC=TAC=234$ mt) and Central AI/Western AI combined (2011 $ABC=TAC=220$ mt).

Fishery: There is no directed fishery for these rockfish species in the BSAI. In the AI, they are primarily taken as incidental catch in the POP trawl fishery, and to a lesser extent the Atka mackerel trawl fishery and the Pacific cod longline fishery. In the EBS, rougheye rockfish are generally caught in the Pacific cod longline fishery and various bottom trawl fisheries.

For More Information:

P. Spencer and C. Rooper.
2010. Assessment of Blackspotted and Rougheye Rockfish Stock Complex in the Bering Sea and Aleutian Islands.

<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Exploitable biomass (mt), pre-season catch specifications (mt), and total catches (mt, including discards) of Blackspotted and Rougheye rockfish in the BSAI, 2004-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2004	208	195	195	259	10,400
2005	90	223	223	298	11,900
2006	203	224	224	299	11,900
2007	167	202	202	269	10,800
2008	214	202	202	269	10,800
2009	195	539	539	660	19,000
2010	232	547	547	669	21,200
2011	-	454	454	549	19,319

*Rougheye Rockfish removed from Other Rockfish catatory 2003.

¹Catch data current through November 2010.

²TAC, ABC and OFL from annual Federal Register.

³Biomass data corresponds to the annual SAFE report projections issued the preceding year.



Shortraker Rockfish


AFSC, NOAA Fisheries

Biology: Shortraker rockfish *Sebastes borealis* are distributed from southeastern Kamchatka, north through the BSAI, the GOA and south to California. Adults are concentrated along the 300-500 m depth interval along the continental slope. Shortraker rockfish predate on shrimps, squids and myctophids. Shortraker rockfish is one of the most long-lived species in the northeast Pacific. Age at 50% maturity is 45 cm, and longevity can exceed 140 years. Natural mortality is estimated to be $M=0.03$. Information on early life history stages of shortaker rockfish is limited.

Catch History: Catches of shortraker and Northern rockfish from 1992-2002 averaged 4,390 mt annually. Catches of shortraker rockfish averaged 3,800 mt annually from 2004-2009.

Stock Assessment: The shortaker rockfish assessment uses a simple surplus production mode (Gompertz-fox). This model incorporates fishery data and fishery independent data from biennial trawl surveys. Shortaker rockfish are managed under Tier 5 of the ABC/OFL control rules. The 2011 projected biomass is 17,452 mt. Catch specifications for 2011 are as follows; OFL=524 mt, ABC= 393 mt, TAC= 393 mt.

Fishery: Shortraker rockfish in the Aleutian Islands are primarily taken in rockfish trawl fisheries and longline fisheries targeting Greenland turbot, sablefish, and Pacific halibut. The central Aleutians comprised 58% of the 2004-2007 AI shortraker catch, followed by the western Aleutians (24%) and eastern Aleutians (18%). In the Eastern Bering Sea, catches of shortraker rockfish largely occur in midwater pollock trawl fisheries and longline fisheries for Pacific cod, Greenland turbot, and halibut. In 2010, 23.5% of the Shortraker rockfish were discarded.



AFSC, NOAA Fisheries



For More Information:


P. Spencer and C. Rooper. 2010. Assessment of the Shortraker Rockfish Stock in the Bering Sea and Aleutian Islands.

<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Total catches, pre-season catch specifications, and exploitable biomass of Shortraker Rockfish* in the BSAI, 2004-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2004	240	526	526	701	23,400
2005	169	596	596	794	26,500
2006	210	580	580	774	25,800
2007	323	424	424	564	18,900
2008	166	424	424	564	18,900
2009	196	387	387	516	17,200
2010	252	387	387	516	17,200
2011	-	393	393	524	17,452

*Shortraker Rockfish removed from Other Rockfish catetory 2003.
¹Catch data current through November 2010.
²TAC, ABC and OFL data from annual Federal Register.
³Biomass from annual SAFE report projections.



Other Rockfish



Shortspine thornyhead rockfish
AFSC, NOAA Fisheries



For More Information:

P. Spencer and C. Rooper. 2010. Assessment of Other Rockfish stock complex in the Bering Sea and Aleutian Islands.

<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Total catches, pre-season catch specifications and exploitable biomass of Other Rockfish in the BSAI, 1995-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1995	849	1,022	1,135	1,135	22,800
1996	642	1,304	1,449	1,449	20,700
1997	468	1,087	1,087	1,449	20,700
1998	588	1,054	1,054	1,492	20,300
1999	765	1,054	1,054	1,405	20,030
2000	840	1,054	1,054	1,405	20,030
2001	906	1,037	1,037	1,383	19,780
2002	952	1,037	1,037	1,383	19,780
2003	737	1,594	1,594	2,126	19,780
2004	655	1,594	1,594	2,126	20,400
2005	464	1,050	1,400	1,870	20,400
2006	579	1,050	1,400	1,870	26,600
2007	652	999	999	1,330	26,700
2008	597	999	999	1,330	36,700
2009	565	1,040	1,040	1,380	39,700
2010	676	1,040	1,040	1,380	39,200
2011	-	1,000	1,280	1,700	44,939

¹Catch data current through November 2010.

²TAC, ABC and OFL from annual Federal Register.

³Biomass from annual SAFE report projections. Biomass includes sharpchin prior to 2003.

Biology: The Other Rockfish complex consists of 7 rockfish species. Shortspine thornyheads and dusky are the two most abundant species for this complex, accounting for about 80% of the survey biomass and fishery catch. Data are limited for many of the "Other Rockfish" complex species.

Dusky rockfish distribution extends from Japan into the BSAI and down to central Oregon. Dusky rockfish are found along the outer continental shelf in patchy distributions. Natural mortality is estimated at $M=0.09$. Dusky rockfish are viviparous. Dusky rockfish longevity is approximately 60 years. Shortspine thornyheads are distributed from Japan to the BSAI down to central California. Shortspine thornyheads are commonly found at depths from 150-450 m. Natural mortality is estimated at $M=0.03$, and shortspine thornyhead longevity extends to 100 years or more. In contrast to many other *Sebastes spp.*, shortspine thornyheads are oviparous.

Catch History: Other Rockfish have been caught in trawl fisheries since Japanese and Soviet fleets began fishing in the BS in the 1960s. Catches of "Other Rockfish" have been tracked since 1977. Catches were relatively high in the BSAI from 1977-1983, ranging annually from 700-2,300 mt. Catches have remained relatively stable from 1993-2009, averaging 677 mt annually.

Stock Assessment: Other Rockfish are managed under Tier 5 of the ABC/OFL control rules. The 2011 projected biomass is 44,939 mt. Catch specifications for 2011 are as follows; OFL=1,700 mt, ABC= 1,280 mt, TAC= 1,000 mt. ABC and TAC are further subdivided into BS and AI limits.

Estimated biomass of Other Rockfish remained relatively stable from the mid-1980s through 2005, and has since increased to about 40,000 mt in recent years.

Fishery: There is no directed fishery for Other Rockfish in the BSAI. Dusky rockfish are primarily taken in the Atka mackerel fishery in the AI and the EBS Pacific cod fishery. Shortspine thornyhead are primarily taken in the AI sablefish and Greenland turbot longline fisheries and EBS pollock trawl fishery.



AFSC, NOAA Fisheries



AFSC, NOAA Fisheries



For More Information:

S. Lowe, J. Ianelli, M. Wilkins, K. Aydin, R. Lauth, and I. Spies. 2010. Assessment of the Atka mackerel stock in the Bering Sea and Aleutian Islands.
<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Atka Mackerel

Biology: Atka mackerel *Pleurogrammus monopterygius* is a schooling, semi-demersal species most commonly found in the AI. Adults occur in large localized aggregations at depths less than 200 m over rough, uneven bottom areas with high tidal currents. Atka mackerel move off the bottom during daylight hours presumably to feed on their main prey items, euphausiids and copepods. They begin to recruit to the fishery at age 3 and longevity can extend to 14 years. Females reach 50% maturity at 31 cm (3.5 years). Natural mortality is estimated at $M=0.30$. Atka mackerel are a substrate spawning fish with male parental care. Behavioral studies have shown that the Atka mackerel mating system is very complex. A significant characteristic is the bright and distinct coloration developed by territorial males during the spawning season. Spawning occurs from July to October, peaking in early September. Atka mackerel have relative low fecundity, with females releasing around 30,000 eggs each year. Eggs are adhesive and deposited in rock crevices in nests guarded by males until hatching, which occurs about 40-45 days later.

Catch History: Beginning in 1970, USSR, Russia and Korea harvested Atka mackerel; foreign catches peaked in 1978 at 24,000 mt. Catches have generally increased since the fishery was fully harvested by U.S. vessels. Peak catch occurred in 1996 (104,000 mt).

Fishery Management: The Atka mackerel fishery is tightly managed to minimize the potential for prey competition with Steller sea lions, including seasonal allowances of TAC and spatial distribution of the fishery away from critical habitat. Additionally, trawling for Atka mackerel was prohibited within 10 nm of all rookeries in areas 542 and 543. Another Steller Sea Lion Biological Opinion (Biop) was released in August 2010, which included fishery closures for mackerel in area 543 and restrictions in 541 and 542.

Since 2008 with the implementation of Amendment 80, the fishery has operated as a catch share fishery, with participants operating as cooperatives. In addition, 2% of the TAC is set aside to vessels using jig gear, and 10.7% is allocated to CDQ groups.

Stock Assessment: Since 2002, the Atka mackerel assessment has used the Assessment Model for Alaska. This model incorporates fishery data and fishery independent data from trawl surveys. Atka mackerel fall under Tier 3a of the ABC/OFL control rules. The 2011 estimated biomass is 438,000 mt. Catch specifications for 2011 are as follows; OFL=101,000, ABC=85,300 mt, TAC=53,080 mt.

Atka mackerel biomass increased during the early 1980s and again in the late 1980s to early 1990s. The most recent peak in female spawning biomass occurred in 2005. Increases in abundance are attributed to series of strong year classes that support the population.

Fishery: Atka mackerel are targeted by trawl catcher processors. Products include whole fish and H&G. One catcher vessel and 12 catcher processors participated in the 2009 BSAI Atka mackerel trawl fishery. Ex-vessel value of catch was \$29.7 million in 2008.

Total catches, pre-season catch specifications, and exploitable biomass of Atka Mackerel in the BSAI, 1998-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1998	57,096	64,300	64,300	134,000	536,000
1999	53,644	66,400	73,300	148,000	595,000
2000	47,229	70,800	70,800	119,000	565,000
2001	61,560	69,300	69,300	138,000	553,000
2002	45,294	49,000	49,000	82,300	439,700
2003	59,350	60,000	63,000	99,700	358,000
2004	60,564	63,000	66,700	99,700	286,000
2005	62,014	63,000	124,000	178,500	486,000
2006	61,883	63,000	110,200	147,000	446,000
2007	58,831	63,000	74,000	86,900	364,000
2008	58,088	60,700	60,700	71,400	323,000
2009	72,273	76,400	83,800	99,400	411,000
2010	68,643	74,000	74,000	88,200	388,000
2011	-	53,080	85,300	101,000	438,000

¹Catch data current through November 2010.

²1988-2010 TAC, ABC and OFL from Federal Register

³Biomass data from SAFE report projections.



AFSC, NOAA Fisheries



For More Information:

O. Ormseth and C. Spital. 2010. Bering Sea and Aleutian Islands Squids. <http://www.afsc.noaa.gov/refm/stocks/assessments.htm>

Biology: There are 14 species in the “Squid” complex in the BSAI. The most abundant species in the BS is the magistrate armhook squid *Berryteuthis magister*. The boreal clubhook squid *Onychoteuthis borealijaponicus* is the most common in the AI. Squid in the BSAI are generally pelagic, however, the North Pacific bobtail squid, magistrate armhook squid and Pacific bobtail squid are often found in close proximity to the bottom. Most species are associated with the slope and basin, with the highest species diversity along the slope region of BS between 200–1500 m. They are active predators, swimming by jet propulsion. Squids prey on euphausiids, zooplankton and small fish and are prey for fish such as Atka mackerel and grenadiers, and marine mammals such as SSLs and fur seals.

Squids are productive, short-lived animals. Squid display rapid growth, patchy distribution and variable recruitment patterns. Populations of the magistrate armhook squid are complex and are made up of multiple cohorts spawned throughout the year. Magistrate squid are dispersed throughout the summer months in the western BS but form large, dense schools over the continental slope between September and October. Three seasonal cohorts are identified in the region; summer-hatched, fall-hatched and winter-hatched. Growth, maturation and mortality rates vary between cohorts. Juvenile and adult magistrate squid also appear to be separated vertically in the water column. Most squid are generally thought to live less than 2-3 years.

Chiroteuthid sp.	<i>Chiroteuthis calyx</i>
Glass squid sp.	<i>Belonella borealis</i>
Glass squid sp.	<i>Galiteuthis phyllura</i>
Minimal armhook squid	<i>Berryteuthis anonychus</i>
Magistrate armhook squid	<i>Berryteuthis magister</i>
Armhook squid	<i>Eogonatus tinro</i>
Boreopacific armhook squid	<i>Gonatopsis borealis</i>
Berry armhook squid	<i>Gonatus berryi</i>
Armhook squid sp.	<i>Gonatus madokai</i>
Armhook squid sp.	<i>Gonatus middendorffi</i>
Clawed armhook squid	<i>Gonatus onyx</i>
Robust clubhook squid	<i>Moroteuthis robusta</i>
Boreal clubhook squid	<i>Onychoteuthis borealijaponicus</i>
North Pacific bobtail squid	<i>Rossia pacifica</i>

Total catches and pre-season catch specifications of Squid in the BSAI, 2007-2011.

Year	Catch	TAC	ABC	OFL
2007	1,188	1,970	1,970	2,620
2008	1,542	1,970	1,970	2,620
2009	353	1,970	1,970	2,620
2010	402	1,970	1,970	2,620
2011	-	425	1,970	2,620

*Squid biomass data unavailable.

Catch history: Japanese and Korean trawl fisheries targeted squid during the 1960s and 1970s; catches peaked in 1978 at 9,000 mt. Catches remained low throughout the late 1980s and early 1990s.

Fishery Management: In 2001, Amendment 66 lifted potential restrictions of allocated squid to CDQ groups, reducing potential for CDQ squid catches to constrain their target fisheries.

Stock Assessment: Squid fall under Tier 6 of the OFL/ABC control rules, and catch specifications are therefore based on the average catch of squid between 1978-1995. Squid estimated biomass is undefined. Catch specifications for squid in 2011 are as follows: OFL=2,620 mt, ABC=1,970 mt, TAC=425 mt.

Fishery: Squid are not a target fishery, and are primarily taken as incidental catch in the pelagic trawl pollock fishery. Discard rates ranged from 16%-88% discarded from 1998-2009.

Ecosystem Components: Squid important components in the diets of many seabirds, fish and marine mammals. Overall fishing removals of squid are low (especially relative to natural predation).



Bigmouth sculpin
AFSC, NOAA Fisheries



For More Information:

O. Ormseth and T. TenBrink. 2010. Bering Sea and Aleutian Islands Sculpins.
<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: There are a total of 48 species of sculpins in the BSAI, with 41 species identified in the Eastern Bering Sea and 22 species in the Aleutian Islands region. Sculpins occupy all benthic habitats and depths. The six species with the highest biomasses include great sculpin (*Myoxocephalus polyacanthocephalus*), threaded sculpin (*Gymnocanthus pistilliger*) plain sculpin (*M. jaok*), warty sculpin (*M. verrucosus*), bigmouth sculpin (*Hemitripterus bolini*), and yellow Irish lord (*H. jordani*).

There is limited BSAI-specific data on age and growth, maturity, or reproductive biology for sculpins identified in this management region. Most if not all sculpins lay adhesive eggs in nests, and many exhibit parental care for eggs.

Catch history: Based on total catch estimates from 1998-2008, sculpins comprised 19-28% of the total Other Species catch during this time period. Sculpins are caught by a wide variety of fisheries, but trawl fisheries for yellowfin sole, Pacific cod, walleye pollock, Atka mackerel and flathead sole, and Pacific cod hook-and-line fishery catch the most. Catches from 2000-2008 ranged from 5,735 mt to 7,670 mt per year.

Fishery Management: Prior to 2011, sculpins were managed as part of the BSAI Other Species complex that included sculpins, skates, sharks, and octopuses. Catch limits for this complex were set by summing the individual ABCs and OFLs for each species group to create an aggregate OFL, ABC, and TAC. Beginning in 2011 the complex was eliminated and each group is managed separately. Sculpins are currently taken only incidentally. For the near future, catch of sculpins will continue to be dependent on the distribution and limitations placed on target fisheries, rather than on any harvest level established for this category.



Blob sculpins
AFSC, NOAA Fisheries

Stock Assessment: Sculpins fall under Tier 5 of the OFL/ABC control rules. Sculpin biomass is estimated to be 208,000 mt in 2011. Catch specifications for sculpins in 2011 are as follows; OFL=58,300 mt, ABC=43,700 mt, TAC=5,200 mt.

Total catches, pre-season catch specifications, and exploitable biomass of Sculpins in the BSAI, 2008-2011.					
Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2008	6,473	-	-	-	-
2009	5,774	-	-	-	-
2010	3,928	-	-	-	-
2011	-	5,200	43,700	58,300	208,000

¹Catch data current through November 2010.
²TAC, ABC and OFL from SAFE
³Biomass data from SAFE report projections.

Fishery: There is currently no target fishery for sculpins in the BSAI, and virtually all are discarded or made into meal. Incidental catches of sculpins are taken in the Pacific cod and Atka mackerel fisheries in the AI, and in the Pacific cod rock sole, and yellowfin sole fisheries in the EBS.



Alaska skate
Beth Matta
AFSC, NOAA Fisheries



For More Information:

O. Ormseth, B. Matta, and J. Hoff. 2010. Bering Sea and Aleutian Islands Skates.

<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: Skates are cartilaginous fishes with large pectoral “wings” attached to the sides of the head. There are 15 species of skates in the BSAI in three genera, *Raja*, *Bathyraja*, and *Amblyraja*. The EBS shelf skate complex is dominated by a single species, the Alaska skate (*Bathyraja parmifera*), occurring at depths of 50 to 200 m. The Bering or sandpaper skate (*B. interrupta*) is the next most common species on the EBS shelf, and is distributed on the outer continental shelf. The dominant species on the EBS slope is the Aleutian skate (*B. aleutica*). A number of other species are found on the EBS slope in significant numbers, including the Alaska skate, Commander skate (*B. lindbergi*), whiteblotched skate (*B. maculata*), whitebrow skate (*B. minispinosa*), rougtail skate (*B. trachura*), and mud skate (*B. taranetzi*). Two rare species, the deepsea skate (*B. abyssicola*) and roughshoulder skate (*Amblyraja badia*), have only recently been reported from EBS slope bottom trawl surveys (Stevenson and Orr 2005). The skate complex in the AI is quite distinct from the EBS shelf and slope complexes, with different species dominating the biomass, as well as at least one endemic species, the recently described butterfly skate, *Bathyraja mariposa*. In the AI, the most abundant species is the whiteblotched skate, *B. maculate*.

Skate life cycles are similar to sharks, with relatively low fecundity, slow growth to large body sizes, and dependence of population stability on high survival rates of a few well developed offspring. Little is known about life history parameters of Alaskan skates. Studies done elsewhere have determined age at maturity and maximum age for big skates and longnose skates to be about 12 to 26 years with maturity occurring at approximately 8 years.



AFSC, NOAA Fisheries

Fishery Management: Until 2011, skate species were managed as part of the “Other species” management category within the BSAI FMP. Beginning in 2011, skates will be managed as a single complex with skate specific ABC and OFL. Currently skates are taken only as bycatch in fisheries directed at target species in the BSAI, so future catches of skates are more dependent on the distribution and limitations placed on target fisheries than on any harvest level established for this category.

Stock Assessment: Catch specifications for the Alaska skate is based on Tier 3a, and for other skates Tier 5 of the ABC/OFL control rules. Skate biomass is projected at 612,000 mt in 2011. Catch specifications for skate in 2011 are as follows; OFL=37,800 mt, ABC=31,500 mt, TAC=16,500 mt.

Total catches, pre-season catch specifications, and exploitable biomass of Skates in the BSAI, 2008-2011.					
Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2008	20,220	-	-	-	-
2009	19,392	-	-	-	634,000
2010	15,190	-	-	-	608,000
2011	-	16,500	31,500	37,800	612,000

¹Catch data current through November 2010.
²TAC, ABC and OFL from SAFE
³Biomass from SAFE report projections.

Fishery: There is currently no target fishery for skates in the BSAI. Most of the skate is caught incidentally in the hook and line fishery for Pacific cod, and trawl fisheries for pollock and flatfish. Between 24% and 39% of the total observed skate catch was retained during the years 2003-2006, primarily consisting of Aleutian skate and Alaska skate.



Salmon shark
AFSC, NOAA Fisheries

Biology: The shark complex consists of 8 species. The species most likely to be encountered in BSAI fisheries and surveys are the Pacific sleeper shark (*Somniosus pacificus*), the spiny dogfish (*Squalus acanthias*), and the salmon shark (*Lamna ditropis*). Sharks are long-lived species with slow growth to maturity, a large maximum size, and low fecundity.

Spiny dogfish tend to segregate by sex and by size. Age-at-50%-maturity for spiny dogfish is estimated to be 35 years for females, and 19 years for males. Spiny dogfish may live up to 100 years, and exhibit very slow growth rates. Spiny dogfish are aplacental viviparous, and embryos are nourished solely by their yolk sac, with a gestation period of 18-24 months.



[For More Information:](#)

C. Tribuzio, K. Echave, C. Rodgveller, J. Heifetz, and K. Goldman. 2010. Assessment of Sharks in the Bering Sea and Aleutian Islands Area. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Pacific sleeper sharks can attain large size. The maximum lengths of captured and measured Pacific sleeper sharks are 440 cm for females and 400 cm for males, although there are reports of individuals of 700 cm in length. Like spiny dogfish, Pacific sleeper sharks are aplacental viviparous.

Salmon sharks also grow relatively large, attaining a maximum length of 215 cm precaudal length for females and about 190 cm for males. Maximum ages for salmon sharks are 17 years for males and 30 years for females. They are thought to live up to 30 years.

Catch history: Incidental catches of shark species in the BSAI fisheries have been very small compared to catches of target species. Sharks have only been reported to species in the catch since 1997 and have made up from 1% to 5% of Other Species catch from 1997 – 2008. Pacific sleeper shark make up 60% of the total shark catch in the BSAI, followed by unidentified sharks at 20%, salmon shark at 9% and spiny dogfish at 2%.

Fishery Management: Shark species were managed as part of the “Other species” management category until 2011, when sharks will be managed as a single complex with shark specific OLF, ABC, and TAC.

Stock Assessment: Sharks fall under Tier 6 of the OFL/ABC control rules, and catch specifications are based on the maximum catch from 1997-2007 (1,362 mt in 2002). Sharks estimated biomass is undefined. Catch specifications for sharks in 2011 are as follows; OFL=1,360 mt, ABC=1,020 mt, TAC= 50 mt. Directed fishing for this species will be prohibited in 2011.

Fishery: There is currently no target fishery for sharks in federally or state managed waters of the BSAI, and most incidentally captured sharks are not retained. Dogfish are at the northern edge of their range in the BSAI but a few are taken in Pacific cod longline fishery (7 mt in 2010). About 90% of the salmon sharks are taken in the Pollock fishery (11 mt in 2010). Sleeper sharks are taken in higher amounts, particularly in the Pacific cod and pollock fisheries, but with some taken in all groundfish fisheries (17 mt total in 2010, but much higher in previous years; 839 mt in 2002).

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2008	186	-	-	-	-
2009	144	-	-	-	-
2010	47	-	-	-	-
2011	-	50	1,020	1,360	Undefined

¹Catch data current through November 2010.
²TAC, ABC and OFL from SAFE
³Biomass from SAFE report projections.



Giant Pacific octopus
Rex Murphy



For More Information:

M.E. Conners and C. Conrath. 2010. Assessment of the Octopus Complex in the Bering Sea and Aleutian Islands Area. <http://www.afsc.noaa.gov/refm/stocks/assessments.htm>

Biology: There are at least 7 species of octopus present in the BSAI, and the species composition both of natural communities and commercial harvest is unknown. Some species, particularly *G. boreopacifica*, are primarily distributed at greater depths than are commonly fished. At depths less than 200 meters *E. dofleini* appears to be the most abundant species.

Octopus life spans are either 1-2 years or 3-5 years depending on the species.


E. dofleini are estimated to mature at 1.5 – 3 years. *E. dofleini* is a terminal spawner,

females die after the eggs hatch while males die shortly after mating. The fecundity of this species in Japanese waters has been estimated at 30,000 to 100,000 eggs per female. Based on larval data, *E. dofleini* is the only octopus in the Bering Sea with a planktonic larval stage.

Giant Pacific octopus	<i>Enteroctopus dofleini</i>
Smoothskin octopus	<i>Benthoctopus leioderma</i>
Flapjack devilfish	<i>Opisthoteuthis californiana</i>
Small pelagic octopus	<i>Japattella diaphana</i>
Stubby octopus	<i>Sasakiopus salebrosus</i>
A deepwater octopus	<i>Graneledone boreopacifica</i>
A deepwater octopus	<i>Benthoctopus oregonesis</i>

Fishery Management: Until 2011, octopus were managed as part of the “Other species” management category within the BSAI FMP. Beginning in 2011, octopuses will be managed as a single complex with specific OFL, ABC, and TAC.

Stock Assessment: Octopus fall under Tier 6 of the OFL/ABC control rules, and catch specifications are based on the average incidental catch 1997-2007. The maximum catch over that time period was 528 tons. There are no historical catch records for octopus, and their biomass has not been estimated. Catch specifications for octopus in 2011 are as follows; OFL=528 mt, ABC=396 mt, TAC=150 mt. Directed fishing for this species will be prohibited in 2011.



Total catches, pre-season catch specifications, and exploitable biomass of Octopus in the BSAI, 2008-2011.

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
2008	191	-	-	-	-
2009	51	-	-	-	-
2010	117	-	-	-	-
2011	-	150	396	528	Undefined

¹Catch data current through November 2010.

²TAC, ABC and OFL from SAFE

³Biomass from SAFE report projections.

Fishery: There is currently no target fishery for octopus in the BSAI. Octopus are taken as incidental catch in trawl, longline, and pot fisheries throughout the BSAI; the highest catch rates are from Pacific cod pot fisheries in the three statistical areas around Unimak Pass. The species composition of the octopus community is not well documented, but recent research indicates that the giant Pacific octopus *Enteroctopus dofleini* is most abundant in shelf waters and predominates in commercial catch.



AFSC, NOAA Fisheries

Walleye Pollock

Biology: Walleye Pollock *Theragra chalcogramma* is an abundant fish species in the GOA, found throughout the shelf regions at depths less than 300 m. Seasonal migrations occur from overwintering areas along the outer shelf to shallower waters (30-140 m) to spawn. Pollock feed on copepods, euphausiids and fish and are prey for other fish, marine mammals and seabirds. Pollock begin to recruit to the fishery at age 3 and longevity extends to 12 years or more (oldest GOA Pollock observed is 22 years). Females reach 50% maturity at approximately 43 cm (ages 4-6), and adults produce 60,000 to 400,000 pelagic eggs. Annual natural mortality is estimated to be $M=0.30$. Peak spawning in the GOA occurs from February to March in the Shumagin Islands and late March in the Shelikof Strait.



For More Information:

M. Dorn, K. Aydin, S. Barbeaux, M. Guttormsen, K. Springer, and M. Wilkins. 2010. Assessment of Walleye Pollock stock in the Gulf of Alaska. <http://www.afsc.noaa.gov/refm/stocks/assessments.htm>

Catch History: Foreign fisheries for pollock developed in the GOA in the early 1970s and peak foreign catches occurred in 1981 at 130,324 mt. A late spawning aggregation was discovered in Shelikof Strait in 1981, and a valuable pollock roe fishery was established in the region. US vessels entered the pollock fishery in 1977 and by 1988, the fishery was fully harvested by the domestic fleet.


Fishery Management: The GOA pollock fishery is regulated under the GOA groundfish FMP through permits and limited entry, catch quotas (TACs), seasons, in-season adjustments, gear restrictions, closed waters, bycatch limits and rates, allocations, regulatory areas, record keeping, reporting requirements and observer monitoring. In 1993, 100% of GOA pollock was apportioned to the inshore sector (vessels that catch fish to deliver to shore based processing plants). In 1998, trawl gear was prohibited east of 140°W, and 100% retention was required for pollock.



AFSC, NOAA Fisheries

Since 1992, GOA pollock catch has been apportioned spatially and temporally to reduce fishery impacts on Steller sea lions (SSLs). Additional SSL protection measures implemented in 2001 established 4 seasons in the Central and Western GOA beginning in January, March, August and October (25% TAC to each season). Additionally, a harvest control rule was

implemented that requires suspension of directed pollock fishing if and when spawning biomass declines below 20%.




Stock Assessment: The GOA pollock assessment is based on an age-structured model. This model incorporates fishery data and fishery independent data from annual bottom trawl surveys and acoustic trawl surveys. GOA Pollock fall under Tier 3b of the ABC/OFL control rules. The 2011 age 3+ biomass is estimated at 893,700 mt. Gulf wide catch specifications for 2011 are as follows; OFL=118,030 mt, ABC=86,970 mt, TAC=86,970 mt. The catch limits are further spatially apportioned into Western, Central area

62, Central area 63, West Yakutat, and Eastern GOA.

Age 3+ GOA pollock model-estimated biomass was high during the early 1980s. Biomass declined through the late 1980s and has remained below target as a result of below average recruitment.

Fishery: The directed fishery is prosecuted by vessels using trawl gear, primarily with pelagic trawls. Small amounts of pollock are also taken



Total catches, pre-season catch specifications, and exploitable biomass of age 3+ Walleye Pollock in the GOA, 1976-2011 (in mt).

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1976	86,527	-	-	-	-
1977	118,356	-	-	-	-
1978	96,935	-	-	-	-
1979	105,748	-	-	-	-
1980	114,622	-	-	-	-
1981	147,744	-	-	-	-
1982	168,740	-	-	-	-
1983	215,608	-	-	-	-
1984	307,401	-	-	-	-
1985	284,826	293,250	-	-	-
1986	87,809	116,600	116,600	-	496,300
1987	69,751	108,000	112,000	-	687,100
1988	65,739	93,000	93,000	-	687,000
1989	78,392	60,200	63,400	-	593,000
1990	90,744	93,000	93,000	-	891,000
1991	100,488	133,400	133,400	-	1,303,000
1992	90,857	87,400	99,400	227,900	838,000
1993	108,908	114,400	160,400	295,020	1,062,000
1994	107,335	109,300	109,300	246,600	726,000
1995	72,618	65,360	65,360	280,400	573,000
1996	51,263	54,810	54,810	86,400	574,000
1997	90,130	79,980	79,980	112,270	1,105,420
1998	125,098	124,730	130,000	186,100	1,156,000
1999	95,590	100,920	100,920	146,000	737,670
2000	73,080	100,000	100,000	139,370	616,710
2001	72,076	95,875	105,810	126,360	727,710
2002	51,937	58,250	58,250	84,090	755,310
2003	50,666	54,350	54,350	78,020	699,120
2004	63,934	71,260	71,260	99,750	769,420
2005	80,846	91,710	91,710	153,030	765,180
2006	71,976	86,807	86,807	118,309	635,732
2007	53,062	68,307	68,307	95,429	861,072
2008	52,500	51,940	51,940	83,150	741,819
2009	44,003	49,900	49,900	69,630	675,749
2010	75,500	84,745	84,745	115,536	797,638
2011	-	86,970	86,970	118,030	893,700

as bycatch in other fisheries. A total of 63 catcher vessels participated in the 2009 GOA directed pollock trawl fishery. About 65% of the catch is landed in Kodiak. Approximately 95% of the catch is pollock in the directed fishery, with incidental catches mainly consisting of arrowtooth flounder, Pacific cod, flathead sole and squid.

Economics: In 2009, ex-vessel value of the catch was \$15.3 million for GOA pollock. Average ex-vessel price paid for GOA Pollock in 2009 was \$0.17/lb. round weight. Primary products were surimi, roe, fillets, H&G, and other products.

Ecosystem Components: In the GOA, the main predators of pollock are arrowtooth flounder, Pacific halibut, Pacific cod, Steller sea lions and the directed pollock fishery. For pollock less than 20 cm, arrowtooth flounder represents close to 50% of total mortality, and the abundance of arrowtooth flounder has increased dramatically in the GOA since the 1980s.



¹Catch data from SAFE report through November 2010.
²1988-2010 TAC, ABC and OFL data from annual Federal Register Harvest Specifications. Does not include EYAK and SEO.
³Biomass from annual SAFE report projections.



Diana Evans, NPFMC

Biology: Pacific cod *Gadus macrocephalus* is a demersal species found in the eastern BS, the AI, and GOA down to central California. Juveniles are typically distributed over the inner continental shelf at depths from 60-150 m. Adults are found at depths from shoreline to 500 m. Mature fish tend to concentrate on the outer continental shelf and prefer muddy or sandy soft sediment substrate. Juveniles feed primarily on small invertebrates and euphausiids. Adult Pacific cod feed on fish such as juvenile pollock, and invertebrates such as polychaetes, amphipods and crangonid shrimp. Predators of Pacific cod include adult Pacific cod, Pacific halibut, salmon sharks and Steller sea lions.

Pacific cod are a relatively fast growing and short-lived fish. Longevity can extend to 19 years. Pacific cod begin to recruit to the fishery around 3 and are 50% recruited by age 7. Natural mortality is estimated at $M=0.38$. Females reach 50% maturity at 50 cm (4-5 years) and larger fish can produce more than 1 million eggs. Adults form spawning aggregations from January to May in the GOA.



For More Information:

G. Thompson, J. Ianelli, and M. Wilkins. 2010. Assessment of Pacific Cod stock in the Gulf of Alaska. <http://www.afsc.noaa.gov/afsc/refm/stocks/assessments.htm>


Catch History: Pacific cod were harvested by foreign fleets targeting higher-value species during the 1970s. By 1976, catches increased to 6,800 mt, and the foreign fishery peaked in 1981 at 35,000 mt. A small joint venture fishery existed through 1988, averaging about 1,400 mt annually. The domestic fishery increased through 1986 and tripled its catch in 1987 to a catch of nearly 31,000 mt. The GOA Pacific cod fishery was fully harvested by domestic vessels in 1987.



SeaAlliance/AGDB

Fishery Management: Pacific cod is regulated under the GOA groundfish FMP through permits, limited entry, catch quotas (TACs), seasons, in-season adjustments, gear restrictions, closed waters, bycatch limits and rates, allocations, regulatory areas, record keeping, reporting requirements and observer monitoring. In 1992, Amendment 23 allocated 90% of GOA Pacific cod to the inshore sector and 10% to the offshore sector. In 1998, trawl gear was prohibited in the East Yakutat/Southeast subareas, and 100% retention of Pacific cod was required. In 2009, the Council passed Amendment 83, which will allocate Western and Central GOA Pacific cod TAC among gear and operation type, based on historic dependency and use by sector, and creates additional entry-level opportunities for jig vessels. If approved, this allocation could be in effect in 2012.

Separate TACs are currently identified for Pacific cod in the Western, Central and Eastern GOA regulatory areas. Within the Central and Western Regulatory Areas, 60% of each component's portion of the TAC is allocated to the A season (January 1 through June 10) and the remainder is allocated to the B season (June 11



through December 31). Longline and trawl fisheries are also associated with a Pacific halibut mortality limit, which can constrain the magnitude and timing of harvests taken by these two gear types.

Stock Assessment: The Pacific cod assessment is based on a Stock Synthesis model that uses both length-structured and age-structured data. This model incorporates fishery data and fishery independent data from the NMFS trawl surveys.

Pacific cod catch limits are set by a Tier 3a ABC/OFL control rule. The 2011 age 3+ biomass is estimated at 428,000 mt for GOA Pacific Cod. Catch specifications for 2011 are as follows; OFL=102,600 mt, ABC=86,800 mt, TAC=65,100 mt. Separate ABCs and TACs are established

for Western, Central, and Eastern GOA. Since 1997, the Council has reduced the


TAC in each area by up to 25% to account for removals in the State waters Pacific cod fishery.

Estimated biomass of Pacific cod peaked in the early 1980s, and then slowly declined as the exceptional 1977 year class gradually exited the population. Estimated biomass appears to be increasing in the short term due to above average recruitment in recent years.

Fishery: The Pacific cod fishery is the second major species (after pollock) targeted in the commercial groundfish catch in the GOA. Pacific cod are taken with trawl, longline, pot and jig gear. Participants in the 2009 GOA directed fishery included 240 vessels using longlines or jig gear, 125 vessels using pot gear, and 64 vessels using trawl gear. Primary bycatch species in the Pacific cod fishery include arrowtooth flounder, and skates and pollock.

Economics: In 2009, ex-vessel value of Pacific cod catch in the GOA was \$23 million, and exvessel price averaged \$0.28/lb round weight. Primary products include whole fish, H&G and fillets.

Ecosystem components: Pacific cod are a prey item for Steller sea lions in the GOA and BSAI.



Total catches, pre-season catch specifications, and exploitable biomass of age 3+ Pacific Cod in the GOA, 1976-2011 (in mt).

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1976	6,764	-	-	-	-
1977	2,267	-	-	-	-
1978	12,190	-	-	-	-
1979	14,904	-	-	-	-
1980	35,345	60,000	-	-	-
1981	36,131	70,000	-	-	-
1982	29,465	60,000	-	-	-
1983	36,540	60,000	-	-	-
1984	23,896	60,000	-	-	-
1985	14,428	60,000	136,000	-	-
1986	25,012	75,000	125,000	-	-
1987	32,939	50,000	125,000	-	-
1988	33,802	80,000	99,000	-	481,700
1989	43,293	71,200	71,200	-	558,700
1990	72,517	90,000	90,000	-	498,044
1991	76,997	77,900	77,900	-	424,100
1992	80,100	63,500	63,500	87,600	363,000
1993	56,488	56,700	56,700	78,100	324,000
1994	47,485	50,400	50,400	71,100	296,000
1995	68,985	69,200	69,200	126,000	573,000
1996	68,280	65,000	65,000	88,000	557,000
1997	77,018	69,115	81,500	180,000	650,000
1998	72,525	66,060	77,900	141,000	785,000
1999	81,785	67,835	84,400	134,000	648,000
2000	66,560	59,800	76,400	102,000	567,000
2001	51,542	52,110	67,800	91,200	526,000
2002	54,483	44,230	57,600	77,100	428,000
2003	52,579	40,540	52,800	70,100	428,000
2004	56,625	48,033	62,810	102,000	484,000
2005	47,585	44,433	58,100	86,200	472,000
2006	47,854	52,264	68,859	95,500	453,000
2007	51,462	52,264	68,859	97,600	375,000
2008	58,963	50,269	64,493	88,660	233,310
2009	52,922	41,807	55,300	66,000	520,000
2010	76,171	59,563	79,100	94,100	701,200
2011	-	65,100	86,800	102,600	428,000


¹Catch includes state waters fishery catch.

²TAC, ABC and OFL data from Federal Register.

³Biomass from annual SAFE report projections issued the preceding year.



Megan Peterson, UAF



AFSC, NOAA Fisheries

Biology: Sablefish *Anoplopoma fimbria* distribution extends from the northern Mexico through the Gulf of Alaska, the AI and into the BS. Adult sablefish are generally found at depths greater than 200 m along the continental slope, shelf gullies and deep fjords. Juvenile sablefish (less than 40 cm) spend the first 2-3 years farther inshore along the continental shelf and begin to move out to the continental slope around age 4. Young-of-the-year sablefish feed primarily on euphausiids and copepods while adults are more opportunistic feeders, relying more heavily on pollock, Pacific herring, Pacific cod, squid and jellyfish. Coho and Chinook salmon are the main predators of young-of-the-year sablefish.



Sablefish are relatively long lived. They begin to recruit to the fishery at age 4 or 5 and longevity often reaches 40 years (the oldest recorded sablefish in Alaska was 94 years old). Female size at 50% maturity is around 65 cm (approximately age 6.5). Females are slightly larger than males, and natural mortality is estimated at $M=0.10$. Alaskan sablefish spawn at pelagic depths near the edges of the continental slope (300-500m) between January and April.



For More Information:

D. Hanselman, C. Lunsford, and C. Rodveller. 2010. Assessment of Sablefish stock in Alaska. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Catch History: US fishermen have harvested sablefish (blackcod) since the end of the 19th century as a byproduct of halibut fisheries. Harvests were relatively small, averaging 1,666 mt from 1930-1957. Japanese longlining began in the EBS around 1958 and expanded into the AI and GOA through the 1970s. Japanese fleet catches increased throughout the 1960s, and peak sablefish catch reached 36,776 mt in 1972. High fishing pressure in the early 1970s by Japanese and USSR vessels may have resulted in a population decline of sablefish in the mid-1970s. By 1988, US fishermen took the majority of the sablefish harvested in the GOA and BSAI. Sablefish was increasingly harvested as a derby-style fishery in the late 1980s and early 1990s until Individual Fishing Quotas were implemented for the hook and line fishery in 1995.




Fishery Management: BSAI and GOA sablefish are managed as one population in federal waters due to their highly migratory behavior during certain life history stages. There are four management areas in the GOA; Western, Central West Yakutat and East Yakutat/Southeast Outside.

In 1985, Amendment 14 to the GOA FMP allocated sablefish TAC by gear type; 80% to fixed gear (including pots) and 20% to trawl in the Western and Central GOA, 95% to fixed gear and 5% to trawl gear in the Eastern GOA. Amendment 20 to the GOA FMP established IFQ management for the GOA sablefish fishery, which began in 1995.

Stock Assessment: The sablefish assessment is based on a statistical sex-specific age-structured model. This model incorporates fishery data and fishery independent data from domestic and Japan-US cooperative longline surveys and the NMFS GOA trawl survey. Sablefish fall under

Tier 3b of the ABC/OFL control rule. The 2011 age 4+ biomass estimated at 149,000 mt for the GOA. Catch specifications for 2011 in the GOA are as follows; OFL=13,340, ABC=11,290 mt, TAC=11,290 mt. Separate ABCs and TACs are established for each GOA subregion Western, Central, West Yakutat, and Southeast Outside.

Fishery: The sablefish IFQ fishery season opening date is concurrent with the halibut fishery for the purposes of reducing bycatch and regulatory discards between the two fisheries. In the GOA, the directed fishery for sablefish is prosecuted with longline gear (pot gear is prohibited for directed sablefish fishing in the GOA). Sablefish are also taken by trawl gear in directed fisheries for rockfish and deepwater flatfish. Primary incidental catch



AFSC, NOAA Fisheries

Total catches, pre-season catch specifications and exploitable biomass of Sablefish in the GOA, 1976-2011 (in mt).

Year	Catch ¹	TAC ²	ABC	OFL	Biomass ³
1976	27,733	-	-	-	-
1977	17,140	-	-	-	-
1978	8,866	-	-	-	-
1979	10,350	13,000	13,000	-	-
1980	8,543	13,000	13,000	-	-
1981	9,917	14,350	14,350	-	-
1982	8,556	12,300	12,300	-	-
1983	9,002	9,480	9,480	-	-
1984	10,230	8,980	8,980	-	-
1985	12,479	8,980	8,980	-	-
1986	21,614	15,000	18,800	-	-
1987	26,325	20,000	25,000	-	383,000
1988	29,903	28,000	35,000	-	520,000
1989	29,842	26,000	30,900	-	426,000
1990	25,701	26,000	26,200	-	312,000
1991	19,580	22,500	22,500	-	194,000
1992	20,451	20,800	20,800	28,200	179,000
1993	22,671	20,900	20,900	27,750	190,400
1994	21,338	25,500	25,500	31,700	218,000
1995	18,631	21,500	21,500	25,730	194,900
1996	15,826	17,080	17,080	22,800	169,500
1997	14,129	14,520	14,520	39,950	199,920
1998	12,758	14,120	14,120	23,450	166,000
1999	13,918	12,700	12,700	19,720	150,000
2000	13,779	13,330	13,330	16,660	169,000
2001	12,127	12,840	12,840	15,720	188,000
2002	12,246	12,820	12,820	19,350	188,000
2003	14,345	14,890	14,890	20,020	182,000
2004	15,630	16,550	16,550	22,160	179,000
2005	13,997	15,940	15,940	19,280	185,000
2006	13,367	14,840	14,840	17,880	152,000
2007	12,265	14,310	14,310	16,906	158,000
2008	12,326	12,730	12,730	15,040	167,000
2009	10,910	11,160	11,160	13,190	149,000
2010	9,998	10,370	10,370	12,270	140,000
2011	-	11,290	11,290	13,340	149,000

¹Catch data through November 2010.

²TAC, ABC and OFL from annual Federal Register.

³Biomass from SAFE report projections for following year.

species in the directed sablefish fishery include shortaker, roughey and thornyhead rockfish.

Economics: In 2009, the ex-vessel value of sablefish catch from the GOA was \$76.5 million. Exvessel prices for GOA sablefish in 2009 averaged \$3.42/lb for fish caught on longline gear and \$2.78/lb for fish taken with trawl gear. For both gear types, the primary product is frozen, head and gutted fish.

Current Issues: Sperm whale and killer whale depredation is problematic for sablefish fisheries in the GOA and BSAI. Depredation occurs when whales remove sablefish from longline gear, damage the fish and/or fishing gear. Killer whale depredation predominates in the BSAI and sperm whale depredation is more common the GOA. Depredation can lead to economic losses in the form of reduced catch, extended travel distances, extended wait times and damaged gear. Depredation may also reduce the accuracy of sablefish stock assessment models.

Additionally, depredating whales may be at greater risk of mortality or injury through vessel strikes or risk of entanglement in gear.



PVOA



AFSC, NOAA Fisheries

Shallow-water Flatfish

Biology: The Shallow-water flatfish complex is comprised of 8 flatfish species. Northern rock sole, southern rock sole, butter sole and yellowfin sole account for the majority of the current biomass of shallow-water flatfish. All flatfish are demersal but have varying depth ranges. Shallow-water flatfish predate on euphausiids, bivalves, polychaetes, amphipods, mollusks and fish. They are prey for Pacific cod, Pacific halibut and skates.

Northern rock sole	<i>Lepidopsetta polyxystra</i>
Southern rock sole	<i>Lepidopsetta bilineata</i>
Butter sole	<i>Pleuronectes isolepis</i>
Yellowfin sole	<i>Pleuronectes asper</i>
Starry flounder	<i>Platichthys stellatus</i>
English sole	<i>Pleuronectes vetulus</i>
Alaska plaice	<i>Pleuronectes quadrituberculatus</i>
Sand sole	<i>Psettichthys melanostictus</i>

Yellowfin sole distribution extends from Sea of Japan,

through the Chuckchi Sea and south to British Columbia. Adult yellowfin sole and rock sole occupy separate winter spawning and summertime feeding distributions on the continental shelf margins. Yellowfin sole are the second most abundant species (after pollock) in Cook Inlet and are also found in Prince William Sound. Yellowfin sole spawning period is protracted and likely extends from May to August, occurring primarily in shallow water. Females are relatively fecund, ranging from 1.3-3.3 million eggs depending on size. Yellowfin sole begin to recruit to the fishery at age 6 and are fully selected by age 13. The estimated age of 50% maturity is 10.5 years for females. Natural mortality is estimated at $M=0.12-0.16$, and longevity extends to 31 years.



For More Information:

B. Turnock W.
Stockhausen. T.
Wilderbuer, and M.
Wilkins. 2010. Assessment of Shallow Water Flatfish in the Gulf of Alaska.
<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>


The rock sole stock in the GOA consists of both northern and southern rock sole. The two species are similar in appearance but have different life history characteristics. Northern rock sole stock spawns beginning in midwinter and peaking during the spring, and the southern rock sole stock spawns during the summer. The estimated age of 50% maturity is 9 years for southern rock sole and 7 years for northern rock sole. Natural mortality is estimated to be $M=0.18-0.20$, and longevity extends to 21 years. Rock sole are most abundant in the Kodiak and Shumagin areas. Adults occupy separate winter spawning and summertime feeding distributions on the continental shelf margins.

Catch History: The flatfish fishery was predominantly a foreign fishery targeting non-flatfish species until 1981. With the cessation of foreign fishing in 1986, joint venture fishing began to account for the majority of flatfish catch, and the fishery was fully domestic by 1988. Shallow-water flatfish catch was 5,455 mt in 1978. Catch declined to a low of 957 mt in 1986 then increased to 9,715 mt in 1993. Shallow-water flatfish catch is often constrained by Pacific halibut bycatch limits.

Fishery Management: The Council divided the “Flatfish” complex into 3 categories (Deep-water flatfish, Shallow-water flatfish, and arrowtooth flounder) in 1990 due to significant differences in halibut bycatch rates, biomass and commercial value in directed fisheries for shallow and Deep-water flatfish. Flathead sole was separated out from the Deep-water flatfish complex in 1991 due to its distributional overlap between both shallow and deep-water groups. In 1993, rex sole was separated from the Deep-water flatfish complex due to concerns regarding POP bycatch.



All flatfish species under the GOA groundfish FMP are regulated through permits, limited entry, catch quotas (TACs), seasons, in-season adjustments, gear restrictions, closed waters, bycatch limits and rates, allocations, regulatory areas, record keeping, reporting requirements and observer monitoring. GOA flatfish species or complexes are managed



with area-specific ABC and TAC apportionments to avoid the potential for localized depletions.

Stock Assessment: The Shallow-water flatfish complex assessment is based on survey biomass estimates. The assessment incorporates fishery data and fishery independent data from annual trawl surveys. Northern rock sole and southern rock sole fall under a Tier 4 of ABC/OFL control rule, and catch limits for the remaining flatfish in the complex are set by a Tier 5 control rule due to limited maturity data. The 2011 projected biomass is 398,961 mt.

Catch specifications for 2011 are as follows; OFL=67,768 mt, ABC=56,242 mt, TAC= 20,062 mt.

Yellowfin sole biomass showed a declining trend from 54,738 t in 2003 to 33,414 t in 2009, and butter sole abundance declined by about 50% from 2007-2009. Northern rock sole, starry flounder and Alaska plaice have been increasing, along with southern rock sole and English sole. Sand sole survey biomass has been variable over time.

Fishery: Since 1988 the majority of Shallow-water flatfish harvest has occurred on the continental shelf and slope east of Kodiak Island in the Central regulatory area. Shallow-water flatfish are generally harvested with trawl gear. Rock sole is the predominant target species in the complex.

Economics: The bottom trawl fishery in the GOA primarily targets rock sole, rex sole and Dover sole. Primary products include whole fish, H&G and fillets. Ex-vessel value of all Flatfish caught in the GOA in 2008 was \$9.2 million. Production in 2008 was 139,150 mt for all flatfish products for a total gross value of \$202.9 million. A total of 33 catcher vessels and 6 catcher processors prosecuted the GOA flatfish fishery.

Total catches, pre-season catch specifications and exploitable biomass of Shallow Water Flatfish* in the GOA, 1991-2011 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass
1991	5,298	12,000	74,000	-	333,900
1992	8,783	11,740	50,480	70,900	257,338
1993	9,715	16,240	50,480	70,860	261,724
1994	9,343	18,630	34,420	44,670	261,720
1995	5,430	18,630	52,270	60,262	355,590
1996	9,350	18,630	52,270	60,262	355,590
1997	7,775	18,630	43,150	59,540	314,960
1998	3,565	18,630	43,150	59,540	315,590
1999	2,577	18,770	43,150	59,540	314,960
2000	6,928	19,400	37,860	45,330	299,100
2001	6,162	19,400	37,860	45,330	299,100
2002	6,195	20,420	49,550	61,810	349,992
2003	4,465	21,620	49,340	61,810	349,990
2004	3,094	20,740	52,070	63,840	375,950
2005	4,769	20,740	52,070	63,840	375,950
2006	7,641	19,972	51,450	62,418	365,766
2007	8,793	19,972	51,450	62,418	103,300
2008	9,708	22,256	60,989	74,364	436,590
2009	8,483	22,256	60,989	74,364	436,590
2010	5,410	20,062	56,242	67,768	398,961
2011	-	20,062	56,242	67,768	398,961

*Separated from Flounders category 1990.

¹Catch data through November 2010.

²Biomass from annual SAFE report projections.



AFSC, NOAA Fisheries

Deepwater Flatfish



For More Information:

W. Stockhausen 2010.
Assessment of Deepwater Flatfish stock in the Gulf of Alaska.

<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: The GOA Deep-water flatfish complex is comprised of 3 flatfish species; Greenland turbot *Reinhardtius hippoglossoides*, Dover sole *Microstomus pacificus*, and deep-sea sole *Embassichthys bathybius*. GOA Dover sole constitutes the majority of the survey biomass and deep-water flatfish catch (generally over 98%). Dover sole are generally found in water deeper than 300 m in the winter but occur at the highest biomass in the 100-200 m depth range during the summer. Dover sole are especially adapted to feeding on small-detrital consuming invertebrates such as polychaetes, amphipods, mollusks, and brittlestars. Dover sole are batch spawners, releasing around 83,000 advanced oocytes in about 9 batches. The peak spawning period occurs from January through May off the Oregon coast. Female Dover sole reach 50% maturity at about 34 cm (6-9 years old). Dover sole recruit to the fishery at 7-10 years, and longevity extends to 55 years. Greenland turbot has a circumpolar distribution in the Atlantic and Pacific. Greenland turbot are typically found from 200-1600 m. Greenland turbot predate on euphausiids, polychaetes and small fish (e.g. pollock) as they mature. Greenland turbot size at 50% maturity is around 60 cm (age 5-10). Greenland turbot begin to recruit to longline fisheries at around 60 cm and are fully recruited at 90 cm. Natural mortality is estimated at $M=0.18$. Biological data is limited for deep-sea sole.

Catch History: Deep-water flatfish catches peaked in 1992 at 11,379 mt, and then declined in 1993, remaining fairly stable from 1993-1999 (average 2,800 mt). After 1999, catches declined, averaging 602 mt annually from 2000-2009.

Stock Assessment: The Deep-water flatfish complex assessment uses a split-sex, age-structured model for Dover sole and mean historical catch data from 1978-1995 for

Greenland turbot and deep-sea sole. Dover sole catch limits are set by a Tier 3a control rule, and Greenland turbot and deep-sea sole fall under Tier 6 due to highly variable survey biomass estimates. The 2011 projected biomass (for Dover sole only) is 89,691 mt. Catch specifications for 2011 are as follows; OFL=7,823 mt, ABC= 6,305 mt, TAC= 6,305 mt.


Total catches, pre-season catch specifications, and exploitable biomass of Deep Water Flatfish* in the GOA, 1990-2011 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass ²
1990	2,380	22,000	108,400	-	-
1991	10,189	15,000	50,500	-	201,500
1992	11,379	19,740	39,280	51,500	169,132
1993	3,823	19,740	45,530	59,650	227,656
1994	3,129	11,080	16,510	19,280	132,030
1995	2,213	11,080	14,590	17,040	116,710
1996	2,193	11,080	14,590	17,040	116,570
1997	3,664	7,170	7,170	9,440	101,430
1998	2,286	7,170	7,170	9,440	101,430
1999	2,285	6,050	6,050	8,070	78,300
2000	985	5,300	5,300	6,980	74,370
2001	804	5,300	5,300	6,980	74,460
2002	559	4,880	4,880	6,430	68,623
2003	946	4,880	4,880	6,430	68,260
2004	680	6,070	6,070	8,010	99,620
2005	412	6,820	6,820	8,490	102,395
2006	405	8,665	8,665	11,008	132,297
2007	287	8,707	8,707	10,431	103,300
2008	563	8,903	8,903	11,343	132,625
2009	466	9,168	9,168	11,578	133,025
2010	502	6,190	6,190	7,680	89,682
2011	-	6,305	6,305	7,823	89,691

*Separated from Flounders category 1990.

¹Catch data through November 2010.

²Biomass from annual SAFE report projections.



Abundance estimates for Greenland turbot and deep-sea sole are highly uncertain. For Dover sole, survey biomass increased throughout the late 1980s, followed by declining estimates through the 1990s. Survey biomass increased again to 99,000 t in 2003.

Fishery: Deep-water flatfish are harvested with trawl gear. Dover sole is the predominant target species in the complex.



AFSC, NOAA Fisheries


Biology: Rex sole *Glyptocephalus zachirus* are distributed from Baja California to the BS, with concentrations in the GOA. Rex sole are closely associated with soft bottom benthic communities and are generally found at depths greater than 300 m. Adult rex sole overwinter near the shelf margins and migrate onto the mid and outer continental shelf each year in April/May. Rex sole exhibit latitudinal changes in growth rates and size at sexual maturity. Size at sexual maturity was greater for rex sole in the GOA than Oregon. Rex sole feed on polychaetes, euphausiids, amphipods and shrimp and are prey for skates, Pacific cod and arrowtooth flounder.



For More Information:

W. Stockhausen. 2010. Assessment of Rex Sole stock in the Gulf of Alaska. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Recruitment to the fishery begins around age 8. Age at 50% maturity for females was estimated at 5.6 years (35.2 cm) in Alaska. Maturity studies from Oregon show males are 50% mature at 16 cm and females at 24 cm. Natural mortality is estimated $M=0.17$, and longevity extends to 27 years. Rex sole are batch spawners with a protracted spawning period in the GOA (peak spawning period occurs April/May).



Total catches, pre-season catch specifications, and exploitable biomass of Rex Sole* in the GOA, 1994-2011 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass ²
1994	3,673	10,140	11,950	13,960	95,630
1995	4,021	9,690	11,210	13,091	89,660
1996	5,874	9,690	11,210	13,091	89,660
1997	3,294	9,150	9,150	11,920	72,330
1998	2,669	9,150	9,150	11,920	72,330
1999	3,060	9,150	9,150	11,920	72,330
2000	3,591	9,440	9,440	12,300	74,600
2001	2,940	9,440	9,440	12,300	81,020
2002	2,941	9,470	9,470	12,320	71,326
2003	3,485	9,470	9,470	12,320	71,330
2004	1,464	12,650	12,650	16,480	99,950
2005	2,176	12,650	12,650	16,480	99,950
2006	3,294	9,200	9,200	12,000	83,600
2007	2,852	9,100	9,100	12,000	82,403
2008	2,703	9,132	9,132	11,933	82,801
2009	4,753	8,996	8,996	11,756	81,572
2010	3,387	9,729	9,729	12,714	88,221
2011	-	9,565	9,565	12,499	86,729

*Separated from Deep Water Flatfish category 1994

¹ Catch through November.

² Biomass data corresponds to the annual SAFE report projections issued the preceding year.

Catch History: Prior to 1981, rex sole was caught incidentally in foreign fisheries targeting higher value species. Catches of rex sole have remained fairly stable since 1994, ranging from 1,464 mt in 2004 to a peak of 5,874 mt in 1996.

Stock Assessment: Rex sole limits are set by a Tier 5 control rule. The 2011 projected biomass is 86,729 mt. and the natural mortality rate (M) = 0.17. Catch specifications for 2011 are as follows; OFL=12,499 mt, ABC= 9,565 mt, TAC=9,565 mt. The ABC and TAC specifications are further subdivided among GOA subareas.

Fishery: GOA rex sole are caught using trawl gear in a directed fishery and fisheries targeting other bottom-dwelling species such as POP, Pacific cod and bottom pollock. Fishing seasons are dictated by seasonal halibut PSC apportionments, with approximately 7 months of fishing occurring between January and November. Catches of rex sole occur primarily in the Western and Central management areas in the GOA.



Arrowtooth Flounder

AFSC, NOAA Fisheries



For More Information:

B. Turnock. 2010.
 Assessment of Arrowtooth
 Flounder stock in the Gulf
 of Alaska.
<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: Arrowtooth flounder (*Atheresthes stomias*) are distributed from the Kamchatka Peninsula, through the BSAI down to central California. Arrowtooth flounder are most abundant at depths from 100-500 m. Adults migrate seasonally from shelf margins in the winter to the inner and middle shelf in April/May with the onset of warmer waters temperatures. Smaller GOA arrowtooth flounder predate on euphausiids, capelin and herring while fish over 40 cm rely primarily on pollock. Predators of arrowtooth flounder include Pacific cod, pollock and skates

Arrowtooth flounder recruitment to the fishery begins at about 3 years, and females are fully recruited by age 10. The estimated length at 50% maturity is 28 cm for males (4 years) and 37 cm for females (5 years) based on samples collected from Washington, and longevity extends to 21 years. Female natural mortality is estimated at $M=0.2$. Male natural mortality has a range estimate ($M=0.27-0.36$). Adult males range in size from 30-50 cm, and females range in size from 30-70 cm. The spawning period for arrowtooth flounder occurs from December to February at depths of 100-360 m. Spawning in the GOA occurs from Kodiak to Yakutat Bay.

Catch History: Prior to 1981, arrowtooth flounder was caught incidentally in foreign fisheries targeting higher value species. From 1991-2000, arrowtooth flounder catches ranged from 10,034 mt-22,583 mt. Catches of arrowtooth flounder were on average greater from 2000-November 2009, peaking in 2008 at 29,293 mt.

Total catches, pre-season catch specifications, and exploitable biomass of Arrowtooth Flounder* in the GOA, 1990-2011 (in mt).


Year	Catch ¹	TAC	ABC	OFL	Biomass ²
1990	7,705	32,000	194,600	-	-
1991	10,035	20,000	340,100	-	2,000,800
1992	15,970	25,000	303,800	427,000	1,787,583
1993	15,560	30,000	321,290	451,690	1,889,922
1994	23,560	30,000	236,240	275,930	1,889,920
1995	18,430	35,000	198,130	231,416	1,585,040
1996	22,183	35,000	198,130	231,416	1,640,000
1997	16,319	35,000	197,840	280,800	1,971,170
1998	12,974	35,000	208,340	295,570	2,062,740
1999	16,209	35,000	217,110	308,880	2,126,714
2000	24,252	35,000	145,360	173,910	1,571,670
2001	19,964	38,000	148,150	173,550	1,586,830
2002	21,230	38,000	146,260	171,060	1,760,000
2003	23,320	38,000	155,140	181,390	1,302,000
2004	15,304	38,000	194,930	228,130	2,453,390
2005	19,770	38,000	216,900	253,900	2,453,390
2006	27,653	38,000	177,844	207,678	2,140,170
2007	25,364	43,000	184,008	214,828	2,146,360
2008	29,293	43,000	226,470	266,914	2,244,870
2009	24,937	43,000	221,512	261,022	1,295,050
2010	23,015	43,000	215,882	254,271	2,139,000
2011	-	43,000	213,150	251,068	2,139,000

*Separated from Flounders category 1990.
¹Catch data through November 2010.
²Biomass from SAFE report projections.

Stock Assessment: The arrowtooth flounder assessment uses an automatic differentiation software developed as a set of libraries under C++ (AD Model Builder). This model incorporates fishery data and fishery independent data from NMFS and IPHC trawl surveys. Arrowtooth flounder catch limits are set by a Tier 3a control rule. The 2011 projected biomass= is 2,121,440 mt. Catch specifications for 2011 are as follows; OFL=251,068 mt, ABC= 213,150 mt, TAC= 43,000 mt.

Arrowtooth flounder biomass has increased steadily since the early 1990s. Estimated biomass averaged 1.7 million mt annually from 2000-2004 and 2 million mt from 2004-2009.

Fishery: There is currently no directed fishery for arrowtooth flounder in the GOA. However, arrowtooth flounder are an important byproduct of more valuable target trawl and longline fisheries, such as Pacific cod and pollock.



Flathead Sole

AFSC, NOAA Fisheries



For More Information:

W. Stockhausen. 2010. Assessment of Flathead Sole stock in the Gulf of Alaska.

<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: Flathead sole *Hippoglossoides elassodon* are distributed in the Kuril Islands, BS, GOA and south to California. Adult flathead sole exhibit a benthic lifestyle and overwinter near the shelf margins before migrating to the mid and outer continental shelf in April or May each year for feeding. They occur primarily on mixed mud and sand bottoms in depths less than 300 m. Pandalid shrimp and brittle stars are the most important prey for adult flathead sole in the GOA, while euphausiids and mysids constitute the most important prey items for juvenile flathead sole. Pacific cod and Pacific halibut are the major predators on adults, while arrowtooth flounder, sculpins, walleye pollock and Pacific cod are the major predators on juveniles.

Flathead sole recruitment to the fishery begins at age 4, and longevity extends to 32 years. Estimated length at 50% maturity is 33 cm (8.7 years). Natural mortality is estimated at $M=0.20$. Flathead sole spawn in March and April, primarily in deeper waters near the margins of the continental shelf. Females release from 70,000-600,000 eggs depending on size.

Catch History: From a high of approximately 2000 t in 1980, annual flathead sole catches declined steadily to a low of around 150 mt in 1986. After 1986, catches increased and reached a peak catch of 3,658 mt in 2009.

Stock Assessment: The flathead sole assessment uses a split-sex, age-based model with age length formulations for fishery and survey selectivity. This model incorporates fishery data and fishery independent data from triennial (1984-1999) and biennial (2001-2009) surveys.


Total catches, pre-season catch specifications, and exploitable biomass of Flathead Sole* in the GOA, 1991-2011 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass ²
1991	1,717	10,000	50,300	-	251,800
1992	2,034	10,000	48,280	63,100	240,615
1993	2,366	10,000	49,450	64,780	247,250
1994	2,580	10,000	35,850	39,310	199,000
1995	2,181	10,000	28,790	31,557	198,470
1996	3,107	9,740	28,790	31,557	198,470
1997	2,446	9,040	26,110	34,010	206,340
1998	1,742	9,040	26,110	34,010	206,340
1999	900	9,040	26,110	34,010	206,340
2000	1,547	9,060	26,270	34,210	207,520
2001	1,911	9,060	26,270	34,210	207,520
2002	2,145	9,280	22,690	29,530	170,915
2003	2,425	11,150	41,390	51,560	132,260
2004	2,390	10,880	51,270	64,750	292,670
2005	2,530	10,390	45,100	56,500	292,670
2006	3,134	9,077	37,820	47,003	291,441
2007	3,163	9,148	39,110	48,658	297,353
2008	3,419	11,054	44,735	55,787	103,300
2009	3,658	11,181	46,464	57,911	323,937
2010	3,458	10,411	47,422	59,295	328,862
2011	-	10,587	49,133	61,412	325,367

¹Catch data from SAFE.

²Biomass from annual SAFE report projections.

Flathead sole catch limits are set by a Tier 3a control rule. The 2011 projected biomass is 325,367 mt. Catch specifications for 2011 are as follows; OFL=61,412 mt, ABC= 49,133 mt, TAC= 10,587 mt. Estimated flathead sole biomass steadily increased from 207,520 mt in 2000 to 328,862 mt in 2010.



Fishery: GOA flathead sole are caught using trawl gear in a directed fishery and fisheries targeting other bottom-dwelling species such as POP, Pacific cod and bottom pollock. Fishing seasons are dictated by seasonal halibut PSC apportionments. The majority of flathead sole in the GOA is taken in the Shelikof Strait and on the Albatross Bank near Kodiak Island and Unimak Island. About 90% of the catch is retained.



Pacific Ocean Perch

AFSC, NOAA Fisheries



For More Information:

D. Hanselman, K. Shotwell, J. Heifetz, and J. Ianelli. 2010. Assessment of Pacific Ocean Perch stock in the Gulf of Alaska. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: Pacific Ocean Perch (POP) *Sebastes alutus* distribution extends from Japan around the Pacific Rim, the BS and south to California. POP are most abundant in AI, GOA and British Columbia and are found primarily offshore along the continental slope at depths from 150-420 m. POP are generally considered a demersal species and are found over cobble substrate. Seasonal changes in depth distribution occur, and adults migrate farther offshore to deeper waters during winter. During late spring and summer, POP migrate to shallower waters inshore for summer feeding. Adults perform diel migrations off the sea floor to feed. POP populations occur in patchy aggregations, and POP are generally planktivorous. Smaller POP feed on calanoid copepods, whereas larger POP rely on euphausiids, shrimp and squids. POP are prey for Pacific halibut, sablefish, Pacific cod and arrowtooth flounder.



SeaAlliance/AGDB

POP is a slow-growing, longlived species. Recruitment to trawl fisheries begins at age 5, and full recruitment to the fishery occurs around age 8. Females reach 50% maturity at 10.5 years in the GOA, and longevity extends to 80 plus years (oldest recorded 84 years in the GOA). Natural mortality is estimated to be $M=0.06$. Females are viviparous, retaining fertilized eggs within the ovary until larval extrusion. Mating takes place in late fall, and larval extrusion occurs in early spring. Females release from 10,000-300,000 eggs each year, depending on size.

Total catches, pre-season catch specifications, and exploitable biomass of Pacific Ocean Perch* in the GOA, 1990-2011 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass ²
1991	6,632	5,800	5,800	-	-
1992	6,158	5,200	5,730	5,730	229,100
1993	2,119	2,560	3,378	3,378	156,300
1994	1,853	2,550	3,030	3,940	101,800
1995	5,742	5,630	6,530	8,232	142,465
1996	8,459	6,960	8,060	10,165	163,220
1997	9,531	9,190	12,990	19,760	301,084
1998	9,266	10,780	12,820	18,090	242,300
1999	10,802	12,590	13,120	18,490	228,190
2000	10,157	13,020	13,020	15,390	200,310
2001	10,860	13,510	13,510	15,390	211,160
2002	11,729	13,190	13,190	15,670	293,240
2003	10,911	13,660	13,660	16,240	298,820
2004	11,528	13,340	13,340	15,840	266,960
2005	11,440	13,575	13,575	16,266	286,367
2006	13,590	14,261	14,261	16,927	312,968
2007	13,046	14,635	14,636	17,158	315,507
2008	12,400	14,999	14,999	17,807	317,511
2009	12,985	15,111	15,111	17,940	318,336
2010	15,520	17,584	17,584	20,243	334,797
2011	-	16,997	16,997	19,566	330,480

* Separated from Slope Rockfish in 1991.

¹Catch data from SAFE.

²Biomass from annual SAFE report projections.

Catch History: POP was harvested in the GOA by the USSR and Japan beginning in the early 1960s. The fishery developed rapidly, and catches peaked in 1965 at 350,000 mt. High fishing effort by the foreign fleet caused a major decline in POP abundance/catches through the late 1960s. Catches continued to decline, and in 1985 foreign trawling in the GOA was prohibited.

The domestic fishery for POP in the GOA began in the early 1980s and expanded each year until 1991. POP catches remained relatively low through the 1990s, averaging 7,072 mt annually from 1991-2000. Catches have increased moderately since 2000, averaging 12,027 mt annually from 2001-November 2009.

Fishery Management: In 1991, POP and the Shortaker/Rougheye complex were separated from the "Slope Rockfish" complex to prevent overfishing. A reduction in TACs after 1991 to promote POP stock rebuilding was also implemented. In 2004, Shortaker and Rougheye rockfish were separated into their own management units due to disproportionately high harvests of

shorttraker rockfish. GOA rockfish stocks and complexes are managed with area-specific

ABC and TAC apportionments to avoid the potential for localized depletions. Amendment 41, effective in 2000, prohibited trawling in the Eastern area east of 140°W longitude, an area previously fished for POP.


The Central GOA Rockfish Pilot Program, effective for 2007 through 2011, rationalized the rockfish and related trawl fisheries. The program provides cooperatives with exclusive catch shares (95% of the CGOA TAC) for target species of POP, Northern rockfish, and Pelagic Shelf rockfish, as well as a allocated a portion of the TAC for suite of secondary species (sablefish, cod, and thornyhead, shortraker and roughey rockfish), and a halibut prohibited species catch limit allocation. Cooperatives receive allocations based on catch history of cooperative member vessels. Sideboard limits for the target rockfish species are established in the Western GOA. A slightly revised program was adopted by the Council in 2010 for implementation in 2012 to 2021.



Mark Fina, NPFC

Stock Assessment: The POP assessment uses an age-structured model using AD Model Builder software. POP catch limits are set under Tier 3a OFL and ABC control rules. This model incorporates fishery data and fishery independent data from biennial trawl surveys. The 2011 projected biomass is 330,480 mt. Catch specifications for 2011 are as follows; OFL=19,566 mt, ABC= 16,997 mt, TAC= 16,997 mt.

Estimated biomass of POP was relatively low during the early 1990s, averaging 158,577 mt from 1991-1995. Since 2000, POP estimated biomass has steadily increased from 211,160 mt in 2000 to 334,797 mt in 2009, averaging 295,567 mt annually from 2000-November 2009.



Fishery: POP are caught primarily in directed bottom trawl fisheries. The percentage of POP in the GOA taken in pelagic trawls increased from 2% in 1990 to 31% in 2008. The majority of POP are caught in the Central regulatory area, and TACs allocated for each area are generally met (except Southeastern area due to prohibited trawling).

Economics: In 2008, production was 20,570 mt for all Alaska rockfish products for a total gross value of \$41.9 million. Ex-vessel value of rockfish catch in the GOA was \$9.5 million.



Northern Rockfish

AFSC, NOAA Fisheries



For More Information:

J. Heifetz, D. Hanselman, K. Shotwell, and J. Ianelli. 2010. Assessment of Northern Rockfish stock in the Gulf of Alaska. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: Northern rockfish *Sebastes polypsinus* distribution extends from the Kamchatka Peninsula, through the BSAI, GOA and British Columbia. The species is most abundant in the central GOA to the western end of the AI. Adults concentrate at discrete sites along the outer continental shelf from 75-150 m. Northern rockfish are demersal and are generally found in aggregations with patchy distributions. Northern rockfish are prey on calanoid copepods, euphausiids and chaetognaths. Based on stomach content data for POP, Pacific halibut and sablefish likely prey on northern rockfish.

Northern rockfish is a slow-growing, longlived species. Age at 50% maturity is 12.8 years in the GOA, and longevity extends to 50 years (oldest recorded 67 in the GOA). GOA northern rockfish grow faster and reach a larger maximum length than the AI northern rockfish. Natural mortality is estimated to be $M=0.06$. Females are viviparous, retaining their fertilized eggs within the ovary until larval extrusion.

Catch History: Northern rockfish were initially harvested by Soviet and Japanese trawlers in the early 1960s. Foreign fishing effort increased quickly in the 1960s, and catches of rockfish in the GOA peaked in 1965 at 350,000 mt. It is likely that GOA northern rockfish comprised some portion of the early foreign catch (exact northern rockfish catch unknown for this period). Northern rockfish was separated from the slope rockfish assemblage in 1993, and catches have remained fairly stable since 1994, ranging from a low of 2,947 mt in 1997 to a high of 5,968 in 1994 (average annual catch equals 4,262 mt from 1994-2009).

Stock Assessment: The northern rockfish assessment uses a separable, age-structured model using AD Model Builder software. This model incorporates fishery data and fishery independent data from biennial trawl surveys. Northern rockfish catch limits are set under Tier 3a of the ABC/OFL control rules. The 2011 projected biomass is 100,463 mt. Catch specifications for 2011 are as follows; OFL=5,784 mt, ABC= 4,854 mt, TAC= 4,854 mt.


Total catches, pre-season catch specifications, and exploitable biomass of Northern Rockfish* in the GOA, 1993-2011 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass ²
1993	4,846	5,760	5,760	10,360	76,800
1994	5,968	5,760	5,760	10,360	76,800
1995	5,634	5,270	5,270	9,926	87,845
1996	3,356	5,270	5,270	9,926	87,850
1997	2,947	5,000	5,000	9,420	83,890
1998	3,058	5,000	5,000	9,420	83,870
1999	5,412	4,990	4,990	9,420	83,870
2000	3,325	5,120	5,120	7,510	85,360
2001	3,150	4,880	4,880	5,780	93,850
2002	3,337	4,980	4,980	5,910	94,350
2003	5,349	5,530	5,530	6,560	108,830
2004	4,806	4,870	4,870	5,790	95,150
2005	4,806	5,091	5,091	6,050	108,274
2006	4,956	5,091	5,091	7,673	136,311
2007	4,187	4,938	4,938	5,890	94,271
2008	4,052	4,549	4,549	5,430	93,391
2009	3,925	4,362	4,362	5,204	90,557
2010	3,871	5,098	5,098	6,070	103,300
2011	-	4,854	4,854	5,784	100,463

Separated from Other Slope Rockfish category 1993.

¹Catch data from the SAFE.

²Biomass from annual SAFE report projections.



Fishery: Northern rockfish are fully allocated as a target species in the CGOA trawl rockfish program, with 95-98% of the CGOA TAC and sideboarded at 74.3% of the WGOA TAC. Important fishing grounds include Portlock Bank, Albatross Bank, Shumagin Bank and Davidson Bank.



Shortraker Rockfish

AFSC, NOAA Fisheries




For More Information:

D. Clausen. 2010. Assessment of Shortraker Rockfish and "Other Slope Rockfish" stocks in the Gulf of Alaska. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: Shortraker rockfish *Sebastes borealis* are distributed from Japan around the Pacific Rim to Southern California, including the BSAI and the GOA. In Alaska, adults are especially concentrated along the continental slope in the 300-500 m depth interval. Shortraker rockfish predate on shrimps, squids, and myctophids. Shortrakers attain the largest size of all *Sebastes*, with a maximum reported length of 120 cm. Shortraker rockfish is one of the most long-lived species in the northeast Pacific, and longevity may exceed 120 years. Natural mortality is estimated to be $M=0.03$. Information on early life history stages of shortraker rockfish is limited.

Catch History: From 1991 to 2004, the NPFMC managed shortraker rockfish in the GOA together with roughey rockfish as an assemblage. Combined catches for the two species ranged from 702 to 2,250 mt, averaging 1,617 mt annually. Shortraker was separated into a single species management unit in 2005, and catches of shortraker rockfish averaged 584 mt annually from 2005-2009.

Stock Assessment: Due to limited biological data, the shortraker rockfish assessment uses a biomass-based approach to calculating ABCs, incorporating fishery independent data from trawl surveys. Shortraker rockfish catch limits are set under Tier 5 ABC/OFL control rules. The 2011 projected biomass is 40,626 mt. Catch specifications for 2011 are as follows; OFL=1,219 mt, ABC=914 mt, TAC= 914 mt.



Fishery: Shortraker rockfish in the GOA are taken in both longline and trawl fisheries; each gear comprises about 50% of the annual catch. Shortrakers in the CGOA are allocated as a secondary species in the CGOA rockfish program. A total of 40% of the CGOA Shortraker TAC is allocated to the catcher processor sector.

Total catches, pre-season catch specifications, and exploitable biomass of Shortraker Rockfish* in the GOA, 2005-2011 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass ²
2005	498	753	753	982	32,723
2006	664	843	843	1,124	37,461
2007	608	843	843	1,124	37,461
2008	598	898	898	1,197	39,905
2009	550	898	898	1,197	39,905
2010	457	914	914	1,219	40,626
2011	-	914	914	1,219	40,626

Separated from Slope Rockfish in 1991 and Shortraker/Roughey in 2004.

¹Catch data from 2009 SAFE

²Biomass from annual SAFE report projections.



Julianne Curry, PVOA



Sharpchin rockfish
AFSC, NOAA Fisheries

Other Slope Rockfish



For More Information:

D. Clausen. 2010. Assessment of Shortracker Rockfish and “Other Slope Rockfish” stocks in the Gulf of Alaska. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: The Other Slope Rockfish (OSR) complex consists of 15 rockfish species, although sharpchin, harlequin, silvergray, redstripe, and redbanded rockfish comprise the majority of the biomass in the GOA. The center of abundance for most of these species is farther south off British Columbia or the US west coast. However, harlequin rockfish are most common in Alaskan waters, and silvergray rockfish appear to be most abundant in southeast Alaska and British Columbia. Within the GOA, OSR are most abundant in the eastern GOA and become increasingly scarce in areas farther west.

Life history data is limited for most OSR species. For sharpchin rockfish, size at 50% maturity is 26.5 cm (10 years). Natural mortality is estimated to be $M=0.05$ for sharpchin and silvergray rockfish, $M=0.10$ for redstripe rockfish, and $M=0.06$ for harlequin and redbanded rockfish and all the minor species in the group.

Blackgill rockfish	<i>Sebastes melanostomus</i>
Bocaccio rockfish	<i>Sebastes paucispinis</i>
Chilipepper rockfish	<i>Sebastes goodei</i>
Darkblotched rockfish	<i>Sebastes crameri</i>
Greenstriped rockfish	<i>Sebastes elongatus</i>
Harlequin rockfish	<i>Sebastes variegatus</i>
Pygmy rockfish	<i>Sebastes wilsoni</i>
Redbanded rockfish	<i>Sebastes babcocki</i>
Redstripe rockfish	<i>Sebastes proriger</i>
Sharpchin rockfish	<i>Sebastes zacentrus</i>
Silvergray rockfish	<i>Sebastes brevispinis</i>
Splitnose rockfish	<i>Sebastes diploproa</i>
Stripetail rockfish	<i>Sebastes saxicola</i>
Vermilion rockfish	<i>Sebastes miniatus</i>
Yellowmouth rockfish	<i>Sebastes reedi</i>

Catch History: Catch data for OSR are only available for the years since 1991, when these 15 species became their own management group in the GOA. Since the mid-1990s, catches for OSR in the GOA have generally been less than 1,000 mt. In particular, the EGOA trawl closure that has been in effect since 1998 has limited the catch of OSR in the GOA.

Total catches, pre-season catch specifications, and exploitable biomass of Other Slope Rockfish* in the GOA, 1993-2010 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass ²
1993	2,810	5,383	8,300	9,850	134,400
1994	1,613	2,235	8,300	9,850	76,500
1995	1,397	2,235	7,110	8,395	112,812
1996	881	2,020	7,110	8,395	112,810
1997	1,217	2,170	5,260	7,560	103,710
1998	861	2,170	5,260	7,560	103,710
1999	788	5,270	5,270	7,560	103,710
2000	577	4,900	4,900	6,390	102,510
2001	559	1,010	4,900	6,390	102,510
2002	774	990	5,040	6,610	107,960
2003	1,078	990	5,050	6,610	107,960
2004	885	670	3,900	5,150	89,460
2005	715	670	3,900	5,150	103,300
2006	931	1,480	4,152	5,394	93,552
2007	690	1,482	4,154	5,394	93,552
2008	809	1,730	4,297	5,624	90,283
2009	881	1,730	4,297	5,624	90,283
2010	798	1,192	3,749	4,881	76,867
2011	-	1,195	3,752	4,881	76,867

*Separated from Slope Rockfish in 1991. Northern Rockfish split from Other rockfish category in 1993.
¹Catch data from SAFE.
²Biomass from annual SAFE report projections.

Stock Assessment: Other Slope Rockfish are managed under Tier 5 of the ABC/OFL control rules (sharpchin rockfish managed under Tier 4). The 2011 projected biomass is 76,867 mt. Catch specifications for 2011 are as follows; OFL=4,881 mt, ABC= 3,752 mt, TAC= 1,195 mt.



Vermilion rockfish
AFSC, NOAA Fisheries

Fishery: There is no directed fishery for OSR in the GOA. Other Slope Rockfish in the GOA are primarily taken in trawl fisheries targeting higher value species. Harlequin and sharpchin rockfish are the predominant OSR species caught. Prior to 1996, more than 90% of the slope rockfish trawl catch was taken by large at-sea factory trawlers. Since then, smaller shore-based trawlers have taken sizeable catches for delivery to processing plants in Kodiak.



Dusky rockfish
AFSC, NOAA Fisheries



For More Information:

C. Lunsford, S. K. Shotwell,
and D. Hanselman. 2010.
Assessment of Pelagic Shelf
Rockfish in the Gulf of
Alaska.
<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Pelagic Shelf Rockfish

Biology: The Pelagic Shelf Rockfish (PSR) complex consists of 3 rockfish species; dusky rockfish *Sebastes variabilis*, yellowtail rockfish *Sebastes flavidus*, and widow rockfish *Sebastes entomelas*. Yellowtail and widow rockfish are less common than dusky rockfish, and life history data is limited for these 2 species. Yellowtail and widow rockfish tend to be concentrated in nearshore areas and offshore banks on the continental shelf. Natural mortality is estimated $M=0.07$ for both yellowtail and widow rockfish.

Dusky rockfish are the most abundant species in the PSR assemblage in the GOA. Adult dusky rockfish are concentrated around offshore banks and near gullies on the outer continental shelf at depths of 100 to 200 m. It is likely that dusky rockfish benthic distribution is associated with hard, rocky bottoms and epibenthic habitats. Dusky rockfish prey on Pacific sandlance and euphausiids. Dusky rockfish age at 50% maturity is approximately 11.3 years. Mortality is estimated to be $M=0.07$, and longevity extends to 60 years. Dusky rockfish are ovoviparous with fertilization, embryonic development, and larval hatching occurring inside the mother. Parturition is believed to occur in the spring in the GOA.



Yellowtail rockfish
AFSC, NOAA Fisheries

Total catches, pre-season catch specifications, and exploitable biomass of Pelagic Shelf Rockfish in the GOA, 1988-2011 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass ²
1988	1,086	3,300	3,300	-	169,700
1989	1,739	3,300	6,600	-	164,300
1990	1,647	8,200	8,200	-	164,000
1991	2,342	4,800	4,800	-	96,300
1992	3,440	6,890	6,890	11,360	75,110
1993	3,193	6,740	6,740	11,300	74,900
1994	2,990	6,890	6,890	11,550	76,500
1995	2,891	5,190	5,190	8,704	57,644
1996	2,302	5,190	5,190	8,704	56,502
1997	2,629	5,140	5,140	8,400	54,220
1998	3,111	5,260	5,260	8,040	55,580
1999	4,826	4,880	4,880	8,190	54,220
2000	3,730	5,980	5,980	9,040	66,440
2001	3,008	5,980	5,980	9,040	66,440
2002	3,318	5,490	5,490	8,220	62,489
2003	2,975	5,490	5,490	8,220	62,500
2004	2,674	4,470	4,470	5,570	57,400
2005	2,235	4,553	4,553	5,680	103,300
2006	2,446	5,436	5,436	6,662	97,368
2007	3,318	5,542	5,542	6,458	99,829
2008	3,634	5,227	5,227	6,400	70,823
2009	3,057	4,781	4,781	5,803	66,603
2010	3,097	5,059	5,059	6,142	66,603
2011	-	4,754	4,754	5,770	66,498

*Separated from Other Rockfish category 1988.

¹Catch data through November 2010.

²Biomass from annual SAFE report projections.

Catch History: PSR catch in the GOA generally increased after the management groups were separated in 1988. From 1998-1995, over 95% of the catch of dusky rockfish was taken by large factory trawler processing fish at sea. In 1996, smaller shore-based trawlers also began taking a portion of the catch in the Central Gulf area for delivery to processing plants in Kodiak. These shore-based trawlers have accounted for 18-74% of the trawl catch in the Central area from 1996-2006. Catches have remained fairly stable since 1994 and peaked in 1999 at 4,826 mt.

Stock Assessment: Black and blue rockfish were removed from the GOA FMP in 1998, and dark rockfish in 2009. PSR are managed under Tier 3 (dusky) and Tier 5 (widow and yellowtail) ABC/OFL control rules. The 2011 projected biomass is 66,498 mt. Catch specifications for 2011 are as follows; OFL=5,770 mt, ABC=4,754 mt, TAC= 4,754 mt.

Fishery: In the CGOA, 95% of the PSR TAC is allocated to the CGOA Rockfish program. The trawl fishery for dusky rockfish begins in May and closes in November. Catches of dusky rockfish are concentrated at a number of offshore banks of the outer continental shelf, west of Yakutat and around Kodiak in areas such as Portlock Bank and Albatross Bank.



Rougheye and Blackspotted Rockfish

Blackspotted rockfish
AFSC, NOAA Fisheries



For More Information:


S. K. Shotwell, D. Hanselman, and D. Clausen. 2010. Assessment of Rougheye and Blackspotted Rockfish stock in the Gulf of Alaska. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: The rougheye and blackspotted (RE/BS) complex consists of 2 species; rougheye rockfish *Sebastes aleutianus* and a species recently identified by genetic research as blackspotted rockfish *Sebastes melanostictus*. These two species are often difficult to differentiate from each other at sea. RE/BS distribution extends from Japan, through the BSAI, GOA to southern California. Adults primarily inhabit a narrow band along the upper continental slope at depths from 300-500 m. Although the two species distributions overlap, blackspotted rockfish are predominant in the AI, while rougheye rockfish are more common in the GOA and southeastern BS.

Rougheye rockfish length at 50% maturity is 44 cm, and longevity may extend to 200 years. Natural mortality for RE/BS is estimated to be $M=0.04$. As with other rockfish, RE/BS are presumed to be viviparous. RE/BS rockfish prey on pandalid shrimps, euphausiids, lanternfish, and crabs. Predators of RE/BS include Pacific halibut, Pacific cod and sablefish.

Catch History: Gulf-wide catches of the rougheye rockfish and blackspotted rockfish ranged from 130-2,418 mt. from 1977-1990. RE/BS rockfish are generally caught with either bottom trawls or longline gear. RE/BS rockfish have been managed as a “bycatch” only species since the creation of the shortraker/rougheye rockfish management subgroup in the Gulf of Alaska in 1991. Catches of rougheye and shortraker rockfish from 1992-2004 ranged from 702 - 2,250 mt, averaged 1,617 mt annually. RE/BS rockfish were separated into their own management unit in 2004, and catches of RE/BS rockfish averaged 345 mt annually from 2005- 2009.

Stock Assessment: The RE/BS rockfish assessment uses a separable age-structured model, which incorporates fishery data and fishery independent data from biennial trawl and annual longline surveys. RE/BS rockfish limits are set by a Tier 3a control rule. The 2011 projected biomass is 45,907 mt. Catch specifications for 2011 are as follows; OFL=1,579 mt, ABC= 1,312 mt, TAC= 1,312 mt.



Total catches, pre-season catch specifications, and exploitable biomass of Rougheye and Blackspotted Rockfish* in the GOA, 2005-2011 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass ²
2005	294	1,007	1,007	1,531	40,281
2006	358	983	983	1,180	37,449
2007	417	988	988	1,148	39,506
2008	389	1,286	1,286	1,548	46,121
2009	280	1,284	1,284	1,545	46,385
2010	447	1,302	1,302	1,568	45,751
2011	-	1,312	1,312	1,579	45,907

*Separated from Slope Rockfish in 1991 and Shortraker/Rougheye in 2004.

¹Catch data from SAFE.

²Biomass from annual SAFE report projections.

Fishery: RE/BS rockfish in the GOA are primarily taken in rockfish bottom trawl fisheries and longline fisheries targeting sablefish and Pacific halibut.



Rougheye rockfish
AFSC, NOAA Fisheries



China rockfish
AFSC, NOAA Fisheries



Demersal Shelf Rockfish

Biology: The Demersal Shelf Rockfish (DSR) complex consists of 7 species. DSR are generally nearshore, bottom-dwelling species, located on the continental shelf and associated with rugged, rocky habitat. DSR species exhibit K-selected life history traits including slow growth and extreme longevity. DSR are viviparous, and parturition occurs from February through September with the majority of the species extruding larvae in spring.

Canary rockfish	<i>Sebastes pinniger</i>
China rockfish	<i>Sebastes nebulosus</i>
Copper rockfish	<i>Sebastes caurimus</i>
Quillback rockfish	<i>Sebastes maliger</i>
Rosethorn rockfish	<i>Sebastes helvomaculatus</i>
Tiger rockfish	<i>Sebastes nigrocinctus</i>
Yelloweye rockfish	<i>Sebastes ruberrimus</i>

For More Information:

K. Green, D. Carlile, M. Jaenicke, S. Meyer, and J. Stahl. 2010. Assessment of the Demersal Shelf Rockfish stock in the SEO District of the Gulf of Alaska.

<http://www.afsc.noaa.gov/refm/stocks/assessments.htm>

The primary species of the fishery is yelloweye rockfish. The oldest recorded yelloweye rockfish is 118 years, and natural mortality is estimated at $M=0.02$. Yelloweye reach a maximum length of about 91 cm with the length at 50% maturity at 45 cm (22 years). Yelloweye feed on shrimp, small crabs and a variety of fishes including small rockfish, herring and sand lance. Yelloweye are in turn prey for larger rockfish, lingcod, salmon and Pacific halibut.

Catch History: The directed fishery for DSR began in 1979 as a small, shore-based, hook and line in Southeast Alaska, which targeted the entire DSR complex. The directed DSR catch increased from 120 mt in 1982 to a peak of 778 mt in 1987.

Fishery Management: DSR are managed jointly by ADF&G and NMFS. The directed DSR season is closed during the halibut IFQ season to prevent over-harvest of DSR, and 33% of DSR quota is allocated pre-halibut season and 67% of DSR quota is allocated post-halibut

season. Directed fishery quotas are set by management area and are based on the remaining ABC after subtracting the estimated DSR incidental catch (landed and at sea discard) in other fisheries.

Exploitable biomass, pre-season catch specifications, and total catches (including discards) of Demersal Shelf Rockfish* in the GOA, 1992-2010 (in mt).

Year	Catch ¹	TAC	ABC	OFL	Biomass ²
1992	511	550	550	732	-
1993	558	800	800	1,600	48,366
1994	540	960	960	1,680	49,280
1995	219	580	580	1,044	26,093
1996	401	950	950	1,702	42,552
1997	406	950	950	1,450	42,552
1998	552	560	560	950	25,031
1999	297	560	560	950	25,031
2000	406	340	340	420	15,100
2001	301	330	330	410	14,695
2002	292	350	350	480	15,615
2003	229	390	390	540	17,510
2004	260	450	450	690	20,168
2003	187	410	410	640	18,508
2006	166	410	410	650	19,558
2007	250	410	410	650	19,558
2008	149	382	382	611	18,329
2009	138	362	362	580	17,390
2010	127	295	295	472	14,321
2011	-	300	300	479	14,395

*Separated from Rockfish in 1991.

¹Catch data through November 2010.

²Biomass from annual SAFE report projections.

Stock

Assessment:

Yelloweye rockfish biomass is estimated from submersible transect density and area estimates of DSR habitat.

DSR catch limits are set by a Tier 4 control rule. The 2011 projected biomass is 14,395 mt. Catch specifications for 2011 are as follows; OFL=479 mt, ABC=300 mt, TAC= 300 mt.

Fishery: The directed fishery for DSR is almost entirely prosecuted by longline gear. Yelloweye accounted for 97% (by weight) and quillback accounted for 1.9% (by weight) of the catch from 2003-2008.



Yelloweye rockfish
Rex Murphy



Thornyhead Rockfish

Shortspine thornyhead
AFSC, NOAA Fisheries



For More Information:

S. Lowe and J. Ianelli.
2010. Assessment of
Thornyheads in the Gulf of
Alaska.
<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: The Thornyhead Rockfish complex consists of 3 species; shortspine *Sebastolobus alascanus*, longspine *Sebastolobus altivelis*, and broadfin *Sebastolobus macrochir* thornyheads. Thornyheads are distinguished from “true” rockfish (*Sebastes*) due to their reproductive biology. Whereas *Sebastes spp.* rockfish are viviparous, thornyheads are oviparous, releasing fertilized eggs in floating gelatinous masses. Thornyheads are also differentiated from *Sebastes spp.* in lacking a swim bladder.

Shortspine thornyheads are distributed in deep-water habitats throughout the North Pacific, and are concentrated between 150-450 m in the cooler, northern part of their range and are generally found in deeper habitats up to 1000 m in the warmer waters of their southern range. Females reach 50% maturity at about 22 cm, and longevity extends to 100 years or more. Natural mortality is estimated to be $M=0.03$.

Shortspine thornyheads feed on shrimps, crabs, zooplankton and amphipods and are in turn prey for arrowtooth flounder, sablefish, sperm whales and sharks. Longspine thornyheads are found only in the eastern north Pacific, around the Shumagin Islands, GOA and south to California. Longspines are generally found in deeper habitats from 200-1,750 m.



NOAA Fisheries Service

Catch History: Foreign rockfish harvests peaked in 1965. The greatest reported harvest of thornheads in the GOA occurred from 1979-1983. Catches declined in 1984 and 1985 due to US management restrictions and a transition to domestic fisheries. US catches continued to increase through 1989, peaking at 3,055 mt. Since then, catches have remained well below the TAC.

Total catches, pre-season catch specifications, and exploitable biomass of age 5+ Thornyhead Rockfish* in GOA, 1992-2011 (mt).


Year	Catch ¹	TAC ²	ABC	OFL	Biomass
1992	2,020	1,800	1,800	2,440	25,700
1993	1,369	1,062	1,180	1,441	26,207
1994	1,320	1,180	1,180	1,440	103,300
1995	1,113	1,900	1,900	2,660	30,341
1996	1,100	1,248	1,560	2,200	26,244
1997	1,240	1,700	1,700	2,400	46,108
1998	1,136	2,000	2,000	2,840	52,271
1999	1,282	1,990	1,990	2,800	53,216
2000	1,307	2,360	2,360	2,820	52,950
2001	1,339	2,310	2,310	2,770	52,100
2002	1,125	1,990	1,990	2,330	77,840
2003	1,159	2,000	2,000	3,050	85,760
2004	818	1,940	1,940	2,590	86,200
2005	719	1,940	1,940	2,590	86,200
2006	779	2,209	2,209	2,945	98,158
2007	701	2,209	2,209	2,945	98,158
2008	741	1,910	1,910	2,540	84,774
2009	666	1,910	1,910	2,540	84,775
2010	553	1,770	1,770	2,360	78,795
2011	-	1,770	1,770	2,360	78,795

* includes longspine and shortspine thornyheads.

¹Catch data through November 2010.

²TAC, ABC and OFL from annual Federal Register.

GOA Trawls survey biomass estimates, 1984-2009



Stock Assessment: Thornyhead rockfish catch limits are set using a Tier 5 control rule. The 2011 projected biomass is 78,795 mt. Catch specifications for 2011 are as follows; OFL=2,360 mt, ABC= 1,770 mt, TAC= 1,770 mt.

Fishery: Thornyheads are caught by bottom trawl as a secondary target species in the CGOA Rockfish program and are also taken incidentally in the sablefish longline fishery. Thornyheads are a valuable rockfish species, and most of the domestic harvest is exported to Japan.



Atka Mackerel

AFSC, NOAA Fisheries

Biology: Atka mackerel *Pleurogrammus monopterygius* are distributed along the continental shelf. Atka mackerel is a schooling, semi-demersal species most commonly found in the AI, but also in the Western and Central GOA. Adult Atka mackerel occur in large localized aggregations at depths less than 200 m over rough, uneven bottom areas with high tidal currents. Atka mackerel feed on euphausiids and copepods and are prey for Pacific cod, arrowtooth flounder and Steller sea lions.



For More Information:

S. Lowe. 2010. Assessment of Atka Mackerel stock in the Gulf of Alaska.

<http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Atka mackerel begin to recruit to the fishery at age 3 and longevity can extend to 14 years. Females reach 50% maturity at 31 cm (3.5 years). Natural mortality is estimated at $M=0.30$. Atka mackerel are a substrate-spawning fish with male parental care. Behavioral studies have shown that the Atka mackerel mating system is very complex. A significant characteristic is the bright and distinct coloration developed by territorial males during the spawning season. Spawning occurs from July to October, peaking in early September. Atka mackerel have relative low fecundity, with females releasing around 30,000 eggs each year. Eggs are adhesive and deposited in rock crevices in nests guarded by males until hatching, which occurs about 40-45 days later.



Jackie Patt, UAF

Catch History: Atka mackerel supported a targeted foreign fishery (primarily Soviet vessels) in the Central GOA during the 1970s and 1980s. Catches peaked in 1975 at about 27,000 mt

then declined dramatically to less than 5 mt in 1986. Joint venture operations participated in the Atka mackerel fishery from 1983-1985, and the fishery was fully domestic by 1986.

Fishery Management: In 1988, Atka mackerel were combined with the Other Species category due to low abundance. In 1994, Atka mackerel were removed from the Other Species category and treated once again as a single species target stock. There has not been a directed Atka mackerel fishery in the GOA since 1996.

Stock Assessment: The existing GOA bottom trawl survey data has limited utility for either absolute abundance estimates or indices for Atka mackerel. Atka mackerel fall under Tier 6 control rule. The 2011 catch specifications for Atka mackerel are as follows; OFL=6,200 mt, ABC=4,700 mt, TAC=2,000 mt.

Fishery: Atka mackerel has been a "bycatch" only fishery since 1996.


Ecosystem Components: Because Atka mackerel is thought to be a common prey item for Steller sea lions, all directed fishing for Atka mackerel is prohibited in the GOA.

Total catches, and pre-season catch specifications of Atka Mackerel* in the GOA, 1994-2010 (in mt).

Year	Catch ¹	TAC	ABC	OFL
1994	3,538	3,500	4,800	19,040
1995	701	3,240	3,240	11,700
1996	1,580	3,240	3,240	9,800
1997	331	1,000	1,000	6,200
1998	317	600	600	6,200
1999	262	600	600	6,200
2000	170	600	600	6,200
2001	76	600	600	6,200
2002	85	600	600	6,200
2003	578	600	600	6,200
2004	819	600	600	6,200
2005	799	600	600	6,200
2006	876	1,500	4,700	6,200
2007	1,453	1,500	4,700	6,200
2008	2,109	1,500	4,700	6,200
2009	2,222	3,328	3,328	6,200
2010	2,409	2,000	4,700	6,200
2011	-	2,000	4,700	6,200

*Added to Other Species category in 1988 and separated from Other Species in 1994.

¹Catch data through November 2010.



Jackie Patt, UAF

Longnose Skates, Big Skates, Other Skates

Biology: The GOA Skate complex is comprised of at least 15 skate species. Big skates and longnose skates dominate the skate biomass in the GOA. *Bathyrāja* sp. compose about a third of total GOA skate biomass, with the majority of these being the Aleutian skate and Bering skate. Skate biomass is concentrated in the Central GOA. Skates feed on bottom invertebrates, such as crustaceans, mollusks and polychaetes and fish. Skates are prey for sharks, Steller sea lions and sperm whales.



For More Information:

O. Ormseth and B. Matta. 2010. Assessment of Gulf of Alaska Skates. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

The highest biomass of skates in the GOA is found in continental shelf waters less than 100 m deep, and is dominated by the big skate. In continental shelf waters from 100-200 m depth, longnose skates dominate skate biomass, and *Bathyrāja* skate species are dominant in the deeper waters extending from 200 to 1000 m or more in depth. Big and longnose skate are generally found in shallower waters in the GOA, and their distribution extends from the Bering Sea to southern Baja California. The Aleutian skate ranges

throughout the north Pacific from northern Japan to northern California and has been found at depths between 16-1602 m. The Alaska skate is restricted to higher latitudes from the Sea of Okhotsk to the eastern GOA at depths from 17-392 m. The range of the Bering skate is undetermined.

Big skate	<i>Raja binoculata</i>
Longnose skate	<i>Raja rhina</i>
Other skates	
Aleutian skate	<i>Bathyrāja aleutica</i>
Bering skate	<i>Bathyrāja interrupta</i>
Alaska skate	<i>Bathyrāja parmifera</i>
Deepsea skate	<i>Bathyrāja abyssicola</i>
Commander skate	<i>Bathyrāja lindbergi</i>
Whiteblotched skate	<i>Bathyrāja maculata</i>
Butterfly skate	<i>Bathyrāja mariposa</i>
Whitebrow skate	<i>Bathyrāja minispinosa</i>
Leopard skate	<i>Bathyrāja pamifera</i> sp.
Mud skate	<i>Bathyrāja taranetzi</i>
Roughtail skate	<i>Bathyrāja trachura</i>
Okhotsk skate	<i>Bathyrāja violacea</i>
Roughsholder skate	<i>Bathyrāja badia</i>



Big Skate
Megan Peterson, UAF

Skates are generally K-selected, with slow-growth, low fecundity and relatively large body size. Skates are oviparous; fertilization is internal, and eggs are deposited in horny cases for incubation. There are 1-7 embryos per egg case in locally occurring *Raja* sp., but little is known about the frequency of breeding or egg deposition for any of the local species. It is estimated that annual fecundity per females may be less than 50 eggs per year. The big skate is the largest skate in the GOA, with maximum sizes observed over 200 cm in the directed fishery in 2003. Observed sizes for the longnose skate range from 165-170 cm. The maximum observed lengths for *Bathyrāja* species from bottom trawl surveys of the GOA range from 86-154 cm. Life history parameter data are limited for GOA skates. The AFSC Age and Growth Program has recently reported a maximum observed age of 25 years for the longnose skate in the GOA and a maximum observed age for GOA big skates of 15 years.

Catch History: Skates were caught as a bycatch only species in the GOA at about 1,000-2,000 mt per year from 1992-1995, principally by the longline Pacific cod and bottom trawl pollock and flatfish fisheries. Most skates during this time period were not retained. A directed skate fishery developed in the GOA in 2003 due to an

increase in the ex-vessel value of skates. The skate fishery was prosecuted generally by longline vessels less than 60 feet around Kodiak Island. Lower ex-vessel prices and a possible reduction in skate catch-per-unit effort resulted in a sharp decline in skate catches in 2004-2005.

Directed fishing for skates in the GOA has been prohibited since 2005. Annual average catches of big skates, longnose skates and other skates from 2005-November 2009 have averaged 996 mt, 638 mt, and 557 mt respectively. Catches are highest in the central GOA regulatory area.

Fishery Management: Since the beginning of domestic fishing in the late 1980s through 2003, all species of skates in the GOA were managed under the Other Species FMP category (skates, sharks, squids, sculpins, and octopuses). Catch limits were determined for all Other Species as 5% of the sum of the TACs for GOA target species. Under Amendment 63 in 2003, GOA skates were removed from the Other Species category in 2004 for separate management in response to a developing fishery. Big and longnose skates were managed together under a single TAC in the Central GOA. The remaining skates were managed as an Other Skates species complex in the Central GOA, and all skates were managed as an Other Skates species complex in the Western and Eastern GOA.

In 2005, big skates and longnose skates were separated into single species management groups due to concerns about disproportionate harvests. The remaining skates (genus *Bathyraja*) continue to be managed as a gulfwide species complex because they were not the targets of the fishery and are more difficult to identify. There has been no directed fishing for skates in the GOA since 2005.

Stock Assessment: The Skates stock assessment used estimated biomass data from NMFS summer bottom trawl surveys from 2003-2009. Skates are managed under Tier 5 of the ABC/OFL control rule, based on an overall natural mortality rate of 0.10 applied to survey biomass estimates for each species group. Gulf wide catch specifications (mt) for 2011 are as follows.

	Biomass	OFL	ABC	TAC	2010 Catch
Big Skates	44,381	4,438	3,328	3,328	2,437
Longnose skates	38,031	3,803	2,852	2,852	1,043
Other skates	28,908	2,791	2,093	2,093	1,464

Note that the ABC and TAC are further broken out into Western, Central, and Eastern Gulf of Alaska for big skate and longnose skates.

Fishery: GOA Skates have been a bycatch only fishery since 2005. Skates are generally caught as bycatch in Pacific halibut and Pacific cod longline fisheries and flatfish trawl fisheries, especially in the GOA Central regulatory area. The incidental catch of Big skates in the Central area has the potential to constrain fisheries.

Ecosystem Components: Skates have few natural predators. In the GOA, skate predators include marine mammals such as Steller sea lions and sperm whales (which may consume adult or juvenile skates), and spiny dogfish (which likely consume juvenile skates).



Spiny dogfish
AFSC, NOAA Fisheries



For More Information:

C. Tribuzio, K. Echave, C. Rodegveiler, J. Heifetz, and K. Goldman. 2010. Assessment of Sharks in the Gulf of Alaska. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Biology: The GOA Shark complex is composed of 8 shark species. The most abundant species in the GOA are the spiny dogfish, the salmon shark and the Pacific sleeper shark. GOA sharks exhibit K-selected life history traits including slow growth to maturity, low fecundity and large size. Spiny dogfish, Pacific sleeper shark and salmon sharks reproduce through aplacental viviparity. Shark diets vary with species and in general sharks are opportunistic feeders, but forage fish, crustaceans, squid and salmon are among the most common prey items.

Spiny dogfish	<i>Squalus acanthias</i>
Salmon shark	<i>Lamna ditropis</i>
Pacific sleeper shark	<i>Somniosus pacificus</i>
Brown cat shark	<i>Apristurus brunneus</i>
White shark	<i>Carcharodon carcharias</i>
Basking shark	<i>Cetorhinus maximus</i>
Sixgill shark	<i>Hexanchus griseus</i>
Blue shark	<i>Prionace glauca</i>

Spiny dogfish are distributed from California to Alaska, through the Aleutian chain to the Asian coast and south to Japan. Spiny dogfish are found at depths ranging from the intertidal to 900 m. Spiny dogfish growth rates are among the slowest of all shark species. Estimates of spiny dogfish age-at-50%-maturity are 20 years for males to 34 years for females. Longevity is estimated to reach between 80 and 100 years. Natural mortality is estimated at $M=0.097$. Spiny dogfish have one of the longest known gestation periods, approximately 18-24 months.



Spiny Dogfish, Mark Fina, NPFMC

Pacific sleeper sharks are found along the North Pacific continental shelf and slope, ranging from Japan to the Bering Sea. Distribution extends as far north as the Chukchi Sea and as far south as Baja California. At higher latitudes, Pacific sleeper sharks are found shallower from littoral zones to surface waters. At lower latitudes, they reside much deeper

and down to 2000 m. Pacific sleeper sharks make extensive, nearly continuous vertical movements. The maximum lengths of captured Pacific sleeper sharks are 440 cm for females and 400 cm for males. Pacific sleeper sharks 150-250 cm in length are most common in Alaska. Pacific sleeper shark age and reproduction data are limited.

Salmon shark distribution in the northern Pacific extends from Japan into the Sea of Okhotsk to the Bering Sea and possibly south as far as Baja California Mexico. Salmon sharks live in areas with sea-surface temperatures between 5°C and 18°C and in depths up to 150 m. However, salmon sharks spend about 72% in waters



Salmon Shark, Ken Goldman, ADF&G

less than 50 m deep. While some salmon sharks migrate south during the winter months, others remain in the GOA throughout the year. Longevity estimates for salmon sharks are between 20-30 years with maturity occurring at 3-5 years for males and 6-9 years for females. Natural mortality is estimated at $M=0.18$.

Catch History: There are currently no directed commercial fisheries for shark species in federal or state managed waters of the GOA, and most incidentally caught sharks are not retained. A small number of spiny dogfish landings in Kodiak were reported in 2004, 2005 and 2007 (approximately 1 mt each year). Spiny dogfish and salmon sharks are also caught in recreational fisheries in the GOA. Estimates of historic catches of sharks range from 308 mt in 1995 to a peak of 2,390 mt in 1998. Catches annually averaged 895 mt from 1992-1999 and 962 mt from 2000-2009.



Spiny dogfish, Cindy Tribuzio, AFSC, NOAA Fisheries

Fishery Management: Until 2011, sharks were managed under the Other Species FMP category (sharks, squids, sculpins, and octopuses). Beginning in 2011, sharks will be managed as a single complex.

Total catches, and pre-season catch specifications of Sharks* in the GOA, 1994-2010 (in mt).

Year	Catch ¹	TAC	ABC	OFL
1994	360	-	-	-
1995	308	-	-	-
1996	484	-	-	-
1997	1,041	-	-	-
1998	2,390	-	-	-
1999	1,036	-	-	-
2000	1,117	-	-	-
2001	853	-	-	-
2002	427	-	-	-
2003	751	-	-	-
2004	573	-	-	-
2005	1,101	-	-	-
2006	1,603	-	-	-
2007	1,406	-	-	-
2008	619	-	-	-
2009	1,167	-	-	-
2010	603	-	-	-
2011		6,197	6,197	8,262

*Split from Other Species in 2011.
¹Catch data through November 2010.

Stock Assessment: Catch specifications for sharks are based on a split Tier system. Tier 5 is used for dogfish sharks, with natural mortality ($M=0.097$) applied to biomass estimate (79,257 mt). Tier 6 is used for other sharks based on average historical catch from 1997-2007. Catch specifications for sharks in 2011 are as follows; OFL=8,262 mt, ABC=6,197 mt.

Fishery: GOA sharks are managed as a bycatch only fishery. In 2010, the catch was 329 mt of spiny dogfish, 159 mt of sleeper sharks, and 107 mt of salmon sharks. On average, over 90% of the sharks are discarded. Spiny dogfish were caught primarily in the longline Pacific cod and bottom trawl flatfish fisheries. Over 90% of Pacific sleeper sharks and salmon sharks were caught in the pollock fishery.



AFSC, NOAA Fisheries

Biology: There are at least 14 species of squid in the Gulf of Alaska and managed as a squid complex. The most common squid near the continental shelf are the minimal armhook squid and the magistrate armhook squid. On the slope, the most common squid species are the boreopacific armhook squid and other *Gonotus* armhook squid. Very little is known about the species of squid in the Gulf of Alaska.



For More Information:

O. Ormseth and C. Spital. 2010. Assessment of Gulf of Alaska Squids. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Squids are short-lived (<4 years), maturing just prior to spawning and dying afterwards. Squid populations consist of multiple cohorts that school with similar sized individuals, and may occupy different areas of the shelf and slope.

Chiroteuthid sp.	<i>Chiroteuthis calyx</i>
Glass squid sp.	<i>Belonella borealis</i>
Glass squid sp.	<i>Galiteuthis phyllura</i>
Minimal armhook squid	<i>Berryteuthis anonychus</i>
Magistrate armhook squid	<i>Berryteuthis magister</i>
Armhook squid	<i>Eogonatus tinro</i>
Boreopacific armhook squid	<i>Gonatopsis borealis</i>
Berry armhook squid	<i>Gonatus berryi</i>
Armhook squid sp.	<i>Gonatus madokai</i>
Armhook squid sp.	<i>Gonatus middendorffi</i>
Clawed armhook squid	<i>Gonatus onyx</i>
Robust clubhook squid	<i>Moroteuthis robusta</i>
Boreal clubhook squid	<i>Onychoteuthis borealijaponicus</i>
North Pacific bobtail squid	<i>Rossia pacifica</i>

Fishery Management:

Squid were defined as an “other species” in the GOA until 2011 when the “other species” complex was separated out into distinct species groupings.

Stock Assessment: Catch specifications for Squid are set using a modified Tier 6 control rule, with catch specifications are based on the highest catch during 1997-2008. Squid estimated biomass in undefined. Catch specifications for squid in 2011 are as follows; OFL=1,530 mt, ABC=1,148 mt, TAC=1,148 mt.

Fishery: There is currently no target fishery for squid in the GOA. GOA squid are primarily (> 90%) taken as incidental catch in the pelagic trawl pollock fishery. They are also taken in smaller numbers in bottom trawl fisheries. About 90% of the squid catch has been retained in recent years.

Total catches, and pre-season catch specifications of Squid* in the GOA, 1997-2010 (in mt).

Year	Catch ¹	TAC	ABC	OFL
1997	98	-	-	-
1998	59	-	-	-
1999	41	-	-	-
2000	19	-	-	-
2001	91	-	-	-
2002	42	-	-	-
2003	92	-	-	-
2004	162	-	-	-
2005	635	-	-	-
2006	1,530	-	-	-
2007	412	-	-	-
2008	84	-	-	-
2009	337	-	-	-
2010	130	-	-	-
2011		1,148	1,148	1,530


*Split from Other Species in 2011.
¹Catch data through November 2010.



Tim Evers

Ecosystem Components:

Squid are not currently a commercially valuable species in the North Pacific, however they play a critical prey role in ecosystems. They are important components in the diets of many seabirds, fish and marine mammals. Overall fishing removals of squid are low (especially relative to natural predation).



Giant Pacific Octopus
Linda Kozak



For More Information :

M.E. Conners and C. Conrath. 2010. Assessment of the Octopus Complex in the GOA.

<http://www.afsc.noaa.gov/refm/stocks/assessments.htm>

Biology: There are at least 7 species of octopus present in federal waters of the GOA, and the species composition both of natural communities and commercial harvest is unknown. At depths less than 200 meters, the giant Pacific octopus *E. dofleini* appears to be the most abundant species. Octopus life spans are either 1-2 years or 3-5 years depending on the species. *E. dofleini* are estimated to mature at 1.5 – 3 years. male *E. dofleini* were found to mature at around 12.5 kg with females thought to mature at larger sizes. *E. dofleini* is a terminal spawner, females die after the eggs hatch while males die shortly after mating. The fecundity of this species in Japanese waters has been estimated at 30,000 to 100,000 eggs per female. There are two other common species of octopus in the GOA: the smoothskin octopus and the flapjack devilfish. The smoothskin octopus occurs from 250-1400 m. and produces few eggs that remain benthic after hatching. The flapjack devilfish is found from 300-1000m deep and spawn up to 2,400 eggs in multiple batches.

Giant Pacific octopus	<i>Enteroctopus dofleini</i>
Smoothskin octopus	<i>Benthoctopus leioderma</i>
Flapjack devilfish	<i>Opisthoteuthis californiana</i>
Pelagic octopus	<i>Japatella diaphana</i>
Red octopus	<i>Octopus californicus</i>
Black octopus	<i>Vampyroteuthis infernalis</i>
a small octopus	<i>Octopus sp. A</i>

Fishery Management: Until 2011, octopus were managed as part of the “Other species” management category within the GOA FMP. Beginning in 2011, octopuses will be managed as a single complex with its own ABC and OFL.

Stock Assessment: Octopus catch limits are specified using a modified Tier 6 control rule, with an estimate of natural mortality ($M=0.53$) applied to the biomass of the 3 most recent NMFS bottom trawl surveys. While the biomass is deemed unreliable for purposes of Tier 5, it does provide a minimum estimate of biomass. Catch



Megan Peterson, UAF

specifications for octopus in 2011 are as follows; OFL=1,272 mt, ABC=954 mt, TAC=954 mt.


Fishery: There is currently no target fishery for octopus in federal waters of the GOA. About 90% of the octopus catch is taken as incidental catch in the Pacific cod pot fisheries in the western and central GOA. In 2010, 271 mt of octopus were retained for human consumption or for bait for the halibut fishery. The species composition of the octopus catch is unknown, but based on research trawl data, the giant Pacific octopus is most abundant in shelf waters and predominates in commercial catch. Preliminary research suggests high survival for octopus released from pot gear.

Total catches, and pre-season catch specifications of Octopus* in the GOA, 1997-2010 (in mt).

Year	Catch ¹	TAC	ABC	OFL
1997	232	-	-	-
1998	112	-	-	-
1999	166	-	-	-
2000	156	--	-	-
2001	88	-	-	-
2002	298	-	-	-
2003	210	-	-	-
2004	286	-	-	-
2005	151	-	-	-
2006	159	-	-	-
2007	262	-	-	-
2008	339	-	-	-
2009	310	-	-	-
2010	324	-	-	-
2011	-	954	954	1,272

*Split from Other Species in 2011.

¹Catch data through November 2010.




Great sculpin
AFSC, NOAA Fisheries

Biology: There are 39 species of sculpins identified in the Gulf of Alaska and managed as a sculpin complex. The most common sculpin species taken incidentally in GOA fisheries are the yellow Irish lord *Hemilepidotus jordani* making up over 60% of the catch, followed by great sculpin *Myoxocephalus polyacanthocephalus*, bigmouth sculpin *Hemitripterus bolini* and plain sculpin *M. joak*. Sculpins lay adhesive eggs in nests, and many exhibit parental care for eggs. Irish lords and great sculpins have an age at 50% maturity of about 7 years.

Catch history: There is no directed fishing for any sculpin species in the GOA at this time. Catch of sculpins in the last 15 years has been averaged about 900 mt per year, reaching a peak in 2008 of 1,943 mt.

Fishery Management: Prior to 2011, sculpins were managed as part of the GOA Other Species complex that included sculpins, skates, sharks, squid and octopus, with an aggregate OFL, ABC, and TAC. Beginning in 2011 sculpins were removed from Other Species and managed as a separate group, as were the remaining species groups. Sculpins are currently taken only as incidental catch in fisheries directed at other target species, and it is likely that catch of sculpins in the near future will continue to be dependent on the distribution and limitations placed on target fisheries, rather than on any harvest level established for this category.

Stock Assessment: Sculpins are managed under Tier 5 of the



For More Information:


O. Ormseth and T. Tenbrink. 2010. Assessment of Gulf of Alaska Sculpins. <http://www.afsc.noaa.gov/ref/m/stocks/assessments.htm>

Catches, pre-season catch specifications and estimated biomass (t) of Sculpins in the GOA, 1997-2011.				
Year	Catch	ABC	OFL	Biomass ²
1997	898	-	-	-
1998	526	-	-	-
1999	544	-	-	30,783
2000	940	-	-	-
2001	587	-	-	30,418
2002	919	-	-	-
2003	629	-	-	26,514
2004	816	-	-	-
2005	626	-	-	33,519
2006	583	-	-	-
2007	960	-	-	32,468
2008	1,943	-	-	-
2009	1,146	-	-	40,726
2010	735	-	-	-
2011	-	5,496	7,328	33,307

*Sculpins removed from Other Species in 2011
¹ Estimated catch data from the SAFE.
² Biomass estimate (t) from trawl surveys.

OFL/ABC guidelines, and catch specifications are based on natural mortality for the complex ($M=0.22$) applied to average survey biomass. Catch specifications for sculpins in 2011 are as follows; OFL=7,328 mt, ABC=5,496 mt, TAC=5,496 mt.

Fishery: There is currently no target fishery for sculpins in the GOA, and virtually all are either discarded or made into meal. Incidental catches of sculpins are taken in the Pacific cod, shallow water flatfish, and rockfish fisheries, as well as the halibut longline fishery.



Bigmouth sculpin,
AFSC, NOAA Fisheries



Plan Team Members and Assessment Authors

Jackie Patt

Bering Sea/Aleutian Islands Groundfish Plan Team

Kerim Aydin (AFSC)
David Barnard (ADF&G)
David Carlile (ADF&G)
Henry Cheng (WDFW)
Bill Clark (IPHC)
Jane DiCosimo (NPFMC)
Lowell Fritz (AFSC)
Mary Furuness (NMFS)
Dana Hanselman (AFSC)
Alan Haynie (AFSC)
Loh-lee Low, Chair (AFSC)
Brenda Norcross (UAF)
Mike Sigler, Vice-Chair (AFSC)
Leslie Slater (USF&W)
Grant Thompson (AFSC)

Gulf of Alaska Groundfish Plan Team

Henry Cheng (WDFW)
Michael Dalton (AFSC)
Robert Foy (UAF)
Nancy Friday (AFSC)
Sarah Gaichas (AFSC)
Ken Goldman (ADF&G)
Kristin Green (ADF&G)
Steven Hare (IPHC)
Jon Heifetz (AFSC)
Jim Ianelli, Co-Chair (AFSC)
Sandra Lowe (AFSC)
Chris Lundsford (AFSC)
Thomas Pearson (NMFS)
Nick Sagalkin (ADFG)
Leslie Slater (USF&W)
Paul Spencer (AFSC)
Diana Stram, Co-Chair (NPFMC)

Bering Sea/Aleutian Islands Stock Assessments

Jim Ianelli (AFSC)
Steve Barbeaux (AFSC)
Grant Thompson (AFSC)
Dana Hanselman (AFSC)
Tom Wilderbuer (AFSC)
William Stockhausen (AFSC)
Paul Spencer (AFSC)
Sandra Lowe (AFSC)
Olav Ormseth (AFSC)
Cindy Tribuzio (AFSC)
Elizabeth Connors (AFSC)

Gulf of Alaska Stock Assessments

Martin Dorn (AFSC)
Grant Thompson (AFSC)
Dana Hanselman (AFSC)
Benjamin Turnock (AFSC)
Teresa A'mar (AFSC)
William Stockhausen (AFSC)
Jon Heifetz (AFSC)
Dave Clausen (AFSC)
Chris Lunsford (AFSC)
Kalei Shotwell (AFSC)
Kristen Green (ADF&G)
Sandra Lowe (AFSC)
Olav Ormseth (AFSC)
Cindy Tribuzio (AFSC)



Council, SSC, AP Members and Staff

Council Members

Eric Olson, Chair, Anchorage, AK
Dave Benson, Vice-Chair, Seattle, WA
Sam Cotton, Eagle River, AK
Ed Dersham, Anchor Point, AK
Duncan Fields, Kodiak, AK
John Henderschedt, Seattle, WA
Dan Hull, Anchorage, AK
Roy Hyder, ODFW, Portland, OR
Jim Balsiger Ph.D., NMFS, Juneau, AK
Phil Anderson/Bill Tweit, WDFW, Seattle, WA
Cora Campbell/Stefanie Moreland, ADF&G, Juneau, AK
Dave Hanson, Ph.D., PSMFC, Lake Oswego, OR
RADM Christopher Colvin/CAPT Mike Cerne, USCG, AK
Denny Lassuy, USFWS, Anchorage, AK
Nicole Ricci, US State Dept, Washington DC

Staff

Chris Oliver, Executive Director
David Witherell, Deputy Director
Gail Bendixen, Administrative Officer
Jane DiCosimo, Senior Plan Coordinator
Diana Evans, Fishery Analyst
Mark Fina, Ph.D., Senior Economist
Jeannie Heltzel, Protected Resources
Nicole Kimball, Fishery Analyst
Peggy Kircher, Administrative Assistant
Jon McCracken, Economist
Sarah Melton, Fishery Analyst
Maria Shawback, Administrative Assistant
Diana Stram, Ph.D., Plan Coordinator

Michael Fey, Data Manager, PSMFC

Scientific and Statistical Committee Members

Pat Livingston, Chair, NOAA Fisheries, AFSC
Farron Wallace, Vice-Chair, WDFW
Jennifer Burns Ph.D., University of Alaska
Susan Hilber, Oregon Department of Fish and Wildlife
Bob Clark, Alaska Dept. of Fish and Game
Anne Hollowed, Ph.D., NOAA Fisheries, AFSC
George Hunt, Ph.D., University of Washington
Gordon Kruse, Ph.D., University of Alaska
Kathy Kuletz, Ph.D., US Fish and Wildlife Service
Seth Macinko, Ph.D., University of Rhode Island
Franz Mueter, Ph.D., University of Alaska
Jim Murphy, Ph.D., University of Alaska
Lew Queirolo, Ph.D., NOAA Fisheries, AKRO
Terry Quinn, Ph.D., University of Alaska
Katherine Reedy-Maschner, Ph.D., Idaho State U.
Raymond Webster, Ph.D., Pacific Halibut Commission
Doug Woodby, Ph.D., Alaska Dept. of Fish and Game

Advisory Panel Members

Tom Enlow, Chair
Lori Swanson, Vice Chair
Matt Moir, Vice Chair
Kurt Cochran
Craig Cross
John Crowley
Julianne Curry
Jerry Downing
Tim Evers
Jeff Farvour
Jan Jacobs
Bob Jacobson
Alexus Kwachka
Chuck McCallum
Teresa Peterson
Ed Poulsen
Becca Robbins Gisclair
Neil Rodriguez
Beth Stewart
Anne Vanderhoeven



North Pacific Fishery Management Council

605 W. 4th Avenue, Suite 306

Anchorage, AK 99501

(907) 271-2809

Fax: (907) 271-2817

www.alaskafisheries.gov/npfmc



Produced by NPFMC under NOAA Award # NA10NMF4410005