

Nonnative Invasive Plants of Southern Forests

A Field Guide for Identification and Control



Author

James H. Miller, Research Ecologist, USDA Forest Service,
Southern Research Station, Auburn University, AL 36849.

Available without charge from
the Southern Research Station

Front Cover

Upper left—Chinese lespedeza (*Lespedeza cuneata*) infestation that developed from dormant seed in the soil seed bank after a forest thinning operation.

Upper right—Kudzu (*Pueraria montana*) infestation on the urban-wildland interface.

Lower left—Chinese privet (*Ligustrum sinense*) and dormant kudzu invading and replacing a pine-hardwood stand.

Lower right—Cogongrass (*Imperata cylindrica*) infestation under mature slash pine (*Pinus elliottii*).

Back Cover

Upper left—Stem injection using a hatchet and spray bottle to apply a hack-and-squirt treatment to control silktree (*Albizia julibrissin*).

Upper right—Hand pulling seedlings of Chinese privet (*Ligustrum sinense*).

Lower left—Prescribed burning can often assist in controlling garlic mustard (*Alliaria petiolata*).

Lower right—Containerized native plants for rehabilitation plantings.

Funding support for 2006 reprinting provided by the
Eastern Forest Environmental Threat Assessment Center, Asheville, NC.

First Printed May 2003
Revised August 2003
Revised December 2004
Revised August 2006

Southern Research Station
P.O. Box 2680
Asheville, NC 28802

**Nonnative Invasive Plants
of Southern Forests**
A Field Guide for Identification and Control

James H. Miller

Acknowledgments

The contributions of Erwin B. Chambliss, USDA Forest Service, Auburn, AL, have been invaluable in image management and layout. Kristine Johnson, Great Smoky Mountains National Park; Johnny Randall, North Carolina Botanical Gardens; Jack Ranney, University of Tennessee; and Fred Nations, Weeks Bay National Estuarine Research Reserve, made comprehensive reviews and provided invaluable recommendations for improvements. Reviews of control recommendations were made by Ron Cornish, Dow AgroScience; Harry Quicke, BASF Corporation; Carroll Walls, UAP Timberlands; and Michael Link, DuPont Corporation. Their knowledgeable comments greatly strengthened content and clarity.

All Plant Images by the Author Except for the Contributions by:

Ted Bodner, Southern Weed Science Society, Champaign, IL

Patrick Breen, Department of Horticulture, Oregon State University, Corvallis, OR

Charles T. Bryson, USDA Agricultural Research Service, Southern Weed Science Research Unit, Stoneville, MS

Thomas Ellis, Jr., Baldwin County Forestry Planning Committee, Bay Minette, AL

Chris Evans, University of Georgia, Tifton, GA

John W. Everest, Department of Agronomy & Soils, Auburn University, Auburn, AL

Jerry Gibson, Deer Park, AL

Kristine Johnson, Great Smoky Mountains National Park, Gatlinburg, TN

Keith Langdon, Great Smoky Mountains National Park, Gatlinburg, TN

John Meade, Department of Plant Science, Rutgers, The State University of New Jersey, New Brunswick, NJ

Fred Nation, Weeks Bay National Estuarine Research Reserve, Fairhope, AL

John M. Randall, The Nature Conservancy, University of California, Davis, CA

Barry A. Rice, The Nature Conservancy, University of California, Davis, CA

Susan Ross, Great Smoky Mountains National Park, Gatlinburg, TN

John Schwegman, Illinois Department of Natural Resources, Springfield, IL

Jody Shimp, Illinois Department of Natural Resources, Springfield, IL

Warner Park Nature Center, Metropolitan Board of Parks and Recreation, Nashville, TN

Hugh Wilson, Texas A&M University, College Station, TX

Plant Names and Plant Distribution Maps from:

USDA Natural Resources Conservation Service's Plants Database: <http://plants.usda.gov>.

Images by Ted Bodner copyrighted by the Southern Weed Science Society for "Forest Plants of the Southeast and Their Wildlife Uses" and used by permission.

ABSTRACT

Invasions of nonnative plants into forests of the Southern United States continue to go unchecked and unmonitored. Invasive nonnative plants infest under and beside forest canopies and dominate small forest openings, increasingly eroding forest productivity, hindering forest use and management activities, and degrading diversity and wildlife habitat. Often called nonnative, exotic, nonindigenous, alien, or noxious weeds, they occur as trees, shrubs, vines, grasses, ferns, and forbs. This book provides information on accurate identification and effective control of the 33 nonnative plants and groups that are currently invading the forests of the 13 Southern States, showing both growing and dormant season traits. It lists other nonnative plants of growing concern, control strategies, and selective herbicide application procedures. Recommendations for preventing and managing invasions on a specific site include maintaining forest vigor with minimal disturbance, constant surveillance and treatment of new unwanted arrivals, and finally rehabilitation following eradication.

Keywords: Alien plants, exotic plant control, exotic weeds, herbicide weed control, integrated vegetation management, invasive exotic plants, invasive nonindigenous plants, noxious plant control.

Contents

Explanation of Codes and Conventions Used in Species Descriptions	iv		
Introduction	1		
Trees			
Tree-of-Heaven	2		
Silktree, Mimosa	4		
Princesstree, Paulownia	6		
Chinaberrytree	8		
Tallowtree, Popcorn tree	10		
Russian Olive	12		
Shrubs			
Silverthorn, Thorny Olive	14		
Autumn Olive	16		
Winged Burning Bush	18		
Chinese / European Privet	20		
Japanese / Glossy Privet	22		
Bush Honeysuckles	24		
Sacred Bamboo, Nandina	26		
Nonnative Roses	28		
Vines			
Oriental Bittersweet	30		
Climbing Yams	32		
Winter Creeper	34		
English Ivy	36		
Japanese Honeysuckle	38		
Kudzu	40		
Vincas, Periwinkles	42		
Nonnative Wisterias	44		
Grasses			
Giant Reed	46		
Tall Fescue	48		
Cogongrass	50		
Nepalese Browntop	52		
Grasses (cont.)			
Chinese Silvergrass	54		
Bamboos	56		
Ferns			
Japanese Climbing Fern	58		
Forbs			
Garlic Mustard	60		
Shrubby Lespedeza	62		
Chinese Lespedeza	64		
Tropical Soda Apple	66		
Other Nonnative Plants Invading Southern Forests and Their Margins	68		
General Principles for Controlling Nonnative Invasive Plants	69		
Effective Treatments	69		
Selective Herbicide Application Methods	69		
Selecting an Effective Herbicide	72		
Other Treatments for an Integrated Approach	73		
The Rehabilitation Phase	74		
Prescriptions for Specific Nonnative Invasive Plants	75		
Nonnative Trees	75		
Nonnative Shrubs	77		
Nonnative Vines	79		
Nonnative Grasses	82		
Nonnative Ferns	83		
Nonnative Forbs and Subshrubs	83		
Sources of Control Information	85		
Glossary of Important Terms	87		
Flower Parts, Flower Types, Inflorescences	91		
Leaf Arrangements, Leaf Divisions, Shapes, Margins	92		
Parts of a Grass Plant	92		

Explanation of Codes and Conventions Used in Species Descriptions

Scientific name



Ailanthus altissima (P. Mill.) Swingle
Synonyms: ailanthus, Chinese sumac, stinking sumac, paradise-tree, copal-tree

International code



AIAL

Common name



Tree-of-Heaven

Forest Inventory and Analysis survey code



0341

Plant. Deciduous tree to 80 feet (25 m) in height and 6 feet (1.8 m) in diameter, with long pinnately compound leaves and circular glands under lobes on leaflet bases. Strong odor from flowers and other parts, sometimes likened to peanuts or cashews.

Stem. Twigs stout, chestnut brown to reddish tan, and smooth-to-velvety with light dots (lenticels) and heart-shaped leaf scars. Buds finely hairy, dome-shaped, and partially hidden by the leaf base. Branches light gray to dark gray, smooth and glossy, with raised dots becoming fissures with age. Bark light gray and rough with areas of light-tan fissures.



Introduction

Invasions of nonnative plants into southern forests continue to go largely unchecked and only partially monitored. Invasive nonnative plants infest under and beside forest canopies and occupy small forest openings, increasingly eroding forest productivity, hindering forest use and management activities, and degrading diversity and wildlife habitat. Often called non-native, exotic, nonindigenous, alien, or noxious weeds, they occur as trees, shrubs, vines, grasses, ferns, and forbs. Some have been introduced into this country accidentally, but most were brought here as ornamentals or for livestock forage. These robust plants arrived without their natural predators of insects and diseases that tend to keep native plants in natural balance. Many have hybridized. Now they increase across the landscape with little opposition, beyond the control and reclamation measures applied by landowners and managers on individual land holdings.

The objective of this book is to provide information on accurate identification and effective control of the 33 plants or groups that are invading the forests of the 13 Southern States at an alarming rate, showing both growing and dormant season traits. It lists other nonnative invasive plants of growing concern and explains control recommendations and selective application procedures. The text and photographs were originally developed to assist in the first region-wide survey and monitoring of these invading species, conducted by the USDA Forest Service's Forest Inventory and Analysis Research Work Unit of the Southern Research Station in collaboration with State forestry management agencies. The four-number survey codes as well as the international plant codes are given for each species (see opposite page).

Integrated vegetation management programs are needed to combat invading nonnative plants. Strategies of surveillance and treatment of new arrivals will safeguard lands, and rehabilitation of existing infestations can be achieved by concerted control measures and reestablishment of native vegetation.



Broadcast treatment of herbicide spray to kudzu.