

**A global simulation of tropospheric ozone and related tracers:
Description and evaluation of MOZART, version 2**

**Larry W. Horowitz
Stacy Walters
Denise L. Mauzerall
Louisa K. Emmons
Phil J. Rasch
Claire Granier
Xuexi Tie
Jean-François Lamarque
Martin Schultz
Guy P. Brasseur**

to be submitted to *Journal of Geophysical Research*

DRAFT -- April 14, 2001


<http://acd.ucar.edu/~larryh/mozart.html>

MOZART Workshop, NCAR

April 16, 2001

MOZART-2

- Model description
 - Overview
 - Recent modifications
- Comparison with observations
- O₃ budget



MOZART-2

Model resolution: 2.8° latitude × 2.8° longitude (≈T42), 20 min timestep,

34 vertical layers (hybrid), surface to ~35 km (5 mb)

Meteorology: Winds, T, P, humidity, surface fluxes

From MACCM-3 (truncated to 34 layers), every 6 hours

Surface emissions: NO_x (fossil fuel, **biomass burning**, soils)

CO, **non-methane hydrocarbons** (FF, vegetation, **biomass burning**, oceans)

Lightning: NO_x source based on convection [*Price et al.*, 1997; *Pickering et al.*, 1998]

Photochemistry: 58 chemical species, 132 kinetic + 31 photolysis reactions

Photolysis rates computed from lookup table

Advection: Flux-form semi-Lagrangian scheme [*Lin and Rood*, 1996]


Convection: Rediagnosed from MACCM-3 data using *Hack* [1994] and

Zhang and MacFarlane [1995] schemes

Boundary layer diffusion: Based on *Holtlag and Boville* [1993]


Dry deposition: Deposition velocities computed with *Wesely* [1989] scheme, using NCEP meteorology and *DeFries and Townshend* [1994] vegetation map

Wet deposition: Based on *Giorgi and Chameides* [1985]






MOZART-2

- Recent modifications
 - Emissions
 - Dry deposition velocities
 - New isoprene scheme
 - Lightning NO_x
 - “Flexible” architecture version
 - Updated reaction rates
 - Improved O_3 upper BC





MOZART-2

NO_x Emissions





MOZART-2

CO Emissions





MOZART-2

Acetone Emissions





MOZART-2

Deposition Velocities (July)



MOZART-2


Lightning NO_x
(Global Annual Total = 2.5 Tg y^{-1})



MOZART-2

Emissions


Species	Fossil fuel combustion	Biomass burning	Biogenic / Soil	Oceans	Total
NO (TgN/y)	23.1	8.7	6.6	0	38.4
CO (Tg/y)	306.9	711.2	181.0	2.0	1201.1
C ₂ H ₆ (TgC/y)	6.4	4.5	0.8	0.1	11.7
C ₃ H ₈ (TgC/y)	10.0	2.2	1.6	0.1	14.00
C ₂ H ₄ (TgC/y)	2.0	12.3	4.3	2.1	20.7
C ₃ H ₆ (TgC/y)	0.9	5.6	0.9	2.5	9.8
C ₄ H ₁₀ (TgC/y)	22.2	23.0	21.4	6.3	72.9
CH ₃ COCH ₃ (Tg/y)	1.0	16.1	10.2	9.9	37.3
ISOP (TgC/y)	0	0	299.0	0	299.0
C ₁₀ H ₁₆ (TgC/y)	0	0	129.1	0	129.1
CH ₄ (Tg/y)	95.0	64.3	145.7	11.3	467.9
N ₂ O (Tg/y)	5.0	1.7	20.7	11.3	38.7
H ₂ (Tg/y)	13.5	14.0	5.3	7.8	40.6



MOZART-2


Emissions (new, old)

Species	Fossil fuel combustion	Biomass burning	Biogenic / Soil	Oceans	Total
NO (TgN/y)	23.1	8.7 10.0	6.6	0	38.4 39.5
CO (Tg/y)	306.9	711.2 858.4	181.0	2.0 20.1	1201.1 1353.8
C ₂ H ₆ (TgC/y)	6.4 3.2	4.5 5.4	0.8	0.1 0.8	11.7 10.2
C ₃ H ₈ (TgC/y)	10.0 5.0	2.2 2.6	1.6	0.1 1.1	14.0 10.2
C ₂ H ₄ (TgC/y)	2.0	12.3 14.1	4.3	2.1 8.4	20.7 28.7
C ₃ H ₆ (TgC/y)	0.9	5.6 6.4	0.9	2.5 10.2	9.8 18.3
C ₄ H ₁₀ (TgC/y)	22.2 44.0	23.0 39.5	21.4 214.3	6.3 63.2	72.9 186.7
Acetone (Tg/y)	1.0	16.1 15.1	10.2	9.9 0	37.3 26.1
Isoprene (TgC/y)	0	0	299.0 500.3	0	299.0 500.3



MOZART-2


- Comparison with observations
 - ~~Ozone sondes~~
 - CMDL (CO)
 - Aircraft measurements
 - DIAL O₃
 - CO
 - Acetone
 - others?



MOZART-2

SUMMARY

- O₃, NO_x, CO, NMHCs, peroxides
 - **very good** agreement with obs
- PAN, CH₂O
 - **good** agreement at most locations
- HNO₃, acetone
 - significant disagreement at some locations



MOZART-2


Ozone budget

Process	Production (Loss) [Tg O ₃ / yr]		
	Global	Northern Hemisphere	Southern Hemisphere
Influx from stratosphere ^a	409 ^b	267	142
Photochemical production	4671	2750	1921
Photochemical loss	-4238	-2463	-1775
Dry deposition	-847	-557	-290

For this budget, the tropopause is defined as the hybrid model level interface corresponding to approximately 100 hPa in the tropics (30°S-30°N) and 250hPa in the extratropics..

a. Includes advection, pressure consistency correction, and convection and vertical diffusion.

b. This term consists of advection (310 Tg/y), pressure consistency correction (91 Tg/y), and convection and vertical diffusion (8 Tg/y).



Ozone fluxes (Tg y^{-1})

