

Appendix II. Summary of the confirmed human-caused mortality and serious injury (SI) events involving baleen whale stocks along the Gulf of Mexico Coast, US East Coast, and adjacent Canadian Maritimes, 2005-2009, with number of events attributed to entanglements or vessel collisions by year.

Stock	Mean annual mortality and SI rate (PBR <sup>1</sup> for reference)	Entanglements			Vessel Collisions		
		Annual rate (US waters / Canadian waters)	Confirmed mortalities (2005, 2006, 2007, 2008, 2009)	Confirmed SIs (2005, 2006, 2007, 2008, 2009)	Annual rate (US waters / Canadian waters)	Confirmed mortalities (2005, 2006, 2007, 2008, 2009)	Confirmed SIs (2005, 2006, 2007, 2008, 2009)
Western North Atlantic right whale ( <i>Eubalaena glacialis</i> )	<b>2.4</b> (0.8)	<b>0.8</b> (0.8/0)	(0, 1, 1, 0, 0)	(0, 0, 0, 0, 2)	<b>1.6</b> (1.2 / 0.4)	(2, 4, 0, 0, 0)	(1, 1, 0, 0, 0)
Gulf of Maine humpback whale ( <i>Megaptera novaeangliae</i> )	<b>5.2</b> (1.1)	<b>3.8</b> (3.4 / 0.4)	(0, 1, 1, 2, 2)	(0, 4, 2, 4, 3)	<b>1.4</b> (1.4 / 0)	(0, 3, 3, 1, 0)	<b>0</b>
Western North Atlantic fin whale ( <i>Balaenoptera physalus</i> )	<b>2.6</b> (6.5)	<b>0.8</b> (0.6 / 0.2)	(0, 0, 2, 0, 0)	(0, 1, 1, 0, 0)	<b>1.8</b> (1.4 / 0.4)	(5, 0, 2, 1, 1)	<b>0</b>
Nova Scotian sei whale ( <i>B. borealis</i> )	<b>1.2</b> (0.4)	<b>0.6</b> (0.4 / 0.2)	(0, 0, 0, 1, 0)	(0, 1, 0, 1, 0)	<b>0.6</b> (0.6 / 0)	(0, 1, 1, 0, 1)	<b>0</b>
Western North Atlantic blue whale <sup>2</sup> ( <i>B. musculus</i> )	<b>0</b> (-)	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>
Canadian East Coast minke whale ( <i>B. acutorostrata</i> )	<b>2.4</b> (69)	<b>2.0</b> (0.8 / 1.2)	(1, 1, 1, 4, 0)	(0, 0, 1, 0, 2)	<b>0.4</b> (0.4 / 0)	(1, 0, 0, 0, 1)	<b>0</b>
Western North Atlantic Bryde's whale ( <i>B. edeni</i> )	<b>0.2</b> (0.1)	<b>0</b>	<b>0</b>	<b>0</b>	<b>0.2</b> (0.2 / 0)	(0, 0, 0, 0, 1)	<b>0</b>

<sup>1</sup> Potential Biological Removal (PBR)

<sup>2</sup> Stock abundance estimates outdated; no PBR established for this stock.