

WHITE-SIDED DOLPHIN (*Lagenorhynchus acutus*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

White-sided dolphins are found in temperate and sub-polar waters of the North Atlantic, primarily in continental shelf waters to the 100m depth contour. The species inhabits waters from central West Greenland to North Carolina (about 35°N) and perhaps as far east as 43°W (Evans 1987; Hamazaki 2002). Distribution of sightings, strandings and incidental takes suggest the possible existence of three stock units: Gulf of Maine, Gulf of St. Lawrence and Labrador Sea stocks (Palka *et al.* 1997). Evidence for a separation between the population in the southern Gulf of Maine and the Gulf of St. Lawrence population comes from a virtual absence of summer sightings along the Atlantic side of Nova Scotia. This was reported in Gaskin (1992), is evident in Smithsonian stranding records, and was obvious during abundance surveys conducted in the summers of 1995 and 1999 which covered waters from Virginia to the Gulf of St. Lawrence. White-sided dolphins were seen frequently in Gulf of Maine waters and in waters at the mouth of the Gulf of St. Lawrence, but only a few sightings were recorded between these two regions.

The Gulf of Maine population of white-sided dolphins is most common in continental shelf waters from Hudson Canyon (approximately 39°N) on to Georges Bank, and in the Gulf of Maine and lower Bay of Fundy. Sightings data indicate seasonal shifts in distribution (Northridge *et al.* 1997). During January to May, low numbers of white-sided dolphins are found from Georges Bank to Jeffreys Ledge (off New Hampshire), with even lower numbers south of Georges Bank, as documented by a few strandings collected on beaches of Virginia and North Carolina. From June through September, large numbers of white-sided dolphins are found from Georges Bank to the lower Bay of Fundy. From October to December, white-sided dolphins occur at intermediate densities from southern Georges Bank to southern Gulf of Maine (Payne and Heinemann 1990). Sightings south of Georges Bank, particularly around Hudson Canyon, occur year round but at low densities. The Virginia and North Carolina observations appear to represent the southern extent of the species' range.

Prior to the 1970s, white-sided dolphins in U.S. waters were found primarily offshore on the continental slope, while white-beaked dolphins (*L. albirostris*) were found on the continental shelf. During the 1970s, there was an apparent switch in habitat use between these two species. This shift may have been a result of the decrease in herring and increase in sand lance in the continental shelf waters (Katona *et al.* 1993; Kenney *et al.* 1996).

POPULATION SIZE

The total number of white-sided dolphins along the eastern U.S. and Canadian Atlantic coast is unknown, although eight estimates from select regions are available from: 1) spring, summer and autumn 1978-1982; 2) July-September 1991-1992; 3) June-July 1993; 4) July-September 1995; 5) July-August 1999; 6) August 2002; 7) June-July 2004; and 8) August 2006. The best available current abundance estimate for white-sided dolphins in the western North Atlantic stock is 63,368 (CV=0.27), an average of the surveys conducted in August within the last 8 years (2002 and 2006). An average is used to account for the large inter-annual variability of the abundance estimates for this species. This variability may be associated with the water temperature and prey patterns.




Figure 1. Distribution of white-sided dolphin sightings from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1998, 1999, 2002, 2004 and 2006. Isobaths are the 100m, 1000m and 4000m depth contours.

Earlier abundance estimates

An abundance estimate of 28,600 white-sided dolphins (CV=0.21) was obtained from an aerial survey program conducted from 1978 to 1982 on the continental shelf and shelf edge waters between Cape Hatteras, North Carolina and Nova Scotia (CETAP 1982). An abundance estimate of 20,400 (CV=0.63) white-sided dolphins was obtained from two shipboard line transect surveys conducted during July to September 1991 and 1992 in the northern Gulf of Maine-lower Bay of Fundy region (Palka *et al.* 1997). An abundance estimate of 729 (CV=0.47) white-sided dolphins was obtained from a June and July 1993 shipboard line-transect sighting survey conducted principally between the 200 and 2,000 m isobaths from the southern edge of Georges Bank, across the Northeast Channel, to the southeastern edge of the Scotian Shelf (NMFS 1993). An abundance estimate of 27,200 (CV=0.43) white-sided dolphins was obtained from a July to September 1995 sighting survey conducted by two ships and an airplane that covered 32,600 km in waters from Virginia to the mouth of the Gulf of St. Lawrence. Kingsley and Reeves (1998) estimated that there were 11,740 (CV=0.47) white-sided dolphins in the Gulf of St. Lawrence during 1995 and 560 (CV=0.89) white-sided dolphins in the northern Gulf of St. Lawrence during 1996. It is assumed these estimates apply to the Gulf of St. Lawrence stock. During the 1995 survey, 8,427 km of track lines were flown in an area of 221,949 km² during August and September. During the 1996 survey, 3,993 km of track lines were flown in an area of 94,665 km² during July and August. As recommended in the GAMMS Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable and should not be used for PBR determinations.

Recent surveys and abundance estimates

An abundance estimate of 51,640 (CV=0.38) white-sided dolphins was obtained from a 28 July to 31 August 1999 line-transect sighting survey conducted from a ship and an airplane covering waters from Georges Bank to the mouth of the Gulf of St. Lawrence. Total track line length was 8,212 km. Shipboard data were analyzed using the modified direct duplicate method (Palka 1995) which accounts for school size bias and for $g(0)$, the probability of detecting a group on the track line. Aerial data were not corrected for $g(0)$ (Palka 2000). The 1999 estimate is larger than the 1995 estimate due to, at least in part, the fact that the 1999 survey covered the upper Bay of Fundy and the northern edge of Georges Bank for the first time and white-sided dolphins were seen in both areas.

An abundance estimate 109,141 (CV=0.30) white-sided dolphins was obtained from an aerial survey conducted in July and August 2002 which covered 7,465 km of trackline over waters from the 1000 m depth contour on the southern edge of Georges Bank to Maine (Table 1). The value of $g(0)$ used for this estimation was derived from the pooled data of 2002, 2004 and 2006 aerial survey data.

An abundance estimate of 2,330 (CV=0.80) white-sided dolphins was obtained from a line-transect sighting survey conducted during 12 June to 4 August 2004 by a ship and plane that surveyed 6,180 km of trackline from the 100m depth contour on the southern Georges Bank to the lower Bay of Fundy. The Scotian shelf south of Nova Scotia was not surveyed. (Table 1). Shipboard data were collected using the two independent team line transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and $g(0)$, the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line transect method (Hiby 1999) and analyzed accounting for $g(0)$ and biases due to school size and other potential covariates (Palka 2005).

An abundance estimate of 17,594 (CV=0.30) white-sided dolphins was generated from an aerial survey conducted in August 2006 which surveyed 10,676 km of trackline in the region from the 2000m depth contour on the southern edge of Georges Bank to the upper Bay of Fundy and to the entrance of the Gulf of St. Lawrence. (Table 1; Palka pers. comm.)

The average abundance estimate of white-sided dolphins from surveys conducted in August during the last 8 years (2002 and 2004) is 63,368 (CV=0.27). An average was used to incorporate the large inter-annual variability and thus provide an average number of white-sided dolphins that could be within the Gulf of Maine-western Scotian shelf region.

Table 1. Summary of recent abundance estimates for western North Atlantic stock of white-sided dolphins. Month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV).			
Month/Year	Area	N_{best}	CV
Jul-Aug 1999	Georges Bank to mouth of Gulf of St. Lawrence	51,640	0.38
Aug 2002	S. Gulf of Maine to Maine	109,141	0.30
Jun-Jul 2004	Gulf of Maine to lower Bay of Fundy	2,330	0.80
Aug 2006	S. Gulf of Maine to upper Bay of Fundy to Gulf of St. Lawrence	17,594	0.30

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for the western North Atlantic stock of white-sided dolphins is 63,368 (CV=0.27). The minimum population estimate for these white-sided dolphins is 50,883.

Current Population Trend

There are insufficient data to determine population trends for this species.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. Life history parameters that could be used to estimate net productivity include: calving interval is 2-3 years; lactation period is 18 months; gestation period is 10-12 months and births occur from May to early August, mainly in June and July; length at birth is 110cm; length at sexual maturity is 230-240 cm for males, and 201-222 cm for females; age at sexual maturity is 8-9 years for males and 6-8 years for females; mean adult length is 250 cm for males and 224 cm for females (Evans 1987); and maximum reported age for males is 22 years and for females, 27 years (Sergeant *et al.* 1980).

For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 50,883. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, threatened, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.5 because because the CV of the average mortality estimate is less than 0.3 (Wade and Angliss 1997). PBR for the western North Atlantic stock of white-sided dolphin is 509.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fishery Information

Detailed fishery information is reported in Appendix III

Earlier Interactions

NMFS observers in the Atlantic foreign mackerel fishery reported 44 takes of Atlantic white-sided dolphins incidental to fishing activities in the continental shelf and continental slope waters between March 1977 and December 1991 (Waring *et al.* 1990; NMFS unpublished data). Of these animals, 96% were taken in the Atlantic mackerel fishery. This total includes 9 documented takes by U.S. vessels involved in joint-venture fishing operations in which U.S. captains transfer their catches to foreign processing vessels. No incidental takes of white-sided dolphins were observed in the Atlantic mackerel JV fishery when it was observed in 1998.

During 1991 to 2005, two white-sided dolphins were observed taken in the Atlantic pelagic drift gillnet fishery, both in 1993. Estimated annual fishery-related mortality and serious injury (CV in parentheses) was 4.4 (.71) in 1989, 6.8 (.71) in 1990, 0.9 (.71) in 1991, 0.8 (.71) in 1992, 2.7 (0.17) in 1993 and 0 from 1994 to 2005. There was no fishery during 1997.

The mid-Atlantic gillnet fishery occurs year round from New York to North Carolina and has been observed since 1993. One white-sided dolphin was observed taken in this fishery during 1997. None were observed taken in other years. The estimated annual mortality (CV in parentheses) attributed to this fishery was 0 for 1993 to 1996, 45 (0.82) for 1997, 0 for 1998 to 2001, unknown in 2002 and 0 in 2003-2005.

U.S.

Northeast Sink Gillnet

This fishery occurs year round from in the Gulf of Maine, Georges Bank and in southern New England waters. Between 1990 and 2005 there were 54 white-sided dolphin mortalities observed in the Northeast sink gillnet fishery. Most were taken in waters south of Cape Ann during April to December. In recent years, the majority of the takes have been east and south of Cape Cod. During 2002, one of the takes was off Maine in the fall Mid-coast Closure Area in a pingered net. Estimated annual fishery-related mortalities (CV in parentheses) were 49 (0.46) in 1991, 154 (0.35) in 1992, 205 (0.31) in 1993, 240 (0.51) in 1994, 80 (1.16) in 1995, 114 (0.61) in 1996 (Bisack 1997), 140 (0.61) in 1997, 34 (0.92) in 1998, 69 (0.70) in 1999, 26 (1.00) in 2000, 26 (1.00) in 2001, 30 (0.74) in 2002, 31 (0.93) in 2003, 7 (0.98) in 2004, and 59 (0.49) in 2005. Average annual estimated fishery-related mortality during 2001-2005 was 31 white-sided dolphins per year (0.35) (Table 2).

Northeast Bottom Trawl

Forty-three mortalities were documented between 1991 and 2005 in the Northeast bottom trawl fishery; 1 during 1992, 0 in 1993, 2 in 1994, 0 in 1995-2001, 1 in 2002, 12 in 2003, 16 in 2004, and 12 in 2005. Estimated annual fishery-related mortalities (CV in parentheses) were 110 (0.97) in 1992, 0 in 1993, 182 (0.71) in 1994, 0 in 1995-1999, 137 (0.34) in 2000, 161 (0.34) in 2001, 70 (0.32) in 2002, 216 (0.27) in 2003, 200 (0.30) in 2004, and 213 (0.28) in 2005. The 2001-2005 average mortality attributed to the northeast bottom trawl was 192 animals (CV=0.13)(Table 2).

Northeast Atlantic (Gulf of Maine/Georges Bank) JV and TALFF Herring Fishery

A U.S. joint venture (JV) mid-water (pelagic) trawl fishery was conducted during 2001 on Georges Bank from August to December. No white-sided dolphins were incidentally captured. Two white-sided dolphins were incidentally captured in a single mid-water trawl during foreign fishing operations (TALFF) (Table 2). During TALFF fishing operations all nets fished by the foreign vessel are observed. The total mortality attributed to the Atlantic herring JV and TALFF mid-water trawl fisheries in 2001 was 2 animals (Table 2).

Northeast Mid-water Trawl Fishery (Including Pair Trawl)

The observer coverage in this fishery was highest after 2003, though a few trips in earlier years were observed (Table 2). A white-sided dolphin was observed taken in the single trawl fishery on the northern edge of Georges Bank (off of Massachusetts) during July 2003 in a haul that was targeting (and primarily caught) herring and 3 white-sided dolphins were taken in 2005 in paired trawls targeting herring. Due to small sample sizes, the bycatch rate model used the 2003 to September 2006 observed mid-water trawl data from paired and single northeast and mid-Atlantic mid-water trawls (Palka, pers. com.). The model that best fit these data was a Poisson logistic regression model that included latitude, bottom depth, and whether a kite panel was used on pair-trawl hauls as significant explanatory variables, and soak duration as the unit of effort. Estimated annual fishery-related mortalities (CV in parentheses) were unknown in 2001-2002, 24 (0.56) in 2003, 19 (0.58) in 2004, and 15(.68) in 2005 (Table 2; Palka pers. com.). The average annual estimated fishery-related mortality during 2001-2005 was 19 (0.35).

Mid-Atlantic Mid-water Trawl Fishery (Including Pair Trawl)

The observer coverage in this fishery was highest after 2003, though a few trips in other years were observed (Table 2). A white-sided dolphin was observed taken in the pair trawl fishery near Hudson Canyon (off New Jersey) during February 2004 in a haul that was targeting mackerel (and landed nothing). In 2005, 5 white-sided dolphins were taken in paired trawls targeting mackerel. Due to small sample sizes, the bycatch rate model used the 2003 to September 2006 observed mid-water trawl data, including paired and single, and Northeast and mid-Atlantic mid-water trawls (Palka, pers. com.). The model that best fit these data was a Poisson logistic regression model that included latitude, bottom depth, and whether a kite panel was used on pair-trawl hauls as significant explanatory variables, and soak duration as the unit of effort. Estimated annual fishery-related mortalities (CV in parentheses) were unknown in 2001-2002, 51 (0.46) in 2003, 105 (0.38) in 2004, and 97(.76) in 2005 (Table 2; Palka pers. com.).

The average annual estimated fishery-related mortality during 2001-2005 was 84 (0.34).

Mid-Atlantic Bottom Trawl Fishery

One white-sided dolphin incidental take was observed in 1997, resulting in a mortality estimate of 161 (CV=1.58) animals. No takes were observed from 1998 through 2004, and one take was observed in 2005. Recently observer coverage for this fishery was around 1%, except for 2004 where it was 3% (Table 2). Estimated annual fishery-related mortalities (CV in parentheses) were 27 (0.17) in 2000, 27 (0.19) in 2001, 25 (0.17) in 2002, 31 (0.25) in 2003, 26 (0.20) in 2004, and 38 (0.29) in 2005. The 2001-2005 average mortality attributed to the mid-Atlantic bottom trawl was 29 animals (CV=0.11).

Table 2. Summary of the incidental mortality of white-sided dolphins (*Lagenorhynchus acutus*) by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the mortalities recorded by on-board observers (Observed Mortality), the estimated annual mortality (Estimated Mortality), the estimated CV of the annual mortality (Estimated CVs) and the mean annual mortality (CV in parentheses).

Fishery	Years	Vessels	Data Type ^a	Observer Coverage ^b	Observed Mortality	Estimated Mortality	Estimated CVs	Mean Annual Mortality
Northeast Sink Gillnet ^d	01-05	1993=349 1998=301	Obs. Data Weighout Trip Logbook	.04, .02, .03, .06, .07	1, 1, 1, 1, 5	26, 30, 31, 7, 59	1.00, .74, .93, .98, .49	31 (0.35)
Northeast Bottom Trawl ^c	01-05	unk	Obs. Data Weighout	.01, .03, .04, .05, .12	0, 1, 12, 16, 47	161, 170, 216, 200, 213	.34, .32, .27, .30, .28	192 (0.13)
GOM/GB Herring Trawl-TALFF ^e	2001	2	Obs. Data	1.00	2	2	0	2 (0)
Northeast Mid-water Trawl - Including Pair Trawl	01-05	24, 27, 28, 22, 25	Obs. Data Weighout Trip Logbook	.001, 0, .031, .126, .199	0,0,1,0,3	unk, unk, 24, 19, 15	unk, unk, .56, .58, .68	19 (0.35)
Mid-Atlantic Mid-water Trawl - Including Pair Trawl	01-05	23, 20, 23, 25, 31	Obs. Data Weighout Trip Logbook	0, .003, .018, .064, .084	0,0,0,1,5	unk, unk, 51, 105, 97	unk, unk, .46, .38, .76	84 (0.34)
Mid-Atlantic Bottom Trawl ^c	01-05	unk	Obs. Data Weighout Trip Logbook	.01, .01, .01, .03, .03	0, 0, 0, 0, 1	27, 25, 31, 26, 38	.19, .17, .25, .20, .29	29 (.11)
Total								357(0.11)

a Observer data (Obs. Data), used to measure bycatch rates, are collected within the Northeast Observer Program. NEFSC collects landings data (Weighout) that are used as a measure of total effort in the Northeast gillnet fishery. Mandatory Vessel Trip Report (VTR) (Trip Logbook) data are used to determine the spatial distribution of fishing effort in the sink gillnet fishery and in the two mid-water trawl fisheries. In addition, the Trip Logbooks are the primary source of the measure of total effort (soak duration) in the two mid-water trawl fisheries.

b Observer coverages for the Northeast sink gillnet are ratios based on metric tons of fish landed. Observer coverages of the trawl fisheries are ratios based on trips.

c A new method was used to develop preliminary estimates of mortality for the mid-Atlantic and Northeast trawl fisheries during 2000-2005. They are a product of bycatch rates predicted by covariates in a model framework and effort reported by commercial fishermen on mandatory vessel logbooks. This method differs from the previous method used to estimate mortality in these fisheries prior to 2000. Therefore, the estimates reported prior to 2000 can not be compared to estimates during 2000-2005. In addition, the fisheries listed in Table 2 reflect new definitions defined by the proposed List of Fisheries for 2005 (FR Vol. 69, No. 231, 2004). The 'North Atlantic bottom trawl' fishery is now referred to as the 'Northeast bottom trawl'. The Illex, Loligo and Mackerel fisheries are now part of the mid-Atlantic and Northeast bottom trawl fisheries.

d After 1998, a weighted bycatch rate was applied to effort from both pingered and non-pingered hauls within the stratum where white-sided dolphins were observed taken. During the years 1997, 1999, 2001, 2002, and 2004, respectively, there were 2, 1, 1, 1, and 1 observed white-sided dolphins taken on pingered trips. No takes were observed on pinger trips during 1995, 1996, 1998, 2000, and 2005.

e There were two foreign vessels that harvested Atlantic herring in the U.S. fishery under a TALFF quota. During TALFF fishing operations all nets fished by the foreign vessel are observed.

CANADA

There is little information available that quantifies fishery interactions involving white-sided dolphins in Canadian waters. Two white-sided dolphins were reported caught in groundfish gillnet sets in the Bay of Fundy during 1985 to 1989, and 9 were reported taken in West Greenland between 1964 and 1966 in the now non-operational salmon drift nets (Gaskin 1992). Several (number not specified) were also taken during the 1960's in the now non-operational Newfoundland and Labrador groundfish gillnets. A few (number not specified) were taken in an experimental drift gillnet fishery for salmon off West Greenland which took place from 1965 to 1982 (Read 1994).

Hooker *et al.* (1997) summarized bycatch data from a Canadian fisheries observer program that placed observers on all foreign fishing vessels operating in Canadian waters, on between 25-40% of large Canadian fishing vessels (greater than 100 feet long), and on approximately 5% of smaller Canadian fishing vessels. Bycaught marine mammals were noted as weight in kilos rather than by the numbers of animals caught. Thus the number of individuals was estimated by dividing the total weight per species per trip by the maximum recorded weight of each species. During 1991 through 1996, an estimated 6 white-sided dolphins were observed taken. One animal was from a longline trip south of the Grand Banks (43° 10'N 53° 08'W) in November 1996 and the other 5 were taken in the bottom trawl fishery off Nova Scotia in the Atlantic Ocean; 1 in July 1991, 1 in April 1992, 1 in May 1992, 1 in April 1993, 1 in June 1993 and 0 in 1994 to 1996.

Estimation of small cetacean bycatch is currently underway for Newfoundland fisheries using data collected during 2001 to 2003 (pers. comm. J. Lawson, DFO). White-sided dolphins were reported to have been caught in the Newfoundland nearshore gillnet fishery and offshore monkfish/skate gillnet fisheries.

Herring Weirs

During the last several years, one white-sided dolphin was released alive and unharmed from a herring weir in the Bay of Fundy (A. Westgate, pers. comm.). Due to the formation of a cooperative program between Canadian fishermen and biologists, it is expected that most dolphins and whales will be able to be released alive. Fishery information is available in Appendix III.

Other Mortality

U.S.

Mass strandings involving up to a hundred or more animals at one time are common for this species. From 1968 to 1995, 349 Atlantic white-sided dolphins were known to have stranded on the New England coast (Hain and Waring 1994; Smithsonian stranding records 1996). The causes of these strandings are not known. Because such strandings have been known since antiquity, it could be presumed that recent strandings are a normal condition (Gaskin 1992). It is unknown whether human causes, such as fishery interactions and pollution, have increased the number of strandings. Stranding data probably underestimate the extent of fishery-related mortality and serious injury because all of the marine mammals that die or are seriously injured may not wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interaction.

White-sided dolphin stranding records from 1997 that are in the NMFS/NE Regional Office strandings and entanglement database have been reviewed and updated. The most recent five years are reported in Table 3. Cause of death was investigated and it was determined that the documented human interactions were as follows: 1 animal possibly killed by a boat collision off Maine during 2001; 2 animals with indications of fishery interactions found in March 2002 in Massachusetts; and 1 animal with indications of fishery interactions found in May 2002 in Virginia, 1 animal with indications of fishery interactions was found in Massachusetts during 2004, and one animal during 2004 was found with twine blocking its esophagus (thus, this is a human interaction, but not necessarily a fishery interaction). In 2005 5 animals had signs of human interaction but in no case was the human interaction able to be determined to be the cause of death. (Table 3).

Mass strandings in Massachusetts occur frequently (Table 3). There were 80 animals in a mass stranding near Wellfleet, Massachusetts, during the week of 29 January to 3 February 1998. Of these, 2 were released alive. Of the 4 found in Massachusetts during the November 1998 mass stranding, 1 was released alive. Fifty-three animals stranded in Wellfleet, Massachusetts during 19-24 March 1999. During 1999, of the 70 strandings, 38 were found alive, and 3 of these animals were released alive. During 2000, 5 were found alive (3 in April and 2 in August), and the 2 in August were released alive. During 2002, there were mass strandings in March and August, of which a few were released alive. During 2003 in Massachusetts 36 white-sided dolphins were involved in mass strandings in

January, April and November, of which 25 were found alive. There were no mass strandings in 2004. In 2005 there were mass strandings in February, April, May and January. A total of 26 white-sided dolphins were involved in mass strandings, of which 11 were successfully released.

CANADA

Small numbers of white-sided dolphins have been taken off southwestern Greenland and they have been taken deliberately by shooting elsewhere in Canada (Reeves *et al.* 1999). The Nova Scotia Stranding Network documented whales and dolphins stranded on the coast of Nova Scotia during 1991 to 1996 (Hooker *et al.* 1997). Researchers with Dept. of Fisheries and Oceans (DFO), Canada documented strandings on the beaches of Sable Island during 1970 to 1998 (Lucas and Hooker 2000). Sable Island is approximately 170km southeast of mainland Nova Scotia. White-sided dolphins stranded at nearly all times of the year on the mainland and on Sable Island. On the mainland of Nova Scotia, a total of 34 stranded white-sided dolphins was recorded between 1991 and 1996: 2 in 1991 (August and October), 26 in July 1992, 1 in Nov 1993, 2 in 1994 (February and November), 2 in 1995 (April and August) and 2 in 1996 (October and December). During July 1992, 26 white-sided dolphins stranded on the Atlantic side of Cape Breton. Of these, 11 were released alive and the rest were found dead. Among the rest of the Nova Scotia strandings, 1 was found in Minas Basin, 2 near Yarmouth and the rest near Halifax. On Sable Island, 10 stranded white-sided dolphins were documented between 1991 and 1998; all were males, 7 were young males (< 200cm), 1 in January 1993, 5 in March 1993, 1 in August 1995, 1 in December 1996, 1 in April 1997 and 1 in February 1998.

Whales and dolphins stranded between 1997 and 2005 on the coast of Nova Scotia as recorded by the Marine Animal Response Society (MARS) and the Nova Scotia Stranding Network are as follows (Table 3): 0 white-sided dolphins stranded in 1997 to 2000, 3 in September 2001 (released alive), 5 in November 2002 (4 were released alive), 0 in 2003, 19-24 in 2004 (15-20 in October (some (unspecified) were released alive) and 4 in November were released alive), and 0 in 2005.

Area	Year					Total
	2001	2002	2003	2004	2005	
Maine ^b	2	4	2	10	3	21
New Hampshire					1	1
Massachusetts ^{a,b}	16	53	59	34	60	222
Rhode Island		2			2	4
Connecticut			1			1
New York		1	2	1		4
New Jersey		1	1	1	6	9
Delaware						0
Maryland					1	1
Virginia ^b		1		4	3	8
North Carolina			1	2	3	6
TOTAL US	18	62	66	52	79	277
Nova Scotia	3	6		2		8
GRAND TOTAL	21	68	66	54	79	285

^a Records of mass strandings in Massachusetts are: March 1999 - 53 animals; April 2000 - 5 animals; August 2000 - 11 animals; April 2001 - 6 animals; March 2002 - 31 animals, of which 7 were released alive; August 2002 - 3 animals, of which 1 was released alive; January 2003 - 4 animals; April 2003 - 28 animals; November 2003 - 4 animals; February 2005 - 8 animals (3 released alive), April 2005 - 6 animals (all released alive), May 2005 strandings of 2 animals (both released alive but one died later), 3 animals (one released alive), and 5 animals, and December 2005 - 2 animals.

^b Strandings that appear to involve a human interaction are: 1 animal from Maine in 2001 that was a possible boat collision; 1 animal from Virginia in May 2002 had signs of fishery interaction; 2 animals from Massachusetts in March 2002 had signs of fishery interactions; 1 animal from Massachusetts in 2004 was a fishery interaction; and 1 other animal from Massachusetts in 2004 was found with twine obstructing its esophagus. In 2005 5 animals had signs of human interaction but in no case was the human interaction able to be determined to be the cause of death.

STATUS OF STOCK

The status of white-sided dolphins, relative to OSP, in the U.S. Atlantic EEZ is unknown. The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine population trends for this species. The total U.S. fishery-related mortality and serious injury for this stock is not less than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. This is a non-strategic stock because the 2001-2005 estimated average annual human related mortality does not exceed PBR.

REFERENCES CITED

- Barlow, J., S.L. Swartz, T.C. Eagle, and P.R. Wade. 1995. U.S. Marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73 pp.
- Bisack, K.D. 1997. Harbor porpoise bycatch estimates in the New England multispecies sink gillnet fishery: 1994 and 1995. Rep. int Whal. Comm 47:705-14.
- CETAP. 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf. Cetacean and Turtle Assessment Program, University of Rhode Island. Final Report, Contract AA51-C78-48, Bureau of Land Management, Washington, DC, 538 pp.
- Evans, P.G.H. 1987. The natural history of whales and dolphins. Facts on File Publications, New York, 343 pp.
- Gaskin, D.E. 1992. Status of Atlantic white-sided dolphin, *Lagenorhynchus acutus*, in Canada. Can.Fld. Nat. 106: 64-72.
- Hain, J. H. W. and G. T. Waring. 1994. Status of and human effects upon marine mammals. Pages. 9-20 *In*: R. W. Langton, J. B. Pearce, and J.A. Gibson (eds). Selected living resources, habitat conditions, and human perturbations of the Gulf of Maine: Environmental and ecological considerations for fishery managements. NOAA Tech. Mem. NMFS-NE-106.
- Hamazaki, T. 2002. Spatiotemporal prediction models of cetacean habitats in the mid-western North Atlantic Ocean (from Cape Hatteras, North Carolina, USA to Nova Scotia, Canada). Mar. Mammal Sci. 18(4):920-939.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. Pages 179-189 *in*: G. W. Garner, S. C. Amstrup, J. L. Laake, B. F. J. Manly, L. L. McDopnald, and D. G. Robertson. (eds). Marine Mammal Survey and Assessment Methods. Balkema, Rotterdam.
- Hooker, S.K., R.W. Baird, and M.A. Showell. 1997. Cetacean strandings and bycatches in Nova Scotia, Eastern Canada, 1991-1996. Meeting document SC/49/O5 submitted to the 1997 International Whaling Commission meeting in Bournemouth, UK.
- Katona, S.K., V. Rough, and D.T. Richardson. 1993. A field guide to whales, porpoises, and seals from Cape Cod to Newfoundland. Smithsonian Institution Press. Washington, DC. 316 pp.
- Kenney, R.D., P.M. Payne, D.W. Heinemann, and H.E. Winn. 1996. Shifts in Northeast shelf cetacean distributions relative to trends in Gulf of Maine/Georges Bank finfish abundance. Pp. 169-196 *in*: K. Sherman, N.A. Jaworski and T. Smada (eds.). The northeast shelf ecosystem: assessment, sustainability, and management. Blackwell Science, Cambridge, MA 02142, USA.
- Kingsley, M.C.S. and R.R. Reeves. 1998. Aerial surveys of cetaceans in the Gulf of St. Lawrence in 1995 and 1996. Can. J. Zool. 76:1529-1550.
- Lucas, Z.N. and S.K. Hooker. 2000. Cetacean strandings on Sable Island, Nova Scotia, 1970-1998. Can. Fld. Nat. 114(1):46-61.
- National Marine Fisheries Service [NMFS]. 1993. Cruise results, NOAA ship DELAWARE II, Cruise No. DEL 93-06, Marine Mammal Survey. 5 pp. [Available from: NMFS, Northeast Fisheries Science Center, 166 Water Street, Woods Hole, MA 02543.]

- Northridge, S., M. Tasker, A. Webb, K. Camphuysen, and M. Leopold. 1997. White-beaked *Lagenorhynchus albirostris* and Atlantic white-sided dolphin *L. acutus* distributions in northwest European and U.S. North Atlantic waters. Rep. int. Whal. Commn 47:797-805.
- Palka, D. 2005. Aerial surveys in the northwest Atlantic: estimation of $g(0)$. In Proceedings of the workshop on Estimation of $g(0)$ in line-transect surveys of cetaceans, ed. F. Thomsen, F. Ugarte, and P.G.H. Evans. ECS Newsletter No. 44 – Special Issue. April 2005. Pgs 12-7
- Palka, D. 2000. Abundance of the Gulf of Maine/Bay of Fundy harbor porpoise based on shipboard and aerial surveys during 1999. NOAA/NMFS/NEFSC Ref. Doc. 00-07. 29 pp.
- Palka, D. and Hammond, P.S. 2001. Accounting for responsive movement in line transect estimates of abundance. Can. J. Fish. Aquat. Sci. 58:777-787.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. pp. 27-50 In: A. Bjørge and G.P. Donovan (eds.) Biology of the Phocoenids. Rep. int Whal. Commn (Special Issue) 16.
- Palka, D., A. Read, and C. Potter. 1997. Summary of knowledge of white-sided dolphins (*Lagenorhynchus acutus*) from the U.S. and Canadian North Atlantic waters. Rep. int Whal. Commn 47:729-34.
- Payne, M. and D.W. Heinemann. 1990. A distributional assessment of cetaceans in the shelf and shelf edge waters of the northeastern United States based on aerial and shipboard surveys, 1978-1988. Report to NMFS. 108p. [Available from National Marine Fisheries Science Center, 166 Water St., Woods Hole, MA 02543.]
- Read, A. J. 1994. Interactions between cetaceans and gillnet and trap fisheries in the northwest Atlantic. Rep. int Whal. Commn (Special Issue) 15: 133-147.
- Reeves, R. R., C. Smeenk, R. L. Brownell, Jr., and C. C. Kinze. 1999. Atlantic white-sided dolphin *Lagenorhynchus acutus* (Gray, 1828). Pp. 31-56 In S. H. Ridgway and Sir R. Harrison (eds.) Handbook of Marine Mammals. Volume 6: The Second Book of Dolphins and the Porpoises. Academic Press, Boston.
- Sergeant, D.E., D.J. St. Aubin, and J.R. Geraci. 1980. Life history and northwest Atlantic status of the Atlantic white-sided dolphin, *Lagenorhynchus acutus*. Cetology 37:1-12.
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Waring, G.T., P. Gerrior, P.M. Payne, B.L. Parry and J.R. Nicolas. 1990. Incidental take of marine mammals in foreign fishery activities off the northeast United States, 1977-1988. Fish. Bull., U.S. 88(2):347-360.