

**Table 1.98A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	8,402	7,949	1,554	1,406	567	604
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1,341	1,373	250	220	89	69
New England	364	408	94	41	22	14
Middle Atlantic	976	964	155	178	67	55
Midwest	1,688	1,643	467	336	186	115
East North Central	1,078	1,071	277	219	115	97
West North Central	611	572	190	117	71 <sup>b</sup>	19
South	2,835	2,779	517	639	178	318
South Atlantic	1,680	1,583	307	322	114	184
East South Central	462	346	106	132	47	59
West South Central	693	850	104	184	17	75
West	2,538	2,153	321	211	114	102
Mountain	640	707	106	102	38	47
Pacific	1,898 <sup>a</sup>	1,446	215	110	76	55
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	4,721	4,445	868	769	388	352
Small Metro	2,643	2,328	483	408	123	158
250K – 1 Mil. Pop.	1,682	1,495	306	232	84	90
< 250K Pop.	961	833	177	176	38	67
Nonmetro	1,038	1,175	203	229	57	94
Urbanized	398	552	65	85	28	21
Less Urbanized	537	536	112	127	27	65
Completely Rural	102	87	26	17	2	8

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.98B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	3.6	3.3	0.7	0.6	0.2	0.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	3.0	3.0	0.6	0.5	0.2	0.2
New England	3.1	3.4	0.8	0.3	0.2	0.1
Middle Atlantic	2.9	2.9	0.5	0.5	0.2	0.2
Midwest	3.2	3.1	0.9	0.6	0.3	0.2
East North Central	2.9	2.8	0.7	0.6	0.3	0.3
West North Central	3.8	3.6	1.2	0.7	0.4 <sup>b</sup>	0.1
South	3.4	3.3	0.6	0.8	0.2	0.4
South Atlantic	3.8	3.6	0.7	0.7	0.3	0.4
East South Central	3.3	2.4	0.8	0.9	0.3	0.4
West South Central	2.7	3.3	0.4	0.7	0.1	0.3
West	4.8	4.0	0.6	0.4	0.2	0.2
Mountain	4.2	4.5	0.7	0.7	0.3	0.3
Pacific	5.1 <sup>a</sup>	3.8	0.6	0.3	0.2	0.1
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	3.8	3.5	0.7	0.6	0.3	0.3
Small Metro	3.8	3.3	0.7	0.6	0.2	0.2
250K – 1 Mil. Pop.	3.7	3.2	0.7	0.5	0.2	0.2
< 250K Pop.	4.0	3.4	0.7	0.7	0.2	0.3
Nonmetro	2.6	2.9	0.5	0.6	0.1	0.2
Urbanized	2.5	3.1	0.4	0.5	0.2	0.1
Less Urbanized	2.7	2.9	0.6	0.7	0.1	0.4
Completely Rural	2.3	2.1	0.6	0.4	0.0	0.2

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.99A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	163	157	100	95	26	32
<b>GEOGRAPHIC DIVISION</b>						
Northeast	20	16	13	10	4	1
New England	10 <sup>a</sup>	3	5	1	3	1
Middle Atlantic	10	14	8	9	1	1
Midwest	26	42	19	27	1 <sup>b</sup>	14
East North Central	15	23	9	14	0 <sup>a</sup>	5
West North Central	11	19	10	13	0 <sup>a</sup>	9
South	60	53	42	34	18	12
South Atlantic	33	22	23	18	12	6
East South Central	10	3	5	2	3	1
West South Central	18	28	14	14	2	5
West	57	45	26	25	4	5
Mountain	21	19	13	15	1	5
Pacific	36	26	13	10	3	*
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	51 <sup>a</sup>	83	30	51	2 <sup>b</sup>	21
Small Metro	59	36	34	21	10	7
250K – 1 Mil. Pop.	37	19	21	11	5	4
< 250K Pop.	21	17	13	11	5	3
Nonmetro	54	37	36	23	14	5
Urbanized	16	12	13	7	8	2
Less Urbanized	33	18	19	12	6	2
Completely Rural	5	8	5	3	*	0

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.99B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.7	0.6	0.4	0.4	0.1	0.1
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.4	0.4	0.3	0.2	0.1	0.0
New England	0.8 <sup>a</sup>	0.2	0.4	0.1	0.3	0.0
Middle Atlantic	0.3	0.4	0.2	0.3	0.0	0.0
Midwest	0.5	0.7	0.3	0.5	0.0 <sup>b</sup>	0.2
East North Central	0.4	0.6	0.2	0.3	0.0 <sup>a</sup>	0.1
West North Central	0.7	1.1	0.6	0.8	0.0 <sup>a</sup>	0.5
South	0.7	0.6	0.5	0.4	0.2	0.1
South Atlantic	0.7	0.5	0.5	0.4	0.3	0.1
East South Central	0.7	0.2	0.4	0.1	0.2	0.1
West South Central	0.6	0.9	0.5	0.5	0.1	0.2
West	1.0	0.8	0.4	0.4	0.1	0.1
Mountain	1.2	1.1	0.8	0.8	0.0	0.3
Pacific	0.9	0.6	0.3	0.2	0.1	*
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.4 <sup>a</sup>	0.6	0.2	0.4	0.0 <sup>b</sup>	0.2
Small Metro	0.8 <sup>a</sup>	0.5	0.5	0.3	0.1	0.1
250K – 1 Mil. Pop.	0.8	0.4	0.4	0.2	0.1	0.1
< 250K Pop.	0.9	0.7	0.6	0.4	0.2	0.1
Nonmetro	1.3	0.9	0.8	0.5	0.3	0.1
Urbanized	0.9	0.6	0.7	0.4	0.5	0.1
Less Urbanized	1.6	0.9	0.9	0.6	0.3	0.1
Completely Rural	1.1	1.9	1.0	0.7	*	0.0

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.100A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	1,182	1,193	266	275	52	55
<b>GEOGRAPHIC DIVISION</b>						
Northeast	128	157	31	42	8	6
New England	48	44	15	14	5	2
Middle Atlantic	81	114	16	28	3	5
Midwest	268	282	66	78	16	13
East North Central	145	175	29 <sup>a</sup>	49	10	8
West North Central	123	106	37	29	6	5
South	373	348	99	81	18	10
South Atlantic	177	153	54	36	10	7
East South Central	76	70	12	16	3	2
West South Central	121	126	33	29	5	1
West	412	406	70	73	10	25
Mountain	168 <sup>a</sup>	124	31	28	5	9
Pacific	244	282	40	46	5	16
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	557	565	119	118	27	29
Small Metro	436	424	92	105	17	19
250K – 1 Mil. Pop.	257	278	52	68	11	11
< 250K Pop.	179	146	40	37	6	8
Nonmetro	188	204	55	51	8	6
Urbanized	62	97	16	26	1	4
Less Urbanized	101	93	23	20	6	2
Completely Rural	24	14	15	5	*	*

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.100B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	3.8	3.8	0.9	0.9	0.2	0.2
<b>GEOGRAPHIC DIVISION</b>						
Northeast	2.3	2.8	0.6	0.7	0.2	0.1
New England	3.3	3.0	1.0	1.0	0.4	0.1
Middle Atlantic	2.0	2.7	0.4	0.7	0.1	0.1
Midwest	3.7	3.8	0.9	1.1	0.2	0.2
East North Central	2.9	3.4	0.6	1.0	0.2	0.2
West North Central	5.5	4.7	1.6	1.3	0.3	0.2
South	3.4	3.1	0.9	0.7	0.2	0.1
South Atlantic	3.3	2.7	1.0	0.6	0.2	0.1
East South Central	4.0	3.6	0.7	0.8	0.2	0.1
West South Central	3.3	3.3	0.9	0.8	0.1	0.0
West	5.6	5.5	1.0	1.0	0.1	0.3
Mountain	7.7 <sup>a</sup>	5.5	1.4	1.2	0.2	0.4
Pacific	4.8	5.5	0.8	0.9	0.1	0.3
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	3.5	3.4	0.8	0.7	0.2	0.2
Small Metro	4.3	4.2	0.9	1.0	0.2	0.2
250K – 1 Mil. Pop.	4.0	4.2	0.8	1.0	0.2	0.2
< 250K Pop.	4.8	4.1	1.1	1.0	0.2	0.2
Nonmetro	3.8	4.1	1.1	1.0	0.2	0.1
Urbanized	2.8	4.1	0.7	1.1	0.0	0.2
Less Urbanized	4.5	4.3	1.0	0.9	0.2	0.1
Completely Rural	5.2	3.9	3.3	1.5	*	*

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.101A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	7,057	6,599	1,188	1,036	488	517
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1,192	1,199	206	168	76	61
New England	307	362	75	26	13	12
Middle Atlantic	885	837	132	142	63	49
Midwest	1,395	1,320	381	231	169 <sup>a</sup>	88
East North Central	918	873	238	156	105	83
West North Central	476	447	143	75	65 <sup>b</sup>	5
South	2,402	2,378	376	524	143	296
South Atlantic	1,471	1,408	230	268	92	171
East South Central	377	273	89	114	42	55
West South Central	554	697	56	142	9 <sup>a</sup>	70
West	2,069	1,702	225	113	100	72
Mountain	451	564	62	60	33	33
Pacific	1,617 <sup>a</sup>	1,139	163	54	67	39
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	4,113	3,797	719	599	359	302
Small Metro	2,148	1,868	357	281	96	132
250K – 1 Mil. Pop.	1,387	1,198	233	153	69	76
< 250K Pop.	761	670	124	128	27	56
Nonmetro	796	934	112	155	34	83
Urbanized	320	444	36	52	19	14
Less Urbanized	403	426	70	95	15	61
Completely Rural	73	64	6	8	*	8

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.101B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	3.9	3.6	0.7	0.6	0.3	0.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	3.4	3.4	0.6	0.5	0.2	0.2
New England	3.3	3.9	0.8	0.3	0.1	0.1
Middle Atlantic	3.4	3.2	0.5	0.5	0.2	0.2
Midwest	3.4	3.2	0.9 <sup>a</sup>	0.6	0.4 <sup>a</sup>	0.2
East North Central	3.2	3.1	0.8	0.5	0.4	0.3
West North Central	3.9	3.7	1.2	0.6	0.5 <sup>b</sup>	0.0
South	3.8	3.7	0.6	0.8	0.2	0.5
South Atlantic	4.3	4.1	0.7	0.8	0.3	0.5
East South Central	3.5	2.5	0.8	1.1	0.4	0.5
West South Central	2.9	3.6	0.3	0.7	0.0 <sup>a</sup>	0.4
West	5.2	4.2	0.6	0.3	0.3	0.2
Mountain	3.9	4.8	0.5	0.5	0.3	0.3
Pacific	5.7 <sup>a</sup>	3.9	0.6	0.2	0.2	0.1
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	4.3	3.9	0.7	0.6	0.4	0.3
Small Metro	4.1	3.5	0.7	0.5	0.2	0.2
250K – 1 Mil. Pop.	4.1	3.4	0.7	0.4	0.2	0.2
< 250K Pop.	4.3	3.6	0.7	0.7	0.2	0.3
Nonmetro	2.6	3.0	0.4	0.5	0.1	0.3
Urbanized	2.7	3.3	0.3	0.4	0.2	0.1
Less Urbanized	2.6	3.0	0.5	0.7	0.1	0.4
Completely Rural	2.1	1.9	0.2	0.3	*	0.3

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.102A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	8,239	7,792	1,454	1,310	541	572
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1,321	1,356	237	210	85	68
New England	355	406	89	40	19	14
Middle Atlantic	966	950	148	170	66	54
Midwest	1,663	1,601	447	309	185 <sup>a</sup>	101
East North Central	1,063	1,048	268	205	115	91
West North Central	600	553	180	104	70 <sup>b</sup>	10
South	2,775	2,726	475	605	161	306
South Atlantic	1,647	1,561	284	304	102	178
East South Central	453	343	101	130	45	58
West South Central	675	822	90	170	14	70
West	2,481	2,108	295	187	110	97
Mountain	619	688	93	87	38	42
Pacific	1,862 <sup>a</sup>	1,421	202	100	73	55
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	4,670	4,362	838	718	386	331
Small Metro	2,585	2,292	449	386	113	151
250K – 1 Mil. Pop.	1,645	1,476	285	221	79	87
< 250K Pop.	940	816	164	165	33	64
Nonmetro	984	1,138	167	206	42	90
Urbanized	382 <sup>a</sup>	541	52	78	20	19
Less Urbanized	505	519	93	115	20	63
Completely Rural	97	78	22	14	2	8

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.102B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	3.9	3.7	0.7	0.6	0.3	0.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	3.3	3.3	0.6	0.5	0.2	0.2
New England	3.3	3.8	0.8	0.4	0.2	0.1
Middle Atlantic	3.2	3.2	0.5	0.6	0.2	0.2
Midwest	3.5	3.3	0.9	0.6	0.4 <sup>a</sup>	0.2
East North Central	3.2	3.1	0.8	0.6	0.3	0.3
West North Central	4.2	3.8	1.3	0.7	0.5 <sup>b</sup>	0.1
South	3.7	3.6	0.6	0.8	0.2	0.4
South Atlantic	4.2	3.9	0.7	0.8	0.3	0.4
East South Central	3.6	2.7	0.8	1.0	0.4	0.5
West South Central	2.9	3.6	0.4	0.7	0.1	0.3
West	5.3	4.4	0.6	0.4	0.2	0.2
Mountain	4.5	5.0	0.7	0.6	0.3	0.3
Pacific	5.6 <sup>a</sup>	4.2	0.6	0.3	0.2	0.2
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	4.1	3.9	0.7	0.6	0.3	0.3
Small Metro	4.2	3.6	0.7	0.6	0.2	0.2
250K – 1 Mil. Pop.	4.1	3.5	0.7	0.5	0.2	0.2
< 250K Pop.	4.4	3.7	0.8	0.7	0.2	0.3
Nonmetro	2.7	3.2	0.5	0.6	0.1	0.2
Urbanized	2.7	3.4	0.4	0.5	0.1	0.1
Less Urbanized	2.9	3.2	0.5	0.7	0.1	0.4
Completely Rural	2.5	2.1	0.5	0.4	0.1	0.2

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.