

Table 1.118C Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 569 | 517 | 300 | 299 | 186 | 207 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 240 | 214 | 112 | 99 | 61 | 71 |
| New England | 141 | 130 | 66 | 65 | 26 | 46 |
| Middle Atlantic | 190 | 173 | 90 | 76 | 54 | 54 |
| Midwest | 208 | 195 | 103 | 110 | 60 | 67 |
| East North Central | 172 | 152 | 86 | 96 | 48 | 60 |
| West North Central | 117 | 122 | 56 | 53 | 36 | 29 |
| South | 351 | 301 | 191 | 166 | 134 | 125 |
| South Atlantic | 266 | 211 | 148 | 122 | 98 | 96 |
| East South Central | 140 | 133 | 79 | 57 | 60 | 43 |
| West South Central | 178 | 167 | 91 | 96 | 68 | 67 |
| West | 312 | 303 | 176 | 199 | 98 | 133 |
| Mountain | 107 | 120 | 55 | 76 | 32 | 55 |
| Pacific | 293 | 277 | 166 | 184 | 92 | 121 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 447 | 406 | 226 | 244 | 140 | 163 |
| Small Metro | 303 | 261 | 164 | 154 | 101 | 104 |
| 250K – 1 Mil. Pop. | 225 | 223 | 125 | 128 | 82 | 77 |
| < 250K Pop. | 191 | 151 | 107 | 89 | 62 | 70 |
| Nonmetro | 206 | 178 | 99 | 105 | 60 | 69 |
| Urbanized | 132 | 121 | 67 | 69 | 35 | 48 |
| Less Urbanized | 143 | 108 | 58 | 73 | 27 | 49 |
| Completely Rural | 80 | 56 | 48 | 22 | 46 | 8 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.118D Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.24 | 0.22 | 0.13 | 0.13 | 0.08 | 0.09 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.53 | 0.47 | 0.25 | 0.22 | 0.13 | 0.16 |
| New England | 1.20 | 1.09 | 0.56 | 0.54 | 0.22 | 0.39 |
| Middle Atlantic | 0.57 | 0.52 | 0.27 | 0.23 | 0.16 | 0.16 |
| Midwest | 0.39 | 0.36 | 0.19 | 0.20 | 0.11 | 0.12 |
| East North Central | 0.46 | 0.40 | 0.23 | 0.25 | 0.13 | 0.16 |
| West North Central | 0.73 | 0.76 | 0.35 | 0.33 | 0.22 | 0.18 |
| South | 0.42 | 0.36 | 0.23 | 0.20 | 0.16 | 0.15 |
| South Atlantic | 0.61 | 0.48 | 0.34 | 0.28 | 0.22 | 0.22 |
| East South Central | 0.99 | 0.93 | 0.56 | 0.40 | 0.43 | 0.30 |
| West South Central | 0.69 | 0.64 | 0.35 | 0.37 | 0.26 | 0.26 |
| West | 0.59 | 0.56 | 0.33 | 0.37 | 0.19 | 0.25 |
| Mountain | 0.70 | 0.77 | 0.36 | 0.49 | 0.21 | 0.35 |
| Pacific | 0.78 | 0.73 | 0.44 | 0.48 | 0.25 | 0.32 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.36 | 0.32 | 0.18 | 0.19 | 0.11 | 0.13 |
| Small Metro | 0.44 | 0.37 | 0.24 | 0.22 | 0.15 | 0.15 |
| 250K – 1 Mil. Pop. | 0.49 | 0.48 | 0.28 | 0.27 | 0.18 | 0.16 |
| < 250K Pop. | 0.80 | 0.62 | 0.45 | 0.36 | 0.26 | 0.29 |
| Nonmetro | 0.51 | 0.44 | 0.25 | 0.26 | 0.15 | 0.17 |
| Urbanized | 0.82 | 0.68 | 0.42 | 0.39 | 0.22 | 0.27 |
| Less Urbanized | 0.73 | 0.59 | 0.30 | 0.40 | 0.14 | 0.27 |
| Completely Rural | 1.79 | 1.37 | 1.09 | 0.55 | 1.04 | 0.20 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.118P Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.0894 | | 0.1726 | | 0.3328 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.4067 | | 0.3624 | | 0.9240 | |
| New England | 0.5095 | | 0.1069 | | 0.3968 | |
| Middle Atlantic | 0.5916 | | 0.9837 | | 0.6611 | |
| Midwest | 0.1547 | | 0.5798 | | 0.1645 | |
| East North Central | 0.2285 | | 0.6506 | | 0.3601 | |
| West North Central | 0.4339 | | 0.0813 | | 0.2666 | |
| South | 0.7761 | | 0.8310 | | 0.6715 | |
| South Atlantic | 0.9967 | | 0.9890 | | 0.8272 | |
| East South Central | 0.8954 | | 0.1007 | | 0.1167 | |
| West South Central | 0.5180 | | 0.3528 | | 0.4435 | |
| West | 0.2642 | | 0.1355 | | 0.2277 | |
| Mountain | 0.1665 | | 0.2077 | | 0.8743 | |
| Pacific | 0.5330 | | 0.2691 | | 0.1706 | |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.1721 | | 0.1736 | | 0.7167 | |
| Small Metro | 0.6527 | | 0.7673 | | 0.6028 | |
| 250K – 1 Mil. Pop. | 0.6236 | | 0.9568 | | 0.5588 | |
| < 250K Pop. | 0.9278 | | 0.6843 | | 0.1591 | |
| Nonmetro | 0.3201 | | 0.1918 | | 0.2299 | |
| Urbanized | 0.9616 | | 0.3932 | | 0.2127 | |
| Less Urbanized | 0.1602 | | 0.0524 | | 0.0477 | |
| Completely Rural | 0.9420 | | 0.0956 | | 0.0387 | |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.119C Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 69 | 70 | 59 | 59 | 39 | 39 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 24 | 30 | 19 | 23 | 13 | 14 |
| New England | 11 | 17 | 8 | 14 | 6 | 7 |
| Middle Atlantic | 20 | 25 | 18 | 18 | 11 | 13 |
| Midwest | 27 | 28 | 23 | 25 | 14 | 15 |
| East North Central | 23 | 24 | 19 | 21 | 12 | 13 |
| West North Central | 14 | 15 | 13 | 13 | 7 | 7 |
| South | 41 | 44 | 38 | 36 | 26 | 24 |
| South Atlantic | 29 | 32 | 29 | 27 | 20 | 15 |
| East South Central | 17 | 20 | 14 | 15 | 10 | 11 |
| West South Central | 23 | 22 | 20 | 19 | 13 | 15 |
| West | 42 | 37 | 34 | 33 | 23 | 22 |
| Mountain | 16 | 21 | 14 | 20 | 10 | 10 |
| Pacific | 39 | 30 | 31 | 27 | 20 | 19 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 54 | 50 | 45 | 41 | 29 | 29 |
| Small Metro | 35 | 37 | 31 | 30 | 23 | 17 |
| 250K – 1 Mil. Pop. | 28 | 32 | 26 | 25 | 21 | 14 |
| < 250K Pop. | 21 | 21 | 18 | 17 | 11 | 10 |
| Nonmetro | 24 | 29 | 23 | 25 | 14 | 17 |
| Urbanized | 16 | 16 | 14 | 15 | 10 | 11 |
| Less Urbanized | 19 | 22 | 15 | 17 | 9 | 12 |
| Completely Rural | 12 | 8 | 11 | 7 | 7 | 5 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.119D Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.28 | 0.28 | 0.24 | 0.24 | 0.16 | 0.15 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.53 | 0.66 | 0.43 | 0.50 | 0.30 | 0.32 |
| New England | 0.99 | 1.41 | 0.66 | 1.17 | 0.55 | 0.57 |
| Middle Atlantic | 0.62 | 0.74 | 0.53 | 0.54 | 0.35 | 0.38 |
| Midwest | 0.48 | 0.49 | 0.40 | 0.44 | 0.24 | 0.27 |
| East North Central | 0.58 | 0.60 | 0.47 | 0.53 | 0.29 | 0.34 |
| West North Central | 0.83 | 0.87 | 0.74 | 0.75 | 0.43 | 0.44 |
| South | 0.47 | 0.49 | 0.43 | 0.40 | 0.30 | 0.27 |
| South Atlantic | 0.67 | 0.71 | 0.66 | 0.60 | 0.46 | 0.34 |
| East South Central | 1.17 | 1.41 | 0.98 | 1.03 | 0.67 | 0.73 |
| West South Central | 0.79 | 0.75 | 0.68 | 0.64 | 0.45 | 0.50 |
| West | 0.73 | 0.63 | 0.58 | 0.57 | 0.39 | 0.38 |
| Mountain | 0.95 | 1.24 | 0.82 | 1.15 | 0.58 | 0.60 |
| Pacific | 0.95 | 0.71 | 0.75 | 0.64 | 0.50 | 0.47 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.41 | 0.37 | 0.34 | 0.31 | 0.22 | 0.22 |
| Small Metro | 0.48 | 0.50 | 0.43 | 0.41 | 0.32 | 0.23 |
| 250K – 1 Mil. Pop. | 0.58 | 0.63 | 0.52 | 0.51 | 0.42 | 0.28 |
| < 250K Pop. | 0.91 | 0.88 | 0.78 | 0.68 | 0.47 | 0.41 |
| Nonmetro | 0.56 | 0.68 | 0.53 | 0.59 | 0.33 | 0.40 |
| Urbanized | 0.96 | 0.89 | 0.80 | 0.83 | 0.59 | 0.58 |
| Less Urbanized | 0.88 | 1.11 | 0.71 | 0.86 | 0.42 | 0.61 |
| Completely Rural | 2.42 | 1.79 | 2.36 | 1.57 | 1.48 | 1.20 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.119P Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.9726 | | 0.7254 | | 0.9876 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.9288 | | 0.3218 | | 0.9740 | |
| New England | 0.3885 | | 0.8143 | | 0.0793 | |
| Middle Atlantic | 0.4419 | | 0.3016 | | 0.3772 | |
| Midwest | 0.0446 | | 0.2710 | | 0.2330 | |
| East North Central | 0.0625 | | 0.2540 | | 0.2001 | |
| West North Central | 0.4064 | | 0.7790 | | 0.8577 | |
| South | 0.4295 | | 0.9633 | | 0.3012 | |
| South Atlantic | 0.8239 | | 0.4087 | | 0.0551 | |
| East South Central | 0.1634 | | 0.2761 | | 0.8611 | |
| West South Central | 0.4870 | | 0.6405 | | 0.6184 | |
| West | 0.7473 | | 0.5150 | | 0.7425 | |
| Mountain | 0.3947 | | 0.5449 | | 0.5714 | |
| Pacific | 0.9328 | | 0.6865 | | 0.9323 | |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.0883 | | 0.4279 | | 0.7829 | |
| Small Metro | 0.1368 | | 0.7686 | | 0.4720 | |
| 250K – 1 Mil. Pop. | 0.5219 | | 0.6866 | | 0.1035 | |
| < 250K Pop. | 0.1093 | | 0.2651 | | 0.1375 | |
| Nonmetro | 0.2122 | | 0.0505 | | 0.1600 | |
| Urbanized | 0.1379 | | 0.0114 | | 0.3385 | |
| Less Urbanized | 0.4719 | | 0.2922 | | 0.2917 | |
| Completely Rural | 0.4761 | | 0.3265 | | 0.9890 | |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.120C Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 119 | 130 | 86 | 97 | 51 | 61 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 44 | 47 | 32 | 39 | 19 | 22 |
| New England | 23 | 24 | 19 | 22 | 11 | 12 |
| Middle Atlantic | 37 | 41 | 25 | 33 | 16 | 19 |
| Midwest | 52 | 53 | 38 | 40 | 22 | 26 |
| East North Central | 43 | 43 | 34 | 31 | 19 | 22 |
| West North Central | 27 | 32 | 17 | 25 | 11 | 14 |
| South | 70 | 80 | 51 | 59 | 33 | 37 |
| South Atlantic | 48 | 57 | 36 | 43 | 25 | 31 |
| East South Central | 34 | 36 | 23 | 22 | 13 | 14 |
| West South Central | 37 | 44 | 27 | 33 | 17 | 16 |
| West | 71 | 73 | 48 | 53 | 25 | 34 |
| Mountain | 22 | 37 | 20 | 28 | 11 | 12 |
| Pacific | 66 | 62 | 42 | 44 | 23 | 32 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 92 | 98 | 64 | 74 | 36 | 46 |
| Small Metro | 67 | 72 | 48 | 52 | 33 | 32 |
| 250K – 1 Mil. Pop. | 52 | 61 | 38 | 40 | 28 | 23 |
| < 250K Pop. | 41 | 40 | 27 | 34 | 18 | 22 |
| Nonmetro | 39 | 42 | 31 | 34 | 17 | 23 |
| Urbanized | 28 | 28 | 24 | 23 | 11 | 16 |
| Less Urbanized | 26 | 27 | 18 | 22 | 11 | 15 |
| Completely Rural | 14 | 9 | 9 | 6 | 7 | 3 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.120D Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.39 | 0.41 | 0.28 | 0.30 | 0.16 | 0.19 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.80 | 0.83 | 0.57 | 0.70 | 0.35 | 0.39 |
| New England | 1.59 | 1.62 | 1.33 | 1.50 | 0.76 | 0.80 |
| Middle Atlantic | 0.92 | 0.98 | 0.62 | 0.80 | 0.39 | 0.45 |
| Midwest | 0.72 | 0.73 | 0.53 | 0.54 | 0.31 | 0.36 |
| East North Central | 0.87 | 0.85 | 0.68 | 0.62 | 0.39 | 0.43 |
| West North Central | 1.22 | 1.40 | 0.78 | 1.08 | 0.48 | 0.64 |
| South | 0.63 | 0.71 | 0.46 | 0.52 | 0.30 | 0.33 |
| South Atlantic | 0.89 | 1.01 | 0.67 | 0.77 | 0.47 | 0.55 |
| East South Central | 1.78 | 1.85 | 1.22 | 1.16 | 0.66 | 0.73 |
| West South Central | 1.00 | 1.16 | 0.72 | 0.88 | 0.45 | 0.43 |
| West | 0.97 | 0.98 | 0.66 | 0.71 | 0.35 | 0.46 |
| Mountain | 1.02 | 1.63 | 0.93 | 1.25 | 0.48 | 0.55 |
| Pacific | 1.29 | 1.20 | 0.82 | 0.85 | 0.45 | 0.62 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.58 | 0.59 | 0.40 | 0.45 | 0.22 | 0.28 |
| Small Metro | 0.66 | 0.71 | 0.47 | 0.51 | 0.32 | 0.32 |
| 250K – 1 Mil. Pop. | 0.81 | 0.92 | 0.60 | 0.60 | 0.43 | 0.34 |
| < 250K Pop. | 1.10 | 1.13 | 0.73 | 0.95 | 0.49 | 0.63 |
| Nonmetro | 0.78 | 0.86 | 0.62 | 0.69 | 0.33 | 0.47 |
| Urbanized | 1.25 | 1.19 | 1.06 | 0.95 | 0.48 | 0.67 |
| Less Urbanized | 1.15 | 1.24 | 0.81 | 1.01 | 0.50 | 0.71 |
| Completely Rural | 2.97 | 2.52 | 1.92 | 1.60 | 1.48 | 0.94 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.120P Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.0030 | | 0.1570 | | 0.0103 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.0593 | | 0.3420 | | 0.3818 | |
| New England | 0.2963 | | 0.4962 | | 0.3621 | |
| Middle Atlantic | 0.1288 | | 0.1028 | | 0.5875 | |
| Midwest | 0.1707 | | 0.5137 | | 0.1287 | |
| East North Central | 0.4360 | | 0.7374 | | 0.4552 | |
| West North Central | 0.2005 | | 0.1126 | | 0.1186 | |
| South | 0.1337 | | 0.8421 | | 0.1559 | |
| South Atlantic | 0.3955 | | 0.8380 | | 0.0838 | |
| East South Central | 0.7759 | | 0.2060 | | 0.9743 | |
| West South Central | 0.0305 | | 0.6179 | | 0.7386 | |
| West | 0.1519 | | 0.1040 | | 0.1835 | |
| Mountain | 0.4721 | | 0.8318 | | 0.4546 | |
| Pacific | 0.2104 | | 0.0423 | | 0.2649 | |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.0266 | | 0.0627 | | 0.0766 | |
| Small Metro | 0.3513 | | 0.4908 | | 0.6514 | |
| 250K – 1 Mil. Pop. | 0.2631 | | 0.3316 | | 0.5824 | |
| < 250K Pop. | 0.9627 | | 0.8582 | | 0.1743 | |
| Nonmetro | 0.0163 | | 0.2858 | | 0.0059 | |
| Urbanized | 0.0870 | | 0.7030 | | 0.0208 | |
| Less Urbanized | 0.0814 | | 0.1180 | | 0.0176 | |
| Completely Rural | 0.9375 | | 0.1885 | | 0.0103 | |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.121C Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 539 | 477 | 275 | 275 | 172 | 193 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 233 | 208 | 106 | 87 | 57 | 61 |
| New England | 143 | 133 | 64 | 63 | 23 | 43 |
| Middle Atlantic | 180 | 163 | 84 | 61 | 52 | 44 |
| Midwest | 188 | 189 | 86 | 97 | 50 | 55 |
| East North Central | 156 | 146 | 72 | 85 | 41 | 48 |
| West North Central | 105 | 121 | 47 | 48 | 29 | 27 |
| South | 343 | 274 | 177 | 149 | 125 | 115 |
| South Atlantic | 263 | 194 | 136 | 111 | 89 | 87 |
| East South Central | 136 | 115 | 74 | 49 | 59 | 42 |
| West South Central | 172 | 155 | 86 | 87 | 65 | 62 |
| West | 283 | 268 | 161 | 188 | 91 | 130 |
| Mountain | 97 | 115 | 48 | 71 | 29 | 54 |
| Pacific | 266 | 241 | 154 | 174 | 86 | 119 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 421 | 376 | 212 | 229 | 132 | 154 |
| Small Metro | 282 | 238 | 146 | 138 | 86 | 95 |
| 250K – 1 Mil. Pop. | 208 | 193 | 108 | 115 | 68 | 70 |
| < 250K Pop. | 179 | 139 | 94 | 77 | 53 | 65 |
| Nonmetro | 192 | 171 | 81 | 90 | 55 | 62 |
| Urbanized | 125 | 112 | 53 | 58 | 30 | 42 |
| Less Urbanized | 131 | 103 | 47 | 63 | 24 | 46 |
| Completely Rural | 64 | 58 | 37 | 21 | 37 | 4 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.121D Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.30 | 0.26 | 0.15 | 0.15 | 0.10 | 0.11 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.66 | 0.59 | 0.30 | 0.25 | 0.16 | 0.17 |
| New England | 1.55 | 1.43 | 0.69 | 0.68 | 0.25 | 0.46 |
| Middle Atlantic | 0.70 | 0.63 | 0.33 | 0.24 | 0.20 | 0.17 |
| Midwest | 0.46 | 0.46 | 0.21 | 0.24 | 0.12 | 0.14 |
| East North Central | 0.55 | 0.51 | 0.25 | 0.30 | 0.14 | 0.17 |
| West North Central | 0.87 | 1.00 | 0.39 | 0.40 | 0.24 | 0.23 |
| South | 0.54 | 0.42 | 0.28 | 0.23 | 0.20 | 0.18 |
| South Atlantic | 0.77 | 0.56 | 0.40 | 0.32 | 0.26 | 0.25 |
| East South Central | 1.26 | 1.06 | 0.68 | 0.45 | 0.55 | 0.38 |
| West South Central | 0.90 | 0.80 | 0.45 | 0.45 | 0.34 | 0.32 |
| West | 0.71 | 0.66 | 0.41 | 0.46 | 0.23 | 0.32 |
| Mountain | 0.85 | 0.99 | 0.42 | 0.61 | 0.26 | 0.46 |
| Pacific | 0.94 | 0.84 | 0.54 | 0.60 | 0.30 | 0.41 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.44 | 0.39 | 0.22 | 0.24 | 0.14 | 0.16 |
| Small Metro | 0.54 | 0.44 | 0.28 | 0.26 | 0.17 | 0.18 |
| 250K – 1 Mil. Pop. | 0.61 | 0.55 | 0.32 | 0.33 | 0.20 | 0.20 |
| < 250K Pop. | 1.01 | 0.75 | 0.53 | 0.41 | 0.30 | 0.35 |
| Nonmetro | 0.62 | 0.55 | 0.26 | 0.29 | 0.18 | 0.20 |
| Urbanized | 1.03 | 0.82 | 0.44 | 0.43 | 0.25 | 0.31 |
| Less Urbanized | 0.86 | 0.73 | 0.31 | 0.45 | 0.16 | 0.33 |
| Completely Rural | 1.82 | 1.76 | 1.07 | 0.65 | 1.07 | 0.13 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.121P Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.3039 | | 0.4035 | | 0.8065 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.6502 | | 0.7611 | | 0.8368 | |
| New England | 0.5701 | | 0.0601 | | 0.3408 | |
| Middle Atlantic | 0.9028 | | 0.3590 | | 0.3693 | |
| Midwest | 0.1882 | | 0.5923 | | 0.5021 | |
| East North Central | 0.2130 | | 0.8604 | | 0.6939 | |
| West North Central | 0.5711 | | 0.2090 | | 0.5467 | |
| South | 0.8248 | | 0.7969 | | 0.4990 | |
| South Atlantic | 0.8442 | | 0.9579 | | 0.7097 | |
| East South Central | 0.7949 | | 0.1123 | | 0.0964 | |
| West South Central | 0.9467 | | 0.5096 | | 0.5746 | |
| West | 0.3989 | | 0.3124 | | 0.3854 | |
| Mountain | 0.2788 | | 0.1960 | | 0.6134 | |
| Pacific | 0.6731 | | 0.5517 | | 0.2516 | |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.2914 | | 0.4216 | | 0.8951 | |
| Small Metro | 0.9137 | | 0.9861 | | 0.5094 | |
| 250K – 1 Mil. Pop. | 0.9083 | | 0.7017 | | 0.9740 | |
| < 250K Pop. | 0.9788 | | 0.5814 | | 0.3037 | |
| Nonmetro | 0.7174 | | 0.5344 | | 0.9674 | |
| Urbanized | 0.6471 | | 0.6552 | | 0.6822 | |
| Less Urbanized | 0.3937 | | 0.2569 | | 0.3686 | |
| Completely Rural | 0.9190 | | 0.1853 | | 0.0472 | |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.122C Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 565 | 510 | 293 | 298 | 182 | 205 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 237 | 215 | 109 | 98 | 58 | 71 |
| New England | 138 | 133 | 68 | 67 | 25 | 48 |
| Middle Atlantic | 189 | 173 | 85 | 73 | 52 | 53 |
| Midwest | 204 | 191 | 98 | 108 | 57 | 64 |
| East North Central | 170 | 149 | 84 | 93 | 46 | 57 |
| West North Central | 113 | 119 | 52 | 53 | 33 | 28 |
| South | 347 | 300 | 186 | 164 | 133 | 122 |
| South Atlantic | 263 | 208 | 142 | 120 | 97 | 93 |
| East South Central | 137 | 136 | 79 | 58 | 60 | 43 |
| West South Central | 178 | 166 | 90 | 95 | 69 | 65 |
| West | 313 | 294 | 172 | 200 | 95 | 134 |
| Mountain | 105 | 114 | 53 | 74 | 31 | 54 |
| Pacific | 295 | 271 | 164 | 186 | 89 | 123 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 446 | 401 | 223 | 247 | 139 | 163 |
| Small Metro | 295 | 259 | 159 | 153 | 96 | 101 |
| 250K – 1 Mil. Pop. | 220 | 219 | 119 | 129 | 76 | 76 |
| < 250K Pop. | 188 | 147 | 106 | 85 | 60 | 68 |
| Nonmetro | 201 | 176 | 92 | 99 | 57 | 68 |
| Urbanized | 131 | 119 | 65 | 67 | 33 | 46 |
| Less Urbanized | 138 | 106 | 52 | 67 | 26 | 48 |
| Completely Rural | 72 | 58 | 39 | 22 | 40 | 5 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.122D Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.27 | 0.24 | 0.14 | 0.14 | 0.09 | 0.10 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.58 | 0.53 | 0.27 | 0.24 | 0.14 | 0.17 |
| New England | 1.30 | 1.24 | 0.63 | 0.63 | 0.23 | 0.45 |
| Middle Atlantic | 0.63 | 0.57 | 0.28 | 0.24 | 0.17 | 0.17 |
| Midwest | 0.43 | 0.40 | 0.21 | 0.22 | 0.12 | 0.13 |
| East North Central | 0.51 | 0.44 | 0.25 | 0.28 | 0.14 | 0.17 |
| West North Central | 0.79 | 0.83 | 0.36 | 0.37 | 0.23 | 0.19 |
| South | 0.46 | 0.39 | 0.25 | 0.22 | 0.18 | 0.16 |
| South Atlantic | 0.67 | 0.52 | 0.36 | 0.30 | 0.25 | 0.23 |
| East South Central | 1.08 | 1.06 | 0.62 | 0.46 | 0.47 | 0.34 |
| West South Central | 0.78 | 0.72 | 0.39 | 0.41 | 0.30 | 0.28 |
| West | 0.67 | 0.61 | 0.37 | 0.42 | 0.20 | 0.28 |
| Mountain | 0.77 | 0.82 | 0.39 | 0.53 | 0.23 | 0.39 |
| Pacific | 0.88 | 0.79 | 0.49 | 0.55 | 0.27 | 0.36 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.40 | 0.36 | 0.20 | 0.22 | 0.12 | 0.14 |
| Small Metro | 0.48 | 0.41 | 0.26 | 0.24 | 0.16 | 0.16 |
| 250K – 1 Mil. Pop. | 0.54 | 0.52 | 0.29 | 0.31 | 0.19 | 0.18 |
| < 250K Pop. | 0.88 | 0.67 | 0.49 | 0.39 | 0.28 | 0.31 |
| Nonmetro | 0.56 | 0.49 | 0.26 | 0.28 | 0.16 | 0.19 |
| Urbanized | 0.91 | 0.74 | 0.45 | 0.42 | 0.23 | 0.29 |
| Less Urbanized | 0.78 | 0.65 | 0.30 | 0.41 | 0.15 | 0.29 |
| Completely Rural | 1.81 | 1.58 | 1.00 | 0.60 | 1.02 | 0.15 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.122P Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.0866 | | 0.1889 | | 0.3265 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.4082 | | 0.4738 | | 0.9194 | |
| New England | 0.4427 | | 0.1215 | | 0.2796 | |
| Middle Atlantic | 0.6545 | | 0.8404 | | 0.5217 | |
| Midwest | 0.0902 | | 0.4276 | | 0.2366 | |
| East North Central | 0.1426 | | 0.8237 | | 0.5200 | |
| West North Central | 0.3742 | | 0.0585 | | 0.2498 | |
| South | 0.8588 | | 0.8135 | | 0.8261 | |
| South Atlantic | 0.9715 | | 0.8649 | | 0.8757 | |
| East South Central | 0.7527 | | 0.0683 | | 0.1061 | |
| West South Central | 0.5817 | | 0.4069 | | 0.5063 | |
| West | 0.2769 | | 0.1647 | | 0.2490 | |
| Mountain | 0.1998 | | 0.2622 | | 0.7710 | |
| Pacific | 0.5222 | | 0.2943 | | 0.1716 | |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.1134 | | 0.1314 | | 0.6781 | |
| Small Metro | 0.7965 | | 0.7278 | | 0.4898 | |
| 250K – 1 Mil. Pop. | 0.6849 | | 0.9676 | | 0.8321 | |
| < 250K Pop. | 0.9056 | | 0.5279 | | 0.2051 | |
| Nonmetro | 0.4035 | | 0.3680 | | 0.3733 | |
| Urbanized | 0.8779 | | 0.7262 | | 0.2865 | |
| Less Urbanized | 0.1936 | | 0.0774 | | 0.0842 | |
| Completely Rural | 0.9872 | | 0.0945 | | 0.0171 | |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.