

Table 1.128C Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug or Method of Administration	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
HALLUCINOGENS								
PCP (Angel Dust, Phencyclidine)	284	271	20	20	45	49	282	262
LSD (Acid)	514	494	34	26	115	108	496	469
Peyote	280	262	15	13	38	37	276	258
Mescaline	340	319	13	10	40	40	340	314
Psilocybin (Mushrooms)	439	433	32	30	104	103	418	403
"Ecstasy" (MDMA)	297	285	40	32	112	109	247	255
INHALANTS	474	441	67	68	111	104	457	407
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	328	317	29	28	46	44	321	309
Correction Fluid, Degreaser, or Cleaning Fluid	127	116	32	31	36	36	116	106
Gasoline or Lighter Fluid	174	156	38	41	46	46	160	142
Glue, Shoe Polish, or Toluene	186	174	43	44	43	42	178	159
Halothane, Ether, or Other Anesthetics	104	101	12	12	24	23	101	97
Lacquer Thinner or Other Paint Solvents	130	126	28	29	34	35	124	114
Lighter Gases (Butane, Propane)	85	63	24	24	28	25	75	55
Nitrous Oxide or "Whippets"	335	310	31	29	96	87	315	289
Spray Paints	116	105	34	37	38	32	101	95
Other Aerosol Sprays	119	105	27	29	38	39	107	92
ANY NEEDLE USE^{1,2}	257	188	13	11	29	30	249	185
Heroin Needle Use ^{1,2}	171	130	6	7	18	19	170	128
Cocaine Needle Use ^{1,2}	170	144	6	4	17	18	169	143
Stimulant Needle Use ^{1,2}	150	117	8	7	19	21	146	115
Methamphetamine Needle Use ^{1,2}	122	102	5	6	17	18	119	100
HEROIN	223	205	14	11	35	37	220	201
Smoke Heroin ²	129	121	8	8	22	24	127	117
Sniff or Snort Heroin ²	170	168	11	8	30	28	166	165

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.128D Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Percentages, 2002 and 2003

Drug or Method of Administration	Total	AGE GROUP (Years)						
		12-17		18-25		26 or Older		
	2002	2003	2002	2003	2002	2003	2002	2003
HALLUCINOGENS								
PCP (Angel Dust, Phencyclidine)	0.26	0.25	0.20	0.19	0.45	0.42	0.32	0.30
LSD (Acid)	0.12	0.11	0.08	0.08	0.15	0.16	0.16	0.14
Peyote	0.22	0.21	0.14	0.10	0.37	0.34	0.28	0.26
Mescaline	0.12	0.11	0.06	0.05	0.12	0.12	0.15	0.14
Psilocybin (Mushrooms)	0.14	0.13	0.05	0.04	0.13	0.12	0.19	0.17
"Ecstasy" (MDMA)	0.19	0.18	0.13	0.12	0.33	0.32	0.23	0.22
	0.13	0.12	0.16	0.13	0.36	0.34	0.14	0.14
INHALANTS								
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	0.20	0.19	0.27	0.27	0.36	0.33	0.25	0.22
Correction Fluid, Degreaser, or Cleaning Fluid	0.14	0.13	0.12	0.11	0.15	0.14	0.18	0.17
Gasoline or Lighter Fluid	0.05	0.05	0.13	0.12	0.12	0.11	0.06	0.06
Glue, Shoe Polish, or Toluene	0.07	0.07	0.15	0.17	0.15	0.15	0.09	0.08
Halothane, Ether, or Other Anesthetics	0.08	0.07	0.17	0.18	0.14	0.13	0.10	0.09
Lacquer Thinner or Other Paint Solvents	0.04	0.04	0.05	0.05	0.08	0.07	0.06	0.05
Lighter Gases (Butane, Propane)	0.06	0.05	0.11	0.12	0.11	0.11	0.07	0.06
Nitrous Oxide or "Whippets"	0.04	0.03	0.10	0.10	0.09	0.08	0.04	0.03
Spray Paints	0.14	0.13	0.13	0.11	0.31	0.27	0.18	0.16
Other Aerosol Sprays	0.05	0.04	0.14	0.15	0.12	0.10	0.06	0.05
	0.05	0.04	0.11	0.12	0.12	0.12	0.06	0.05
ANY NEEDLE USE^{1,2}								
Heroin Needle Use ^{1,2}	0.11	0.08	0.05	0.04	0.09	0.10	0.14	0.10
Cocaine Needle Use ^{1,2}	0.07	0.05	0.03	0.03	0.06	0.06	0.09	0.07
Stimulant Needle Use ^{1,2}	0.07	0.06	0.02	0.02	0.06	0.06	0.09	0.08
Methamphetamine Needle Use ^{1,2}	0.06	0.05	0.03	0.03	0.06	0.07	0.08	0.06
	0.06	0.05	0.02	0.02	0.06	0.06	0.07	0.06
HEROIN								
Smoke Heroin ²	0.10	0.09	0.05	0.04	0.11	0.12	0.12	0.11
Sniff or Snort Heroin ²	0.05	0.05	0.03	0.03	0.07	0.08	0.07	0.06
	0.07	0.07	0.04	0.03	0.10	0.09	0.09	0.09

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.128P Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug or Method of Administration	Total	AGE GROUP (Years)					
		12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002
HALLUCINOGENS							
PCP (Angel Dust, Phencyclidine)	0.6951		0.0066		0.0978		0.8669
LSD (Acid)	0.3193		0.2649		0.1595		0.2355
Peyote	0.6127		0.0000		0.0001		0.4706
Mescaline	0.9656		0.9590		0.8532		0.9502
Psilocybin (Mushrooms)	0.6033		0.0060		0.0249		0.3683
"Ecstasy" (MDMA)	0.1014		0.9330		0.8673		0.0859
	0.1102		0.0000		0.5186		0.0093
INHALANTS	0.8484		0.6767		0.1087		0.9175
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	0.8987		0.6896		0.2721		0.9940
Correction Fluid, Degreaser, or Cleaning Fluid	0.9501		0.6073		0.1935		0.8416
Gasoline or Lighter Fluid	0.3614		0.7304		0.2615		0.4546
Glue, Shoe Polish, or Toluene	0.2017		0.9591		0.5773		0.2253
Halothane, Ether, or Other Anesthetics	0.0885		0.2649		0.4458		0.0748
Lacquer Thinner or Other Paint Solvents	0.7504		0.7453		0.3357		0.9947
Lighter Gases (Butane, Propane)	0.3490		0.5436		0.2554		0.3793
Nitrous Oxide or "Whippets"	0.6610		0.0811		0.2442		0.9954
Spray Paints	0.7395		0.0547		0.3190		0.9885
Other Aerosol Sprays	0.9366		0.0326		0.8482		0.6524
ANY NEEDLE USE^{1,2}	0.2730		0.7636		0.5358		0.2381
Heroin Needle Use ^{1,2}	0.8810		0.7265		0.7708		0.8437
Cocaine Needle Use ^{1,2}	0.1032		0.3674		0.2874		0.0855
Stimulant Needle Use ^{1,2}	0.3691		0.5895		0.6254		0.3429
Methamphetamine Needle Use ^{1,2}	0.6624		0.8573		0.8756		0.6356
HEROIN	0.9050		0.4102		0.7162		0.9105
Smoke Heroin ²	0.4794		0.7742		0.9972		0.4646
Sniff or Snort Heroin ²	0.9935		0.2128		0.8573		0.9711

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.129C Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

Pain Reliever	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Darvocet®, Darvon®, or Tylenol® with Codeine	445	424	51	49	88	100	428	398
Percocet®, Percodan®, or Tylox®	312	320	27	28	70	77	297	299
Vicodin®, Lortab®, or Loracet®	354	389	48	47	102	106	321	352
Codeine	249	248	31	31	70	74	229	231
Demerol®	168	156	16	13	41	40	161	148
Dilaudid®	156	94	3	4	16	12	156	93
Fioricet®	61	69	4	5	12	9	61	67
Fiorinal®	96	79	7	5	8	10	96	78
Hydrocodone	172	196	25	28	57	73	155	176
Methadone	76	93	15	14	25	31	72	86
Morphine	128	111	23	19	39	41	121	101
Oxycontin®	116	140	21	20	40	56	103	125
Phenaphen® with Codeine	102	79	13	10	20	24	99	74
Propoxyphene	51	42	8	7	8	11	51	40
SK-65®	12	13	5	5	4	5	10	11
Stadol®	33	37	5	2	6	5	32	37
Talacen®	16	15	6	3	5	5	15	14
Talwin®	84	66	5	2	6	6	83	66
Talwin NX®	29	21	5	4	4	8	28	19
Tramadol	15	36	5	4	5	12	13	33
Ultram®	86	96	14	11	23	27	81	88

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.129D Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Percentages, 2002 and 2003

Pain Reliever	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Darvocet®, Darvon®, or Tylenol® with Codeine	0.19	0.18	0.21	0.19	0.28	0.32	0.24	0.22
Percocet®, Percodan®, or Tylox®	0.13	0.13	0.11	0.11	0.23	0.24	0.17	0.17
Vicodin®, Lortab®, or Loracet®	0.15	0.16	0.20	0.19	0.33	0.34	0.18	0.19
Codeine	0.11	0.10	0.13	0.12	0.23	0.23	0.13	0.13
Demerol®	0.07	0.07	0.06	0.05	0.13	0.13	0.09	0.08
Dilaudid®	0.07	0.04	0.01	0.02	0.05	0.04	0.09	0.05
Fioricet®	0.03	0.03	0.02	0.02	0.04	0.03	0.03	0.04
Fiorinal®	0.04	0.03	0.03	0.02	0.03	0.03	0.05	0.04
Hydrocodone	0.07	0.08	0.10	0.11	0.18	0.23	0.09	0.10
Methadone	0.03	0.04	0.06	0.06	0.08	0.10	0.04	0.05
Morphine	0.05	0.05	0.09	0.08	0.13	0.13	0.07	0.06
Oxycontin®	0.05	0.06	0.09	0.08	0.13	0.18	0.06	0.07
Phenaphen® with Codeine	0.04	0.03	0.05	0.04	0.06	0.08	0.05	0.04
Propoxyphene	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.02
SK-65®	0.01	0.01	0.02	0.02	0.01	0.02	0.01	0.01
Stadol®	0.01	0.02	0.02	0.01	0.02	0.02	0.02	0.02
Talacen®	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01
Talwin®	0.04	0.03	0.02	0.01	0.02	0.02	0.05	0.04
Talwin NX®	0.01	0.01	0.02	0.02	0.01	0.02	0.02	0.01
Tramadol	0.01	0.02	0.02	0.02	0.02	0.04	0.01	0.02
Ultram®	0.04	0.04	0.06	0.04	0.07	0.09	0.05	0.05

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.129P Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Pain Reliever	Total	AGE GROUP (Years)					
		12-17		18-25		26 or Older	
		2002	2003	2002	2003	2002	2003
Darvocet®, Darvon®, or Tylenol® with Codeine	0.3578		0.3624		0.3132		0.4199
Percocet®, Percodan®, or Tylox®	0.0184		0.1155		0.1928		0.0463
Vicodin®, Lortab®, or Loracet®	0.0000		0.9693		0.0002		0.0000
Codeine	0.9703		0.8604		0.1228		0.6300
Demerol®	0.8669		0.7114		0.8148		0.8978
Dilaudid®	0.2359		0.2674		0.2493		0.2746
Fioricet®	0.8576		0.6345		0.2660		0.7471
Fiorinal®	0.6356		0.4815		0.8702		0.6588
Hydrocodone	0.0000		0.0506		0.0000		0.0231
Methadone	0.0589		0.6992		0.0299		0.1873
Morphine	0.8369		0.1620		0.1822		0.6531
Oxycontin®	0.0000		0.3522		0.0000		0.0011
Phenaphen® with Codeine	0.8420		0.8758		0.2488		0.6099
Propoxyphene	0.9450		0.8676		0.1228		0.7702
SK-65®	0.7702		0.7163		0.5670		0.9182
Stadol®	0.6351		0.8241		0.3294		0.5000
Talacen®	0.6045		0.2823		0.9699		0.7876
Talwin®	0.0353		0.5166		0.8736		0.0368
Talwin NX®	0.1456		0.8000		0.7915		0.1180
Tramadol	0.0006		0.6005		0.0002		0.0211
Ultram®	0.3025		0.0651		0.4994		0.2553

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.130C Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

Tranquilizer	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Klonopin® or Clonazepam	147	151	19	17	41	47	139	139
Xanax®, Alprazolam, Ativan®, or Lorazepam	271	274	32	27	72	76	248	254
Valium® or Diazepam	437	379	27	27	77	78	427	361
Atarax®	73	49	8	6	11	9	71	47
BuSpar®	71	56	9	9	22	23	68	49
Equanil®	20	38	6	4	4	4	19	38
Flexeril®	143	136	9	9	24	31	140	129
Librium®	149	110	5	6	12	13	149	110
Limbital®	26	11	4	3	9	4	24	11
Meprobamate	52	47	4	4	7	5	51	46
Miltown®	52	36	5	3	3	3	52	36
Rohypnol®	48	42	10	11	17	23	42	37
Serax®	59	40	7	4	8	5	58	40
Soma®	137	145	17	17	36	47	135	134
Tranxene®	44	35	4	4	4	7	43	34
Vistaril®	59	43	7	4	11	9	57	42

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.130D Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Percentages, 2002 and 2003

Tranquilizer	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Klonopin® or Clonazepam	0.06	0.06	0.08	0.07	0.13	0.15	0.08	0.08
Xanax®, Alprazolam, Ativan®, or Lorazepam	0.12	0.12	0.13	0.11	0.23	0.24	0.14	0.14
Valium® or Diazepam	0.19	0.16	0.11	0.11	0.25	0.25	0.24	0.20
Atarax®	0.03	0.02	0.03	0.02	0.04	0.03	0.04	0.03
BuSpar®	0.03	0.02	0.04	0.04	0.07	0.07	0.04	0.03
Equanil®	0.01	0.02	0.02	0.01	0.01	0.01	0.01	0.02
Flexeril®	0.06	0.06	0.04	0.04	0.08	0.10	0.08	0.07
Librium®	0.06	0.05	0.02	0.02	0.04	0.04	0.08	0.06
Limbital®	0.01	0.00	0.02	0.01	0.03	0.01	0.01	0.01
Meprobamate	0.02	0.02	0.01	0.02	0.02	0.02	0.03	0.03
Miltown®	0.02	0.02	0.02	0.01	0.01	0.01	0.03	0.02
Rohypnol®	0.02	0.02	0.04	0.04	0.06	0.07	0.02	0.02
Serax®	0.03	0.02	0.03	0.02	0.03	0.01	0.03	0.02
Soma®	0.06	0.06	0.07	0.07	0.11	0.15	0.07	0.07
Tranxene®	0.02	0.01	0.02	0.02	0.01	0.02	0.02	0.02
Vistaril®	0.02	0.02	0.03	0.02	0.04	0.03	0.03	0.02

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.130P Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Tranquilizer	Total	AGE GROUP (Years)					
		12-17		18-25		26 or Older	
		2002	2003	2002	2003	2002	2003
Klonopin® or Clonazepam	0.0296		0.3864		0.0003		0.2217
Xanax®, Alprazolam, Ativan®, or Lorazepam	0.0013		0.9235		0.0196		0.0065
Valium® or Diazepam	0.8262		0.8595		0.3123		0.9693
Atarax®	0.3481		0.3605		0.8140		0.4133
BuSpar®	0.2118		0.9351		0.5172		0.2547
Equanil®	0.2671		0.8187		0.7322		0.2617
Flexeril®	0.3279		0.9049		0.0622		0.5366
Librium®	0.0537		0.7612		0.8630		0.0503
Limbitrol®	0.0079		0.6705		0.0177		0.0500
Meprobamate	0.9142		0.6215		0.2045		0.8191
Miltown®	0.2573		0.5476		0.7355		0.2710
Rohypnol®	0.6326		0.6892		0.3671		0.9488
Serax®	0.5074		0.3163		0.1564		0.7051
Soma®	0.1787		0.7186		0.4193		0.2818
Tranxene®	0.6308		0.6545		0.3950		0.5082
Vistaril®	0.1633		0.1797		0.0392		0.3872

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.131C Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

Stimulant	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Methamphetamine, ¹ Desoxyn®, or Methedrine	369	369	27	22	61	65	356	365
Prescription Diet Pills ²	342	295	23	21	36	43	342	293
Ritalin® or Methylphenidate	195	160	33	29	67	68	178	137
Cylert®	41	37	6	6	9	7	38	36
Dexedrine®	201	168	11	11	28	28	197	163
Dextroamphetamine	75	62	5	8	11	13	74	61
Didrex®	36	38	7	6	8	7	34	37
Eskatrol®	82	28	3	2	7	2	81	28
Ionamin®	86	79	6	6	8	6	86	78
Mazanor®	18	10	6	4	2	4	17	8
Obedrin-LA®	39	6	3	5	3	3	39	*
Plegine®	23	18	5	2	4	2	22	18
Preludin®	122	85	4	4	8	11	122	84
Sanorex®	37	23	3	6	6	4	37	22
Tenuate®	75	32	4	5	4	6	74	31

*Low precision; no estimate reported.

¹ Crank, crystal, ice, or speed.

² For example, Amphetamines, Benzedrine®, Biphetamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.131D Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Percentages, 2002 and 2003

Stimulant	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Methamphetamine, ¹ Desoxyn®, or Methedrine	0.16	0.16	0.11	0.09	0.20	0.20	0.20	0.20
Prescription Diet Pills ²	0.15	0.12	0.09	0.08	0.12	0.13	0.19	0.16
Ritalin® or Methylphenidate	0.08	0.07	0.13	0.11	0.22	0.21	0.10	0.08
Cylert®	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.02
Dexedrine®	0.09	0.07	0.04	0.04	0.09	0.09	0.11	0.09
Dextroamphetamine	0.03	0.03	0.02	0.03	0.03	0.04	0.04	0.03
Didrex®	0.02	0.02	0.03	0.02	0.03	0.02	0.02	0.02
Eskatrol®	0.03	0.01	0.01	0.01	0.02	0.01	0.05	0.02
Ionamin®	0.04	0.03	0.02	0.03	0.03	0.02	0.05	0.04
Mazanor®	0.01	0.00	0.02	0.02	0.01	0.01	0.01	0.00
Obedrin-LA®	0.02	0.00	0.01	0.02	0.01	0.01	0.02	*
Plegine®	0.01	0.01	0.02	0.01	0.01	0.00	0.01	0.01
Preludin®	0.05	0.04	0.02	0.02	0.03	0.04	0.07	0.05
Sanorex®	0.02	0.01	0.01	0.03	0.02	0.01	0.02	0.01
Tenuate®	0.03	0.01	0.02	0.02	0.01	0.02	0.04	0.02

*Low precision; no estimate reported.

¹ Crank, crystal, ice, or speed.

² For example, Amphetamines, Benzedrine®, Biphetamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.131P Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Stimulant	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Methamphetamine, ¹ Desoxyn®, or Methedrine	0.6807		0.2342		0.0843		0.9580	
Prescription Diet Pills ²	0.3588		0.5581		0.5317		0.3452	
Ritalin® or Methylphenidate	0.0822		0.2757		0.2526		0.0244	
Cylert®	1.0000		0.5781		0.7117		1.0000	
Dexedrine®	0.0032		0.9370		0.3779		0.0038	
Dextroamphetamine	0.5397		0.3496		1.0000		0.4702	
Didrex®	0.7510		0.7865		0.5268		0.6027	
Eskatrol®	0.0137		0.1230		0.1158		0.0229	
Ionamin®	0.4715		0.4142		0.5843		0.4738	
Mazanor®	0.0913		0.3677		0.3203		0.0821	
Obedrin-LA®	0.1178		0.2802		0.8608		*	
Plegine®	0.4330		0.1823		0.1674		0.6917	
Preludin®	0.2756		0.6959		0.2488		0.2248	
Sanorex®	0.4472		0.2687		0.4055		0.4062	
Tenuate®	0.1100		0.6668		0.3632		0.0801	

*Low precision; no estimate reported.

¹ Crank, crystal, ice, or speed.

² For example, Amphetamines, Benzedrine®, Biphetamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.132C Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

Sedative	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Methaqualone, Sopor®, or Quaalude®	325	283	12	11	28	26	324	279
Barbiturates ¹	220	200	6	8	16	17	220	197
Restoril® or Temazepam	96	93	10	9	14	13	93	91
Amytal®	71	43	2	5	6	6	71	42
Butisol®	36	27	3	4	5	5	35	27
Chloral Hydrate	52	38	3	4	4	7	52	37
Dalmane®	102	63	6	6	6	7	101	62
Halcion®	116	88	6	6	11	11	115	86
Phenobarbital	145	116	10	7	15	15	144	115
Placidyl®	118	88	6	5	8	5	118	87
Tuinal®	144	105	5	6	6	4	144	105

*Low precision; no estimate reported.

¹ For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.132D Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Percentages, 2002 and 2003

Sedative	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Methaqualone, Sopor®, or Quaalude®	0.14	0.12	0.05	0.04	0.09	0.08	0.18	0.15
Barbiturates ¹	0.09	0.08	0.02	0.03	0.05	0.05	0.12	0.11
Restoril® or Temazepam	0.04	0.04	0.04	0.04	0.05	0.04	0.05	0.05
Amytal®	0.03	0.02	0.01	0.02	0.02	0.02	0.04	0.02
Butisol®	0.02	0.01	0.01	0.02	0.01	0.02	0.02	0.01
Chloral Hydrate	0.02	0.02	0.01	0.02	0.01	0.02	0.03	0.02
Dalmane®	0.04	0.03	0.03	0.02	0.02	0.02	0.06	0.03
Halcion®	0.05	0.04	0.03	0.02	0.03	0.03	0.06	0.05
Phenobarbital	0.06	0.05	0.04	0.03	0.05	0.05	0.08	0.06
Placidyl®	0.05	0.04	0.02	0.02	0.02	0.01	0.07	0.05
Tuinal®	0.06	0.04	0.02	0.02	0.02	0.01	0.08	0.06

*Low precision; no estimate reported.

¹ For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.132P Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Sedative	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Methaqualone, Sopor®, or Quaalude®	0.3736		0.8331		0.1486		0.4465	
Barbiturates ¹	0.7084		0.1489		0.3521		0.7369	
Restoril® or Temazepam	0.4988		0.9187		0.2967		0.5877	
Amytal®	0.4181		0.0001		0.7820		0.2913	
Butisol®	0.8745		0.2055		0.7766		0.7813	
Chloral Hydrate	0.3048		0.4487		0.0390		0.1831	
Dalmane®	0.0843		0.8977		0.7951		0.0796	
Halcion®	0.0292		0.5173		0.4111		0.0326	
Phenobarbital	0.1242		0.5093		0.4689		0.1586	
Placidyl®	0.0641		0.8999		0.0190		0.0891	
Tuinal®	0.0653		0.4442		0.2476		0.0669	

*Low precision; no estimate reported.

¹ For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.