

Table 2.19C Any Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	797	783	883	828	832	803
12	25	23	19	17	13	11
13	36	33	27	26	19	16
14	40	37	34	31	27	26
15	42	48	38	40	33	32
16	44	46	40	42	35	39
17	42	46	40	42	41	41
18	46	52	51	54	51	51
19	50	48	53	50	52	49
20	47	48	53	51	49	50
21	42	47	49	48	51	47
22	40	41	45	47	46	46
23	40	42	49	47	48	47
24	39	37	44	45	42	46
25	36	43	43	48	42	47
26-29	184	157	202	188	198	190
30-34	185	186	205	199	204	202
35-39	211	211	243	239	237	238
40-44	176	195	240	240	236	236
45-49	204	217	237	240	222	225
50-54	266	278	301	299	287	293
55-59	238	212	258	254	236	248
60-64	221	199	241	200	233	199
65 or Older	451	380	343	297	336	279

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.19D Any Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.34	0.33	0.38	0.35	0.35	0.34
12	0.62	0.56	0.47	0.42	0.31	0.27
13	0.86	0.77	0.65	0.59	0.46	0.36
14	0.92	0.91	0.77	0.76	0.61	0.62
15	1.02	1.16	0.92	0.96	0.80	0.77
16	1.08	1.09	0.99	1.01	0.87	0.93
17	1.07	1.12	1.01	1.02	1.04	1.01
18	1.06	1.14	1.17	1.17	1.16	1.10
19	1.22	1.23	1.29	1.30	1.28	1.26
20	1.17	1.19	1.32	1.26	1.22	1.24
21	1.04	1.19	1.22	1.22	1.25	1.19
22	1.05	1.03	1.16	1.16	1.18	1.14
23	1.05	1.05	1.28	1.19	1.27	1.17
24	1.11	1.03	1.24	1.23	1.19	1.26
25	1.07	1.19	1.28	1.33	1.26	1.29
26-29	1.21	1.07	1.32	1.27	1.30	1.29
30-34	0.93	0.92	1.03	0.99	1.03	1.00
35-39	0.96	1.01	1.10	1.14	1.08	1.14
40-44	0.80	0.86	1.10	1.06	1.08	1.04
45-49	0.96	1.01	1.12	1.12	1.05	1.05
50-54	1.41	1.39	1.60	1.50	1.53	1.47
55-59	1.62	1.42	1.76	1.70	1.61	1.66
60-64	1.90	1.69	2.07	1.70	2.00	1.70
65 or Older	1.33	1.11	1.01	0.87	0.99	0.81

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.19P Any Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.4499		0.0635		0.1893	
12	0.1327		0.3673		0.9889	
13	0.0017		0.0130		0.0610	
14	0.0980		0.6489		0.6851	
15	0.1541		0.7776		0.5175	
16	0.0848		0.3846		0.1875	
17	0.0645		0.1144		0.1111	
18	0.0042		0.0544		0.0994	
19	0.5173		0.4415		0.3063	
20	0.9326		0.7933		0.1883	
21	0.0267		0.0052		0.0219	
22	0.3193		0.9003		0.6170	
23	0.3045		0.4413		0.2599	
24	0.3489		0.1341		0.0309	
25	0.5748		0.8580		0.8056	
26-29	0.2580		0.3492		0.3120	
30-34	0.2420		0.4223		0.8593	
35-39	0.7550		0.0536		0.0249	
40-44	0.1984		0.7877		0.8759	
45-49	0.8468		0.8968		0.6932	
50-54	0.2868		0.4121		0.7338	
55-59	0.3111		0.6308		0.6367	
60-64	0.8860		0.0738		0.1319	
65 or Older	0.5673		0.8065		0.5286	

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.20C Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	843	825	835	819	795	789
12	24	22	18	15	11	10
13	34	31	26	23	18	15
14	40	35	32	28	25	23
15	41	45	36	37	30	29
16	44	46	39	41	33	36
17	43	48	38	41	37	40
18	50	55	51	53	49	48
19	52	48	54	50	50	49
20	47	50	52	51	47	50
21	44	49	51	49	51	48
22	43	44	45	48	43	46
23	42	44	48	48	48	46
24	40	40	43	46	42	47
25	38	47	43	46	42	45
26-29	189	170	196	191	190	189
30-34	199	205	211	201	209	198
35-39	227	230	230	234	225	227
40-44	195	218	233	237	225	228
45-49	213	229	219	231	210	222
50-54	281	284	285	286	271	277
55-59	243	237	247	229	230	224
60-64	229	209	226	184	217	184
65 or Older	474	407	314	267	297	246

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.20D Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.36	0.35	0.36	0.34	0.34	0.33
12	0.58	0.53	0.43	0.36	0.26	0.25
13	0.82	0.72	0.61	0.54	0.43	0.34
14	0.92	0.84	0.74	0.68	0.58	0.55
15	0.99	1.08	0.88	0.89	0.73	0.71
16	1.09	1.09	0.96	0.99	0.82	0.86
17	1.08	1.17	0.96	1.01	0.93	0.97
18	1.16	1.20	1.18	1.15	1.13	1.05
19	1.27	1.24	1.32	1.28	1.23	1.25
20	1.19	1.24	1.30	1.26	1.17	1.24
21	1.08	1.25	1.25	1.25	1.26	1.23
22	1.11	1.10	1.17	1.20	1.12	1.14
23	1.10	1.11	1.26	1.19	1.25	1.15
24	1.15	1.10	1.21	1.28	1.19	1.30
25	1.14	1.28	1.30	1.26	1.26	1.25
26-29	1.24	1.15	1.28	1.29	1.24	1.28
30-34	1.00	1.02	1.06	1.00	1.05	0.98
35-39	1.03	1.10	1.04	1.12	1.02	1.08
40-44	0.89	0.96	1.06	1.05	1.02	1.01
45-49	1.00	1.07	1.03	1.08	0.99	1.04
50-54	1.50	1.42	1.52	1.43	1.44	1.39
55-59	1.65	1.59	1.68	1.53	1.57	1.50
60-64	1.97	1.78	1.94	1.56	1.87	1.57
65 or Older	1.39	1.19	0.92	0.78	0.88	0.72

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.20P Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.3636		0.0538		0.2244	
12	0.0729		0.1289		0.8606	
13	0.0018		0.0023		0.0115	
14	0.0183		0.1868		0.9222	
15	0.1093		0.4585		0.6186	
16	0.2520		0.5508		0.1184	
17	0.0986		0.0519		0.2037	
18	0.0012		0.0389		0.1609	
19	0.4503		0.3273		0.3952	
20	0.9736		0.7746		0.1708	
21	0.0299		0.0012		0.0173	
22	0.3565		0.8564		0.3627	
23	0.3166		0.2774		0.1361	
24	0.4405		0.2327		0.1176	
25	0.6559		0.8775		0.7890	
26-29	0.5240		0.4084		0.2530	
30-34	0.1132		0.7537		0.8722	
35-39	0.8958		0.0748		0.0444	
40-44	0.3097		0.6542		0.7647	
45-49	0.4478		0.8886		0.8738	
50-54	0.4626		0.8153		0.7970	
55-59	0.9652		0.4552		0.6629	
60-64	0.8122		0.0792		0.1656	
65 or Older	0.4073		0.9329		0.8280	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.21C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	691	695	318	314	279	284
12	12	10	7	6	5	3
13	15	18	11	12	8	6
14	21	19	16	14	10	10
15	23	21	17	17	12	12
16	25	25	19	18	14	13
17	29	29	22	22	13	16
18	38	39	27	28	19	19
19	41	38	28	27	22	19
20	48	39	35	27	24	19
21	43	40	26	25	22	19
22	38	41	25	23	19	18
23	44	42	26	26	21	21
24	40	41	19	25	14	21
25	36	36	16	19	15	15
26-29	164	169	92	95	85	80
30-34	195	186	107	105	97	95
35-39	206	202	109	111	97	98
40-44	201	200	96	88	76	84
45-49	170	200	75	85	69	78
50-54	232	230	94	80	87	74
55-59	198	210	60	105	40	104
60-64	172	169	86	71	82	68
65 or Older	328	316	142	136	130	135

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.21D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.29	0.29	0.14	0.13	0.12	0.12
12	0.28	0.24	0.18	0.14	0.11	0.08
13	0.36	0.41	0.26	0.27	0.19	0.15
14	0.49	0.45	0.37	0.35	0.22	0.25
15	0.55	0.50	0.40	0.40	0.29	0.29
16	0.62	0.60	0.46	0.44	0.35	0.31
17	0.74	0.70	0.55	0.53	0.34	0.40
18	0.87	0.84	0.62	0.61	0.44	0.41
19	0.99	0.98	0.67	0.70	0.54	0.50
20	1.21	0.98	0.87	0.66	0.61	0.48
21	1.07	1.03	0.64	0.64	0.55	0.48
22	0.99	1.02	0.65	0.56	0.48	0.44
23	1.15	1.04	0.68	0.66	0.56	0.51
24	1.15	1.11	0.55	0.68	0.41	0.57
25	1.09	1.01	0.49	0.52	0.44	0.42
26-29	1.07	1.14	0.60	0.65	0.56	0.54
30-34	0.98	0.92	0.54	0.52	0.49	0.47
35-39	0.94	0.96	0.49	0.53	0.44	0.47
40-44	0.92	0.88	0.44	0.39	0.35	0.37
45-49	0.80	0.93	0.35	0.40	0.33	0.36
50-54	1.24	1.15	0.50	0.40	0.46	0.37
55-59	1.35	1.41	0.41	0.71	0.27	0.69
60-64	1.48	1.44	0.74	0.61	0.70	0.57
65 or Older	0.97	0.92	0.42	0.40	0.38	0.39

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.21P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.1752		0.4411		0.7189	
12	0.6255		0.4216		0.7829	
13	0.4650		0.8546		0.9276	
14	0.3237		0.7954		0.9175	
15	0.9926		0.3544		0.5006	
16	0.2887		0.2681		0.1263	
17	0.2040		0.3721		0.1045	
18	0.0392		0.3108		0.4017	
19	0.1132		0.3841		0.2876	
20	0.2953		0.5916		0.9170	
21	0.1425		1.0000		0.9293	
22	0.4845		0.4173		0.7363	
23	0.1891		0.5904		0.5184	
24	0.4450		0.3648		0.0323	
25	0.3943		0.5384		0.5163	
26-29	0.9545		0.9467		0.8027	
30-34	0.0992		0.8426		0.8821	
35-39	0.5601		0.6291		0.7888	
40-44	0.7049		0.6349		0.7759	
45-49	0.0822		0.6069		0.5485	
50-54	0.3084		0.2758		0.2421	
55-59	0.1417		0.0652		0.0093	
60-64	0.1035		0.1782		0.1257	
65 or Older	0.2523		0.5262		0.7574	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.22C Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	817	873	483	463	350	331
12	15	14	10	11	7	6
13	23	20	16	14	7	8
14	28	27	23	22	17	15
15	35	34	29	28	17	18
16	37	38	31	33	22	22
17	39	39	34	34	27	24
18	47	51	42	44	31	32
19	52	51	43	46	33	34
20	54	51	46	38	34	31
21	50	47	40	36	30	30
22	50	52	40	42	30	31
23	49	48	40	37	29	26
24	46	47	31	36	24	28
25	43	47	30	37	21	26
26-29	206	184	136	145	103	116
30-34	196	204	143	144	102	97
35-39	236	224	172	139	121	99
40-44	232	249	158	143	123	105
45-49	238	235	155	133	117	100
50-54	307	321	187	178	143	130
55-59	266	281	129	162	97	122
60-64	251	237	77	90	52	74
65 or Older	424	421	145	130	113	104

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.22D Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.35	0.37	0.21	0.19	0.15	0.14
12	0.35	0.34	0.23	0.27	0.17	0.14
13	0.54	0.47	0.37	0.33	0.16	0.18
14	0.64	0.64	0.52	0.52	0.39	0.37
15	0.84	0.83	0.69	0.69	0.43	0.45
16	0.91	0.91	0.77	0.79	0.55	0.52
17	0.98	0.96	0.85	0.82	0.69	0.59
18	1.08	1.10	0.97	0.95	0.72	0.70
19	1.26	1.31	1.05	1.18	0.81	0.88
20	1.36	1.25	1.16	0.95	0.86	0.77
21	1.24	1.19	0.99	0.93	0.74	0.78
22	1.28	1.30	1.04	1.05	0.77	0.76
23	1.27	1.19	1.05	0.92	0.75	0.66
24	1.31	1.28	0.88	0.99	0.67	0.77
25	1.29	1.28	0.90	1.03	0.64	0.72
26-29	1.35	1.25	0.89	0.98	0.67	0.79
30-34	0.98	1.01	0.72	0.71	0.51	0.48
35-39	1.07	1.07	0.78	0.66	0.55	0.47
40-44	1.06	1.10	0.72	0.63	0.56	0.46
45-49	1.12	1.10	0.73	0.62	0.55	0.47
50-54	1.63	1.60	0.99	0.89	0.76	0.65
55-59	1.81	1.88	0.88	1.08	0.66	0.82
60-64	2.16	2.02	0.66	0.77	0.45	0.63
65 or Older	1.25	1.23	0.43	0.38	0.33	0.30

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.22P Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.4274		0.2256		0.9156	
12	0.9538		0.3447		0.8856	
13	0.0763		0.3693		0.3820	
14	0.7881		0.4655		1.0000	
15	0.0174		0.5256		0.7613	
16	0.0042		0.1508		0.7629	
17	0.9174		0.8909		0.4112	
18	0.0324		0.1277		0.8950	
19	0.3759		0.7939		0.6393	
20	0.6226		0.8267		0.2575	
21	0.0131		0.1230		0.9226	
22	0.2250		0.1085		0.1791	
23	0.7699		0.2200		0.4627	
24	0.4221		0.0316		0.1353	
25	0.0451		0.0981		0.2174	
26-29	0.1053		0.7107		0.5781	
30-34	0.6880		0.4377		0.8226	
35-39	0.3301		0.1399		0.2007	
40-44	0.0083		0.6243		0.7638	
45-49	0.9749		0.2466		0.4822	
50-54	0.4723		0.8155		0.3986	
55-59	0.1098		0.2299		0.5174	
60-64	0.6004		0.8590		0.4548	
65 or Older	0.6811		0.8111		0.7103	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.23C Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	660	603	922	891	989	919
12	25	29	21	23	11	14
13	38	37	33	33	20	22
14	42	41	39	38	31	28
15	42	44	42	42	34	35
16	44	45	41	46	39	40
17	36	38	40	44	41	41
18	43	44	49	51	49	52
19	43	39	47	48	54	53
20	42	33	51	43	59	49
21	28	31	35	36	47	46
22	31	28	37	38	48	47
23	28	30	40	39	45	48
24	31	26	37	36	43	43
25	27	27	36	37	41	45
26-29	138	115	171	157	188	191
30-34	143	132	187	192	209	209
35-39	142	136	221	205	245	238
40-44	150	148	210	232	239	259
45-49	148	171	215	230	264	246
50-54	193	198	325	290	352	333
55-59	191	170	254	265	267	281
60-64	166	164	243	250	260	258
65 or Older	385	377	502	481	476	467

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.23D Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.28	0.25	0.39	0.37	0.42	0.39
12	0.62	0.71	0.52	0.55	0.27	0.34
13	0.92	0.86	0.79	0.77	0.49	0.50
14	0.96	0.99	0.89	0.91	0.70	0.69
15	1.01	1.07	1.03	1.01	0.82	0.85
16	1.08	1.06	1.01	1.09	0.95	0.96
17	0.90	0.93	1.02	1.08	1.03	1.01
18	0.98	0.96	1.12	1.10	1.11	1.13
19	1.04	1.00	1.14	1.24	1.32	1.36
20	1.06	0.82	1.29	1.08	1.47	1.23
21	0.69	0.79	0.87	0.92	1.15	1.18
22	0.80	0.70	0.95	0.95	1.23	1.17
23	0.73	0.75	1.05	0.97	1.17	1.19
24	0.87	0.72	1.06	0.99	1.21	1.18
25	0.80	0.73	1.09	1.02	1.23	1.24
26-29	0.91	0.78	1.12	1.06	1.23	1.29
30-34	0.72	0.65	0.94	0.95	1.05	1.03
35-39	0.64	0.65	1.01	0.98	1.12	1.14
40-44	0.68	0.65	0.95	1.02	1.09	1.14
45-49	0.70	0.80	1.02	1.07	1.25	1.15
50-54	1.02	0.99	1.73	1.45	1.88	1.66
55-59	1.30	1.14	1.73	1.78	1.82	1.88
60-64	1.43	1.40	2.09	2.12	2.24	2.20
65 or Older	1.13	1.10	1.48	1.40	1.40	1.36

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.23P Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.9739		0.0385		0.1065	
12	0.1616		0.3877		0.0476	
13	0.3073		0.5363		0.4898	
14	0.0881		0.4952		0.7860	
15	0.9292		0.8910		0.4123	
16	0.8335		0.9104		0.7044	
17	0.1749		0.1032		0.5028	
18	0.0850		0.0874		0.0934	
19	0.4008		0.4215		0.6460	
20	0.2366		0.1969		0.0258	
21	0.2191		0.2136		0.4174	
22	0.1566		0.7354		0.0911	
23	0.7691		0.2772		0.4428	
24	0.1683		0.8732		0.7223	
25	0.4996		0.5669		0.3196	
26-29	0.8364		0.7449		0.9419	
30-34	0.9675		0.5536		0.1540	
35-39	1.0000		0.6995		0.7471	
40-44	0.7347		0.1732		0.2813	
45-49	0.5495		0.0413		0.4869	
50-54	0.6440		0.4127		0.6053	
55-59	0.4431		0.5281		0.4497	
60-64	0.5314		0.2124		0.2084	
65 or Older	0.6512		0.0194		0.0440	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.24C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TYPE OF ALCOHOL USE					
	Any Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002	2003	2002	2003	2002	2003
TOTAL	989	919	732	682	392	380
12	11	14	7	7	1	3
13	20	22	14	12	6	3
14	31	28	25	22	12	10
15	34	35	27	27	11	15
16	39	40	34	32	15	17
17	41	41	34	39	21	21
18	49	52	44	50	31	29
19	54	53	51	49	32	30
20	59	49	55	50	38	38
21	47	46	48	48	41	38
22	48	47	51	52	38	40
23	45	48	45	52	34	37
24	43	43	44	46	28	34
25	41	45	42	46	28	32
26-29	188	191	177	183	107	120
30-34	209	209	206	192	113	107
35-39	245	238	227	194	131	124
40-44	239	259	224	224	111	125
45-49	264	246	212	207	143	120
50-54	352	333	240	249	132	137
55-59	267	281	195	206	108	113
60-64	260	258	184	168	113	75
65 or Older	476	467	261	226	101	107

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.24D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TYPE OF ALCOHOL USE					
	Any Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002	2003	2002	2003	2002	2003
TOTAL	0.42	0.39	0.31	0.29	0.17	0.16
12	0.27	0.34	0.16	0.18	0.03	0.08
13	0.49	0.50	0.34	0.28	0.13	0.07
14	0.70	0.69	0.58	0.54	0.28	0.25
15	0.82	0.85	0.66	0.66	0.27	0.37
16	0.95	0.96	0.83	0.77	0.37	0.41
17	1.03	1.01	0.87	0.95	0.53	0.50
18	1.11	1.13	1.02	1.09	0.70	0.64
19	1.32	1.36	1.26	1.26	0.79	0.78
20	1.47	1.23	1.38	1.23	0.94	0.95
21	1.15	1.18	1.17	1.23	1.02	0.97
22	1.23	1.17	1.33	1.29	0.97	0.99
23	1.17	1.19	1.19	1.31	0.89	0.93
24	1.21	1.18	1.25	1.26	0.81	0.94
25	1.23	1.24	1.27	1.26	0.85	0.87
26-29	1.23	1.29	1.16	1.24	0.70	0.81
30-34	1.05	1.03	1.03	0.95	0.57	0.53
35-39	1.12	1.14	1.03	0.93	0.60	0.59
40-44	1.09	1.14	1.02	0.99	0.50	0.55
45-49	1.25	1.15	1.00	0.97	0.68	0.56
50-54	1.88	1.66	1.28	1.25	0.70	0.69
55-59	1.82	1.88	1.33	1.38	0.74	0.76
60-64	2.24	2.20	1.59	1.43	0.97	0.64
65 or Older	1.40	1.36	0.77	0.66	0.30	0.31

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.24P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TYPE OF ALCOHOL USE					
	Any Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2002	2003	2002	2003	2002	2003
TOTAL	0.1065		0.5430		0.8287	
12	0.0476		0.7863		0.3534	
13	0.4898		0.1780		0.0373	
14	0.7860		0.9170		0.7956	
15	0.4123		0.9282		0.1459	
16	0.7044		0.9304		0.6592	
17	0.5028		0.6718		0.5864	
18	0.0934		0.4939		0.2142	
19	0.6460		0.2431		0.7656	
20	0.0258		0.1516		0.4903	
21	0.4174		0.1479		0.2934	
22	0.0911		0.6445		0.7521	
23	0.4428		0.1244		0.9488	
24	0.7223		0.0086		0.0890	
25	0.3196		0.6694		0.1061	
26-29	0.9419		0.1352		0.3717	
30-34	0.1540		0.1354		0.9962	
35-39	0.7471		0.4391		0.8892	
40-44	0.2813		0.3998		0.0902	
45-49	0.4869		0.6213		0.3230	
50-54	0.6053		0.9297		0.4008	
55-59	0.4497		0.3335		0.7003	
60-64	0.2084		0.1369		0.0696	
65 or Older	0.0440		0.7159		0.3368	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.