

**Table 2.1C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	797	783	883	828	832	803
Cigarettes	843	825	835	819	795	789
Smokeless Tobacco	691	695	318	314	279	284
Cigars	817	873	483	463	350	331
Pipe Tobacco <sup>2</sup>	726	801	--	--	166	131
<b>ALCOHOL</b>	660	603	922	891	989	919
Binge Alcohol Use <sup>3</sup>	--	--	--	--	732	682
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	392	380

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.1D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.34	0.33	0.38	0.35	0.35	0.34
Cigarettes	0.36	0.35	0.36	0.34	0.34	0.33
Smokeless Tobacco	0.29	0.29	0.14	0.13	0.12	0.12
Cigars	0.35	0.37	0.21	0.19	0.15	0.14
Pipe Tobacco <sup>2</sup>	0.31	0.34	--	--	0.07	0.06
<b>ALCOHOL</b>	0.28	0.25	0.39	0.37	0.42	0.39
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.31	0.29
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.17	0.16

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.1P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.4499		0.0635		0.1893	
Cigarettes	0.3636		0.0538		0.2244	
Smokeless Tobacco	0.1752		0.4411		0.7189	
Cigars	0.4274		0.2256		0.9156	
Pipe Tobacco <sup>2</sup>	0.7389		--		0.2797	
<b>ALCOHOL</b>	0.9739		0.0385		0.1065	
Binge Alcohol Use <sup>3</sup>	--		--		0.5430	
Heavy Alcohol Use <sup>3</sup>	--		--		0.8287	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.2C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	108	113	92	93	81	80
Cigarettes	109	111	87	90	74	73
Smokeless Tobacco	61	55	41	40	27	30
Cigars	79	82	65	65	47	44
Pipe Tobacco <sup>2</sup>	35	33	--	--	14	18
<b>ALCOHOL</b>	107	112	105	104	80	82
Binge Alcohol Use <sup>3</sup>	--	--	--	--	67	67
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	31	33

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.2D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.44	0.45	0.37	0.37	0.33	0.32
Cigarettes	0.44	0.45	0.35	0.36	0.30	0.29
Smokeless Tobacco	0.25	0.22	0.17	0.16	0.11	0.12
Cigars	0.32	0.33	0.26	0.26	0.19	0.17
Pipe Tobacco <sup>2</sup>	0.14	0.13	--	--	0.06	0.07
<b>ALCOHOL</b>	0.43	0.45	0.42	0.42	0.32	0.33
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.27	0.27
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.12	0.13

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.2P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.0002		0.0438		0.0797	
Cigarettes	0.0002		0.0051		0.0511	
Smokeless Tobacco	0.2370		0.5002		0.6430	
Cigars	0.0105		0.6166		0.8391	
Pipe Tobacco <sup>2</sup>	0.2646		--		0.3391	
<b>ALCOHOL</b>	0.3519		0.5577		0.8725	
Binge Alcohol Use <sup>3</sup>	--		--		0.9274	
Heavy Alcohol Use <sup>3</sup>	--		--		0.7209	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.3C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	132	142	154	152	150	151
Cigarettes	139	152	155	147	150	151
Smokeless Tobacco	133	119	78	76	59	56
Cigars	149	151	111	115	83	84
Pipe Tobacco <sup>2</sup>	82	75	--	--	24	25
<b>ALCOHOL</b>	104	101	127	131	164	160
Binge Alcohol Use <sup>3</sup>	--	--	--	--	161	155
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	112	114

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.3D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.43	0.45	0.50	0.48	0.48	0.48
Cigarettes	0.45	0.48	0.50	0.46	0.48	0.47
Smokeless Tobacco	0.43	0.37	0.25	0.24	0.19	0.18
Cigars	0.48	0.48	0.36	0.36	0.27	0.26
Pipe Tobacco <sup>2</sup>	0.27	0.24	--	--	0.08	0.08
<b>ALCOHOL</b>	0.34	0.32	0.41	0.41	0.53	0.50
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.52	0.49
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.36	0.36

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.3P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.2277		0.0977		0.4260	
Cigarettes	0.1001		0.0331		0.4180	
Smokeless Tobacco	0.0016		0.5694		0.5084	
Cigars	0.5072		0.9881		0.2913	
Pipe Tobacco <sup>2</sup>	0.3119		--		0.1779	
<b>ALCOHOL</b>	0.3714		0.6491		0.1799	
Binge Alcohol Use <sup>3</sup>	--		--		0.3381	
Heavy Alcohol Use <sup>3</sup>	--		--		0.7023	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.4C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	763	729	848	774	797	748
Cigarettes	808	773	792	764	754	738
Smokeless Tobacco	654	668	302	295	270	274
Cigars	793	847	462	443	331	332
Pipe Tobacco <sup>2</sup>	714	775	--	--	160	126
<b>ALCOHOL</b>	621	566	894	865	953	887
Binge Alcohol Use <sup>3</sup>	--	--	--	--	702	639
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	355	349

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.4D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.43	0.40	0.47	0.43	0.44	0.41
Cigarettes	0.45	0.43	0.44	0.42	0.42	0.41
Smokeless Tobacco	0.36	0.37	0.17	0.16	0.15	0.15
Cigars	0.44	0.47	0.26	0.24	0.18	0.18
Pipe Tobacco <sup>2</sup>	0.40	0.43	--	--	0.09	0.07
<b>ALCOHOL</b>	0.35	0.31	0.50	0.48	0.53	0.49
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.39	0.35
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.20	0.19

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.4P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.9643		0.1433		0.2724	
Cigarettes	0.8807		0.1735		0.3266	
Smokeless Tobacco	0.4716		0.5315		0.7570	
Cigars	0.6531		0.2023		0.6931	
Pipe Tobacco <sup>2</sup>	0.8789		--		0.3208	
<b>ALCOHOL</b>	0.9806		0.0305		0.0542	
Binge Alcohol Use <sup>3</sup>	--		--		0.3613	
Heavy Alcohol Use <sup>3</sup>	--		--		0.9935	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.5C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	45	41	34	30	24	20
Cigarettes	44	40	31	28	21	18
Smokeless Tobacco	19	19	13	13	9	7
Cigars	27	25	18	18	10	10
Pipe Tobacco <sup>2</sup>	13	13	--	--	7	8
<b>ALCOHOL</b>	47	51	40	42	24	26
Binge Alcohol Use <sup>3</sup>	--	--	--	--	16	14
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	6	5

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.5D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.54	0.49	0.41	0.35	0.29	0.23
Cigarettes	0.53	0.47	0.38	0.33	0.26	0.21
Smokeless Tobacco	0.23	0.22	0.16	0.15	0.11	0.08
Cigars	0.32	0.29	0.22	0.21	0.12	0.11
Pipe Tobacco <sup>2</sup>	0.16	0.15	--	--	0.09	0.09
<b>ALCOHOL</b>	0.57	0.61	0.48	0.50	0.29	0.30
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.19	0.17
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.07	0.05

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.5P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.0010		0.0111		0.1385	
Cigarettes	0.0007		0.0008		0.0461	
Smokeless Tobacco	0.6870		0.8387		0.9747	
Cigars	0.1375		0.8534		0.5592	
Pipe Tobacco <sup>2</sup>	0.7269		--		0.7281	
<b>ALCOHOL</b>	0.9265		0.9489		0.5901	
Binge Alcohol Use <sup>3</sup>	--		--		0.3042	
Heavy Alcohol Use <sup>3</sup>	--		--		0.1747	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.6C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	60	63	52	52	42	40
Cigarettes	60	60	50	48	39	37
Smokeless Tobacco	32	29	22	22	16	17
Cigars	44	44	37	35	24	24
Pipe Tobacco <sup>2</sup>	21	18	--	--	9	7
<b>ALCOHOL</b>	60	62	57	56	45	45
Binge Alcohol Use <sup>3</sup>	--	--	--	--	37	36
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	16	19

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.6D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.71	0.77	0.61	0.62	0.49	0.49
Cigarettes	0.71	0.72	0.59	0.58	0.45	0.45
Smokeless Tobacco	0.38	0.35	0.26	0.27	0.18	0.20
Cigars	0.51	0.53	0.44	0.43	0.29	0.30
Pipe Tobacco <sup>2</sup>	0.25	0.22	--	--	0.11	0.09
<b>ALCOHOL</b>	0.71	0.75	0.68	0.68	0.53	0.54
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.43	0.44
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.18	0.23

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.6P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.0585		0.7382		0.9127	
Cigarettes	0.0137		0.2196		0.7409	
Smokeless Tobacco	0.5618		0.5931		0.6349	
Cigars	0.1147		0.9401		0.7654	
Pipe Tobacco <sup>2</sup>	0.1428		--		0.4217	
<b>ALCOHOL</b>	0.3162		0.8953		0.5603	
Binge Alcohol Use <sup>3</sup>	--		--		0.8061	
Heavy Alcohol Use <sup>3</sup>	--		--		0.2835	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.7C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	60	66	57	61	55	57
Cigarettes	62	67	56	59	50	55
Smokeless Tobacco	41	39	29	28	20	21
Cigars	55	57	48	48	36	34
Pipe Tobacco <sup>2</sup>	23	24	--	--	8	15
<b>ALCOHOL</b>	55	54	57	60	55	58
Binge Alcohol Use <sup>3</sup>	--	--	--	--	50	52
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	25	28

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.7D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.75	0.79	0.71	0.73	0.69	0.69
Cigarettes	0.77	0.80	0.70	0.71	0.63	0.66
Smokeless Tobacco	0.51	0.47	0.36	0.34	0.25	0.26
Cigars	0.68	0.68	0.60	0.58	0.45	0.41
Pipe Tobacco <sup>2</sup>	0.28	0.29	--	--	0.10	0.18
<b>ALCOHOL</b>	0.69	0.65	0.72	0.72	0.69	0.70
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.62	0.62
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.31	0.34

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.7P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.0140		0.0971		0.0382	
Cigarettes	0.0540		0.0854		0.0501	
Smokeless Tobacco	0.1036		0.1517		0.8613	
Cigars	0.0456		0.3789		0.4188	
Pipe Tobacco <sup>2</sup>	0.5493		--		0.0963	
<b>ALCOHOL</b>	0.2837		0.2087		0.4469	
Binge Alcohol Use <sup>3</sup>	--		--		0.7997	
Heavy Alcohol Use <sup>3</sup>	--		--		0.8738	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.8C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	83	88	99	91	95	88
Cigarettes	89	92	101	89	93	87
Smokeless Tobacco	78	68	52	48	38	32
Cigars	94	92	79	76	57	57
Pipe Tobacco <sup>2</sup>	45	44	--	--	18	19
<b>ALCOHOL</b>	76	72	90	87	102	99
Binge Alcohol Use <sup>3</sup>	--	--	--	--	93	97
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	61	61

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.8D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.67	0.70	0.80	0.73	0.76	0.71
Cigarettes	0.72	0.73	0.81	0.71	0.75	0.70
Smokeless Tobacco	0.62	0.54	0.42	0.39	0.31	0.25
Cigars	0.75	0.73	0.64	0.60	0.45	0.46
Pipe Tobacco <sup>2</sup>	0.36	0.35	--	--	0.15	0.15
<b>ALCOHOL</b>	0.61	0.58	0.73	0.69	0.82	0.79
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.75	0.77
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.49	0.49

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.8P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.1417		0.1754		0.3969	
Cigarettes	0.1081		0.1264		0.5770	
Smokeless Tobacco	0.0084		0.4993		0.2392	
Cigars	0.0497		0.3707		0.8016	
Pipe Tobacco <sup>2</sup>	0.0926		--		0.2921	
<b>ALCOHOL</b>	0.9830		0.9356		0.6328	
Binge Alcohol Use <sup>3</sup>	--		--		0.8156	
Heavy Alcohol Use <sup>3</sup>	--		--		0.6659	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.9C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	95	110	115	119	111	113
Cigarettes	97	118	115	119	111	114
Smokeless Tobacco	96	93	53	55	43	45
Cigars	116	112	80	87	62	62
Pipe Tobacco <sup>2</sup>	64	61	--	--	16	16
<b>ALCOHOL</b>	69	64	91	89	110	111
Binge Alcohol Use <sup>3</sup>	--	--	--	--	120	114
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	85	89

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.9D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.51	0.57	0.62	0.62	0.60	0.59
Cigarettes	0.52	0.61	0.62	0.62	0.60	0.59
Smokeless Tobacco	0.51	0.48	0.28	0.29	0.23	0.24
Cigars	0.62	0.58	0.43	0.45	0.34	0.32
Pipe Tobacco <sup>2</sup>	0.35	0.32	--	--	0.08	0.08
<b>ALCOHOL</b>	0.37	0.33	0.49	0.46	0.59	0.58
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.65	0.59
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.46	0.46

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.9P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.6536		0.3166		0.6753	
Cigarettes	0.3673		0.1295		0.4858	
Smokeless Tobacco	0.0263		0.9322		0.8775	
Cigars	0.5097		0.3291		0.1915	
Pipe Tobacco <sup>2</sup>	0.9292		--		0.4569	
<b>ALCOHOL</b>	0.2484		0.7132		0.2151	
Binge Alcohol Use <sup>3</sup>	--		--		0.1957	
Heavy Alcohol Use <sup>3</sup>	--		--		0.4531	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.10C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	246	246	277	270	283	283
Cigarettes	259	275	281	278	281	278
Smokeless Tobacco	252	250	142	144	126	124
Cigars	270	273	193	200	146	153
Pipe Tobacco <sup>2</sup>	159	165	--	--	35	37
<b>ALCOHOL</b>	198	175	260	243	292	285
Binge Alcohol Use <sup>3</sup>	--	--	--	--	267	273
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	156	157

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.10D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.70	0.70	0.79	0.77	0.81	0.81
Cigarettes	0.74	0.79	0.80	0.80	0.80	0.80
Smokeless Tobacco	0.72	0.72	0.40	0.41	0.36	0.35
Cigars	0.77	0.78	0.55	0.57	0.41	0.44
Pipe Tobacco <sup>2</sup>	0.45	0.47	--	--	0.10	0.11
<b>ALCOHOL</b>	0.56	0.50	0.74	0.69	0.83	0.81
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.76	0.78
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.44	0.45

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.10P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.9943		0.9791		0.5987	
Cigarettes	0.5010		0.7836		0.5474	
Smokeless Tobacco	0.2217		0.8537		0.9558	
Cigars	0.4185		0.3889		0.6041	
Pipe Tobacco <sup>2</sup>	0.7555		--		0.9631	
<b>ALCOHOL</b>	0.9024		0.8443		0.3146	
Binge Alcohol Use <sup>3</sup>	--		--		0.8372	
Heavy Alcohol Use <sup>3</sup>	--		--		0.5628	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.11C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	725	686	780	696	735	680
Cigarettes	754	718	724	669	693	654
Smokeless Tobacco	588	629	264	260	233	245
Cigars	741	806	403	394	296	288
Pipe Tobacco <sup>2</sup>	675	720	--	--	156	121
<b>ALCOHOL</b>	583	534	814	799	880	830
Binge Alcohol Use <sup>3</sup>	--	--	--	--	624	568
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	314	308

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.11D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.50	0.47	0.54	0.48	0.51	0.47
Cigarettes	0.52	0.49	0.50	0.46	0.48	0.45
Smokeless Tobacco	0.41	0.43	0.18	0.18	0.16	0.17
Cigars	0.51	0.55	0.28	0.27	0.21	0.20
Pipe Tobacco <sup>2</sup>	0.47	0.49	--	--	0.11	0.08
<b>ALCOHOL</b>	0.40	0.37	0.56	0.55	0.61	0.57
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.43	0.39
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.22	0.21

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.11P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.9642		0.1280		0.1808	
Cigarettes	0.9397		0.1221		0.2036	
Smokeless Tobacco	0.8255		0.4472		0.7737	
Cigars	0.4648		0.3397		0.4906	
Pipe Tobacco <sup>2</sup>	0.8706		--		0.3094	
<b>ALCOHOL</b>	0.9613		0.0187		0.0932	
Binge Alcohol Use <sup>3</sup>	--		--		0.3964	
Heavy Alcohol Use <sup>3</sup>	--		--		0.8086	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.12C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	780	755	876	810	829	784
Cigarettes	827	799	826	802	790	774
Smokeless Tobacco	678	687	312	310	276	283
Cigars	814	855	483	456	346	333
Pipe Tobacco <sup>2</sup>	719	791	--	--	163	130
<b>ALCOHOL</b>	627	578	910	884	972	909
Binge Alcohol Use <sup>3</sup>	--	--	--	--	730	675
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	389	376

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.12D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.37	0.35	0.42	0.38	0.39	0.37
Cigarettes	0.39	0.38	0.39	0.38	0.38	0.36
Smokeless Tobacco	0.32	0.32	0.15	0.15	0.13	0.13
Cigars	0.39	0.40	0.23	0.21	0.16	0.16
Pipe Tobacco <sup>2</sup>	0.34	0.37	--	--	0.08	0.06
<b>ALCOHOL</b>	0.30	0.27	0.43	0.42	0.46	0.43
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.35	0.32
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.19	0.18

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.12P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.7895		0.0973		0.2451	
Cigarettes	0.6554		0.1001		0.2930	
Smokeless Tobacco	0.2015		0.4865		0.6821	
Cigars	0.5816		0.2509		0.9369	
Pipe Tobacco <sup>2</sup>	0.7701		--		0.2325	
<b>ALCOHOL</b>	0.9052		0.0414		0.0971	
Binge Alcohol Use <sup>3</sup>	--		--		0.5452	
Heavy Alcohol Use <sup>3</sup>	--		--		0.8533	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.13C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	454	474	593	596	572	573
Cigarettes	509	528	548	582	527	545
Smokeless Tobacco	567	567	302	286	264	256
Cigars	631	621	437	408	327	305
Pipe Tobacco <sup>2</sup>	620	681	--	--	139	127
<b>ALCOHOL</b>	379	373	586	563	656	599
Binge Alcohol Use <sup>3</sup>	--	--	--	--	540	517
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	346	321

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.13D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.40	0.41	0.52	0.52	0.50	0.50
Cigarettes	0.45	0.46	0.48	0.51	0.46	0.47
Smokeless Tobacco	0.50	0.49	0.27	0.25	0.23	0.22
Cigars	0.56	0.54	0.38	0.35	0.29	0.27
Pipe Tobacco <sup>2</sup>	0.55	0.59	--	--	0.12	0.11
<b>ALCOHOL</b>	0.33	0.32	0.52	0.49	0.58	0.52
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.48	0.45
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.30	0.28

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.13P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.6300		0.1026		0.1377	
Cigarettes	0.4151		0.1823		0.3168	
Smokeless Tobacco	0.1065		0.2919		0.5227	
Cigars	0.2760		0.0623		0.3449	
Pipe Tobacco <sup>2</sup>	0.3948		--		0.5493	
<b>ALCOHOL</b>	0.9716		0.1095		0.9318	
Binge Alcohol Use <sup>3</sup>	--		--		0.6632	
Heavy Alcohol Use <sup>3</sup>	--		--		0.3729	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.14C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	593	589	537	520	513	507
Cigarettes	627	586	525	503	501	488
Smokeless Tobacco	311	296	101	130	100	126
Cigars	447	446	178	174	114	123
Pipe Tobacco <sup>2</sup>	267	286	--	--	90	35
<b>ALCOHOL</b>	501	471	644	646	644	600
Binge Alcohol Use <sup>3</sup>	--	--	--	--	415	382
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	177	174

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.14D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.49	0.48	0.44	0.42	0.42	0.41
Cigarettes	0.52	0.48	0.43	0.41	0.41	0.40
Smokeless Tobacco	0.26	0.24	0.08	0.11	0.08	0.10
Cigars	0.37	0.36	0.15	0.14	0.09	0.10
Pipe Tobacco <sup>2</sup>	0.22	0.23	--	--	0.07	0.03
<b>ALCOHOL</b>	0.41	0.38	0.53	0.53	0.53	0.49
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.34	0.31
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.15	0.14

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.14P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.5244		0.2295		0.6577	
Cigarettes	0.5856		0.1019		0.3865	
Smokeless Tobacco	0.7925		0.6591		0.6295	
Cigars	0.9572		0.2191		0.0335	
Pipe Tobacco <sup>2</sup>	0.3154		--		0.2440	
<b>ALCOHOL</b>	0.9762		0.1321		0.0169	
Binge Alcohol Use <sup>3</sup>	--		--		0.5977	
Heavy Alcohol Use <sup>3</sup>	--		--		0.0456	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.15C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	76	78	68	64	59	56
Cigarettes	76	78	61	60	53	47
Smokeless Tobacco	53	49	38	38	26	29
Cigars	62	63	51	50	37	36
Pipe Tobacco <sup>2</sup>	30	28	--	--	13	17
<b>ALCOHOL</b>	77	74	74	73	55	57
Binge Alcohol Use <sup>3</sup>	--	--	--	--	46	46
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	24	23

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.15D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.60	0.61	0.53	0.50	0.47	0.44
Cigarettes	0.60	0.61	0.48	0.47	0.42	0.37
Smokeless Tobacco	0.42	0.39	0.30	0.30	0.21	0.23
Cigars	0.49	0.49	0.40	0.39	0.30	0.28
Pipe Tobacco <sup>2</sup>	0.24	0.22	--	--	0.10	0.13
<b>ALCOHOL</b>	0.61	0.58	0.59	0.57	0.43	0.45
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.37	0.36
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.19	0.18

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.15P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.0067		0.1777		0.5264	
Cigarettes	0.0075		0.0542		0.4055	
Smokeless Tobacco	0.4139		0.9364		0.3174	
Cigars	0.0080		0.3235		0.8613	
Pipe Tobacco <sup>2</sup>	0.5541		--		0.1328	
<b>ALCOHOL</b>	0.0507		0.1377		0.6271	
Binge Alcohol Use <sup>3</sup>	--		--		0.5837	
Heavy Alcohol Use <sup>3</sup>	--		--		0.4515	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.16C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	71	73	59	64	51	52
Cigarettes	71	72	57	63	49	50
Smokeless Tobacco	25	25	18	15	9	6
Cigars	46	50	35	39	26	23
Pipe Tobacco <sup>2</sup>	19	17	--	--	9	7
<b>ALCOHOL</b>	70	77	69	73	58	60
Binge Alcohol Use <sup>3</sup>	--	--	--	--	46	46
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	19	23

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.16D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.59	0.60	0.49	0.52	0.42	0.42
Cigarettes	0.58	0.59	0.47	0.51	0.40	0.41
Smokeless Tobacco	0.21	0.20	0.15	0.12	0.08	0.05
Cigars	0.38	0.40	0.29	0.32	0.21	0.19
Pipe Tobacco <sup>2</sup>	0.15	0.14	--	--	0.07	0.06
<b>ALCOHOL</b>	0.58	0.63	0.57	0.60	0.48	0.49
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.38	0.38
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.16	0.19

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.16P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.0063		0.1095		0.0484	
Cigarettes	0.0054		0.0301		0.0450	
Smokeless Tobacco	0.2755		0.1439		0.0769	
Cigars	0.3457		0.6638		0.9074	
Pipe Tobacco <sup>2</sup>	0.2578		--		0.4088	
<b>ALCOHOL</b>	0.5729		0.5330		0.5090	
Binge Alcohol Use <sup>3</sup>	--		--		0.6917	
Heavy Alcohol Use <sup>3</sup>	--		--		0.1886	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.17C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	91	87	102	107	100	106
Cigarettes	97	98	105	106	99	109
Smokeless Tobacco	103	106	69	72	56	55
Cigars	106	114	92	98	73	72
Pipe Tobacco <sup>2</sup>	72	67	--	--	22	21
<b>ALCOHOL</b>	70	68	86	88	111	100
Binge Alcohol Use <sup>3</sup>	--	--	--	--	110	105
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	87	91

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.17D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.58	0.54	0.66	0.67	0.64	0.67
Cigarettes	0.63	0.62	0.67	0.67	0.64	0.69
Smokeless Tobacco	0.67	0.67	0.44	0.45	0.36	0.34
Cigars	0.68	0.72	0.59	0.62	0.47	0.45
Pipe Tobacco <sup>2</sup>	0.47	0.42	--	--	0.14	0.13
<b>ALCOHOL</b>	0.45	0.43	0.55	0.55	0.71	0.63
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.71	0.66
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.56	0.57

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.17P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.3523		0.4744		0.6052	
Cigarettes	0.2452		0.2173		0.8404	
Smokeless Tobacco	0.0011		0.3848		0.3210	
Cigars	0.7345		0.7895		0.3986	
Pipe Tobacco <sup>2</sup>	0.4469		--		0.1790	
<b>ALCOHOL</b>	0.5155		0.7319		0.0602	
Binge Alcohol Use <sup>3</sup>	--		--		0.2165	
Heavy Alcohol Use <sup>3</sup>	--		--		0.9594	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.18C Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	93	100	103	100	101	93
Cigarettes	94	101	105	96	102	92
Smokeless Tobacco	62	51	24	22	8	11
Cigars	88	90	58	60	38	43
Pipe Tobacco <sup>2</sup>	30	32	--	--	12	12
<b>ALCOHOL</b>	74	73	89	93	102	114
Binge Alcohol Use <sup>3</sup>	--	--	--	--	99	100
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	55	59

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.18D Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: Standard Errors of Percentages, 2002 and 2003**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.60	0.63	0.67	0.63	0.65	0.59
Cigarettes	0.60	0.64	0.68	0.61	0.66	0.58
Smokeless Tobacco	0.40	0.32	0.16	0.14	0.05	0.07
Cigars	0.57	0.57	0.38	0.38	0.25	0.27
Pipe Tobacco <sup>2</sup>	0.19	0.20	--	--	0.08	0.08
<b>ALCOHOL</b>	0.48	0.46	0.57	0.59	0.66	0.72
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.64	0.63
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.35	0.37

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 2.18P Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	0.3498		0.0788		0.4659	
Cigarettes	0.1768		0.0596		0.2794	
Smokeless Tobacco	0.2144		0.5673		0.1368	
Cigars	0.4468		0.7274		0.5289	
Pipe Tobacco <sup>2</sup>	0.3574		--		0.7311	
<b>ALCOHOL</b>	0.0740		0.7636		0.9413	
Binge Alcohol Use <sup>3</sup>	--		--		0.9107	
Heavy Alcohol Use <sup>3</sup>	--		--		0.5429	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.